1 **Clinical *Burkholderia pseudomallei* isolates from Queensland carry diverse**
2 ***bimA_{Bm}* genes that are associated with central nervous system disease and**
3 **are phylogenomically distinct from other Australian strains**

4

5 Delaney Burnard¹, Michelle J Bauer¹, Caitlin Falconer¹, Ian Gassie^{1,2}, Robert E Norton³,
6 David L Paterson¹, Patrick N A Harris⁴

7 ¹University of Queensland Centre for Clinical Research, Royal Brisbane and Woman's Hospital,
8 Herston, Queensland, 4029, Australia.

9 ²Department of Infectious Diseases, Mater Hospital Brisbane, South Brisbane, Queensland, 4101,
10 Australia.

11 ³Pathology Queensland, Townsville University Hospital, Townsville, Queensland, 4814, Australia.

12 ⁴Pathology Queensland, Royal Brisbane & Women's Hospital, Herston, Queensland, 4029, Australia.

13 ***Corresponding author:** Dr Patrick N A Harris

14 University of Queensland Centre for Clinical Research, Building 71/918 Royal Brisbane &
15 Women's Hospital Campus, Herston, QLD, 4029

16 **Email:** p.harris@uq.edu.au; **Tel:** +61 (0) 7 3346 6081

17 **Keywords:** *Burkholderia pseudomallei*, melioidosis, central nervous system disease,
18 virulence factor

19

20

21 Abstract

22 *Burkholderia pseudomallei* is an environmental gram-negative bacterium that causes the
 23 disease melioidosis and is endemic in many countries of the Asia-Pacific region. In Australia,
 24 the mortality rate remains high at approximately 10%. The bacterium is almost exclusively
 25 found in the endemic region, which spans the Northern Territory, Western Australia and
 26 North Queensland. However, Queensland infections remain understudied compared to
 27 those of the Northern Territory. This pilot study aimed to assess the prevalence of central
 28 nervous system disease associated variant *bimA_{Bm}*, identify circulating antimicrobial
 29 resistance mutations and genetically distinct strains from Queensland, via comparative
 30 genomics. From 76 clinical isolates, we identified the *bimA_{Bm}* variant in 20 (26.3%) isolates
 31 and in 9 (45%) of the isolates with documented central nervous system infection (n=18).
 32 Explorative analysis suggests a significant association between isolates carrying the *bimA_{Bm}*
 33 variant and central nervous system disease (OR 2.8, 95% CI 1.3-6.0, P=0.009) compared with
 34 isolates carrying the wildtype *bimA_{Bp}*, with the possibility of additional driving factors.
 35 Furthermore, 50% of isolates were identified as novel multi-locus sequence types (STs),
 36 while the *bimA_{Bm}* variant was more commonly identified in isolates with novel sequence
 37 STs, compared to those of previously described STs. Additionally, mutations associated with
 38 acquired antimicrobial resistance were only identified in 14.5% of all genomes. The findings
 39 of this research have provided clinically relevant genomic data of *B. pseudomallei* in
 40 Queensland and suggest that the *bimA_{Bm}* variant may enable risk stratification for the
 41 development CNS complications and be a potential therapeutic target.

42 Introduction

43 *Burkholderia pseudomallei* is an environmental gram-negative pathogen present in Asia,
 44 South America, Africa and the Pacific. Infections cause the disease melioidosis, with an
 45 extensive range of disease manifestations including, but not limited to, pneumonia, sepsis,
 46 and skin and soft tissue abscesses[1]. Successful treatment can be compromised by the
 47 diagnosis of central nervous system disease (e.g. encephalomyelitis, abscess, meningitis,
 48 cranial nerve impairment)[2]. As these infections are often sub-clinical, exhibit varying times
 49 to clinical presentation, have diverse clinical presentations and affect multiple other organs;
 50 they can be challenging cases to manage[2,3].

51 Central nervous system infections have been associated with the virulence factor *bimA_{Bp}*
 52 mutant, *bimA_{Bm}*. These *bimA* genes encode as intracellular motility factor A, type V effector
 53 proteins that utilise host cellular actin. This motility results in the ability to evade the host
 54 immune system and invade the central nervous system via inter-cellular host cell
 55 migration[4]. Patients with infections possessing the *bimA_{Bm}* gene are suggested to be 14
 56 times more likely to present with neurological disease, while the wild-type variant *bimA_{Bp}* is
 57 linked to pneumonia[5,6].

58 This gene has been identified in both clinical and environmental isolates from the Northern
 59 Territory; yet isolates from North Queensland have not been as extensively characterised,
 60 despite both states being endemic regions[5,6]. Clinical presentations of neurological
 61 disease have been documented in a retrospective Queensland isolate collection[7];
 62 however, the diversity of *bimA_{Bm}* and the association with central nervous system disease in
 63 Queensland remains unknown.

64 This study aimed to use comparative genomics to screen for the presence of the virulence
65 factor *bimA_{Bm}*, which has been previously linked to neurological disease in *B. pseudomallei*
66 infections in patients of the Northern Territory. Furthermore, genomic diversity and
67 antimicrobial resistance profiles are described.

68 **Methods**

69 *Isolates*

70 Clinical *B. pseudomallei* isolates have been collected prospectively over the last 22 years
 71 (1996-2018) from participating hospital pathology facilities (Cairns, Townsville and Central
 72 laboratories of Pathology Queensland). A total of 76 isolates were selected between
 73 postcodes 4895-4500 to include isolates from all of Queensland, from a collection of 400
 74 clinical isolates. All isolates with documented central nervous system presentation were
 75 included (n=18) and the remaining 57 isolates were selected to include a variety of other
 76 common *B. pseudomallei* clinical presentations (Table 1). An additional 24 publicly available
 77 reference genomes were included in this study, of these, 13 were derived from the
 78 Northern Territory and one representative each from Papa New Guinea, China and Thailand
 79 (Table S1).

80 *DNA extraction and WGS*

81 The 76 isolates were recovered from -80°C storage and subcultured twice to ensure purity.
 82 DNA extraction was performed with the QIAGEN DNAeasy ultra-pure DNA extraction kit
 83 according to manufacturer's instructions. Sequencing libraries were generated using the
 84 Nextera Flex DNA library preparation kit and sequenced on the MiniSeq™ System (Illumina
 85 Inc., San Diego, CA, USA) on a high output 300 cycle cartridge according to the
 86 manufacturer's instructions. Five strains with unique *BimA_{Bm}* sequence and/or genomic
 87 diversity were prepped for long read sequencing using the MinION (Oxford Nanopore
 88 Technologies, Oxford, UK), where sequencing libraries were generated using the Rapid
 89 Barcoding Sequencing Kit (SQK-RBK004) and run on a flow cell R9.4.1 for 72 hrs.

90 *Genomic Analysis*

91 Illumina reads were trimmed with Trimmomatic v0.36[8] and quality assessed with
92 multiQC[9], genomes were assembled with SPAdes v3.14.0[10] and annotated with Prokka
93 v1.13[11]. Long reads were filtered, quality checked, assembled, and polished with Illumina
94 reads where applicable using the MicroPIPE pipeline[12]. Genomes derived from this study
95 are publicly available at NCBI BioProject ID: PRJNA717363.

96 All read mapping was performed with BWA-MEM[13]. MLST sequence types were
97 determined with MLST[14] (<https://github.com/tseemann/mlst>) and read mapped to alleles
98 retrieved from pubMLST, where SNPs were suspected[14]. Genotypic antimicrobial
99 resistance was determined with ArDaP[15]. Whole genome alignment (4,950,632bp) and
100 phylogenomic analysis of the 24 reference genomes and 76 genomes derived from this
101 study ($n=100$) were achieved using parSNP[16] JModelTest[17] (TVM+F+I+G4) and
102 IQtree[18] (1000 bootstrap (BS) replicates). Reference genomes were downloaded from
103 NCBI, under the “complete” genome filter, where Australian reference strains were included
104 if they were annotated with significant metadata (Table S1). The reference strains for
105 virulence factors *bimA_{Bm}* and *bimA_{Bp}* were MSHR0668 (locus BH02_RS34290)
106 NZ_CP009546.1 and K96243 (locus BPSS1492) NC_006351.1, respectively.

107 *Statistical analysis*

108 Using Stata and the csi command for unstratified cumulative incidence data[19], and
109 isolates with *bimA_{Bm}* or CNS presentation ($n=29$), Odds Ratios (ORs) and Chi-Square were
110 calculated, with a P value <0.05 considered significant.

111 *Ethics*

This study received ethical approval from the Royal Brisbane & Women's Hospital Ethics Committee as a low or negligible-risk study (LNR/2020/QRBW/65573), with site specific authority obtained from the Townsville Hospital and Health Service and approval under the Queensland Public Health Act.

Results

In total, we identified the *bimA_{Bm}* virulence factor in 20/76 Queensland isolates (26.3%), geographically limited to the northern half of Queensland (Figure 1). Of the 18 isolates with known CNS disease, nine were found to carry the *bimA_{Bm}* virulence factor and nine carried the wild type *bimA_{Bp}*. A total of 20 isolates were confirmed to carry the *bimA_{Bm}* virulence factor, of which eleven had no clinical record of CNS disease. The Stata csi analysis supported *bimA_{Bm}* being associated with CNS infection (Odds Ratio 2.8, 95% CI 1.3-6.1, P=0.009).

The variation observed in the 20 *bimA_{Bm}* protein sequences was significant amongst isolates, with 18 proteins possessing unique sequences, while none of the sequences produced in this study were identical to that of the reference strain MSHR668. The variation was observed almost only in the proline-rich region of the protein (90-160aa) (Figure 2). However, five isolates carried an amino acid variant ΔN213S (TSV1, TSV152, TSV141, TSV164, CAM60), of which four isolates had documented CNS disease (one individual died before diagnosis, CAM60), and four cases resulted in death (one individual made an unexpected recovery while in intensive care, TSV 152). These isolates were collected from as early as 1996, with the latest in 2012 and were significantly geographically dispersed (350-850 km apart). Additionally, four of these same isolates (TSV1, TSV152, TSV164, CAM60) and isolate TSV294 also possessed an upstream mutation ΔR11H. Additionally, all isolates

carrying *bimA_{Bm}* also possess a truncated *bimC* gene which plays a role in intracellular spread and lies upstream of *bimA_{Bm}*.

From the 76 isolates we were able to identify 27 known multi-locus sequence types (STs) from 38 isolates (50%) (Table 3). From these, at least three sequence types (109, 151, 1667) have also been identified in the Northern Territory. Of the 38 novel sequence types, 26 isolates were comprised of previously described alleles in a novel combination, and 12 were due to a SNP present in one of the seven alleles (Figure 3). Novel allele variants were identified across six of the alleles, with ace the only allele without any variants (n=12). In each novel case, sequence types were confirmed via read mapping to all seven alleles. In total 34 of the 38 isolates carried unique novel STs, as two of the STs occurred twice and one ST occurred three times in the sample set. Among the 20 isolates carrying the *bimA_{Bm}* variant, 13 (65%) carried novel STs, over previously described STs (n=7, 35%; Figure 3).

Phylogenomic analysis revealed the majority of Queensland isolates with novel STs branched significantly closer to the Queensland *bimA_{Bm}* reference genome MSHR668, than to the Northern Territory and Asian genomes. This was also true for the *bimA_{Bm}* variant. Two separate lineages can be identified in the phylogeny (point of divergence is approximately in the middle of the phylogeny), with the three clades at the bottom half of the phylogeny carrying 33 of the 38 novel STs identified (86.8%) and 18 of the 20 *bimA_{Bm}* variants present in the sample set (90%). MLST diversity did not appear to be influenced by or correlated with the date the isolate was collected (Figure 3). No Queensland isolate was identical to that of a Northern Territory isolate, with limited clustering with Northern Territory or other Australian isolates. Additionally, five of the Queensland isolates included in this study clustered within the Australian/Asian clade. In this clade the reference genome K96243 from

Thailand is the most basal, with BPC006 from China branching within the Australian isolates CAM2, 112, 189, TSV38 and 95. All of the isolates within this clade were of known ST and did not carry the *bimA_{Bm}* variant.

Approximately one third of the genomes did not contain any AMR related mutations at all (35.5%). The pre-cursor mutation to imipenem resistance (*penA* Δ T153A) was identified in 57.9% of all isolates. Only mutations associated with ceftazidime and meropenem resistance were predicted within the sample set. Ceftazidime resistance (loss of the PBP3 homolog BPSS1219) was only predicted from one isolate (1.3%), while meropenem resistance was predicted to be caused by the loss of function of the *amrR* efflux pump (BPSL1805) in 11 of the 76 isolates (14.5%). Meropenem resistance was encoded in 15% of *bimA_{Bm}* isolates and 16% of *bimA_{Bp}* isolates, respectively (Table 2).

Discussion

Variation in the *bimA_{Bm}* gene was present and extensive. Twenty isolates carried the virulence factor and only two of the sequences produced from this study were identical (TSV39, TSV25). No sequence generated from this study was identical to any of the previously described reference sequences, suggesting depth of variation in the *bimA_{Bm}* gene is extensive and yet to be seen (Figure 2). Since the 2000's sequence variation of *BimA_{Bm}* in *B. pseudomallei* has not been discussed in great detail, despite being studied recently[20]. For example, five isolates possessed a ΔN213S substitution, positioned between the proline rich region and the YadA-like head domain, a highly conserved region[21]. Interestingly, the isolates with this mutation had a mortality rate of 80% and documented CNS disease at a rate of 80% (one patient died before CNS disease could be diagnosed). Unfortunately, similar to previous studies the effect these variants have upon the function of the protein and resulting virulence remain unclear[21]. Reference sequences from MSHR33, 491 and 172 all possess the mutation as well, suggesting this is not a recent event[21]. However, such high mortality rates and disease severity suggests that isolates with this type of mutation may be more virulent than those without.

Previous studies have suggested that *bimA_{Bm}* carrying isolates are 14 times more likely to develop CNS disease than the wild-type *bimA_{Bp}* isolates[5]. Explorative analysis of this data suggested a significant association between the *bimA_{Bm}* variant and the development of CNS disease, compared to the wild type. However, as this dataset was selected for particular factors (CNS disease and postcode) these numbers should be viewed with caution. Perhaps what is more representative of the Queensland population is the 11 *bimA_{Bm}* variants identified from the 58 non-CNS disease isolates, suggesting 19% of *B. pseudomallei*

infections will carry the virulence factor. Indeed, larger sample sizes collected at random will be needed to confirm these numbers and this should be considered for future studies. Sarovich *et al.*, also assessed the likelihood of clinical co-variants that may be associated with CNS disease, but did not find other associations[5]. Due to the lack of complete clinically relevant data surrounding the isolates in this study, we were unable to calculate if this was applicable to our results.

In this study, CNS disease isolates were evenly split between *bimA_{Bp}* and *bimA_{Bm}* genes and more isolates presented without CNS disease and *BimA_{Bm}* (n=11), than with both of these variables (n=9). Therefore, it is likely this gene is not the only factor driving the development of CNS disease. This was also evident in the Northern Territory, with 85.5% of isolates carrying the variant, not diagnosed with CNS disease. Additionally, 1% presented with CNS disease, but no variant[5]. In both studies this is a significant proportion of *bimA_{Bm}* isolates that have not developed CNS disease. An environmental study[6] of both *bimA_{Bm}* and another virulence factor lipopolysaccharide (LPS) suggested the *bimA_{Bm}* virulence factor was more likely to occur with LPS genotype B, the more prevalent genotype in Australia. Genotype A, is more prevalent in Thailand and Southeast Asia, where CNS disease is less frequently reported[5,6]. Furthermore, the gene upstream of *bimA*, *bimC* was also assessed in this study. As both *bimA* and *bimC* genes have been shown to play a role in the intracellular spread of *B. pseudomallei*[22]. As all *bimA_{Bm}* isolates in this study possessed a truncated *bimC* gene, the affect this truncation has upon virulence and the role of *bimC* in intracellular spread remains unknown. However, it suggests the variant is significant to virulence in these strains and should be further investigated. There is no way to show a definitive correlation between these two virulence factors and *bimA_{Bm}* in Queensland without including both clinical and environmental isolates in future studies. The authors

suggest a pan-genomic analysis coupled with extensive clinical metadata from as many isolates as possible representing both states evenly, to be the best approach at identifying all genes associated with CNS disease. It is possible that there may be some combination of genes derived only from the unique genetic diversity of Australian isolates responsible for the increased incidences of CNS disease observed here in Australia compared to those overseas.

Although Queensland has been included in MLST diversity analyses before, novel STs have not been described, which is significant given 50% of isolates in this study were novel STs (Figure 3, Table 3)[23,24]. A large number of isolates (34%) were comprised of a previously undescribed combination of alleles to generate a novel ST. It also appears that isolates with the *bimA_{Bm}* virulence factor are more likely to be novel STs. This implies that Queensland isolates are exchanging genetic material[24]. The exchange of genetic material was also evidenced by the lack of clonality seen in both previously described and novel STs in Queensland. ST 70 was the most commonly described (6.5%) and was identified in both Brisbane and Cairns isolates suggesting one ST does not dominate certain spatial areas in this study, despite this being reported for the Northern Territory[6,23]. Furthermore, the same studies reported that Queensland and the Northern Territory do not share any common STs. However, in this study we identified at least three STs shared between the states (STs 109, 151, 1667), with the potential of more, as the exact location was not recorded for all STs in pubMLST. The identification of shared STs here, may be due to the previous under-representation, or exclusion of Queensland in *B. pseudomallei* isolates in previous studies[23–26].

Genomic analyses also identified the loss of function of the *amrR* regulator (*amrAB-OprA* efflux pump), associated with meropenem resistance in 14.5% of isolates, however this has not been validated phenotypically[15]. A pre-cursor mutation in *penA* Δ T153A, was common amongst the isolates and in combination with other missense and promoter mutations can cause imipenem and amoxicillin-clavulanic acid resistance[15]. This is not overly worrisome as imipenem is not used to treat *B. pseudomallei* in Australia, meropenem is the preferred carbapenem, however amoxicillin-clavulanic acid is sometimes used in eradication therapy[3]. Phenotypically, antimicrobial resistance circulating in the Northern Territory comprises of approximately 3% for doxycycline and 0.9 % for trimethoprim-sulfamethoxazole[27], while isolates remain susceptible to meropenem and ceftazidime[1,27,28]. This data suggests AMR may be rarely encountered in Queensland isolates and the current selection of antimicrobials for use against *B. pseudomallei* infections will be effective.

Conclusion

This study has revealed a significant amount of genetic diversity in Queensland *B. pseudomallei* isolates, such as novel MLST sequence types and unique *bimA_{Bm}* gene sequences. The virulence factor *bimA_{Bm}* is linked to CNS disease, yet it is suspected there are multiple drivers for this type of infection. Further exploration into the virulence factors responsible for CNS disease should focus on maximum sample size, pan-genomics, detailed clinical metadata and include environmental samples. Identification of CNS disease drivers may provide an excellent therapeutic target or act as a screening test to warn clinicians and prompt additional investigations such as a lumbar puncture or MRI.

263 **Funding**

264 This study/project has been funded by a Research Grant 2020 of the European Society of
265 Clinical Microbiology and Infectious Diseases (ESCMID) to DB.

267 Acknowledgements

268 The authors would like to thank Valentine Murigneux for the use of, and their assistance in
269 their long-read assembly pipeline MicroPIPE.

271 **Disclosures**

DLP has received research support from Shionogi, MSD and Pfizer and honoraria for speaking or advisory boards from MSD, Pfizer, Sumitomo, Biomerieux, Accelerate and Lyosvant. PNAH reports grants from Shionogi, Merck (MSD) and Sandoz and honoraria for speaking or advisory boards from Pfizer, Sandoz and Sumitomo. All other authors declare no conflicts of interest.

277 **Author contribution**

278 DB: conceptualization, methodology, analysis, writing. MJB: methodology, analysis, writing-
 279 review and edits. CF: methodology, analysis, writing-review and edits. IG: methodology,
 280 writing-review and edits. REN: data curation, writing-review and edits. PNAH: supervisor,
 281 writing-review and edits. DLP: supervisor, writing-review and edits.

282 Tables

283 **Table 1.** Metadata regarding the *B. pseudomallei* isolates included in this study.

Date range of collection	1996-2018
Geographical range	4000-4895 Queensland, Australia
Gender	31 (40.8%) Female, 45 (59.2%) Male
Age range	6-84 years
Central Nervous System Disease	18 (23.7%)
Mortality*	15 (19.7%)
Isolation sites:	
Blood	42
Pus	8
Sputum	7
Cerebral spinal fluid	4
Tissue	4
Brain	2
Endotracheal aspirate	2
Bronchoalveolar Lavage	1
Gastrointestinal Tract	1
Liver	1
Lung aspirate	1
Lymph	1
Urine	1
Unknown	1
Total:	76

284 *only information for 23/76 isolates was available.

285 **Table 2.** Comparison of clinical presentation, sequence type and antimicrobial resistance of

Variable	<i>bimA_{Bm}</i> Isolates (%) (n=20)	<i>bimA_{Bp}</i> Isolates (%) (n=56)	286 isolates (%) 287 carrying the 288 <i>bimA_{Bm}</i> or <i>bimA_{Ps}</i> 289 virulence factor. 290 291
CNS presentation	45	16	
Novel ST	40	53.6	
Known ST	60	46.4	
<i>amrR</i> mutation	15	16	
Pre-cursor mutation	45	64.3	
No AMR mutation	35	35.7	
Mortality*	40	12.5	

292

293

294

295

296 *Information was only available for 23/76 isolates

297 **Table 3.** Multilocus sequence types of clinical *Burkholderia pseudomallei* from Queensland,
298 included in this study.

Sequence Type	Occurrence
24	2
35	1
70	5
109	1
151	1
235	1
252	3
254	1
257	2
283	1
286	1
591	2
593	2
594	1
958	1
1041	1
1042	1

1453	1
1667	2
1707	1
1711	1
1712	1
1754	1
1756	1
1757	1
1760	1
1769	1
Novel	38 (26 = novel allele combination, 12 = SNP in allele)

299

300 Figure Legends

301 **Figure 1.** Distribution of *B. pseudomallei* isolates with the *bimA_{Bm}* virulence factor in north
302 Queensland.

303

304 **Figure 2.** Variation of the *BimA_{Bm}* proline-rich region identified in Queensland *B.*
305 *pseudomallei* isolates compared to the Queensland *BimA_{Bm}* reference genome MSHR668.

306

307 **Figure 3.** Maximum likelihood phylogenomic tree of Australian *B. pseudomallei* isolates,
308 rooted to reference MSHR668. Bootstrap support ≥ 80 is shown at nodes in red and

309 reference genomes are labelled with strain name stated in NCBI. Tree was built using IQtree
310 100 genomes and a 4,950,632 bp alignment generated from parSNP.

- [1] Gassiep I, Armstrong M, Norton R. Human Melioidosis. Clinical Microbiology Reviews 2020;33. <https://doi.org/10.1128/CMR.00006-19>.
- [2] Wongwandee M, Linasmita P. Central nervous system melioidosis: A systematic review of individual participant data of case reports and case series. PLOS Neglected Tropical Diseases 2019;13:e0007320. <https://doi.org/10.1371/journal.pntd.0007320>.
- [3] eTG complete (digital). Melbourne: Therapeutic Guidelines Limited; 2019 Jun. (eTG August 2020 edition) <https://www.tg.org.au> n.d.
- [4] Stevens MP, Stevens JM, Jeng RL, Taylor LA, Wood MW, Hawes P, et al. Identification of a bacterial factor required for actin-based motility of *Burkholderia pseudomallei*. Molecular Microbiology 2005;56:40–53. <https://doi.org/10.1111/j.1365-2958.2004.04528.x>.
- [5] Sarovich DS, Price EP, Webb JR, Ward LM, Voutsinos MY, Tuanyok A, et al. Variable Virulence Factors in *Burkholderia pseudomallei* (Melioidosis) Associated with Human Disease. PLOS ONE 2014;9:e91682. <https://doi.org/10.1371/journal.pone.0091682>.
- [6] Webb JR, Rachlin A, Rigas V, Sarovich DS, Price EP, Kaestli M, et al. Tracing the environmental footprint of the *Burkholderia pseudomallei* lipopolysaccharide genotypes in the tropical “Top End” of the Northern Territory, Australia. PLOS Neglected Tropical Diseases 2019;13:e0007369. <https://doi.org/10.1371/journal.pntd.0007369>.
- [7] Deuble M, Aquilina C, Norton R. Neurologic Melioidosis. Am J Trop Med Hyg 2013;89:535–9. <https://doi.org/10.4269/ajtmh.12-0559>.
- [8] Bolger, A. M., Lohse, M., & Usadel, B. Trimmomatic: A flexible trimmer for Illumina Sequence Data. Bioinformatics, Btu170 2014.
- [9] Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for multiple tools and samples in a single report. Bioinformatics 2016;32:3047–8. <https://doi.org/10.1093/bioinformatics/btw354>.
- [10] Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. J Comput Biol 2012;19:455–77. <https://doi.org/10.1089/cmb.2012.0021>.
- [11] Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics 2014;30:2068–9. <https://doi.org/10.1093/bioinformatics/btu153>.
- [12] Murigneux V, Roberts LW, Forde BM, Phan M-D, Nhu NTK, Irwin AD, et al. MicroPIPE: An end-to-end solution for high-quality complete bacterial genome construction. BioRxiv 2021:2021.02.02.429319. <https://doi.org/10.1101/2021.02.02.429319>.
- [13] Li H, Durbin R. Fast and accurate long-read alignment with Burrows-Wheeler transform. Bioinformatics 2010;26:589–95. <https://doi.org/10.1093/bioinformatics/btp698>.
- [14] Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics: BIGSdb software, the PubMLST.org website and their applications. Wellcome Open Res 2018;3:124. <https://doi.org/10.12688/wellcomeopenres.14826.1>.
- [15] Madden DE, Webb JR, Steinig EJ, Mayo M, Currie BJ, Price EP, et al. Taking the next-gen step: comprehensive antibiotic resistance detection from *Burkholderia pseudomallei* genomes. BioRxiv 2019:720607. <https://doi.org/10.1101/720607>.
- [16] Treangen TJ, Ondov BD, Koren S, Phillippy AM. The Harvest suite for rapid core-genome alignment and visualization of thousands of intraspecific microbial genomes. Genome Biology 2014;15:524. <https://doi.org/10.1186/s13059-014-0524-x>.

- [17] Darriba D, Taboada GL, Doallo R, Posada D. jModelTest 2: more models, new heuristics and high-performance computing. Nat Methods 2012;9:772. <https://doi.org/10.1038/nmeth.2109>.
- [18] Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol 2015;32:268–74. <https://doi.org/10.1093/molbev/msu300>.
- [19] Stata Statistical Software. College Station, TX: StataCorp LLC; 2019.
- [20] Morris JL, Fane A, Sarovich DS, Price EP, Rush CM, Govan BL, et al. Increased Neurotropic Threat from *Burkholderia pseudomallei* Strains with a *B. mallei*-like Variation in the bimA Motility Gene, Australia. Emerg Infect Dis 2017;23:740–9. <https://doi.org/10.3201/eid2305.151417>.
- [21] Sitthidet C, Stevens JM, Chantratita N, Currie BJ, Peacock SJ, Korbsrisate S, et al. Prevalence and Sequence Diversity of a Factor Required for Actin-Based Motility in Natural Populations of *Burkholderia Species*. Journal of Clinical Microbiology 2008;46:2418–22. <https://doi.org/10.1128/JCM.00368-08>.
- [22] Srinon V, Chaiwattananrungruengpaisan S, Korbsrisate S, Stevens JM. *Burkholderia pseudomallei* BimC Is Required for Actin-Based Motility, Intracellular Survival, and Virulence. Front Cell Infect Microbiol 2019;9. <https://doi.org/10.3389/fcimb.2019.00063>.
- [23] McRobb E, Kaestli M, Price EP, Sarovich DS, Mayo M, Warner J, et al. Distribution of *Burkholderia pseudomallei* in Northern Australia, a Land of Diversity. Appl Environ Microbiol 2014;80:3463–8. <https://doi.org/10.1128/AEM.00128-14>.
- [24] Cheng AC, Ward L, Godoy D, Norton R, Mayo M, Gal D, et al. Genetic Diversity of *Burkholderia pseudomallei* Isolates in Australia. J Clin Microbiol 2008;46:249–54. <https://doi.org/10.1128/JCM.01725-07>.
- [25] Johnson SL, Baker AL, Chain PS, Currie BJ, Daligault HE, Davenport KW, et al. Whole-Genome Sequences of 80 Environmental and Clinical Isolates of *Burkholderia pseudomallei*. Genome Announc 2015;3:e01282-14. <https://doi.org/10.1128/genomeA.01282-14>.
- [26] Smet BD, Sarovich DS, Price EP, Mayo M, Theobald V, Kham C, et al. Whole-Genome Sequencing Confirms that *Burkholderia pseudomallei* Multilocus Sequence Types Common to Both Cambodia and Australia Are Due to Homoplasy. Journal of Clinical Microbiology 2015;53:323–6. <https://doi.org/10.1128/JCM.02574-14>.
- [27] Crowe A, McMahon N, Currie BJ, Baird RW. Current antimicrobial susceptibility of first-episode melioidosis *Burkholderia pseudomallei* isolates from the Northern Territory, Australia. Int J Antimicrob Agents 2014;44:160–2. <https://doi.org/10.1016/j.ijantimicag.2014.04.012>.
- [28] Jenney AWJ, Lum G, Fisher DA, Currie BJ. Antibiotic susceptibility of *Burkholderia pseudomallei* from tropical northern Australia and implications for therapy of melioidosis. International Journal of Antimicrobial Agents 2001;17:109–13. [https://doi.org/10.1016/S0924-8579\(00\)00334-4](https://doi.org/10.1016/S0924-8579(00)00334-4).

[illegible]