Analysis of the Second COVID-19 Wave in India Using a Birth-Death Model

Narayanan C. Viswanath

Government Engineering College, Thrissur, Kerala, India, 680009.

Email: narayanan_viswanath@yahoo.com

Mobile: +91 9495851753

ORCID ID: <https://orcid.org/0000-0001-9191-7436>

Abstract

India is witnessing the second wave of the COVID-19 disease from the first half of February 2021. The method in [5] is applied here to analyze the second wave in India. We start with fitting a birth-death model to the active and total cases data for the period from 13th to 28th February 2021. This initial dataset is expanded step by step by adding the two future week's data to it until 14th May 2021. This resulted in six models in total. The efficacy of each model is tested in terms of predictions made for the next two weeks. The infectivity rates are found to be ever-increasing in the case of the five initial models. The infectivity rate for the sixth model, which is based on the data from 13th February to 14th May 2021, shows a decreasing nature with an increase in time. This indicates a decline in the second wave, which may start from 4th June 2021 according to the fitted parameters.

Keywords: Second COVID-19 wave; birth-death model; India

Introduction

According to Ranjan et al. [3], a rise in the number of COVID-19 cases in India started from 13th February 2021 which indicated the start of the second wave of the disease progression. Using a SIR model [2], they predict a peaking of the same in mid-May 2021.

[4] applied a generalized birth-death model [1] for modeling the total and actively infected COVID-19 cases in several countries. This turned out to be a special case, of the standard SIR model [2], in which the susceptible cases variable has no explicit role. [5] discussed application of the model in [4] when a rise in the number of cases occurs after a fall. Here we check the efficacy of the method in [5] in analyzing the second wave that occurred in India.

Methods

By taking the infection birth rate as $\lambda(t) = ae^{-bt}$ and the recovery rate as μ in a generalized birth-death model [1], [4] derives the following equations for the actively infected $I(t)$ and total $M(t)$ cases as:

$$I(t) = I(0)e^{\left(\frac{a}{b}[1-e^{-bt}] - \mu t\right)}. \quad \dots (1)$$

$$M(t) = I(0) \left(1 + \int_0^t a e^{-b\tau} e^{\left(\frac{a}{b}[1-e^{-b\tau}]-\mu\tau\right)} d\tau \right). \quad \dots (2)$$

[5] discussed application of the above model when a rise in the number of cases occurs after a fall. Their idea was to fit equations (1) and (2) to a dataset after adjusting the total cases so that the total and active cases remain the same at the starting epoch. We follow the method in [5] for fitting equations (1) and (2) to the active and total cases data [7]. Fitting was done using the *nlinfit* function available in the MATLAB R2019b [6] software.

Results and Discussion

We consider 13th February 2021 as a starting point. We fitted equations (1) and (2) using the method discussed in [5] for the data from 13th to 28th February 2021. Figure 1 shows the fit. The parameters for this fit are: $a = 0.0794821535$; $b = -0.0157707070$; $I(0) = 132167$; $\mu = 0.0739635918$. Here notice that the negative value for parameter b indicates that the infection rate is going to increase forever. This according to us points to a failure of the model as far as a long-term prediction is concerned. The absence of the number of susceptible cases in the model could be the reason for this phenomenon. Figure 2 is the comparison of the predictions by the fitted model with actual data for the next two weeks (from 1st to 14th March 2021). Figure 2 shows that the accuracy of the prediction decreases as time increases.

Next, we expanded the prediction dataset by adding the actual data of the next two weeks to it. Figure 3 shows the fit for the data from 13th February to 14th March 2021. The parameters for this fit are: $a = 0.0895802716$; $b = -0.0046661410$; $I(0) = 128946$; $\mu = 0.0782528444$. Again the parameter b turns out to be negative. Prediction for the next two weeks, which is from 15th to 28th March 2021, is given in Figure 4. It shows a decent prediction, whose accuracy begins to decrease as time increases.

We then considered the data from 13th February to 31st March 2021. Figure 5 shows the fit whose parameters are given by: $a = 0.0784394479$; $b = -0.0135156063$; $I(0) = 135863$; $\mu = 0.0778148298$. Figure 6 shows the predictions for the next two weeks: from 1st to 14th April 2021. The parameters for Figure 7, which shows the fit for the data from 13th February to 14th April 2021, are given by: $a = 0.0868287781$; $b = -0.0090513907$; $I(0) = 122037$; $\mu = 0.0750109712$. Figure 8 shows the quality of predictions for the next two weeks (from 15th to 28th April 2021). Figures 5 to 8 reveal that while the quality of fit (Figures 5 and 7) remains the same, the quality of future prediction decreases (Figures 6 and 8).

Figure 9 shows that the fitted counts are slightly greater than the actual values towards the end of the fit, when fitting was done on the data from 13th February to 30th April 2021. The parameters for this fit are: a

$= 0.12608518091$; $b = -0.00129869234$; $I(0) = 58896$; $\mu = 0.07907911057$. The predicted values for the period of 1st to 14th of May 2021 are much higher than the actual values as can be observed from Figure 10. More specifically, the actual and predicted active cases for May 14th are 3679691 and 8594118 respectively. The actual and predicted total cases for the same date are 24372243 and 30651576 respectively.

A fit of the data from 13th February to 14th May 2021 gives the parameters as $a = 0.236576657262$; $b = 0.008817517656$; $I(0) = 4806$; $\mu = 0.088195225005$. Notice that parameter b is positive for the first time since 13th February 2021, indicating a decrease in the infection rate.

According to [4], a point of inflection on the curve (1) of actively infected cases where its decline starts is given by

$$t_s = \frac{1}{b} \log \left(\frac{a}{\mu} \right) \dots (3)$$

The parameters for the final fit give t_s as equal to 111.9, which indicates the decline of active cases may start from 4th June 2021.

Figure 1. Comparison of the fitted values of active and total cases with the actual values for India based on the data from 13th to 28th February 2021. The parameters for this fit are: $a = 0.0794821535$; $b = -0.0157707070$; $I(0) = 132167$; $\mu = 0.0739635918$.

Figure 2. Comparison of the predicted active and total cases with their actual values for India from 1st to 14th March 2021, when fitting was done based on the data from 13th to 28th February 2021.

Figure 3. Comparison of the fitted values of active and total cases with the actual values for India based on the data from 13th February to 14th March 2021. The parameters for this fit are: $a = 0.0895802716$; $b = -0.0046661410$; $I(0) = 128946$; $\mu = 0.0782528444$.

Figure 4. Comparison of the predicted active and total cases with their actual values for India from 15th to 28th March 2021, when fitting was done based on the data from 13th February to 14th March 2021.

Figure 5. Comparison of the fitted values of active and total cases with the actual values for India based on the data from 13th February to 31st March 2021. The parameters for this fit are: $a = 0.0784394479$; $b = -0.0135156063$; $I(0) = 135863$; $\mu = 0.0778148298$.

Figure 6. Comparison of the predicted active and total cases with their actual values for India from 1st to 14th April 2021, when fitting was done based on the data from 13th February to 31st March 2021.

Figure 7. Comparison of the fitted values of active and total cases with the actual values for India based on the data from 13th February to 14th April 2021. The parameters for this fit are: $a = 0.0868287781$; $b = -0.0090513907$; $I(0) = 122037$; $\mu = 0.0750109712$.

Figure 8. Comparison of the predicted active and total cases with their actual values for India from 15th to 28th April 2021, when fitting was done based on the data from 13th February to 14th April 2021.

Figure 9. Comparison of the fitted values of active and total cases with the actual values for India based on the data from 13th February to 30th April 2021. The parameters for this fit are: $a = 0.12608518091$; $b = -0.00129869234$; $I(0) = 58896$; $\mu = 0.07907911057$.

Figure 10. Comparison of the predicted active and total cases with their actual values for India from 1st to 14th May 2021, when fitting was done based on the data from 13th February to 30th April 2021.

Figure 11. Comparison of the fitted values of active and total cases with the actual values and future prediction for India based on the data from 13th February to 14th May 2021. The parameters for this fit are: $a = 0.236576657262$; $b = 0.008817517656$; $I(0) = 4806$; $\mu = 0.088195225005$.

References

1. Kendall, D.G. On the generalized birth-and-death process. *Ann. Math. Stat.* 19(1):1-15; (1948).

2. Kermack, W.O., McKendrick, A.G. A Contribution to the mathematical theory of epidemics. Proc. Roy. Soc. Lond A 115, 700-721 (1927).
3. Ranjan, R., Sharma, A., Verma, M.K. Characterization of the Second Wave of COVID-19 in India. doi: <https://doi.org/10.1101/2021.04.17.21255665> (2021)
4. Viswanath, N.C. Analysis and Prediction of COVID-19 characteristics using a birth-and-death model. doi: <https://doi.org/10.1101/2020.06.23.20138719> (2020)
5. Viswanath, N.C. Prediction of COVID-19 Active and Total Cases After a Fall and Rise of Cases. doi: <https://doi.org/10.1101/2020.07.02.20145045> (2020)
6. MATLAB. Version 9.7.0.1261785 (R2019b Update 3). Natick, Massachusetts: The MathWorks Inc. (2019)
7. Worldometers. Total Coronavirus Cases in India. <https://www.worldometers.info/coronavirus/country/india>. (May 15, 2021)