

Kaplan-Meier plot for treatment groups





Figure 2: Kaplan-Meier plot for treatment groups time to clinical cure (log-rank=16.98).



Kaplan-Meier plot for treatment groups considering the overall patient length of stay

Figure 3: Kaplan-Meier plot for treatment groups considering the overall patient length of stay.

Pt.Group	(N=250)	2.31 (1.54 - 3.5)			-	→ <0.001 ***
Age_Yrs	(N=250)	0.99 (0.97 - 1.0)				0.033 *
Fever	(N=250)	1.48 (0.96 - 2.3)		⊢■		0.075
Dyspnea	(N=250)	0.74 (0.45 - 1.2)	F	B		0.241
S_throat	(N=250)	0.93 (0.33 - 2.6)				0.884
Pneumonia	(N=250)	1.01 (0.82 - 1.3)		↓ ₽ ↓		0.897
Cough	(N=250)	0.98 (0.63 - 1.5)		⊢−−− ∎		0.857
ECG_Find	(N=250)	1.03 (0.87 - 1.2)		⊧ ₩ 1		0.737
CT_Base	(N=250)	0.98 (0.87 - 1.1)		H -		0.719
# Events: 128; Global p-val AIC: 1182.32; Concordance		39982 0.2	0.5	1	2	

Figure 4: Cox regression between the treatments groups regarding suspected covariates at baseline.




Figure 5: Kaplan-Meier plot for treatment groups considering the time to undetectable SARS-COV-2 RNA on two consecutive nasopharyngeal swabs.