Lockdown babies- Birth and new parenting experiences during the 2020 Covid-19 lockdown in South Africa

Elise Farley¹*, Amanda Edwards¹, Emma Numanoglu², Tamsin K. Phillips³

¹ Nudibrink Consulting, Cape Town, South Africa;

² Breastfeeding Matters, Cape Town, South Africa;

³ Division of Epidemiology & Biostatistics, School of Public Health and Family Medicine, University of Cape Town, South Africa.

*Corresponding author: Elise Farley, elisefarley@gmail.com ; +27795601251. 10 Princes Rd, Claremont, Cape Town, South Africa.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

ABSTRACT

Background

Perceived birth experiences of parents can have a lasting impact on children. We explored the birth and new parenting experiences of South African parents during the Covid-19 lockdown.

Methods

We conducted an online cross-sectional survey with consenting parents of babies born in South Africa during 2020. Factors associated with negative birth emotions and probable depression were estimated using logistic regression.

Results

Most of the 520 respondents were females (n= 496, 95%) who gave birth at private hospitals (n=426, 86%). Mothers reported having overall positive birth emotions (n= 399, 80%). Multivariable analysis showed that having the baby during lockdown (adjusted odds ratio (aOR) 5.02; CI 1.28 – 19.66); being diagnosed with Covid-19 (aOR 3.17; CI 1.07 – 9.42); having negative new parenting emotions (aOR 6.07; CI 3.27 – 11.29); a preterm baby (aOR 3.02; CI 1.36 – 6.70) and lockdown related barring of preferred in hospital support (aOR 2.45; CI 1.35 – 4.43) were associated with mothers reporting predominately negative emotions about the birth. Having their chosen delivery method reduced the odds of negative birth emotions (aOR 0.4; CI 0.22 – 0.72). Multivariable analysis showed that having predominantly negative new parenting emotions (aOR 10.75; CI 5.41- 21.37), breastfeeding struggles (aOR 2.16; CI 1.36 – 3.46); lockdown preventing health care access (aOR 2.06; CI 1.20 – 3.54) and creating financial strain (aOR 2.58; CI 1.08 – 6.18) were associated with probable minor depression

Conclusions

Lockdown exacerbated many birth and parenting challenges including mental health and health care access. However, overall experiences were positive and there was a strong sense of resilience amongst parents.

INTRODUCTION

The first 1000 days of a baby's life are the most crucial in terms of development [1]. As part of these 1000 days, the birth of a baby and the perceived birth experiences of the parents can have a lasting impact on the child's life. Subjective birth-experiences have been shown to be associated with the development of post-traumatic stress symptoms in the mother [2, 3] and the mental health of parents can have a long-lasting impact on the child's behavioural and emotional state [4, 5].

The Covid-19 pandemic and ensuing national lockdowns have had widespread global ramifications on every aspect of life, including population wide mental health [6, 7]. These "lockdowns" have severely altered people's social networks and support structures due to the restrictions on seeing family and friends, drastic changes to routines and, for some, impacted economic livelihoods [7]. South Africa entered one of the world's strictest lockdowns on 26 March 2020, and followed a tiered approach in attempts to curb the spread of COVID-19 nationally.

Both having a baby and living through lockdown are stressful situations where support is imperative to ensure positive outcomes for both parents and children. A United Kingdom based study assessing the maternal experiences and infant feeding during the Covid-19 period showed that lockdown has had an impact on parents, resulting in difficulties including changes in delivery plan, a lack of breastfeeding support, no maternal healthcare, mental health challenges and a lack of social support [8]. In a similar manner, we explored the unique birth and new parenting experiences in South Africa during the Covid-19 lockdown.

PARTICIPANTS, ETHICS AND METHODS

STUDY DESIGN, SETTING AND STUDY PARTICIPANTS

We conducted cross-sectional online survey with parents of babies born in South Africa during 2020. Data collection took place from 09 September to 10 October 2020. Participants were aged 18 years or older, living in South Africa at the time of the baby's birth and had a baby born between 1 January 2020 until 10 October 2020.

SAMPLING

A mixture of convenience and snowball sampling was used for the study. We initially sent the survey out to networks of parents who attended antenatal courses run by co-author EN (through her organisation Breastfeeding Matters). This grouping was chosen as it was anticipated that these parents had children born during our study period. The co-authors also sent the survey out on WhatsApp

parenting support groups. Respondents were asked to share the survey with other parents who met the inclusion criteria. Due to the time sensitive nature of the study and restrictions in reaching potential respondents via alternative mechanisms under ongoing lockdown conditions, this recruitment strategy was chosen as it was thought to be the most appropriate for the respondent population who are frequently connected through multiple online formats.

DATA COLLECTION

The data collection tool was created in English using KoBoCollect (KoboToolBox©). A link to the survey was generated and sent out to respondents over email, WhatsApp and via social media. The questionnaire includes a selection of questions derived from a similar United Kingdom study [8] (Supplementary material, Table S1).

Data collected included sociodemographic characteristics, details of the birth and baby, information on feeding. Respondents were asked about their predominant emotions during birth and the new parenting phase. A selection of emotions were listed under the positive (joyful, grateful, relieved, hopeful) and negative (fearful, resentful, inadequate, anxious, sad, overwhelmed) response to guide respondents. Respondents were asked about how Covid-19 had impacted them. Both closed and open response questions were included to explore the full range of emotions experienced.

The 10 point Edinburgh Postnatal Depression Scale [9] was used to assess which respondents showed symptoms of probable depression. It must be noted that this tool was not used as a diagnostic tool. Minor depression was indicated for respondents who scored 10 or more and major depression was indicated for those who scored 13 or more [10]. Minor depression was used in our analysis as this cut off is more sensitive than that for major depression. Participants who scored above 10 were prompted to reach out for support and all participants were provided with resources for psychosocial support services.

ANALYSIS

Due to the small sample size of male respondents (n=24), we conducted separate analyses on responses from female and male respondents. For the male cohort, we performed a descriptive analyses. Categorical variables are reported as frequencies and proportions. Depending on their distribution, continuous variables are summarised using medians and interquartile ranges (IQR) or means and standard deviations (SD).

For the female cohort, we performed a similar descriptive analysis. In addition, we compared the responses between those who reported having predominately positive emotions about their birth, and

those who had predominantly negative emotions about their birth. Comparisons were also made by lockdown level at the time of delivery (5 categories: Alert Level 5: 26 March -30 April, Level 4: 1 – 31 May, Level 3: 1 June – 17 August, Level 2: 18 August – 20 September, Level 1: 21 September – 28 December, after which time adjusted levels were implemented), and in binary between those whose babies were born prior to lockdown (1 January – 25 March 2020) and those whose babies were born during lockdown (25 March – end of data collection on 10 October 2020). Bivariate associations were examined using chi-squared tests, t-tests and Kruskal Wallis tests as appropriate. Missing data numbers are recorded in each table.

Univariable and multivariable analysis were conducted using logistic regression, to estimate factors associated with predominant birth emotions (positive and negative) and probable minor depression (yes or no). Variables chosen for inclusion in the multivariable analysis were those with ten or more cases and a univariable strength of association equivalent to a p-value < 0.2, after assessing collinearity among variables [11]. Odds ratios (OR), adjusted odds ratios (aOR) and their respective 95% confidence intervals (CI) are reported. All quantitative data analysis was conducted with Stata 15 (StataCorp LP, College Station, TX, USA).

To complement quantitative analyses, free text responses were analysed by coinvestigator AE using thematic content analysis [5]. Coding was conducted by hand, allowing for the addition of new concepts in two rounds of analysis. The frequency of each concept was then catalogued and quantified in MS Excel, before high-frequency related concepts were merged into core themes.

ETHICAL STATEMENT

Ethical approval was obtained from the University of Cape Town Human Research Ethics Committee (385/2020). Informed consent was obtained from all respondents by asking them to first read the information sheet and then provide consent by selecting the ‘Yes’ button on the online survey. If respondents declined to participate after reading the information sheet, this was recorded and they were directed to a page with parenting and mental health resources. The survey was anonymous and no identifying information such as names or addresses were collected.

RESULTS

In total, 520 parents completed the survey, 496 (95%) were females and 24 (5%) were males.

FEMALES

SOCIO-DEMOGRAPHIC CHARACTERISTICS

The mean age of female respondents was 32 years (SD 4 years); 327 (66%) were from the Western Cape; 183 (37%) highest qualification was an undergraduate degree; 339 (68%) were employed full-time at the time of the survey; 474 (96%) were married, in a civil partnership or cohabiting with a partner and 426 (86%) gave birth at a private hospital (Table 1). Mothers in our study gave birth to 506 babies in total, 255 were males (50%). For just over half ($n=260$; 53%) of the mothers in the study, this was their first child. The majority were born at a normal gestational age range (37-42 completed weeks of pregnancy [12], $n=436$, 88%). The mean birth weight was 3.2kg (SD 0.6kg), 47 (10%) babies were classified as having a low birth weight (0-2499g), 414 (84%) as normal birthweight (2500 – 3999g) and 32 (7%) as having a high birthweight (>4000g).

LOCKDOWN IMPACT

The Covid-19 lockdown led to mothers reporting delays in registering their babies ($n=231$, 47%) and difficulties in purchasing baby goods ($n=229$, 46%). The main non-medical activities that mothers missed due to lockdown were visiting friends and family ($n=441$, 89%), outdoor exercise ($n=197$, 40%) and going to restaurants ($n=179$, 36%). The majority of babies in the study cohort were born during lockdown (pre-lockdown $n=78$, 16%; level 5 $n=95$, 19%; level 4 $n=88$, 18%; level 3 $n=168$, 34%; level 2 $n=64$, 13% and level 1 $n=3$, 1%). Mothers with babies born prior to lockdown were more likely to have their preferred method of birth ($n=51$; 65%) in comparison to those with babies born during level 4 ($n=48$; 55%), level 3 ($n=97$; 58%) and level 2 ($n=37$; 58%) lockdown ($p=0.009$). There were fewer babies placed skin to skin after birth during lockdown (pre-lockdown $n=67$, 86%; lockdown $n=313$, 75%; $p=0.040$). More mothers reported having a negative birth experience during lockdown (pre-lockdown $n=3$, 4%; lockdown $n=94$, 23%; $p<0.001$). A vast majority of mothers who gave birth during lockdown reported that the pandemic had affected their birth experience (pre-lockdown $n=16$, 21%; lockdown $n=325$, 78%; $p<0.001$) and nearly half reported that someone they had wanted to be at the birth was not able to be there because of the Covid-19 pandemic (pre-lockdown $n=6$, 8%; lockdown $n=198$, 47%; $p<0.001$). A third as many mothers saw a lactation consultant in hospital during lockdown ($n=53$, 13%) in comparison to prior to lockdown ($n=30$, 39%; $p<0.001$). Lockdown did not affect the number of babies born preterm (pre-lockdown $n=9$, 12%; lockdown $n=48$, 12%) (Supplementary material, Table S2).

REGRESSION

Predominant emotions about birth

The majority of mothers reported having overall positive emotions about their birth (positive $n=399$, 80%; negative $n=97$, 20%). The main reasons for negative birth experiences were a traumatic experience ($n=42$, 8%), feeling disconnected from what was happening ($n=37$, 7%), and not having a partner at the birth ($n=34$, 7%). Figure 1 illustrates the words mothers used most commonly to describe their birth experience. The risk factors associated with having predominately negative emotions at birth are shown in Table 2. These include having the baby during lockdown (aOR 5.02; CI 1.28 – 19.66); being diagnosed with Covid-19 (aOR 3.17; CI 1.07 – 9.42); having predominantly negative emotions during the new parenting phase (aOR 6.07; CI 3.27 – 11.29); having a preterm baby (aOR 3.02; CI 1.36 – 6.70) and lockdown related barring of preferred in hospital support for the birth (aOR 2.45; CI 1.35 – 4.43). Being able to have their delivery method reduced the odds of having negative birth emotions (aOR 0.4; CI 0.22 – 0.72) (Table 2).

Mental health

Multivariable analysis showed that respondents aged between 25 and 34 years (aOR 0.25; CI 0.06 – 1.01) and aged 35 years or older (aOR 0.16; CI 0.04– 0.68) were less likely to have symptoms of probable minor depression in comparison to those aged between 18 and 24 years. Those who self-identified as having predominantly negative emotions in the new parenting phase were more likely to show symptoms of probable minor depression (aOR 10.75; CI 5.41– 21.37). Further factors associated with probable minor depression were struggling with breastfeeding (aOR 2.16; CI 1.36 – 3.46); lockdown preventing health care access (aOR 2.06; CI 1.20 – 3.54) and lockdown putting financial strain on families, causing issues paying for living expenses such as rent (aOR 2.58; CI 1.08 – 6.18) (Table 3).

SUPPORT

Very few mothers ($n=69$; 14%) reported accessing professional support for mental health after their birth. However, several reported accessing other forms of support via social media ($n=328$; 66%), websites ($n=309$; 62%) and on WhatsApp moms groups ($n=226$; 46%). Most ($n=474$; 96%) moms are able to access a reliable internet connection (Table 4).

MALES

The median age of male respondents ($n=24$) was 37 years (IQR 29, 39). Male respondents mainly reported having the preferred method of birth ($n=15$; 63%); feeling positive about the birth experience

(n=23; 95%); and that the birth did not affect their parenting (n=17; 71%). The Covid-19 pandemic and resultant lockdown affected 30% (n=7) of fathers birth experiences, mainly due to the father not being able to visit the mother and child in hospital after the birth (n=5; 21%) and because the pandemic led to a change in expected birth plan (n=4; 17%). Twenty-five percent of (n=6) of fathers showed symptoms of major depression. The most widely used forms of online support for fathers are websites (n=20; 83%) and social media (n=11; 46%) (Table 5).

THEMATIC ANALYSIS

Thematic content analysis of open responses indicated five core themes that confirm and deepen the quantitative findings of birth and new parenting experiences under COVID-19.

Social exclusion and isolation leading to increased stress and anxiety

Overwhelmingly the most common theme (n=169), new parents reported strong feelings of social exclusion and isolation, particularly during the birthing experience that impacted on their levels of stress and anxiety as new parents. The lack of social support due to the forced absence of a partner, close family members and friends were the most cited reasons for feelings of social isolation and loneliness.

"Not being able to be with my parents. My mom for help and of course my sister. I never envisioned not getting help, love and support. The celebration I envisioned was taken away."

Interestingly, for some parents the absence of outside family and friends was seen as a positive opportunity for the new family to bond without interruption. Although, this was the case for a significant minority of respondents and typically when at least one significant other was allowed to be present:

"No visitors allowed at the hospital. I saw this as a positive so my husband and I could bond with new-born and become a little family unit"

Rapidly modified health services impact experiences of quality and quantity of care

The second most commonly reported theme (n=88) indicated how rapidly modified health services impacted new parents' experiences of both the quality and quantity of care received during lockdown. New COVID-19 testing protocols, shorter hospital stays and reduced availability of expected services

and appointments (e.g. lactation consultants, midwives, gynaecologists) reduced the quantity of care reported by new parents. While, unclear policies and procedures, conflicting medical advice, unexpected medical procedures and reduced availability and attentiveness of medical personnel impacted perceptions on the quality of care received. These issues are summarised by one new mother who reported her hospital experience during lockdown:

“The Covid test was done 30 minutes after being booked into the hospital, but the test was never done. So when my baby was born 16 hours later, I wasn't allowed near her as my test results were not back yet. I only held my baby 7 hours after she was born - she was given formula without my consent. Dad wasn't allowed in, so he only saw her when we were discharged. I was also served food and drink in take away containers as my test results were not back yet. Only after tests came back negative, were I given food in a plate.”

Anxiety and the virus: increasing the “mental load of motherhood”

Feelings of fear and anxiety linked to COVID-19 permeated the pre-, peri- and post-natal experiences reported by new mothers (n=64). Anxiety was specifically described due to fear of infection, the implications of COVID-19 regulations on access to healthcare and social support during birth, managing work and additional childcare responsibilities, and the social reintegration for both family and baby once lockdown eased. For some, the anxiety of COVID-19 prevented them from seeking the healthcare and support they needed as new parents:

“I'm scared to take my baby anywhere because of Covid-19. I should have done a breastfeeding course prior to delivery but I did not because I was too scared of contracting the virus.”

While others reported the significant impact of COVID-19 and the ensuing lockdown on their mental health and wellbeing:

“Increased mental load of motherhood. Working from home with pre-schooler and baby. Struggle with post-partum anxiety and depression.”

These findings highlight quantitative results that emphasize the need for increased access to mental health support services for new parents, especially under COVID-19 conditions.

The impact of changing work and economic circumstances

Changing work and economic circumstances due to COVID-19 lockdown added additional stress to birth and new parenting experiences (n=51). The economic impact of lockdown introduced higher levels of uncertainty with reports of reduced income, delayed UIF payments, retrenchments and job losses. Combined with reduced access to childcare and the challenge of looking for new work while also adjusting to the arrival of a new baby, many parents were forced to make significant changes to their living circumstances:

"Due to my husband losing his income we are financially in a crisis (even still). We had to move when my baby was just 2 weeks old due to COVID. We cannot afford any help from nanny or schooling. I had to go back to work sooner in order to receive a salary to keep us barely afloat"

A positive birth experience can support a positive transition to parenthood

Despite the rapid changes required by COVID-19 conditions, women were more likely to report a positive transition to parenting if their birthing experience had gone according to plan (n=60):

"I was so happy my delivery went according to plan. I felt empowered and therefore I think I went into those first few weeks much more confidently."

Together with feelings of empowerment and the ability to cope, a positive, supportive birthing experience was also linked to new mothers' experiences of bonding with their babies:

"The calm birth set the scene for a calm baby and mom, and ensured a proper bond between us too."

These findings, although not universal, correspond with the overall positive emotions (80.4%) reported by new mothers in this study. They also highlight the importance of positive birthing experiences as well as the resilience of new mothers giving birth under unique Covid-19 conditions.

DISCUSSION

Our study showed that the Covid-19 lockdown impacted the birth and new parenting experiences of parents whose children were born in 2020. Parents with children born under lockdown were less likely to have their preferred choice of birth method, had worse self-reported birth experiences, were less likely to have skin-to-skin contact with their babies after birth and were frequently unable to have

someone at their birth that they had wanted to be there. Health services were modified due to the pandemic and this impacted new parents' experiences of both the quality and quantity of care received. Despite these challenges, parents reported positive birth and new parenting experiences, showing resilience.

Our results were similar to two studies, one from the United Kingdom [8] and the other from Italy [13], which showed that the Covid-19 lockdown led to distress for parents who reported social isolation as a key challenge. The United Kingdom study also reported that lockdown led to shorter hospital stays, changes to the delivery plans, a lack of breastfeeding support, challenges accessing healthcare, limited social support and mental health difficulties [8], findings corroborated by our study.

Our findings on the prominence of probable depression (63% mothers, 29% fathers), are higher than the usual in this context (33%), which are higher than worldwide pre-Covid-19 estimates (13%) [5]. This finding is mirrored by Canadian study which found increased levels of depression amongst new fathers [14] and a report from an Australian postnatal support hotline that calls increased 75% during the first week of lockdown [15], indicating an increase in new parents requiring mental health support. Our study showed that mental health issues affected many respondents who were younger, had negative emotions about their birth, struggled with breastfeeding, could not access health care and had lockdown related financial concerns. Parents need support to overcome these challenges. Other studies have shown that digital methods of care provision during the post-partum period are effective [16]. These forms of care are reported in our study to be acceptable (specifically social media, websites and WhatsApp moms groups) and should form the basis of interventions for this population.

Mothers in our cohort reported that lockdown restrictions made the new parenting phase more challenging. In particular there was the ban on the sale of baby products which was introduced with Level 5 lockdown and only overturned after the restriction was ruled unconstitutional in early April 2020 [17]. A further challenge was the registration of babies, which by law should happen within 30 days of their birth. Prior to lockdown, registrations took place within hospitals, this was cancelled during lockdown, as were many services at Home Affairs, making registration difficult. The ban on exercise, which was a feature of several lockdown levels, was noted by respondents as something that impact their postpartum quality of life. Exercise in the postpartum period has been linked to several positive changes for the mother including improved psychosocial well-being in other settings [18]. Future lockdown restriction implementers and policy makers should take these challenges into consideration. Baby products should always be included in essential items lists, and be available for sale in shops and online; plans should be made to assist new parents to register their babies in a timely manner and outdoor exercise with precautions such as masks, should be allowed.

Not specific to the Covid-19 lockdown, but a noteworthy finding is the caesarean section rate in our study cohort (63%), a finding similar to other studies of private healthcare users in South Africa (60% [19]; 73% [20], 77% [21]). The WHO states that caesarean section are medically necessary for 10 - 15% of all births [22]. While undoubtedly lifesaving, caesarean sections can lead to severe and permanent physical complications [22, 23], increased post-traumatic stress [23], mental health issues [23] and impairments in quality of life [23] for the mother. Caesarean sections have also been reported to lead to challenges with infant-mother bonding [24], breastfeeding issues [23, 25], an increase risk of non-communicable diseases [26] and adverse effects on children's sensory perception and neuropsychiatric development [24]. Our findings corroborate these reports and show that most mothers wanted to have a natural birth, and that a change in delivery method was directly associated with a negative birth experience which was associated with probable depression. Due to these potential issues, caesarean sections should only be used when medically necessary [22]. The problematically high caesarean section rate in South African private health care needs to be addressed. One such solution would be initiating a midwife led birthing process as opposed to the current Gynaecologist led process [15].

One of the main limitations of our study is that the sample of parents who responded to the survey do not represent the vast majority of South African citizens. The use of convenience sampling in order to rapidly recruit participants in a short window of time resulted in our study sample being mainly degreed, married, middle-to high-income earners who had their babies in private hospitals. The findings from our study can thus only be generalised to this portion of the population. Future studies need to investigate the birth and new parenting experiences in populations of lower socioeconomic status and of public sector patients. The use of convenience sampling, further limiting the generalisability of our results [27]. There is also the potential for recall bias in this study due to the differing lengths of time between the birth and completion of survey, and it is difficult to tease out the independent effects of the lockdown levels of different postpartum periods.

Covid-19 and the ensuing lockdown exacerbated many of the usual challenges of birth and new parenting. Mental health was a key challenge, along with difficulties accessing support. However, overall experiences were positive and there was a strong sense of resilience amongst parents. As the world continues to battle the Covid-19 pandemic on all fronts, efforts to support new parents during this critical phase should not be neglected.

REFERENCES

1. Wrottesley S V., Lamper C, Pisa PT. Review of the importance of nutrition during the first 1000 days: Maternal nutritional status and its associations with fetal growth and birth, neonatal and infant outcomes among African women. *J Dev Orig Health Dis.* 2015;7(2):144–62. DOI: 10.1017/S2040174415001439
2. Garthus-Niegel S, von Soest T, Vollrath M, Eberhard-Gran M. The impact of subjective birth experiences on post-traumatic stress symptoms: a longitudinal study. *Arch Womens Ment Heal.* 2013;16:1–10. DOI: <http://dx.doi.org.ezproxy.uct.ac.za/10.1007/s00737-012-0301-3>
3. Garthus-Niegel S, Knoph C, von Soest T, Nielsen CS, Eberhard-Gran M. The Role of Labor Pain and Overall Birth Experience in the Development of Posttraumatic Stress Symptoms: A Longitudinal Cohort Study. *Birth.* 2014;41(1):108–15. DOI: 10.1111/birt.12093
4. Latva R, Korja R, Salmelin RK, Lehtonen L, Tamminen T. How is maternal recollection of the birth experience related to the behavioral and emotional outcome of preterm infants? *Early Hum Dev.* 2008;84(9):587–94. DOI: 10.1016/j.earlhumdev.2008.02.002
5. Gutierrez-Galve L, Stein A, Hanington L, Heron J, Ramchandani P. Paternal depression in the postnatal period and child development: Mediators and moderators. *Pediatrics.* 2015;135(2):e339–47. DOI: 10.1542/peds.2014-2411
6. Fiorillo A, Gorwood P. The consequences of the COVID-19 pandemic on mental health and implications for clinical practice. *Eur Psychiatry.* 2020;63(1):e32. DOI: 10.1192/j.eurpsy.2020.35
7. Venkatesh A, Edirappuli S. Social distancing in covid-19: What are the mental health implications? *BMJ.* 2020;369(April):2020. DOI: 10.1136/bmj.m1379
8. Vazquez-Vazquez A, Dib S, Rougeaux E, Wells J, Fewtrell M. The impact of the Covid-19 lockdown on the experiences and feeding practices of new mothers in the UK: Preliminary data from the COVID-19 New Mum Study. *medRxiv.* 2020; DOI: doi.org/10.1101/2020.06.17.20133868
9. Cox J, Holden J, Sagovsky R. Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. *Br J Psychiatry* 150782-786. 1987;150:782–6.
10. Matthey S, Henshaw C, Elliott S, Barnett B. Variability in use of cut-off scores and formats on

- the Edinburgh Postnatal Depression Scale - Implications for clinical and research practice. Arch Womens Ment Health. 2006;9(6):309–15. DOI: 10.1007/s00737-006-0152-x
11. Chen Q, Nian H, Zhu Y, Talbot H, Griffin M, Harrell F. Too many covariates and too few cases? – A comparative study. Stat Med. 2017;35(25):4546–58. DOI: 10.1002/sim.7021.
 12. WHO. Preterm birth: Key facts [Internet]. World Health Organization. 2018 [cited 2021 May 8].
 13. Stampini V, Monzani A, Amadori R. A Survey Among Italian Pregnant Women and New-mothers During the COVID-19 Pandemic Lockdown. :1–16.
 14. Cameron EE, Joyce KM, Rollins K, Roos LE. Paternal Depression & Anxiety During the COVID-19 Pandemic. Preprint. 2020; DOI: 10.31234/osf.io/drs9u
 15. Matthews S. More Midwife-based Interventions Could Save Millions of Lives. New Security Beat. 2020 Dec 9;
 16. Gibson L, Hanson VL. “Digital motherhood”: How does technology support new mothers? Conf Hum Factors Comput Syst - Proc. 2013;313–22. DOI: 10.1145/2470654.2470700
 17. Rabkin F. Products for the care of babies and toddlers now essential items. Mail and Guardian. 2020 Apr 6;
 18. Alvar J, Vélez ID, Bern C, Herrero M, Desjeux P, Cano J, Jannin J, de Boer M. Leishmaniasis worldwide and global estimates of its incidence. PLoS One. 2012;7(5). DOI: 10.1371/journal.pone.0035671
 19. R.P. N, J. M. Rising rates of Caesarean sections: An audit of Caesarean sections in a specialist private practice. South African Fam Pract. 2009;51(3):254–8.
 20. Solanki GC, Cornell JE, Daviaud E, Fawcus S. Caesarean section rates in South Africa: A case study of the health systems challenges for the proposed National Health Insurance. South African Med J. 2020;110(8):747–50. DOI: 10.7196/SAMJ.2020.v110i8.14699
 21. Council for Medical Schemes. Epidemiology and trends of caesarean section births in the medical schemes '. 2020.
 22. World Health Organisation. WHO statement on caesarean section rates. World Health Organisation. 2015. DOI: 10.1111/1471-0528.13526

23. Benton M, Salter A, Tape N, Wilkinson C, Turnbull D. Women's psychosocial outcomes following an emergency caesarean section: A systematic literature review. *BMC Pregnancy Childbirth*. 2019;19(1). DOI: 10.1186/s12884-019-2687-7
24. Chen H, Tan D. Cesarean section or natural childbirth? Cesarean birth may damage your health. *Front Psychol*. 2019;10(FEB):1–7. DOI: 10.3389/fpsyg.2019.00351
25. Hobbs AJ, Mannion CA, McDonald SW, Brockway M, Tough SC. The impact of caesarean section on breastfeeding initiation, duration and difficulties in the first four months postpartum. *BMC Pregnancy Childbirth*. 2016;16(1):1–9. DOI: 10.1186/s12884-016-0876-1
26. Stinson LF, Payne MS, Keelan JA. A critical review of the bacterial baptism hypothesis and the impact of cesarean delivery on the infant microbiome. *Front Med*. 2018;5(MAY). DOI: 10.3389/fmed.2018.00135
27. Etikan I, Musa S, Alkassim R. Comparison of convenience sampling and purposive sampling. *Am J Theor Appl Stat*. 2016;5(1):1–4. DOI: 10.11648/j.ajtas.20160501.11

AUTHOR STATEMENTS

ACKNOWLEDGEMENTS

The authors are grateful to the parents of the babies born during lockdown who agreed to be a part of this study.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ETHICAL STATEMENT

Ethical approval was obtained from the University of Cape Town Human Research Ethics Committee (385/2020). Informed consent was obtained from all respondents by asking them to first read the information sheet and then provide consent by selecting the ‘Yes’ button on the online survey. If respondents declined to participate after reading the information sheet, this was recorded and they were directed to a page with parenting and mental health resources. The survey was anonymous and no identifying information such as names or addresses were collected.

FUNDING

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors. The authors had full access to all data and final responsibility for the decision to submit for publication.

AUTHOR CONTRIBUTIONS

All authors were responsible for the study concept, design, data collection, analysis, and interpretation of data. All authors drafted and critically reviewed the manuscript.

TABLES

Table 1- Female respondent's socio-demographic characteristics and the impact of Covid-19 lockdown

		Total	
Variable	Categories	N=496	%
SOCIODEMOGRAPHIC CHARACTERISTICS			
Age respondent (years)	Mean, SD	32.38	4.3
Province of residence	Eastern Cape	15	3.0%
	Free State	4	0.8%
	Gauteng	126	25.4%
	KwaZulu-Natal	16	3.2%
	Limpopo	1	0.2%
	Mpumalanga	3	0.6%
	North West	3	0.6%
	Northern Cape	1	0.2%
	Western Cape	327	65.9%
Highest education	Grade 10	7	1.4%
	Matric	98	19.8%
	Undergraduate degree	183	36.9%
	Honours degree	141	28.4%
	Masters degree	55	11.1%
	PhD	12	2.4%
Employment	Full-time	339	68.3%
	Part-time	49	9.9%
	Unemployed	108	21.8%
How many children do you have?	1	260	52.4%
	2	184	37.1%
	3 or more	47	9.5%
	Missing	5	1.0%
Household status	Married/ civil partnership/ cohabitation	474	95.6%
	Single parent living alone	8	1.6%
	Single parent living with family	14	2.8%
Place of birth	Government clinic	2	0.4%
	Government hospital	37	7.5%
	Home birth	24	4.8%
	Private hospital	426	85.9%
	Other	7	1.4%

Table 2: Factors associated with predominantly negative emotions about the birth, variables with p<0.2 in the univariable analysis include in multivariable analysis									
	Study population			Unadjusted analysis			Adjusted analysis		
	Subjects (N)	Predominant birth emotion negative (N)	Predominant birth emotion negative (%)	OR	95% CI	P	OR	95% CI	P
All subjects	496	97	(19.6)						
Study variables									
Age (years)						0.010			0.699
18-24	21	8	(38.1)	1.00	(Ref)		1.00	(Ref)	
25-24	326	70	(21.5)	0.44	(0.18, 1.11)		0.67	(0.20, 2.27)	
35andabove	149	19	(12.8)	0.24	(0.09, 0.65)		0.57	(0.15, 2.15)	
Highest education level completed						<0.001			<0.001
High school	105	38	(36.2)	1.00	(Ref)		1.00	(Ref)	
Undergrad	183	31	(16.9)	0.36	(0.21, 0.63)		0.29	(0.14, 0.59)	
Postgrad	208	28	(13.5)	0.27	(0.16, 0.48)		0.30	(0.14, 0.63)	
Household status						0.046			0.274
Married/ civil partnership/ cohabitation	474	88	(18.6)	1.00	(Ref)		1.00	(Ref)	
Single parent living on my own	8	3	(37.5)	2.63	(0.62, 11.22)		3.12	(0.59, 16.68)	
Single parent living with family	14	6	(42.9)	3.29	(1.11, 9.72)		2.01	(0.48, 8.51)	
Predominant negative emotions during the new parenting phase						<0.001			<0.001
Positive	345	36	(10.4)	1.00	(Ref)		1.00	(Ref)	
Negative	151	61	(40.4)	5.82	(3.62, 9.35)		6.07	(3.27, 11.29)	
Edinburgh scale, minor depression						<0.001			0.061
No depression	182	14	(7.7)	1.00	(Ref)		1.00	(Ref)	

Probable depression	314	83	(26.4)	4.31	(2.37, 7.86)		2.09	(0.97, 4.52)	
Gestational age of child/ren at birth (weeks)						<0.001			0.006
Normal	436	74	(17.0)	1.00	(Ref)		1.00	(Ref)	
Premature	57	21	(36.8)	2.85	(1.58, 5.17)		3.02	(1.36, 6.70)	
Prolonged	1	1	(100.0)	1.00	(Ref)		1.00	(Ref)	
Did you end up having your preferred choice of delivery method?						<0.001			0.002
No	192	55	(28.6)	1.00	(Ref)		1.00	(Ref)	
Yes	304	42	(13.8)	0.40	(0.25, 0.63)		0.40	(0.22, 0.72)	
Someone I wanted to be at the birth was not able to be there because of Covid-19						<0.001			0.003
No	292	32	(11.0)	1.00	(Ref)		1.00	(Ref)	
Yes	204	65	(31.9)	3.80	(2.37, 6.08)		2.45	(1.35, 4.43)	
Was your baby placed on you skin-to-skin soon after birth?						<0.001			0.080
No	116	44	(37.9)	1.00	(Ref)		1.00	(Ref)	
Yes	380	53	(13.9)	0.27	(0.17, 0.43)		0.57	(0.30, 1.07)	
Were you discharged from hospital after the birth earlier than expected because						0.117			0.411
No	359	64	(17.8)	1.00	(Ref)		1.00	(Ref)	
Yes	137	33	(24.1)	1.46	(0.91, 2.35)		1.30	(0.69, 2.45)	
Did you have enough support around breastfeeding in hospital?						<0.001			0.175
No	175	49	(28.0)	1.00	(Ref)		1.00	(Ref)	
Yes	321	48	(15.0)	0.45	(0.29, 0.71)		0.67	(0.37, 1.20)	
Lockdown prevented me reaching out to healthcare providers for help with issues						0.040			0.420
No	368	64	(17.4)	1.00	(Ref)		1.00	(Ref)	
Yes	128	33	(25.8)	1.65	(1.02, 2.66)		1.31	(0.68, 2.54)	
Lockdown at time of birth						<0.001			0.021
Pre-lockdown	78	3	(3.8)	1.00	(Ref)		1.00	(Ref)	

Lockdown	418	94	(22.5)	7.25	(2.24, 23.53)		5.02	(1.28, 19.66)	
Have you been diagnosed with Covid-19?						0.039			0.038
No	471	88	(18.7)	1.00	(Ref)		1.00	(Ref)	
Yes	25	9	(36.0)	2.45	(1.05, 5.72)		3.17	(1.07, 9.42)	
Covid-19 has affected my ability to pay rent						0.019			0.304
No	425	77	(18.1)	1.00	(Ref)		1.00	(Ref)	
Yes	65	20	(30.8)	2.01	(1.12, 3.59)		1.74	(0.60, 5.03)	
Covid-19 has affected my ability to pay for food						0.135			0.033
No	431	81	(18.8)	1.00	(Ref)		1.00	(Ref)	
Yes	59	16	(27.1)	1.61	(0.86, 3.00)		0.28	(0.09, 0.90)	
Covid-19 has affected my ability to pay for other essentials						0.004			0.700
No	374	63	(16.8)	1.00	(Ref)		1.00	(Ref)	
Yes	116	34	(29.3)	2.05	(1.26, 3.32)		1.15	(0.57, 2.31)	
OR = odds ratio; CI = confidence interval; P = p value from logistic regression model									

Table 3: Factors associated with probable minor depression, variables with p<0.2 in the univariable analysis include in multivariable analysis									
	Study population			Unadjusted analysis			Adjusted analysis		
	Subjects (N)	Edinburgh scale, minor depression N	Edinburgh scale, minor depression (%)	OR	95% CI	P	OR	95% CI	P
All subjects	496	314	(63.3)						
Study variables									
Age (years)						0.007			0.023
18-24	21	18	(85.7)	1.00	(Ref)		1.00	(Ref)	
25-24	326	215	(66.0)	0.32	(0.09, 1.12)		0.25	(0.06, 1.01)	
35 and above	149	81	(54.4)	0.20	(0.06, 0.70)		0.16	(0.04, 0.68)	
Highest education level completed						0.048			0.729
High school	105	77	(73.3)	1.00	(Ref)		1.00	(Ref)	
Undergrad	183	114	(62.3)	0.60	(0.36, 1.02)		0.79	(0.42, 1.51)	
Postgrad	208	123	(59.1)	0.53	(0.31, 0.88)		0.78	(0.41, 1.48)	
Predominant birth emotion negative						<0.001			0.151
Positive	399	231	(57.9)	1.00	(Ref)		1.00	(Ref)	
Negative	97	83	(85.6)	4.31	(2.37, 7.86)		1.70	(0.82, 3.52)	
Predominant negative emotions during the new parenting phase						<0.001			<0.001
Positive	345	174	(50.4)	1.00	(Ref)		1.00	(Ref)	
Negative	151	140	(92.7)	12.51	(6.54, 23.94)		10.75	(5.41, 21.37)	
Actual birth method						0.118			0.133
C-section	311	205	(65.9)	1.00	(Ref)		1.00	(Ref)	
Vaginal	185	109	(58.9)	0.74	(0.51, 1.08)		0.69	(0.43, 1.12)	
Was your baby placed on you skin-to skin soon after birth?						0.036			0.979
No	116	83	(71.6)	1.00	(Ref)		1.00	(Ref)	
Yes	380	231	(60.8)	0.62	(0.39, 0.97)		0.99	(0.56, 1.76)	
Were you discharged from hospital after the birth earlier than expected because						0.086			0.149
No	359	219	(61.0)	1.00	(Ref)		1.00	(Ref)	
Yes	137	95	(69.3)	1.45	(0.95, 2.20)		1.45	(0.87, 2.42)	
Struggled with breastfeeding						<0.001			0.001
No	221	122	(55.2)	1.00	(Ref)		1.00	(Ref)	

Yes	275	192	(69.8)	1.88	(1.30, 2.72)		2.16	(1.36, 3.46)	
Did you have enough support around breastfeeding in hospital?						0.110			0.713
No	175	119	(68.0)	1.00	(Ref)		1.00	(Ref)	
Yes	321	195	(60.7)	0.73	(0.49, 1.07)		1.10	(0.67, 1.80)	
Did you access any mental health support services after your 2020 birth?						0.014			0.111
No	427	261	(61.1)	1.00	(Ref)		1.00	(Ref)	
Yes	69	53	(76.8)	2.11	(1.17, 3.81)		1.74	(0.88, 3.44)	
Lockdown prevented me reaching out to healthcare providers for help with issues						<0.001			0.009
No	368	213	(57.9)	1.00	(Ref)		1.00	(Ref)	
Yes	128	101	(78.9)	2.72	(1.70, 4.37)		2.06	(1.20, 3.54)	
Have you been diagnosed with Covid-19?						0.084			0.226
No	471	294	(62.4)	1.00	(Ref)		1.00	(Ref)	
Yes	25	20	(80.0)	2.41	(0.89, 6.53)		1.98	(0.66, 5.95)	
Covid-19 has affected my ability to pay rent						0.004			0.033
No	425	259	(60.9)	1.00	(Ref)		1.00	(Ref)	
Yes	65	52	(80.0)	2.56	(1.35, 4.85)		2.58	(1.08, 6.18)	
Covid-19 has affected my ability to pay for food						0.007			0.582
No	431	264	(61.3)	1.00	(Ref)		1.00	(Ref)	
Yes	59	47	(79.7)	2.48	(1.28, 4.81)		0.75	(0.27, 2.09)	
Covid-19 has affected my ability to pay for other essentials						<0.001			0.063
No	374	221	(59.1)	1.00	(Ref)		1.00	(Ref)	
Yes	116	90	(77.6)	2.40	(1.48, 3.88)		1.79	(0.97, 3.31)	
OR = odds ratio; CI = confidence interval; P = p value from logistic regression model									

Table 4: Support accessed by mothers after birth in 2020

		N=496	%
Did you access any mental health support services after your 2020 birth?	Yes	69	13.9%
	No	427	86.1%
Did you find any of these resources helpful during this time?	Social media	328	66.1%
	Websites	309	62.3%
	WhatsApp moms groups	226	45.6%
	Blogs	173	34.9%
	Other	46	9.3%
Did you have a reliable internet connection to be able to access support online?	Yes	474	95.6%
	No	22	4.4%
Were internet data charges too expensive for you to be able to use the internet to access support?	Yes	55	11.1%
	No	441	88.9%

Table 5: Fathers of babies born during 2020, birth and new parenting experiences			
Variable	Categories	N=24	%
SOCIODEMOGRAPHIC CHARACTERISTICS			
Age respondent (years)	Median, IQR	36.5 years	29, 38.5 years
BIRTH			
Preferred method of birth occurred	Yes	15	62.5%
	No	9	37.5%
Predominant birth emotion	Negative	1	4.17%
	Positive	23	95.83%
	Missing	0	0.00%
Happy with birth experience	Yes	21	87.50%
	No	3	12.50%
NEW PARENTING			
Birth affected new parenting	Yes	2	29.17%
	No	17	70.83%
	Missing	4	
Predominant new parenting emotions	Negative	2	8.33%
	Positive	22	91.67%
COVID-19			
Covid-19 affected birth experience	Yes	7	29.17%
	No	17	70.83%
If yes, how has Covid-19 affected birth experience	I could not visit us after the birth	5	20.83%
	Our birth plan changed due to the pandemic	4	16.67%
	Friends and family not able to support	4	16.67%
	I was not allowed to be at the birth	3	12.50%
	Caesarean section was offered earlier than expected	2	8.33%
	Treated as Covid-19 positive until test results came back negative	2	8.33%
Has the Covid-19 pandemic affected the employment status of you or your partner?	Yes	5	20.83%
	No	19	79.17%
Has the Covid-19 pandemic affected any of the following	My ability to pay rent	1	4.17%
	My ability to pay for other essentials (including medicine, utility bills etc.)	3	12.50%
MENTAL HEALTH			
Edinburgh scale, minor depression	Likely no depression (0-9)	17	70.83%
	Probable minor depression	7	29.17%
Edinburgh scale, major depression	Likely no major depression	18	75.00%
	Probable major depression	6	25.00%
Did you find any of these resources helpful during this time?	Blogs	7	29.17%
	Websites	20	83.33%
	Social media	11	45.83%
	WhatsApp moms groups	8	33.33%
	Other	3	12.50%

FIGURES

Figure 1: Words mothers used to describe their birth experience

SUPPLEMENTARY MATERIAL

S1- STUDY QUESTIONNAIRE

ID respondent	Automatically generated
Date and time of interview	Automatically generated
Eligibility questions	
Are you aged 18 years or older?	Yes No (not eligible- end survey)
Are you mother/father of a baby born in 2020?	Yes No (not eligible- end survey)
Do you live in South Africa?	Yes No
If yes, in which province?	Eastern Cape Free State Gauteng KwaZulu-Natal Limpopo

	Mpumalanga Northern Cape North West Western Cape
If no, which country do you live in?	Country: (not eligible- end survey)
Sociodemographic Questions	
Age (in years)	
Gender	Female
	Male
Highest education level completed	None
	Primary school (grade 7/ standard 5)
	Grade 10/ standard 8
	Matric
	Undergraduate degree
	Honours degree
	Masters degree

	PhD
Employment	Unemployed
	Employed part-time
	Employed full-time
Employment type	Government
	Non-profit
	For-profit/ business
	Self-employed
	Public institution (academia etc.)
How many children do you have?	
What is your relationship to the child in question	Mother
	Father
Household status	Single parent living on my own
	Single parent living with family
	Married/ civil partnership/ cohabitation

Before birth	
Which service-providers did you consult with prior to your birth?	Mid-wife
	Doula
	Lactation consultant
	Gynaecologist
	Public clinic
	Private clinic
	Pediatrician
	None
	Other:
Were these service providers helpful?	Always
	Sometimes
	Never
How did you intend to feed your baby before you gave birth?	Breastfeed Formula feed Mixed (formula and breastfed) Did not have a preference

Before birth, were you planning on vaccinating your child according to the government advised vaccination schedule?	Yes No
If no, why not?	
Birth	
How many children did you give birth to in 2020?	One Twins Triplets
When was your child born?	DD/MM/20_
Where was the child in question born? (The child born in 2020).	Private hospital
	Government hospital
	Government clinic
	Home birth
	Other:
Gestational age of child at birth (weeks and days)	
Baby's birth weight (kg)	

*Repeat this question according to number of children born	
Baby's sex	Female
*Repeat this question according to number of children born	Male
What was your birth plan prior to delivery?	Vaginal no medication
	Vaginal with epidural
	Vaginal with other pain medication
	Vaginal – induced and no other medication
	Vaginal – induced and epidural or other pain medication
	Planned C-section
Did you end up having your preferred choice of delivery method?	Yes
	No
If no, what delivery method did you have?	Vaginal no medication
	Vaginal with epidural
	Vaginal with other pain medication
	Vaginal – induced and no other medication
	Vaginal – induced and epidural or other pain medication
	Unplanned C-section

What one word would you use to describe your birth experience?	
What are the predominant emotions you feel about your birth experience?	Positive (joyful, grateful, relieved, hopeful) Negative (fearful, resentful, inadequate, anxious, sad, overwhelmed)
Was your baby placed on you skin-to skin soon after birth?	
How many nights did you stay in hospital?	
Were you happy with your birth experience?	Yes
	No
If no, why? *Select all that apply	I was unhappy with the delivery method
	I was unhappy with the care provided at the hospital
	I was unhappy with the care provided by the doctor
	I found the experience traumatic
	I was in too much pain
	The hospital staff did not show compassion or care during the birth
	I could not have my partner with me during the birth
	I feel I was not in control of the choices made during the birth process

	My labor was too long
	I was unhappy as I ended up using pain medication
	I did not want the medical interventions that were used (forceps, suction etc.)
	My birth changed from fine too dangerous in a short amount of time
	I felt disconnected from what was happening
	I had complications
	I had to have a general anaesthetic
	No-one explained what was happening to me
	My baby was in danger
	Other:
Do you feel your delivery experience affected your parenting during the first few weeks of your baby's life?	Yes
	No
If yes, how?	Free text:
Did you have enough support around breastfeeding in hospital?	Yes
	No
Did you see a lactation consultant while in hospital?	Yes

	No
If no, would you have liked to see a lactation consultant in hospital?	Yes
	No
2020-parenting	
What one word would you use to describe your 2020 parenting experience?	
What are the predominant emotions you feel about your 2020 parenting experience?	Positive (joyful, grateful, relieved, hopeful) Negative (fearful, resentful, inadequate, anxious, sad, overwhelmed)
How are you currently feeding your baby?	Exclusive breastfeeding (breastfeeding only) Breastfeeding with solid food Formula feeding Formula feeding and solid foods Mixed (breastfeeding and formula) Mixed feeding (breastfeeding and formula) with solid foods

Have you struggled at all with breastfeeding?	Yes No Never breastfed
If yes, why?	I could not get my baby to latch I feel my milk supply is too low I did not know if my baby was getting enough milk My breasts were engorged I got mastitis My baby has an anatomical mouth problem which makes breastfeeding (such as a tongue tie or cleft) I have flat or inverted nipples My nipples are very painful Clogged milk ducts My baby got nipple confusion My breast implants affected my milk ducts Other:
Which service providers did you consult with after your birth?	Mid-wife

	Doula
	Lactation consultant
	Gynecologist
	Public clinic
	Private clinic
	Pediatrician
	None
	Other:
<p>Were these service providers helpful?</p> <p>*Answer for each service provider listed above</p>	Always
	Sometimes
	Never
<p>Did you see a lactation consultant after discharge from hospital?</p>	Yes
	No
<p>If no, would seeing a lactation consultant after discharge from hospital have improved your breastfeeding experience?</p>	Yes
	No
<p>Did you ever get conflicting advice around breastfeeding from healthcare professionals?</p>	Yes
	No

Have you vaccinated your children at the government suggested times?	☐ Yes ☐ No
Covid-19	
If no, was Covid-19 the reason for the delays in vaccinating your child?	☐ Yes ☐ No
Which, if any, of these options might make you feel safer to get your child vaccinated??	☐ Drive-through ☐ Home visit ☐ An option for vaccinations at a site away from the main clinic (in local church, school or community centre) ☐ None of these
Was there someone you had wanted to be with your during your delivery who was not able to be there because of Covid-19?	☐ Yes ☐ No
Who was unable to be with you during your labour and delivery because of Covid-19?	☐ Partner/ husband/ wife ☐ Mother ☐ Other family member ☐ Doula ☐ Midwife

	Other
Were you discharged from hospital after the birth earlier than expected because of Covid-19?	Yes No
Have you been diagnosed with Covid-19?	Yes No
If yes, how did this affect you?	I had to be isolated from my partner
	My partner could not help with the baby
	I was sick during pregnancy
	I was sick and had to look after the baby
	I had no symptoms
	I was very worried about passing the infection on to others
	Other:
Was your partner diagnosed with Covid-19?	Yes No
If yes, how did this affect you?	My partner had to be isolated from me
	My partner could not help with the baby
	I was worried about becoming infected

	I was worried about my partner spreading the infection on to others
	Other:
Have you been in close contact with someone diagnosed with Covid-19?	Yes
	No
If yes, how did this affect you?	I was worried about becoming infected
	I was worried my baby would become infected
	I was frustrated as I could not take care of the Covid-19 infected person
	Other:
Do you feel the Covid-19 pandemic affected your birth experience?	Yes
	No
If yes, in what way? *Tick all that apply	Me/my partner was not allowed to be at the birth
	Me/my partner could not visit us after the birth
	My baby was taken away from me after birth (as I was Covid-19 positive)
	My birth plan changed due to the pandemic
	Delivery method was changed by doctor
	Doctor was unavailable to assist at birth

	Induction was suggested earlier than expected
	C-section was offered earlier than expected
	Doctor advised me to contact other service providers who offer home births
	My doula was not allowed to attend the delivery
	Other:
Has the Covid-19 pandemic affected you after delivery during the 2020 parenting phase?	Yes
	No
If yes, in what way?	Limited social visits
	Limited support available from family due to Covid-19 restrictions
	Limited support available from medical service providers due to Covid-19 restrictions
	Not able to purchase necessary goods
	Anxiety about going to shops to buy groceries
	Anxiety about your baby getting sick
	Anxiety over other family member getting sick
	Not being able to have the help (domestic worker, night nurse, nanny etc.) you had planned on during this time

	Worrying about going back to work and not being able to find childcare
Did you or other members of the family (not the baby) have health issues during lockdown?	Yes
	No
If yes, did you seek care?	Yes
	No
If no, was that because of lockdown?	Yes
	No
Were there any health issues unrelated to the baby that you or your family members wanted to seek care for but didn't because of Covid-19?	Yes
	No
Were there baby items you needed that you couldn't buy during lockdown?	Yes
	No
Which non-medical services, activities and goods did you miss the most because of lockdown? (Select top 3)	Restaurants
	Hairdressers
	Home improvement shops (Builders Warehouse etc.)
	Outdoor exercise
	Online clothes shopping

	Alcohol
	Travel between provinces
	Travel in your city
	Shopping at grocery store
	Visiting friends and family
	Other:
Have you faced delays in registering the birth of your baby because of Covid-19?	Yes
	No
Has the Covid-19 pandemic affected the employment status of you or your partner?	Yes
	No
If yes, how?	I lost my job
	My partner lost their job
	We had to take salary cuts
	My business has closed
	My business has lost many clients
	Other:

<p>Has the Covid-19 pandemic affected any of the following:</p> <p>*Select all that apply</p>	<p>My ability to pay rent</p> <p>My ability to pay for food</p> <p>My ability to pay for other essentials (including medicine, utility bills etc.)</p>
<p>Please tell us about any other ways the Covid-19 pandemic influences your birth or 2020-parenting experiences</p>	<p>Free text</p>
<p>Mental health [9]</p>	
<p>For this section, please check the answer that comes closest to how you have felt IN THE PAST 7 DAYS, not just how you feel today.</p>	
<p>I have been able to laugh and see the funny side of things</p>	<p>As much as I always could</p> <p>Not quite so much now</p> <p>Definitely not so much now</p> <p>Not at all</p>
<p>I have looked forward with enjoyment to things</p>	<p>As much as I ever did</p> <p>Rather less than I used to</p> <p>Definitely less than I used to</p> <p>Hardly at all</p>
<p>I have blamed myself unnecessarily when things went wrong</p>	<p>Yes, most of the time</p>

	Yes, some of the time Not very often No, never
I have been anxious or worried for no good reason	No, not at all Hardly ever Yes, sometimes Yes, very often
I have felt scared or panicky for no very good reason	Yes, quite a lot Yes, sometimes No, not much No, not at all
Things have been getting on top of me	Yes, most of the time I haven't been able to cope at all Yes, sometimes I haven't been coping as well as usual No, most of the time I have coped quite well No, I have been coping as well as ever
I have been so unhappy that I have had difficulty sleeping	Yes, most of the time Yes, sometimes

	Not very often No, not at all
I have felt sad or miserable	Yes, most of the time Yes, quite often Not very often No, not at all
I have been so unhappy that I have been crying	Yes, most of the time Yes, quite often Only occasionally No, never
The thought of harming myself has occurred to me	Yes, quite often Sometimes Hardly ever Never
Support	
Did you access any mental health support services after your 2020 birth?	Yes No

Was this support online or face to face?	☐ Online ☐ Face-to-face ☐ Telephonically
Was this support helpful?	☐ Yes ☐ No
Did you have a reliable internet connection to be able to access support online?	☐ Yes
	☐ No
Were internet data charges too expensive for you to be able to use the internet to access support?	☐ Yes
	☐ No
Did you find any of these resources helpful during this time?	☐ Blogs
	☐ Websites
	☐ Social media
	☐ WhatsApp moms groups
	☐ Other
Any other comments?	
Thank you for your participation in this survey.	

If you require counselling or mental health assistance, please contact:

Suicide Helpline

National help-line: 0800 567 567

The South African Depression and Anxiety Group

National help-line: 0800 21 22 23 or 0800 456 789 or 0800 20 5026

Live online counselling service: www.sadag.org

Lifeline

Helpline: 0861 322 322

Website: <http://lifelinesa.co.za/>

Perinatal Mental Health Project

Website: <https://pmhp.za.org/messages-for-mothers/>

Resources: https://pmhp.za.org/wpcontent/uploads/PMHP_MaternalMentalHealth_Resource-Directory_2017.pdf

Cipla free mental health services:

Helpline: 0800 4567 789

WhatsApp 076 88 22 775

Website: <https://www.cipla.co.za/mental-health/free-mental-health-resources-in-south-africa/>

Coronavirus SA

Coronavirus Hotline: 0800 029 999

WhatsApp: 060 012 3456

Website: <https://sacoronavirus.co.za/> or

<https://coronavirus.datafree.co/>

Supplementary material, Table S2: Comparison of respondents' experiences of birth and the new parenting phase between those with babies born prior to lock down and those born during lockdown								
		Total		Pre-lockdown		Lockdown		Ch sq
Variable	Categories	N=496	%	N=78	%	N=413	%	
BIRTH								
Covid-19 affected birth experience	Yes	341	68.75%	16	20.52%	325	77.75%	<0.001
	No	155	31.25%	62	79.49%	93	22.25%	
How did the lockdown affect your birth experience	My partner could not visit us after the birth	152	30.60%	1	6.25%	151	46.46%	0.002
	My birth plan changed due to the pandemic	60	12.10%	1	6.25%	59	18.15%	0.22
	My partner was not allowed to be at the birth	53	10.70%	1	6.25%	52	16.00%	0.29
	My doula was not allowed to attend the delivery	52	10.50%	1	6.25%	51	15.69%	0.31
	C-section was offered earlier than expected	29	5.80%	1	6.25%	28	8.62%	0.74
	Induction was suggested earlier than expected	24	4.80%	1	6.25%	23	7.08%	0.9
	Delivery method was changed by doctor	19	3.80%	0	0.00%	19	5.57%	0.32
	My baby was taken away from me after birth (as I was Covid-19 positive)	4	0.80%	0	0.00%	4	1.23%	0.66
Birth plan prior to birth (binary)	Vaginal	334	67.34%	49	62.82%	285	68.18%	0.35
	C-section	162	32.66%	29	37.18%	133	31.82%	
Preferred method of birth occurred	Yes	304	61.29%	51	65.38%	253	60.53%	0.42
	No	192	38.71%	27	34.62%	165	39.47%	
Actual birth method (binary)	Vaginal	185	37.30%	26	33.33%	159	38.04%	0.43
	C-section	311	62.70%	52	66.67%	259	61.96%	
Skin to skin after birth	Yes	380	76.61%	67	85.90%	313	74.88%	0.04
	No	116	23.39%	11	14.10%	105	25.12%	
Predominant birth emotion	Negative	97	19.56%	3	3.85%	94	22.49%	<0.001
	Positive	399	80.44%	75	96.15%	324	77.51%	

Feeding plan before birth	Breastfeed	438	88.31%	65	83.33%	373	89.23%	0.33
	Formula feed	22	4.44%	6	7.69%	16	3.83%	
	Mixed (formula and breastfed)	34	6.85%	7	8.97%	27	6.46%	
	Did not have a preference	2	0.40%	0	0.00%	2	0.48%	
Saw lactation consultant in hospital	Yes	83	16.73%	30	38.46%	53	12.68%	<0.001
	No	413	83.27%	48	61.54%	365	87.32%	
Gestational age of baby at birth	Premature (0-36 weeks)	57	11.5%	9	11.50%	48	11.50%	0.91
	Normal (37-42 weeks)	436	87.9%	69	88.50%	367	87.80%	
	Prolonged (43 or more weeks)	1	0.2%	0	0.00%	1	0.20%	
	Missing	2	0.4%	0	0.00%	2	0.50%	
Someone I wanted to be at the birth was not able to be there because of Covid-19	Yes			6	7.69%	198	47.37%	<0.001
	No	204	41.13%	72	92.31%	220	52.63%	
NEW PARENTING								
Covid-19 affected me after delivery	Yes	412	83.06%	67	85.90%	345	82.54%	0.47
	No	84	16.94%	11	14.10%	73	17.46%	
How Covid-19 affected me after delivery	Limited social visits	339	68.30%	57	85.07%	282	81.74%	0.51
	Limited support available from family due to Covid-19 restrictions	321	64.70%	54	80.60%	267	77.39%	0.56
	Anxiety about your baby getting sick	305	61.50%	48	71.64%	257	74.49%	0.63
	Anxiety about going to shops to buy groceries	304	61.30%	49	73.13%	255	73.91%	0.89
	Anxiety over other family member getting sick	267	53.80%	43	64.18%	224	64.93%	0.91
	Not being able to have the help (domestic worker, night nurse, nanny etc.) you had planned on during this time	232	46.80%	49	73.13%	183	53.04%	0.002
	Worrying about going back to work and not being able to find childcare	173	34.90%	38	56.72%	135	39.13%	0.008
	Not able to purchase necessary goods	126	25.40%	23	34.33%	103	29.86%	0.47

	Limited support available from medical service providers due to Covid-19 restrictions	109	22.00%	24	35.82%	85	24.64%	0.06
Predominant new parenting emotions	Negative	151	30.44%	25	32.05%	126	30.14%	0.74
	Positive	345	69.56%	53	67.95%	292	69.86%	
Struggled with breastfeeding	Yes	275	55.44%	40	51.28%	235	56.22%	0.11
	No	203	40.93%	32	41.03%	171	50.91%	
	Never breastfed	18	3.63%	6	7.69%	12	2.87%	
Reason for struggling with breastfeeding	My nipples were painful and cracked	122	24.60%	20	25.64%	102	24.40%	0.55
	I could not get my baby to latch	113	22.80%	13	16.67%	100	23.92%	0.37
	I feel my milk supply is too low	99	20.00%	13	16.67%	86	20.57%	0.64
	I did not know if my baby was getting enough milk	84	16.90%	8	10.26%	76	18.18%	0.23
	My breasts were engorged	60	12.10%	7	8.97%	53	12.68%	0.57
	I have flat or inverted nipples	43	8.70%	2	2.56%	41	9.81%	0.11
	Clogged milk ducts	38	7.70%	10	12.82%	28	6.70%	0.07
	My baby has an anatomical mouth problem which makes breastfeeding (such as a tongue tie or cleft)	31	6.30%	5	6.41%	26	6.22%	0.69
	I got mastitis	24	4.80%	5	6.41%	19	4.55%	0.49
	My baby got nipple confusion	16	3.20%	1	1.28%	15	3.59%	0.47
Conflicting breastfeeding advice	Yes	209	42.14%	33	42.31%	176	42.11%	0.97
	No	287	57.86%	45	57.69%	242	57.89%	
MENTAL HEALTH								
Edinburgh scale (*More sensitive)	Likely no depression (0-9)	182	36.69%	28	35.90%	154	36.84%	0.87
	Probable minor depression (10 or more)	314	63.31%	50	64.10%	264	63.16%	
Edinburgh scale	Likely no major depression (0-12)	282	56.85%	49	62.82%	233	55.74%	0.25
	Probable major depression (13 or more)	214	43.15%	29	37.18%	185	44.26%	