

Māori and Pacific People in New Zealand have higher risk of hospitalisation for COVID-19


Supplementary Material

Contents

- Supplementary Table 1. Ethnicity breakdown of the data.
- Supplementary Table 2 (see supplementary_table2.xlsx). Results of sensitivity analysis for the model for probability of hospitalisation.
- Supplementary Table 3 (see supplementary_table3.xlsx). Results of sensitivity analysis for the model for length of hospital stay.
- Supplementary Figure 1. Model results including 90% confidence intervals for probability of hospitalisation by age and ethnicity.

	Māori	Pacific	Asian	NZ Euro/other
Māori	99	13	1	59
Pacific	13	166	8	15
Asian	1	8	283	6
NZ Euro/other	59	15	6	1171
Total	172	202	298	1251

Supplementary Table 1. Breakdown of the number of cases in the data set by ethnicity showing cases with a single ethnicity (bold numbers) and cases with two ethnicities (non-bold numbers). In addition to the 1,821 single and dual ethnicity cases represented in the table, there were 8 cases with three ethnicities, of which 6 were Māori, Pacific and NZ European/other, and 2 were Pacific, Asian and NZ European/other.



Supplementary Figure 1. Model results including 95% confidence intervals for probability of hospitalisation by age and ethnicity for individuals without underlying health conditions (top row of plots) and with underlying health conditions (bottom row of plots).