

Supplemental Information

S1: Intervention Procedures

Nutrition Interventions

The Community Nutrition component used evidence-based integrated nutrition interventions for the “first 1,000 days” of life. Village Health Support Groups (VHSGs), supervised by health workers and Commune Councils for Women and Children (CCWC), sought to improve childcare and development at multiple levels: individual, family, and community. Five core activities comprised the community initiative designed to prevent malnutrition:

- 1) **Community Dialogues:** This activity was led quarterly by the Village Chief and VHSGs. The community gathered to talk, decide, and take action together to support all children to grow healthy. Communities reviewed progress of creating a healthy environment, discussed one key action to jointly address challenges, and decided together how everyone can come together to achieve this action.
- 2) **Caregiver Group Education Sessions:** Caregiver Groups were peer-led groups of women who use a 13-session experiential learning manual following each of the key behaviors promoted by NOURISH. NOURISH trains two members per group to facilitate monthly sessions for their group, with support from elder women in the community and trained Community Agents.
- 3) **Growth Monitoring and Promotion:** VHSGs monitored every child every month. Children who were sick or not growing well were referred to health centers or referral hospitals, as appropriate, and followed up at home after treatment.
- 4) **Home visits:** VHSGs and Mother Support Group (MSG) members provided tailored interpersonal communication during home visits to promote childcare and feeding practices, home hygiene, and proper use of latrines and handwashing stations. Home visits were conducted for pregnant women, caregivers of children 9-11 months old and caregivers of children not growing well.
- 5) **Village Fairs:** This activity was held twice a year for each village. Village fairs offered women and their families’ hands-on learning experiences that bring together health/nutrition, WASH and agriculture using games, demonstrations and practice, interactive discussions and latrine marketing and sales by local participating sanitation suppliers.

CCT acted as a social safety net mechanism for poor “first 1,000 days” families, serving as an incentive for women to access services, practice specific behaviors, and overcome constraints related to poverty. Eligible families (based on poverty status) could receive up to six payments, for a total of \$65 over the first 1,000 days of a child’s life, which were transferred directly into women’s bank accounts after completed use of health and nutrition services.

- 1) **First transfer:** \$12.50 at 1 month postpartum. Conditions: At least four antenatal care visits, delivery in a health center, and at least two postnatal care visits.
- 2) **Remaining five transfers:** \$10 for the second to fifth transfers and \$12.50 for the last transfer over the next 23 months postpartum. Conditions: Monthly monitoring of children’s growth through Growth Monitoring and Promotion (GMP) at the health centers or in the community, and handwashing station at home.

Vouchers served as another mechanism to encourage demand and overcome access constraints related to poverty. Vouchers were distributed to poor “first 1,000 days” families in communes where the CCT is implemented and is redeemable for discounts on water filters (\$5 co-payment) and two food baskets (\$5 co-payment). Vouchers were only distributed in combined intervention groups.

SBCC consisted of media and materials to promote key behaviors in health/nutrition, sanitation/hygiene, and agriculture. The project's SBCC framework was grounded in evidence of what works in social and behavior change and foundational work done by NOURISH in its first year. On the nutrition side, SBCC supported all the Community Nutrition activities, described above, and was implemented by community change agents (VHSGs and caregiver peer groups).

- 1) **Grow Together:** The campaign focused on 13 key stunting prevention behaviors spanning health, nutrition, WASH, and agriculture to stimulate relevant actions for children to grow and reach their full potential. It was not possible to exclude the WASH messaging from the Grow Together campaign, so caregivers received information on all 13 behaviors as part of the nutrition programming.
- 2) To complement the print materials, the SBCC media plan included three television spots including the foundational Grow Together TV spot, latrine construction and Small Fish Powder; 13 “soundbites”; an advocacy package for local leaders; and more than 20 print materials carrying the same “look and feel” to link to core values and motivations to take action.
- 3) The “first 1,000 days” family SBCC package was centered on a Family Commitment Card enumerating the critical practices and allowing families to prioritize behaviors and visualize successes and gaps. As the Family Card filled with accomplishments, the family was recognized as a growth champion with a child book and other incentives to mark its accomplishment. A behavior wheel checklist to guide home visits showing health/nutrition and sanitation/hygiene practices supplemented the Family Card.

Sanitation Interventions

CLTS aimed to achieve sustained behavior change through the process of community “triggering” leading to spontaneous and long-term abandonment of open defecation practices. This one-time triggering event was conducted in collaboration with the Ministry of Rural Development and provincial and district departments of rural development. In alignment with national Open Defecation Free certification guidelines, CLTS covered entire villages to minimize the risk of fecal-oral contamination for all children. Following CLTS triggering, NOURISH monitored the commitments of families and communities through door-to-door visits. These visits took place at least 5 times per village and were used to also raise awareness and create demand for sanitation/latrines.

Latrine vouchers were a targeted subsidy to poor households in villages that reached 75% sanitation coverage. Vouchers were redeemable for a discount on latrine materials (\$15 co-payment). In the combined-intervention villages, latrine vouchers were initially linked to the CCT program, and so were only offered to those beneficiaries. However, this requirement was phased out and latrine vouchers were eventually available to all poor households in eligible villages.

Supply-side support consisted of collaborating with private and public sector actors to develop locally sensitive market-oriented approaches for the integrated business service centers around “first 1,000 days” products and services. NOURISH encouraged knowledge sharing across small- and medium-sized enterprises (SMEs) as well as utilized existing resource centers and agencies to develop the capacity of SMEs for effective service delivery and to increase their outreach to poor and relatively remote areas. NOURISH identified a number of successful businesses within or outside the project area and organized interfaces between new and existing businesses to give mutual learning opportunities between SMEs and develop linkages for possible collaboration. In addition, suppliers were linked to communes where CLTS triggering had occurred so they could follow-up with households that committed to purchasing or building latrines.

SBCC on the sanitation side consisted of sanitation campaigns in primary schools to ensure children become agents of change and carry new behaviors home. As change agents, children have the potential to convince their families to construct latrines or purchase a handwashing station, and to use them.


S2: Participant inclusion and exclusion

The universe of communes targeted by the program was selected based on two criteria:

- 1) Communes where at least 30% of the population was living below the poverty line according to the 2011 Cambodia Ministry of Planning's Commune Database; and
- 2) Communes where latrine subsidies were not then in place.

From this list of eligible communes, 24 were excluded due to commitments with the provincial governments. The remaining 58 communes were part of the initial universe that the evaluation team randomized in September 2015 into treatment and control arms using Stata. In June 2016, three communes from the list of 58 were excluded due to objections by local government and a perceived overlap with other development programming. Ultimately, the evaluation included 55 communes.

S3: Theory of Change



S4: Anthropometry measurement guidelines

Anthropometric measurement is comprised of weight and length. Weight was measured using Uniscale (UNICEF recommended scale) in Kilogram with precision to one decimal point. Length was measured using a length board (UNICEF / WFP recommended) in Centimetre with precision to one decimal point. Two data measurements were required, one from the measurement taker and another one from an assistant. The measurement procedure followed FANTA Guidelines:

- Weight measurement:
 - *Preparation:* Ensure enough material is available for measurement (scale, battery, pen, tissue, record form, and age calculation form) with proper function. Ensure that the scale is positioned in a plate and smooth surface. Measurement taker is on the right hand of mother/caregiver while assistant is in front of mother/caregiver. Ensure that children dress light clothes with no cap or shoes. Assistant helps mother/caregiver in carrying the child and asks mother/caregiver to go on to the scale after proper functioning.
 - *During Measurement:*
 - If the child is less than 24 months old, do tared weighing:
 - Request mother/caregiver to remove her shoes and stand on the scale (without the child) with head held high and standing still; read the measurement result loudly; press button to tare scale; hand the child to mother/caregiver after scale functioning; read weight of child out loud so that assistant can record the measurement.
 - If the child is 24 – 28 months old, and will stand still, weight the child alone:
 - Ask the mother to help the child remove shoes and outer clothing; talk with the child about the need to stand still; ask the child to stand in the middle of the scale, feet slightly apart, and to remain still until you say so; read weight of child out loud so that assistant can record the measurement.
 - *Second Measurement:* Request mother/caregiver to step off the scale. Repeat the measurement steps. Record second measurement.

- Length measurement:

Depending on a child's age and ability to stand, measure the child's length or height.

If the child is less than 24 months, measure the child's length lying down:

- *Preparation:* Prepare length board on a plate and smooth surface. Ensure length board stability, take off shoe and cap from child. Check measurement level on the length board, and ensure the record form. The assistant should kneel with both knees behind the base of the board; the measurer should kneel with both knees on the right side of the child to hold the foot piece with their right hand.
- *During Measurement:* With the caregiver's help, lay child on his back on the board. The assistant should cup their hands over the child's ears to place child's

head against base of board so that the child is looking straight up. The measurer should make sure the child is laying flat and in the center of the board; place their left hand over the child's shins or over the knees and press them firmly against the board; with their right hand make sure the feet are perpendicular to the floor and place the sliding foot piece firmly against the child's heels; check head, eye, shoulder, hand, buttock, knees and heel for proper positioning. Make sure body is in proper position and still. Measurement must be read to the nearest of 0.1 cm.

- *Second Measurement:* Remove the sliding foot piece and release left hand from the shins or knees. Make sure body is in proper position and still. Repeat the measurement one more time to ensure accuracy of reading. If the two measurements are different by more than 1.0 cm, then a third measurement is taken.

If the child is 24 – 28 months old, measure standing height unless the child is unable to stand:

- *Preparation:* Place the measuring board at a right angle between a level floor and against a straight, hard vertical surface such as a wall or pillar. Make sure the board is not moving. Ask the caregiver to walk the child to the board and to kneel in front of the child. The assistant should place record form nearby on the ground and kneel with both knees on the right side of the child. The measurer should kneel on their right knee on the child's left side.
 - *During Measurement:* The assistant should place the child's feet flat and together in the center of and against the back and base of the board; place right hand just above the child's ankles on the shins, left hand on the child's knees and push against the board; make sure the child's legs are straight and the heels and calves are against the board. The measurer will ask the child to look straight ahead at the mother who should stand in front of the child; place left hand under the child's chin; make sure the shoulders are level, and the head, shoulder blades and buttocks are against the board. With right hand, lower the headpiece on top of the child's head. Measurement must be read to the nearest of 0.1 cm; assistant should record the measurement.
 - *Second Measurement:* Remove the sliding head piece and release hands from chin, shins or knees. Make sure body is in proper position and still. Repeat the measurement one more time to ensure accuracy of reading. If the two measurements are different by more than 1.0 cm, then a third measurement is taken.
- Following the weighing and length measurements, any child who is classified as severely malnourished is referred to the health clinic.
 - Training: The enumerators were trained on the protocols to follow and how to calibrate equipment. They were also tested on accurate reading and recording of weight and length measurements. We conducted hands-on practice in pairs and then we did a standardization exercise where the entire team was tested on their ability to measure child length accurately and precisely. Measurers had to meet the accuracy and precision threshold to pass and be hired as enumerators for data collection.

- Field supervision: The anthropometry specialist was present in the field during the entire baseline phase, accompanying enumerators to ensure proper technique with the height and weight measurements and recording and conducting random repeat measurements as quality control for 5% of the sample.

S5: Sanitation facility validation

To validate sanitation-related data collection, enumerators performed structured observation and recorded WASH conditions in the household and took photographs of household latrines, which were tagged with the unique household identification number for verification of proper classification. A random sample of 20% of the photos were cross-checked by collaborators from MSI with the recorded survey to ensure proper classification

S6: Sanitation coverage prior to intervention rollout

	Any sanitation	SD	95% CI
Nutrition-only	68%	2%	(65%, 72%)
Sanitation-only	59%	2%	(56%, 63%)
Combined	67%	1%	(64%, 70%)
Control	66%	1%	(63%, 68%)

S7: CONSORT checklist for randomized trials

Section/Topic	Item No	Checklist item	Reported on page No
Title and abstract			
	1a	Identification as a randomised trial in the title	_____
	1b	Structured summary of trial design, methods, results, and conclusions (for specific guidance see CONSORT for abstracts)	_____
Introduction			
Background and objectives	2a	Scientific background and explanation of rationale	_____
	2b	Specific objectives or hypotheses	_____
Methods			
Trial design	3a	Description of trial design (such as parallel, factorial) including allocation ratio	_____
	3b	Important changes to methods after trial commencement (such as eligibility criteria), with reasons	_____
Participants	4a	Eligibility criteria for participants	_____
	4b	Settings and locations where the data were collected	_____
Interventions	5	The interventions for each group with sufficient details to allow replication, including how and when they were actually administered	_____
Outcomes	6a	Completely defined pre-specified primary and secondary outcome measures, including how and when they were assessed	_____
	6b	Any changes to trial outcomes after the trial commenced, with reasons	_____
Sample size	7a	How sample size was determined	_____
	7b	When applicable, explanation of any interim analyses and stopping guidelines	_____
Randomisation:			
Sequence generation	8a	Method used to generate the random allocation sequence	_____
	8b	Type of randomisation; details of any restriction (such as blocking and block size)	_____
Allocation concealment mechanism	9	Mechanism used to implement the random allocation sequence (such as sequentially numbered containers), describing any steps taken to conceal the sequence until interventions were assigned	_____

Implementation	10	Who generated the random allocation sequence, who enrolled participants, and who assigned participants to interventions	<hr/>
Blinding	11a	If done, who was blinded after assignment to interventions (for example, participants, care providers, those assessing outcomes) and how	<hr/>
	11b	If relevant, description of the similarity of interventions	<hr/>
Statistical methods	12a	Statistical methods used to compare groups for primary and secondary outcomes	<hr/>
	12b	Methods for additional analyses, such as subgroup analyses and adjusted analyses	<hr/>
Results			
Participant flow (a diagram is strongly recommended)	13a	For each group, the numbers of participants who were randomly assigned, received intended treatment, and were analysed for the primary outcome	<hr/>
	13b	For each group, losses and exclusions after randomisation, together with reasons	<hr/>
Recruitment	14a	Dates defining the periods of recruitment and follow-up	<hr/>
	14b	Why the trial ended or was stopped	<hr/>
Baseline data	15	A table showing baseline demographic and clinical characteristics for each group	<hr/>
Numbers analysed	16	For each group, number of participants (denominator) included in each analysis and whether the analysis was by original assigned groups	<hr/>
Outcomes and estimation	17a	For each primary and secondary outcome, results for each group, and the estimated effect size and its precision (such as 95% confidence interval)	<hr/>
	17b	For binary outcomes, presentation of both absolute and relative effect sizes is recommended	<hr/>
Ancillary analyses	18	Results of any other analyses performed, including subgroup analyses and adjusted analyses, distinguishing pre-specified from exploratory	<hr/>
Harms	19	All important harms or unintended effects in each group (for specific guidance see CONSORT for harms)	<hr/>
Discussion			
Limitations	20	Trial limitations, addressing sources of potential bias, imprecision, and, if relevant, multiplicity of analyses	<hr/>
Generalisability	21	Generalisability (external validity, applicability) of the trial findings	<hr/>

Interpretation	22	Interpretation consistent with results, balancing benefits and harms, and considering other relevant evidence	_____
Other information			_____
Registration	23	Registration number and name of trial registry	_____
Protocol	24	Where the full trial protocol can be accessed, if available	_____
Funding	25	Sources of funding and other support (such as supply of drugs), role of funders	_____
