***Plasmodium falciparum* hrp2 and hrp3 gene deletion status in Africa and South America by highly sensitive and specific digital PCR**

Short title: ***hrp2/hrp3* deletion typing by digital PCR**

Claudia A. Vera-Arias¹, Aurel Holzschuh^{1,2}, Colins O. Oduma^{3,4}, Kingsley Badu⁵, Mutala Abdul-Hakim⁵, Joshua Yukich⁶, Manuel W. Hetzel^{2,7}, Bakar S. Fakhri^{2,7,8}, Abdullah Ali⁹, Marcelo U. Ferreira¹⁰, Simone Ladeia-Andrade¹¹, Fabián E. Sáenz¹², Yaw Afrane¹³, Endalew Zemene¹⁴, Delenasaw Yewhalaw¹⁴, James W. Kazura¹⁵, Guiyun Yan¹⁶, Cristian Koepfli¹

¹ University of Notre Dame, Notre Dame, IN, United States

² Swiss Tropical and Public Health Institute, Switzerland

³ Kenya Medical Research Institute-Centre for Global Health Research, Kisumu, Kenya

⁴ Department of Biochemistry and Molecular Biology, Egerton University, Nakuru, Kenya

⁵ Kwame Nkrumah University of Science and Technology, Ghana

⁶ Tulane University, USA

⁷ University of Basel, Basel, Switzerland

⁸ Ifakara Health Institute, Dar es Salaam, United Republic of Tanzania

⁹ Zanzibar Malaria Elimination Programme, Zanzibar, United Republic of Tanzania

¹⁰ University of Sao Paulo, Brazil

¹¹ Laboratory of Parasitic Diseases, Fiocruz, Rio de Janeiro, Brazil

¹² Centro de Investigación para la Salud en América Latina, Facultad de Ciencias Exactas y Naturales,

Pontificia Universidad Católica del Ecuador, Quito, Ecuador

¹³ Department of Medical Microbiology, University of Ghana Medical School, Accra, Ghana

25 ¹⁴ Tropical and Infectious Diseases Research Center, Jimma University, Ethiopia

26 ¹⁵ Case Western Reserve University, Cleveland, USA

27 ¹⁶ Program in Public Health, University of California, Irvine, CA 92617, USA

28

29 * Corresponding author: ckoepfli@nd.edu

30

31

32

Abstract

Background: The most commonly used *Plasmodium falciparum* rapid diagnostic tests target the Histidine-Rich Proteins 2 and 3 (HRP2, HRP3). An increasing number of countries report parasites that carry *hrp2* and/or *hrp3* gene deletions, resulting in false negative test results. Molecular surveillance of *hrp2* and *hrp3* deletions is crucial but adequate protocols have been lacking.

Methods and Findings: We have developed novel assays for deletion typing based on droplet digital PCR (ddPCR), targeting *hrp2* exon1, *hrp2* exon 2, and *hrp3*. In the ddPCR assay, *hrp2* or *hrp3* and a control gene were quantified with very high accuracy in a single tube. The theoretical limit of detection of the ddPCR assay was 0.33 parasites/uL, and thus well suited for typing of low-density asymptomatic infections. The deletion was reliably detected in mixed infections with wild-type and *hrp2*-deleted parasites when the proportion of parasites carrying the deletion was >40%. For a side-by-side comparison with the conventional nested PCR (nPCR) assay, 248 samples from asymptomatic individuals from western Kenya were screened in triplicate by ddPCR and nPCR. No deletions were observed by ddPCR, while by nPCR no band for *hrp2* was observed in 8% of samples. The ddPCR assay was applied to screen 830 samples from six countries in Africa and South America. No or very few deletions were observed in Kenya (n=241), Zanzibar/Tanzania (n=91), and Ghana (n=223). In southwestern Ethiopia, 1/47 (2.1%) samples carried *hrp2* deletion, and 35/47 (74.5%) *hrp3* deletions. In Brazil, 87/187 (46.5%) samples carried *hrp2* deletions, and 116/187 (62%) *hrp3* deletions. In Ecuador, no *hrp2* deletions were observed, but 22/41 (53.7%) samples carried *hrp3* deletions.

Conclusions: Compared to nPCR, the ddPCR assay minimizes the risk of false-negative results (i.e. *hrp2* deletion observed when the sample is wild type), increases sensitivity, and greatly reduces the number of reactions that need to be run. Pronounced differences in the prevalence of deletion were observed among sites, with more *hrp3* than *hrp2* deletions.

57

58

Introduction

In 2019, over 200 million cases of malaria and over 400,000 deaths were recorded [1]. *Plasmodium falciparum* remains the primary cause of malaria in humans. Fast and accurate diagnosis and treatment of clinical episodes are key components of malaria control. Diagnosis is commonly performed either by light microscopy, or rapid diagnostic tests (RDTs). RDTs are lateral flow devices that detect parasite proteins in human blood through immunohistochemistry. Light microscopy requires basic lab infrastructure and skilled microscopists. In contrast, RDTs require minimal infrastructure and training, and results are available within approximately 15 minutes at a cost of less than 1 USD per test. RDTs are the only field-deployable diagnostic tool available at peripheral health centers and for community screening to diagnose asymptomatic infections, for example through reactive case detection [2]. In 2016, over 300 million RDTs were used by malaria control programs [3].

The most sensitive RDTs for *P. falciparum* rely on the detection of Histidine Rich Protein 2 (HRP2) [4]. HRP2 is a highly expressed secreted protein, and thus an ideal target for diagnosis. It is also the target for a new generation of ultra-sensitive RDTs with a limit of detection of <100 parasites/ μ L [5, 6]. While alternative RDTs detecting other proteins, e.g. parasite lactase dehydrogenase (pLDH), or aldolase, are available, they are generally less sensitive [7, 8]. HRP2-based RDTs can also detect HRP3, a structurally similar protein sharing multiple epitopes with HRP2.

In 2010, a report revealed that a large proportion of *P. falciparum* parasites in Peru did not carry the *hrp2* gene [9], and thus could not be detected by HRP2-based RDTs. Since then, an increasing number of reports from Latin America [10-14], Africa [15-22], and Asia [23, 24] found varying proportions of parasites with *hrp2*-deletion, reaching up to 80% of clinical cases in certain hospitals in Eritrea [18]. In addition, *hrp3* can

be deleted. The deletion of either *hrp2* or *hrp3*, or both genes, has no known impact on parasite fitness. RDTs can yield a positive result if *hrp2* is deleted but *hrp3* is expressed, but sensitivity of the RDT is lower in this case [25].

Molecular surveillance to assess the frequency of *hrp2* and *hrp3* deletion is crucial to decide whether HRP2-based RDTs can be used [26]. The WHO recommends to use alternative diagnostics if the prevalence of *hrp2* deletion is above 5% [27]. At this level, the number of false-negative tests due to *hrp2* deletion will exceed the number of false-negative tests because of lower sensitivity of the alternative diagnostics. The prevalence of *hrp2* deletion has been found to differ substantially within countries [12, 17], thus the choice of diagnostics might need to be adapted at sub-national level. Where low levels of deletions are present, HRP2-based RDTs remain a highly useful tool for diagnosis.

Deletion screening has been classically done using nested PCR (nPCR) followed by gel electrophoresis [26]. Absence of a band is interpreted as deletion. False-negative results could occur when PCR conditions are suboptimal, or when parasite density is low and amplification is stochastic. To overcome this limitation, 3-fold repetition of the nested *hrp2* PCR and a control PCR (e.g. *msp2*, or *glurp*) is recommended [26]. As a result, for each sample twelve PCRs need to be run. Deletion status might remain unresolved if results differ among replicates. As an additional problem, multiple clone *P. falciparum* infections are common in most endemic settings [28]. In case of a multiple clone infection with a wild-type parasite and a parasite carrying the deletion, the wild type parasite will result in a band on the gel when using the nPCR assay. Multiple clone infections can thus mask the presence of deletions, resulting in an underestimation of the frequency of deletion [29]. More recently, quantitative PCR (qPCR) protocols for *hrp2/hrp3* deletion typing were published [30], greatly enhancing throughput. However, when parasite densities are low,

considerable variation in quantification is observed between replicates [31]. As a result of the technical challenges for accurate typing, maps of *hrp2* deletion frequency remain scattered and incomplete.

We have developed a novel method for typing of *hrp2* and *hrp3* deletions based on droplet digital PCR (ddPCR). ddPCR yields highly accurate quantification of parasites [31]. In a ddPCR experiment, the reaction volume is partitioned into approximately 15,000 microdroplets, which are then subject to end-point PCR. Each droplet functions as an individual PCR reaction, with amplification occurring if the droplet contains template DNA. The number of positive droplets corresponds to the number of template DNA copies. Using two different probes, two targets can be quantified in a single reaction well, e.g. a control gene and a target gene. As each droplet with template can be considered a ‘within-well replicate’, a single well offers the sensitivity and specificity of a large number of replicates by nPCR or qPCR. The risk of false negative results (i.e. no amplification when template is present) is thus minimal compared to nPCR or qPCR. The novel assay greatly reduces the number of reactions to be run, increase sensitivity and accuracy, and can detect the deletion in mixed infections. The assay was extensively validated using culture strains, and field isolates from Kenya, Zanzibar/Tanzania, Ethiopia, Ghana, Brazil, and Ecuador.

Methods

Ethics statement

Informed written consent was obtained from all study participants or their parents or legal guardians prior to sample collection. The study was approved by the University of Notre Dame Institutional Review Board (approvals 18-08-4803, 19-04-5321, 18-12-5029), the Institutional Scientific and Ethical Review boards of the Noguchi Memorial Institute of Medical Research, University of Ghana, the Committee on Human Research, Publication and Ethics, School of Medical Science, Kwame Nkrumah University of Science and Technology, Kumasi (CHRPE/AP/375/20), the Zanzibar Medical Research Ethics Committee (ZAMREC/0001/Feb/17), the Institutional Review Board of Tulane University (17-993573), the Institutional Review Board of the Ifakara Health Institute (003-2017), the Ethics Commission of North-western and Central Switzerland (Req-2017-00162), the Institutional Review Board of Institute of Health, Jimma University, Ethiopia (RPGC/486/06), Maseno University Ethics Review Committee (MUERC protocol number 00456), the Ethics Committee for Research in Human Beings of the Pontificia Universidad Católica del Ecuador (CEISH-571-2018), the Ministry of Public Health of Ecuador (MSP-DIS-2019-004-O), and the institutional review board of Oswaldo Cruz Foundation, Brazil (no. 022/2009).

Digital PCR assays

Four novel ddPCR assays were developed. One assay targets the conserved first 120 bp of *hrp2* exon 2, and thus is located directly adjacent to the histidine-rich repeats. Different breakpoint for the *hrp2* deletion have been described [26]. The novel primers for exon 2 are located in a region that is deleted in all known deletion variants, thus irrespective of the specific breakpoint, *hrp2* deletion will be detected. The second assay targets *hrp2* exon 1. While exon 1 does not contain antigens that are recognized by RDTs, the deletion of this exon 1 would prevent proper expression of the protein. The third assay targets

hrp3. Each assay was multiplexed with an assay targeting *serine-tRNA ligase* (PF3D7_0717700, herein referred to as '*tRNA*'). *tRNA* is a conserved, essential single copy gene, that is frequently used as reference for gene expression assays [32]. In wild type infections not carrying a deletion, the copy numbers of *tRNA* and *hrp2* or *hrp3* are identical.

Novel primers and probes were developed for all assays (Table 1). Assay conditions are given in Supplementary File S1. Across over >3000 genomes available through MalariaGen and PlasmoDB, no SNPs were recorded in primer and probe sequences, thus the assay can be used for the screening of isolates of global origin. Assay conditions were optimized to achieve maximal separation between positive and negative droplets (Figure 1, Supplementary File S2, Figure S1).

Table 1: Primer and probe sequences

Assay	Sequence 5'-3'
<i>hrp2</i> exon 2	<i>hrp2</i> exon 2 forward
	<i>hrp2</i> exon 2 reverse
	<i>hrp2</i> exon 2 probe
<i>hrp2</i> exon 1	<i>hrp2</i> exon 1 forward
	<i>hrp2</i> exon 1 reverse
	<i>hrp2</i> exon 1 probe
<i>hrp3</i>	<i>hrp3</i> forward
	<i>hrp3</i> reverse
	<i>hrp3</i> probe
<i>tRNA</i>	<i>tRNA</i> forward
	<i>tRNA</i> reverse
	<i>tRNA</i> probe

162

163 **3D7 and Dd2 parasite culture strain mixtures**

164 In order to determine the ability to detect mixed clone infections with only one strain carrying the
 165 deletion, experimental mixtures were made from two well-characterized laboratory strains, 3D7 (no
 166 deletions), and Dd2 (carrying the *hrp2* deletion). Each strain was cultured separately, DNA extracted, and
 167 quantified by ddPCR using the *hrp2/tRNA* ligase assay. No *hrp2* was detected in Dd2. Mixtures were
 168 prepared with a concentration (of both strains combined) of 10, 50, 100, and 500 genomes/μL, and with
 169 a 3D7 to Dd2 ratio of 100:0, 80:20, 60:40, 40:60, 20:80, and 0:100. Each mixture was run in triplicate using
 170 the *hrp2* exon2/ *tRNA* ligase assay. The ratio of *hrp2* to *tRNA* copy numbers was compared to expected
 171 values.

172

173 **Nested PCR assay for *hrp2***

174 In order to compare the ddPCR assay directly to the established nPCR assay, 248 samples from
 175 asymptomatic carriers in western Kenya were run by the *hrp2* exon 2 assay, and the *hrp2* nsPCR in
 176 triplicate. Assay conditions for the nPCR followed published protocols [33, 34] and are given in
 177 Supplementary File S1.

178

179 **Field isolates**

180 Field isolates were screened from Kenya, Zanzibar in the United Republic of Tanzania, Ethiopia, Ghana,
 181 Brazil, and Ecuador. Samples from Kenya (n=248) were from Chulaimbo and Homa Bay in western Kenya
 182 close to Lake Victoria. Samples were collected in a cross-sectional survey including individuals of all ages
 183 from January-August 2019. Finger-prick samples were collected in EDTA microtainers and infections
 184 detected by qPCR. Overall *P. falciparum* prevalence across both sites was 16% [35]. No diagnosis by RDT
 185 or microscopy was done.

186

187 Samples from Zanzibar (n=91) had been collected in the frame of a study on reactive case detection (RCD)

188 from May 2017 to October 2018 [2]. Asymptomatic infections identified through RCD were typed. During

189 RCD, infections were diagnosed by HRP2/pLDH-based RDT (SD BIOLINE Malaria Ag Pf HRP2/pLHD), a blood

190 spot was collected on filter paper, and infections diagnosed by qPCR [2]. Prevalence (not including index

191 cases) was 0.8% by RDT and 2.4% by qPCR. All samples with a density (by qPCR) of >100 parasites/ μ L were

192 selected for *hrp2/hrp3* deletion typing, irrespective of RDT result.

193

194 Samples from Ethiopia (n=47) included clinical cases and asymptomatic individuals sampled in June to

195 November 2016 from Jimma Zone, Oromia Region. *P. falciparum* prevalence was 4% by microscopy and

196 8.3% by qPCR [36]. No RDT screening was conducted.

197

198 From Ghana, 2 sets of samples were typed. The first set (n=11) was collected in June-September of 2017

199 from febrile school children aged 5-14 years [37]. The second set (n=212) was collected in Mankranso and

200 Agona Hospitals in the Ashanti region from febrile patients from September to December 2020.

201

202 In Brazil, samples (n=187) were collected in Cruzeiro do Sul, Upper Juruá Valley, northwestern Brazil, in

203 2010-2013. This is the country's main malaria hotspot, which accounted for nearly 15% of all *P. falciparum*

204 infections in Brazil at the time of the study. Samples were collected from clinical patients 4 to 73 years of

205 age (mean, 26.6) enrolled for a drug efficacy trial [38]. Only baseline samples from patients with

206 microscopy- and PCR-confirmed *P. falciparum* infection were included in this study. Even though these

207 samples had been collected nearly a decade ago and might not reflect the current status of *hrp2/hrp3*

208 deletion, they were included in light of the known high levels of *hrp2/hrp3* deletion in Latin America to

confirm the ability of the assays to reliably detect deletions in field isolates [39]. Because of little template volume available, Brazilian samples were not screened for *hrp2* exon 1 deletion.

In Ecuador, samples (n=41) were collected from clinical patients from March 2019 to April 2020. The samples were collected in Esmeraldas and Carchi Provinces in the north-west of the country, where most *P. falciparum* cases of Ecuador are reported. All infections were confirmed by microscopy and collected as blood spots in filter paper. Three samples were collected from travelers coming from the Pacific coast in Colombia but diagnosed in Ecuador.

Data analysis

For the analysis of the ddPCR data, the following criteria were applied: A minimum of 2 droplets positive for *tRNA* were required to include a sample in data analysis. Samples were repeated if a deletion was observed but ≤ 5 droplets were positive for *tRNA*. At a density of >5 droplets positive for *tRNA*, the probability of a false-negative result for *hrp2* or *hrp3* (i.e. no positive droplet in a wild-type infection) is less than 1:500. Data for all samples and assays are provided in Supplementary File S3.

Results

Assay development and validation

New primers and probes were developed for *hrp2* exon 1, *hrp2* exon 2, *hrp3*, and *tRNA* (Table 1). Upon optimization of assay conditions, clear separation between negative and positive droplets was obtained across a wide range of parasite densities (Figure 1, Supplementary Figure S1). No positive droplets for *hrp2* or *hrp3* were observed in case of deletion, while the separation between negative droplets and those positive for *tRNA* remained clear (Figure 1, Supplementary Figure S1).

A: Wild-type, medium density

B: Wild-type, low density

C: *hrp2* deletion

D: Mixed infection

Figure 1: Examples of *hrp2* exon 2 deletion typing by ddPCR. Droplets positive for *hrp2* are shown in blue (top left of each panel). Droplets positive for *tRNA* are shown in green (bottom right). Droplets positive for *hrp2* and *tRNA* are shown in orange (top right). Negative droplets (for both *hrp2* and *tRNA*) are shown in gray (bottom left). A) Wild-type infection of medium density. Approximately 600 droplets are positive each for *hrp2* and *tRNA*, and 36 for both targets. B) Wild-type sample of low-density: 32 and 29 droplets are positive for *hrp2* and *tRNA*, respectively. C) *hrp2* deletion: Droplets are positive for *tRNA*, but no droplets are

positive for *hrp2*. D) Mixed infection with wild-type parasites and parasites carrying *hrp2* deletion. Only 15 droplets are positive for *hrp2*, but 598 droplets are positive for *tRNA*.

To evaluate reproducibility and the limit of reliable detection, 248 isolates from asymptomatic carriers in Kenya were typed for the *hrp2* exon 2 assay in triplicate. Geometric mean density was 95 genomes/ μ L, and 47/248 samples were at densities <10 genomes/ μ L. By ddPCR, 235/248 (94.6%) of samples met inclusion criteria of ≥ 2 droplets positive for *tRNA* in all 3 replicates. Highly similar quantification among replicates was observed (Figure 2A). For each sample, the highest and lowest value of *hrp2* copies/ μ L was recorded. Correlation among technical replicates was very high ($R^2=0.990$). Likewise, for each sample, the highest and lowest value of *tRNA* copies were recorded, and correlation was very high ($R^2=0.991$).

Figure 2: Validation of assay. A) Samples typed in triplicate for the *hrp2* exon 2/*tRNA* assay. Representative examples of different parasite densities are shown. For each sample, the quantification of *hrp2* exon 2, and of *tRNA* is shown. Results from the same run are connected by a dashed line. B) Mixtures of 3D7 (wild type) and Dd2 (*hrp2* deletion). Mixtures were run in triplicate at densities of 10-500 genomes/reaction, and at ratios of 0-100% Dd2. The expected proportion of *hrp2* to *tRNA* copies corresponds to the proportion of 3D7 in the mixture. The observed proportion reflects the expected proportion closely for all mixtures at 500 and 100 genome/reaction.

No deletions were detected in the 248 samples from western Kenya. I.e., no samples with droplets for *tRNA* but no droplets for *hrp2* were observed, even though density was low in many samples. In the ddPCR, approximately two-thirds of the reaction volume was transformed into droplets. Applying the threshold of 2 droplets positive for *tRNA*, three template genomes were required per reaction well to reach that threshold. Up to 9 μ L of template DNA could be added to one reaction when primers and probes are kept at 100 μ M. Thus, the theoretical limit of detection was 0.33 parasites/ μ L. If ≤ 5 droplets are positive for *tRNA* and a deletion is observed, it is recommended to repeat the sample.

Detection of mixed infections with *hrp2*-negative and wild type parasites

To test the ability of the assay to detect deletions when only part of all parasites in an isolate carry the deletion, experimental mixtures were prepared with DNA from parasite culture of 3D7 (wild-type), and Dd2 (*hrp2* deletion). Mixtures were prepared at ratios from 0% to 100% Dd2, and at densities of 10-500 genomes/reaction. At densities of 100 to 500 genomes/ μ L, the quantification by ddPCR represented the mixture ratio with high accuracy (Figure 2B). Whenever 40% or more of parasites carried the deletion, the observed ratio was clearly below 100%. In cases where only a small proportion of all parasites carried the deletion (20% Dd2 vs. 80% 3D7), the difference in quantification of *hrp2* and *tRNA* was too small to observe the deletion. Likewise, at densities < 50 genomes/reaction the ratio did not accurately reflect the experimental mixture.

The results were corroborated by screening of 739 field samples from five countries. In wild type isolates, a very similar quantification of *hrp2* and *tRNA* was expected. Unless samples carried a clear deletion (i.e.

no or very little *hrp2* signal), in all samples with densities of >50 copies/ μ L, the ratio of *hrp2* to *tRNA* copies was above 0.6 (Figure 3). With increasing parasite density, the ratio got closer to 1.

Figure 3: Ratios of *hrp2* to *tRNA* copies in 684 field isolates. With increasing parasite density (X-axis), the ratio becomes close to 1. Deletions (with no wild type parasites present) have a ratio of 0. Dashed lines show a ratio of *hrp2* to *tRNA* copies of 0.6, and 50 genomes/ μ L. Mixed infection can be reliably detected at densities >50 genomes/ μ L, and if >40% of parasites carry the deletion.

A lower ratio of *hrp2* to *tRNA* copies was observed in samples from Zanzibar, where the mean ratio of *hrp2*/*tRNA* copies was 0.82 in absence of any samples that carried *hrp2* deletion (Supplementary Figure S2). The reason is not known, but might be caused by sampling and storage procedures. Blood samples from Zanzibar were collected on filter paper, and stored at ambient temperature for over three years

prior to extraction. Possibly, this could have resulted in DNA degradation, that affected *hrp2* more than *tRNA*.

Comparison of ddPCR to gel-based nested PCR

The *hrp2* exon 2 ddPCR assay was compared to the classical nPCR assay with visualization of products on agarose gel in 248 asymptomatic infections from western Kenya. The density of asymptomatic infections is often low and thus amplification by PCR can be stochastic. All assays were run in triplicate, i.e. *hrp2*-exon2/*tRNA* by ddPCR, *hrp2* nested PCR, and *msh2* nested PCR.

Samples were included in analysis if the PCR for the control gene was positive in all three replicates, i.e. if >2 droplets were positive for *tRNA* in the ddPCR assay, or a band was detected for *msh2* in all three replicates. For the gel-based assay, 85.5% (212/248) samples met inclusion criteria (Table 2). Among those positive, for 144/212 samples a band for *hrp2* was observed in all three replicates. A band in two replicates was observed for 34 samples, and a band in a single replicate for 17 samples. For 17 samples, despite obtaining a band for *msh2* in all three replicates, no band for *hrp2* was observed. These 17 samples would thus be classified as *hrp2* deletion, resulting in a prevalence of deletion of 8.0% (17/212). Figure 3 shows representative examples of results of the nPCR and ddPCR assays.

Table 2: Comparison between nested PCR and ddPCR-based *hrp2* deletion typing

	nested PCR	ddPCR
Total samples	248	248

Met inclusion criteria	212 (85.5%)	235 (94.8%)
Deletion in 3/3 replicates	17	0
Deletion in 2/3 replicates	17	0
Deletion in 1/3 replicates	34	2
No deletion	144	233
Prevalence of deletion	8.0% (17/212)	0% (0/235)

319

320

321 **Figure 3:** Comparison of nPCR and ddPCR for *hrp2* deletion typing. Representative examples of results obtained by *hrp2* and *msp2*
322 nPCR, and by ddPCR. The expected size of the *hrp2* band is 228 bp. A band is visible in samples 3, 4, and 5, but no band is visible
323 in samples 1, 2, 6, and 7. *msp2* was run as control for the nPCR assay. *msp2* is a size polymorphic gene with amplicons ranging
324 from approximately 200 to 500 bp. Bands are observed for all samples, and multiple bands are observed in case of polyclonal
325 infections. L = 100 bp DNA ladder (New England BioLabs). Samples were run in triplicate, and the same results was obtained all
326 three times. By ddPCR, no deletions were observed in any samples. For samples 1 and 7, the *hrp2* exon 2/*tRNA* ddPCR plot is
327 shown. Droplets are visible for both targets, thus no deletion is observed.

328

329

By ddPCR, the criteria for inclusion (≥ 2 droplets for *tRNA*) was met by 94.8% (235/248) of samples. Among them, ≥ 2 droplets for *hrp2* were detected in all three replicates in 233/235 samples. In 2 samples in only two of the three replicates ≥ 2 droplets were positive for *hrp2*. Both of these samples had one replicate with 1 positive droplet for *hrp2*, and 5 positive droplets for *tRNA*. In none of the samples all three or two out of three replicates were negative for *hrp2*. Thus, the prevalence of deletion by ddPCR was 0%.

***hrp2* and *hrp3* deletions in Africa and South America**

The new ddPCR assay was applied to screen for deletions in 830 samples from Kenya, Zanzibar, Ethiopia, Ghana, Brazil, and Ecuador. The frequency of deletions for all loci and sites is given in Table 3.

Table 3: *hrp2* and *hrp3* deletions in Africa and South America

Site	N	Deletions				
		<i>hrp2</i> exon 2	<i>hrp2</i> exon 1	<i>hrp3</i>	<i>hrp2</i> + <i>hrp3</i> ¹	Mixed ²
Kenya	241	0% (0/241)	0% (0/241)	0% (0/241)	0% (0/241)	
Zanzibar	91	0% (0/91)	0% (0/91)	0 % (0/91)	0% (0/91)	
Ethiopia	47	2.1% (1/47)	2.1% (1/47)	74.5% (35/47)	2.1% (1/47)	1 x <i>hrp3</i>
Ghana	223	0% (0/226)	0% (0/223)	0.4% (1/223)	0% (0/170)	1 x <i>hrp2</i> , 3 x <i>hrp3</i>
Brazil	187	46.5% (87/187)	NA	62.0% (116/187)	46.0% (86/187)	2 x <i>hrp2</i> , 2 x <i>hrp3</i>
Ecuador	41	0% (0/41)	0% (0/41)	53.7% (22/41)	0% (0/39)	

¹ Isolates with deletions of *hrp2* and *hrp3*. Note that these isolates are also counted as deletions in the columns for *hrp2*, and for *hrp3* (e.g. in Brazil 87 isolates carried *hrp2* deletion, of which 86 also carried *hrp3* deletion).

² Isolates with only a proportion of all parasites carrying the deletion.

From Kenya, 241 samples were typed and no deletions of *hrp2* or *hrp3* were observed. In Zanzibar, no deletions were observed among 91 samples. Among 223 samples from Ghana, 1 *hrp2* deletion/wild type mixed sample was detected (ratio of *hrp2* to *tRNA* copies was <0.6, and parasite density >50 genomes/ μ L), 1 *hrp3* deletion, and 3 samples with *hrp3* deletion/wild type mixes.

In Ethiopia, 47 samples met inclusion criteria. One sample carried a deletion of *hrp2* exons 1 and 2, resulting in a frequency of deletion of 2.1%. *hrp3* deletion was observed in 35/47 (74.5%) samples, and one sample carried a mixed infection with wild type/*hrp3* deletion. The sample with *hrp2* deletion was among the samples with *hrp3* deletion, i.e. both genes were deleted.

From Brazil, 187 samples were screened. 87 samples carried a deletion of *hrp2*. Two additional samples carried mixed infections with wild type parasites and *hrp2* deletion. Deletion of *hrp3* was observed in 116/187 (62.0%) samples, and 86/187 (46.0%) samples carried deletions of *hrp2* and *hrp3*. No *hrp2* deletions were observed in Ecuador, but 22/41 (53.7%) samples carried *hrp3* deletions.

Discussion

Molecular surveillance of the extent of *hrp2* and *hrp3* deletions is a high priority task to select the optimal tools for *P. falciparum* diagnosis. The novel assay based on ddPCR yielded highly accurate results, and was able to detect mixed infections with wild-type parasites and parasites carrying *hrp2* deletion. The good performance of the assay was shown by typing of samples from six countries. The samples reflected a range of parasite densities, from low-density asymptomatic infections to high-density clinical infections, and sites represented a range in the frequency of *hrp2* and/or *hrp3* deletions. Across over 800 field samples screened, <10 samples needed to be repeated because of poor separation between negative and positive droplets, or failure of droplet generation.

High sensitivity is required for accurate typing of low-density infections. We validated a threshold of two droplets positive for *tRNA* as limit of detection. When 9 µL of DNA are used as template, this results in a theoretical limit of detection of 0.33 parasites/µL. Sensitivity can be further increased by concentrating the DNA prior to typing.

The ddPCR assay showed increased sensitivity and specificity to type low-density infections compared to nPCR. Out of 248 asymptomatic samples from western Kenya, 95% could be analyzed by ddPCR, but only 85% by nPCR. More importantly, results on deletion status differed substantially. By ddPCR, no deletions were observed. By nPCR, no band was observed for *hrp2* in 8% of samples that had a positive band for the control gene (*msh2*) in all three replicates. Thus, the frequency of deletion was above the 5% threshold, and it would be erroneously recommended to discontinue the use of HRP2-based RDTs. The frequency of deletion by nPCR was similar to a previous study conducted in a nearby site in western Kenya that found *hrp2* deletion in 8/89 samples using nPCR [40]. The direct comparison of ddPCR and nPCR in a large

number of samples points to a possible overestimation of *hrp2* deletion frequency by studies relying on nPCR.

In almost all transmission settings, polyclonal *P. falciparum* infections are frequent [28]. Using a gel-based assay for deletion typing, in a mixed infection with a wild type parasite and a parasite carrying a deletion, the wild type parasite will produce a band and mask the deletion. These infections can be detected by RDT, but depending on the proportion of polyclonal infections, they can result in pronounced underestimation of the true frequency of deletion [29]. The highly accurate quantification by ddPCR allows detection of mixed infections. Based on experimental mixtures of 3D7 and Dd2, and field samples, mixed infections were reliably detected when at least 40% of parasites carried the deletion, and at densities above 100 genomes per reaction. Using a well-working qPCR assay with an amplification efficacy of 100%, a similar difference between *hrp2* and *tRNA* copy numbers would result in less than half a cycle difference. This is within the normal variation of technical replicates [31, 41], and thus such mixed infections could not be detected by qPCR.

Twenty-nine isolates from Zanzibar and 16 from Ghana had tested negative by HRP2-based RDT despite high density of 100 to >10,000 parasites/ μ L [2], yet no *hrp2* deletions were observed in these populations. False-negative RDT results might be caused due to incorrect handling, prozone effect [42], or sequence variation without deletion of the *hrp2* gene [43]. The finding corroborates the importance of molecular typing. Studies comparing microscopy and RDT results can give important clues for the presence of deletions [44], but molecular typing is required for confirmation [18].

No *hrp2* deletions were found in Zanzibar, and Kenya, and one mixed infection in Ghana. The data from Ghana contrasts an earlier study that reported a frequency of deletion of >30% [15]. A single *hrp2* deletion

was found in southwestern Ethiopia. This is in stark contrast to very high levels of deletion in western Ethiopia [45], Eritrea [18], and Sudan [46]. The results corroborate the need for studies assessing *hrp2* deletion and selection of diagnostic tools at sub-national level.

Contrasting findings were obtained from the sample sets from South America, with very high levels of deletion in Brazil, and no deletions in Ecuador. Brazil and Ecuador share no borders, and the amount of human migration is limited. The results add to the heterogenous pattern of *hrp2/hrp3* deletion in South America, with high frequency of *hrp2* deletion in the Amazon [9, 12, 39], but low levels among the Pacific coast [11, 12]. An outbreak of parasites with *hrp2* deletion at the Peruvian Pacific coast was caused by infections imported from the Amazon [47]. Brazil is committed to *P. falciparum* elimination, and the samples typed originated from the main hotspot of transmission [48]. Rapid diagnostic tests (RDTs) remain relatively little used in Brazil. Microscopy remains the diagnostic method of choice, and RDTs are mostly used in remote areas, e.g., populations from Amerindian Reserves and some traditional riverine communities with no access to conventional microscopy. The frequency of *hrp2* deletion in Brazil clearly exceeds the 5% threshold, thus it is recommended that no HRP2-based RDTs are used.

To determine whether the frequency of the deletion exceeds the threshold of 5%, the WHO recommends that for each site a minimum of 370 is typed [49]. Limitations of the molecular assays available for typing have been a major hindrance to type that number of samples in many sites where *hrp2* or *hrp3* deletion is suspected. Even a low proportion of false-negative results could impact the decision to discontinue HRP2-based RDTs. As a result of the scarcity of field data, the spatiotemporal dynamics of *hrp2* deletion in parasite populations, drivers of the deletion, potential fitness costs, and clinical consequences remain poorly understood. Results from simulation studies suggest that the use of *hrp2*-based RDTs selects for

hrp2-negative parasites, in particular if transmission levels are low and a large proportion of all infections turn clinical and result in treatment seeking [50, 51].

In conclusion, the novel, high-throughput, highly sensitive and specific ddPCR assay will facilitate molecular surveillance for *hrp2* and *hrp3* deletion, and thus aid selection of diagnostic tests to accelerate malaria control and elimination. Data obtained using this assay will help to understand the evolutionary processes underlying the de-novo emergence and spread of the deletion.

Acknowledgements

We thank all study participants providing blood samples and the study teams and health center personnel who supported sample collection. We thank Michael T. Ferdig and Katelyn M. Vendrely for providing culture strain DNA for the 3D7/Dd2 experimental mixtures.

Supplementary Files

Supplementary File S1: Assay conditions
Supplementary File S2: Supplementary Figures S1, S2
Supplementary File S3: Database

Financial Disclosure Statement

This work was supported by NIH grants R21AI137891 awarded to CK, and U19 AI129326, D43 TW001505 awarded to GY (<https://www.nih.gov>). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests

The authors state that no competing interests exist.

References

1. World Health Organisation. World Malaria Report 2020. 2020.
2. Stuck L, Fakih BS, Al-Mafazy AH, Hofmann NE, Holzschuh A, Grossenbacher B, et al. Malaria infection prevalence and sensitivity of reactive case detection in Zanzibar. *Int J Infect Dis.* 2020;97:337-46. Epub 2020/06/14. doi: 10.1016/j.ijid.2020.06.017. PubMed PMID: 32534138.
3. WHO. World Malaria Report 2017. 2017.
4. Li B, Sun Z, Li X, Li X, Wang H, Chen W, et al. Performance of pfHRP2 versus pLDH antigen rapid diagnostic tests for the detection of *Plasmodium falciparum*: a systematic review and meta-analysis. *Arch Med Sci.* 2017;13(3):541-9. Epub 2017/05/17. doi: 10.5114/aoms.2017.67279. PubMed PMID: 28507567; PubMed Central PMCID: PMC5420633.
5. Acquah FK, Donu D, Obboh EK, Bredu D, Mawuli B, Amponsah JA, et al. Diagnostic performance of an ultrasensitive HRP2-based malaria rapid diagnostic test kit used in surveys of afebrile people living in Southern Ghana. *Malaria J.* 2021;20(1). doi: ARTN 125 10.1186/s12936-021-03665-7. PubMed PMID: WOS:000625426400002.
6. Hofmann NE, Antunes Moniz C, Holzschuh A, Keitel K, Boillat-Blanco N, Kagoro F, et al. Diagnostic Performance of Conventional and Ultrasensitive Rapid Diagnostic Tests for Malaria in Febrile Outpatients in Tanzania. *J Infect Dis.* 2019;219(9):1490-8. Epub 2018/11/27. doi: 10.1093/infdis/jiy676. PubMed PMID: 30476111; PubMed Central PMCID: PMC6467194.

- 485 7. World Health Organisation. False-negative RDT results and implications of new reports
486 of *P. falciparum* histidine-rich protein 2/3 gene deletions. 2016.
- 487 8. World Health Organisation. Malaria Rapid Diagnostic Test Performance. Results of WHO
488 product testing of malaria RDTs: round 8 (2016–2018)2018.
- 489 9. Gamboa D, Ho MF, Bendezu J, Torres K, Chiodini PL, Barnwell JW, et al. A large
490 proportion of *P. falciparum* isolates in the Amazon region of Peru lack pfhrp2 and pfhrp3:
491 implications for malaria rapid diagnostic tests. PLoS One. 2010;5(1):e8091. Epub 2010/01/30.
492 doi: 10.1371/journal.pone.0008091. PubMed PMID: 20111602; PubMed Central PMCID:
493 PMCPMC2810332.
- 494 10. Maltha J, Gamboa D, Bendezu J, Sanchez L, Cnops L, Gillet P, et al. Rapid diagnostic tests
495 for malaria diagnosis in the Peruvian Amazon: impact of pfhrp2 gene deletions and cross-
496 reactions. PLoS One. 2012;7(8):e43094. doi: 10.1371/journal.pone.0043094. PubMed PMID:
497 22952633; PubMed Central PMCID: PMCPMC3429466.
- 498 11. Saenz FE, Morton LC, Okoth SA, Valenzuela G, Vera-Arias CA, Velez-Alvarez E, et al.
499 Clonal population expansion in an outbreak of *Plasmodium falciparum* on the northwest coast
500 of Ecuador. Malar J. 2015;13 Suppl 1:497. Epub 2015/12/15. doi: 10.1186/s12936-015-1019-2.
501 PubMed PMID: 26651993; PubMed Central PMCID: PMCPMC4676133.
- 502 12. Murillo Solano C, Akinyi Okoth S, Abdallah JF, Pava Z, Dorado E, Incardona S, et al.
503 Deletion of *Plasmodium falciparum* Histidine-Rich Protein 2 (pfhrp2) and Histidine-Rich Protein
504 3 (pfhrp3) Genes in Colombian Parasites. PLoS One. 2015;10(7):e0131576. Epub 2015/07/08.
505 doi: 10.1371/journal.pone.0131576. PubMed PMID: 26151448; PubMed Central PMCID:
506 PMCPMC4494814.

- 507 13. Dorado EJ, Okoth SA, Montenegro LM, Diaz G, Barnwell JW, Udhayakumar V, et al.
508 Genetic Characterisation of Plasmodium falciparum Isolates with Deletion of the pfhrp2 and/or
509 pfhrp3 Genes in Colombia: The Amazon Region, a Challenge for Malaria Diagnosis and Control.
510 PLoS One. 2016;11(9):e0163137. doi: 10.1371/journal.pone.0163137. PubMed PMID:
511 27636709; PubMed Central PMCID: PMC5026348.
- 512 14. Akinyi Okoth S, Abdallah JF, Ceron N, Adhin MR, Chandrabose J, Krishnalall K, et al.
513 Variation in Plasmodium falciparum Histidine-Rich Protein 2 (Pfhrp2) and Plasmodium
514 falciparum Histidine-Rich Protein 3 (Pfhrp3) Gene Deletions in Guyana and Suriname. PLoS One.
515 2015;10(5):e0126805. doi: 10.1371/journal.pone.0126805. PubMed PMID: 25978499; PubMed
516 Central PMCID: PMC4433255.
- 517 15. Amoah LE, Abankwa J, Oppong A. Plasmodium falciparum histidine rich protein-2
518 diversity and the implications for PfHRP 2: based malaria rapid diagnostic tests in Ghana. Malar
519 J. 2016;15:101. Epub 2016/02/20. doi: 10.1186/s12936-016-1159-z. PubMed PMID: 26891848;
520 PubMed Central PMCID: PMC4759916.
- 521 16. Koita OA, Doumbo OK, Ouattara A, Tall LK, Konare A, Diakite M, et al. False-negative
522 rapid diagnostic tests for malaria and deletion of the histidine-rich repeat region of the hrp2
523 gene. Am J Trop Med Hyg. 2012;86(2):194-8. Epub 2012/02/04. doi: 10.4269/ajtmh.2012.10-
524 0665. PubMed PMID: 22302847; PubMed Central PMCID: PMC3269266.
- 525 17. Parr JB, Verity R, Doctor SM, Janko M, Carey-Ewend K, Turman BJ, et al. Pfhrp2-Deleted
526 Plasmodium falciparum Parasites in the Democratic Republic of the Congo: A National Cross-
527 sectional Survey. J Infect Dis. 2016. doi: 10.1093/infdis/jiw538. PubMed PMID: 28177502.

- 528 18. Berhane A, Anderson K, Mihreteab S, Gresty K, Rogier E, Mohamed S, et al. Major Threat
529 to Malaria Control Programs by Plasmodium falciparum Lacking Histidine-Rich Protein 2,
530 Eritrea. Emerg Infect Dis. 2018;24(3):462-70. Epub 2018/02/21. doi: 10.3201/eid2403.171723.
531 PubMed PMID: 29460730; PubMed Central PMCID: PMC5823352.
- 532 19. Menegon M, L'Episcopia M, Nurahmed AM, Talha AA, Nour BYM, Severini C.
533 Identification of Plasmodium falciparum isolates lacking histidine-rich protein 2 and 3 in Eritrea.
534 Infect Genet Evol. 2017;55:131-4. Epub 2017/09/12. doi: 10.1016/j.meegid.2017.09.004.
535 PubMed PMID: 28889944.
- 536 20. Bharti PK, Chandel HS, Ahmad A, Krishna S, Udhayakumar V, Singh N. Prevalence of
537 pfhrp2 and/or pfhrp3 Gene Deletion in Plasmodium falciparum Population in Eight Highly
538 Endemic States in India. PLoS One. 2016;11(8):e0157949. doi: 10.1371/journal.pone.0157949.
539 PubMed PMID: 27518538; PubMed Central PMCID: PMC4982695.
- 540 21. Li P, Xing H, Zhao Z, Yang Z, Cao Y, Li W, et al. Genetic diversity of Plasmodium
541 falciparum histidine-rich protein 2 in the China-Myanmar border area. Acta Trop. 2015;152:26-
542 31. doi: 10.1016/j.actatropica.2015.08.003. PubMed PMID: 26297799; PubMed Central PMCID:
543 PMC4918506.
- 544 22. Kumar N, Pande V, Bhatt RM, Shah NK, Mishra N, Srivastava B, et al. Genetic deletion of
545 HRP2 and HRP3 in Indian Plasmodium falciparum population and false negative malaria rapid
546 diagnostic test. Acta Trop. 2013;125(1):119-21. doi: 10.1016/j.actatropica.2012.09.015.
547 PubMed PMID: 23041541.
- 548 23. Nima MK, Hougard T, Hossain ME, Kibria MG, Mohon AN, Johora FT, et al. Case Report:
549 A Case of Plasmodium falciparum hrp2 and hrp3 Gene Mutation in Bangladesh. American

550 Journal of Tropical Medicine and Hygiene. 2017;97(4):1155-8. doi: 10.4269/ajtmh.16-0884.

551 PubMed PMID: WOS:000423208300030.

552 24. Pati P, Dhangadamajhi G, Bal M, Ranjit M. High proportions of pfhrp2 gene deletion and

553 performance of HRP2-based rapid diagnostic test in Plasmodium falciparum field isolates of

554 Odisha. Malar J. 2018;17(1):394. Epub 2018/10/31. doi: 10.1186/s12936-018-2502-3. PubMed

555 PMID: 30373573; PubMed Central PMCID: PMC6206925.

556 25. Kong A, Wilson SA, Ah Y, Nace D, Rogier E, Aidoo M. HRP2 and HRP3 cross-reactivity and

557 implications for HRP2-based RDT use in regions with Plasmodium falciparum hrp2 gene

558 deletions. Malaria J. 2021;20(1). doi: ARTN 207

559 10.1186/s12936-021-03739-6. PubMed PMID: WOS:000651405700003.

560 26. Cheng Q, Gatton ML, Barnwell J, Chiodini P, McCarthy J, Bell D, et al. Plasmodium

561 falciparum parasites lacking histidine-rich protein 2 and 3: a review and recommendations for

562 accurate reporting. Malar J. 2014;13:283. Epub 2014/07/24. doi: 10.1186/1475-2875-13-283.

563 PubMed PMID: 25052298; PubMed Central PMCID: PMC4115471.

564 27. World Health Organisation. P. falciparum hrp2/3 gene deletions. Conclusions and

565 recommendations of a Technical Consultation. 2016.

566 28. Lopez L, Koepfli C. Systematic review of Plasmodium falciparum and Plasmodium vivax

567 polyclonal infections: Impact of prevalence, study population characteristics, and laboratory

568 procedures. 2021. doi: 10.1101/2021.03.18.436023.

569 29. Watson OJ, Verity R, Ghani AC, Garske T, Cunningham J, Tshefu A, et al. Impact of

570 seasonal variations in Plasmodium falciparum malaria transmission on the surveillance of

571 pfhrp2 gene deletions. Elife. 2019;8. Epub 2019/05/03. doi: 10.7554/eLife.40339. PubMed
572 PMID: 31045490; PubMed Central PMCID: PMCPMC6533063.

573 30. Schindler T, Deal AC, Fink M, Guirou E, Moser KA, Mwakasungula SM, et al. A multiplex
574 qPCR approach for detection of pfhrp2 and pfhrp3 gene deletions in multiple strain infections
575 of Plasmodium falciparum. Sci Rep-Uk. 2019;9. doi: ARTN 13107
576 10.1038/s41598-019-49389-2. PubMed PMID: WOS:000485215700016.

577 31. Koepfli C, Nguitragool W, Hofmann NE, Robinson LJ, Ome-Kaius M, Sattabongkot J, et al.
578 Sensitive and accurate quantification of human malaria parasites using droplet digital PCR
579 (ddPCR). Sci Rep. 2016;6:39183. Epub 2016/12/17. doi: 10.1038/srep39183. PubMed PMID:
580 27982132; PubMed Central PMCID: PMCPMC5159915.

581 32. Friedrich O, Reiling SJ, Wunderlich J, Rohrbach P. Assessment of Plasmodium falciparum
582 PfMDR1 transport rates using Fluo-4. J Cell Mol Med. 2014;18(9):1851-62. Epub 2014/06/04.
583 doi: 10.1111/jcmm.12313. PubMed PMID: 24889967; PubMed Central PMCID:
584 PMCPMC4196660.

585 33. Akinyi S, Hayden T, Gamboa D, Torres K, Bendezu J, Abdallah JF, et al. Multiple genetic
586 origins of histidine-rich protein 2 gene deletion in Plasmodium falciparum parasites from Peru.
587 Sci Rep. 2013;3:2797. Epub 2013/10/01. doi: 10.1038/srep02797. PubMed PMID: 24077522;
588 PubMed Central PMCID: PMCPMC3786299.

589 34. Abdallah JF, Okoth SA, Fontecha GA, Torres RE, Banegas EI, Matute ML, et al. Prevalence
590 of pfhrp2 and pfhrp3 gene deletions in Puerto Lempira, Honduras. Malar J. 2015;14:19. Epub
591 2015/01/22. doi: 10.1186/s12936-014-0537-7. PubMed PMID: 25604310; PubMed Central
592 PMCID: PMCPMC4308922.

35. Oduma CO, Ogolla S, Atieli H, Ondigo BN, Lee MC, Githeko AK, et al. Increased investment in gametocytes in asymptomatic Plasmodium falciparum infections in the wet season. BMC Infect Dis. 2021;21(1):44. Epub 2021/01/11. doi: 10.1186/s12879-020-05761-6. PubMed PMID: 33422001; PubMed Central PMCID: PMC7797145.
36. Zemene E, Koepfli C, Tiruneh A, Yeshiwondim AK, Seyoum D, Lee MC, et al. Detection of foci of residual malaria transmission through reactive case detection in Ethiopia. Malar J. 2018;17(1):390. Epub 2018/10/28. doi: 10.1186/s12936-018-2537-5. PubMed PMID: 30367636; PubMed Central PMCID: PMC6203988.
37. Dieng CC, Gonzalez L, Pestana K, Dhikrullahi SB, Amoah LE, Afrane YA, et al. Contrasting Asymptomatic and Drug Resistance Gene Prevalence of Plasmodium falciparum in Ghana: Implications on Seasonal Malaria Chemoprevention. Genes (Basel). 2019;10(7). Epub 2019/07/19. doi: 10.3390/genes10070538. PubMed PMID: 31315304; PubMed Central PMCID: PMC6678124.
38. Ladeia-Andrade S, de Melo GNP, de Souza-Lima RD, Salla LC, Bastos MS, Rodrigues PT, et al. No Clinical or Molecular Evidence of Plasmodium falciparum Resistance to Artesunate Mefloquine in Northwestern Brazil. American Journal of Tropical Medicine and Hygiene. 2016;95(1):148-54. doi: 10.4269/ajtmh.16-0017. PubMed PMID: WOS:000400206100029.
39. Goes L, Chamma-Siqueira N, Peres JM, Nascimento JM, Valle S, Arcanjo AR, et al. Evaluation of Histidine-Rich Proteins 2 and 3 Gene Deletions in Plasmodium falciparum in Endemic Areas of the Brazilian Amazon. Int J Environ Res Public Health. 2020;18(1). Epub 2020/12/31. doi: 10.3390/ijerph18010123. PubMed PMID: 33375379; PubMed Central PMCID: PMC7795390.

- 615 40. Beshir KB, Sepulveda N, Bharmal J, Robinson A, Mwanguzi J, Busula AO, et al.
616 Plasmodium falciparum parasites with histidine-rich protein 2 (pfhrp2) and pfhrp3 gene
617 deletions in two endemic regions of Kenya. Sci Rep. 2017;7(1):14718. Epub 2017/11/09. doi:
618 10.1038/s41598-017-15031-2. PubMed PMID: 29116127; PubMed Central PMCID:
619 PMCPMC5677122.
- 620 41. Hindson CM, Chevillet JR, Briggs HA, Gallichotte EN, Ruf IK, Hindson BJ, et al. Absolute
621 quantification by droplet digital PCR versus analog real-time PCR. Nat Methods.
622 2013;10(10):1003-5. Epub 2013/09/03. doi: 10.1038/nmeth.2633. PubMed PMID: 23995387;
623 PubMed Central PMCID: PMCPMC4118677.
- 624 42. Gillet P, Mori M, Van Esbroeck M, Van den Ende J, Jacobs J. Assessment of the prozone
625 effect in malaria rapid diagnostic tests. Malar J. 2009;8:271. Epub 2009/12/02. doi:
626 10.1186/1475-2875-8-271. PubMed PMID: 19948018; PubMed Central PMCID:
627 PMCPMC2789093.
- 628 43. Nderu D, Kimani F, Thiong'o K, Karanja E, Akinyi M, Too E, et al. Plasmodium falciparum
629 histidine-rich protein (PfHRP2 and 3) diversity in Western and Coastal Kenya. Sci Rep.
630 2019;9(1):1709. Epub 2019/02/10. doi: 10.1038/s41598-018-38175-1. PubMed PMID:
631 30737461; PubMed Central PMCID: PMCPMC6368535.
- 632 44. Berhane A, Russom M, Bahta I, Hagos F, Ghirmai M, Uqubay S. Rapid diagnostic tests
633 failing to detect Plasmodium falciparum infections in Eritrea: an investigation of reported false
634 negative RDT results. Malar J. 2017;16(1):105. Epub 2017/03/08. doi: 10.1186/s12936-017-
635 1752-9. PubMed PMID: 28264689; PubMed Central PMCID: PMCPMC5339986.

45. Alemayehu GS, Blackburn K, Lopez K, Cambel Dieng C, Lo E, Janies D, et al. Detection of high prevalence of Plasmodium falciparum histidine-rich protein 2/3 gene deletions in Assosa zone, Ethiopia: implication for malaria diagnosis. Malar J. 2021;20(1):109. Epub 2021/02/25. doi: 10.1186/s12936-021-03629-x. PubMed PMID: 33622309.
46. Prosser C, Gresty K, Ellis J, Meyer W, Anderson K, Lee R, et al. Plasmodium falciparum Histidine-Rich Protein 2 and 3 Gene Deletions in Strains from Nigeria, Sudan, and South Sudan. Emerg Infect Dis. 2021;27(2):471-9. Epub 2021/01/27. doi: 10.3201/eid2702.191410. PubMed PMID: 33496220; PubMed Central PMCID: PMC7853540.
47. Baldeviano GC, Okoth SA, Arrospide N, Gonzalez RV, Sanchez JF, Macedo S, et al. Molecular Epidemiology of Plasmodium falciparum Malaria Outbreak, Tumbes, Peru, 2010-2012. Emerg Infect Dis. 2015;21(5):797-803. Epub 2015/04/22. doi: 10.3201/eid2105.141427. PubMed PMID: 25897626; PubMed Central PMCID: PMC4412223.
48. Ferreira MU, Castro MC. Challenges for malaria elimination in Brazil. Malar J. 2016;15(1):284. Epub 2016/05/22. doi: 10.1186/s12936-016-1335-1. PubMed PMID: 27206924; PubMed Central PMCID: PMC4875681.
49. World Health Organization. Surveillance template protocol for pfhrp2/pfhrp3 gene deletions. 2020.
50. Gatton ML, Dunn J, Chaudhry A, Ciketic S, Cunningham J, Cheng Q. Implications of Parasites Lacking Plasmodium falciparum Histidine-Rich Protein 2 on Malaria Morbidity and Control When Rapid Diagnostic Tests Are Used for Diagnosis. J Infect Dis. 2017;215(7):1156-66. Epub 2017/03/23. doi: 10.1093/infdis/jix094. PubMed PMID: 28329034.

657 51. Watson OJ, Slater HC, Verity R, Parr JB, Mwandagalirwa MK, Tshefu A, et al. Modelling
658 the drivers of the spread of Plasmodium falciparum hrp2 gene deletions in sub-Saharan Africa.
659 Elife. 2017;6. Epub 2017/08/25. doi: 10.7554/eLife.25008. PubMed PMID: 28837020; PubMed
660 Central PMCID: PMCPMC5602420.
661