Prevalence of laboratory-related musculoskeletal disorders among biomedical scientists

Nasar Alwahaibi¹, Mallak Al Sadairi ¹, Ibrahim Al Abri ¹, Samira Al Rawahi ².

¹Department of Allied Health Sciences, College of Medicine and Health Sciences, Sultan Qaboos University.
Oman.

²Department of Pathology, Sultan Qaboos University Hospital, Sultan Qaboos University. Oman.

For correspondence: nasar@squ.edu.om

Abstract

Background: Laboratory-related musculoskeletal disorders (LMSDs) are injuries resulted from working in the laboratory. Biomedical scientists (BMSs) play an important role in any health care system. However, they are at high risk of exposure to LMSDs. This study aimed to estimate the prevalence and the associated risk factors of LMSDs among this group of healthcare professionals. Methods: A cross-sectional survey (Nordic musculoskeletal) was used to estimate the prevalence of LMSDs among BMSs. Data were analyzed using Statistical Package for Social Science software version 25. Chi-square was performed to find the significant association between LMSDs and different risk factors. Results: The study included 83 BMSs. Females represented 63.9% and 36.1% were in the age group of 35–44. The overall prevalence of LMSDs was 77.1%. The most prevalent LMSDs were neck, shoulders, and lower back with 50.6%, 49.4%, and 43.4%, respectively. Neck complaints and upper back complaints were found statistically significant with the female gender. Shoulders complaints were associated with pipetting and microscopy. Lower back complaints were associated with pipetting and heavy work at home. A total of 65.57% of BMSs had irregular symptoms of LMSDs, 54.10% experienced moderate pain due to these symptoms, and 44.26% had symptoms that persisted from hours to days. Conclusion: The study found that the prevalence of LMSDs among BMSs was high. Good knowledge, attitude, practice, and training of ergonomics may minimize the prevalence of LMSDs among BMSs.

Key words: Biomedical scientists; Ergonomics; Laboratory-related musculoskeletal disorders; Prevalence; Risk factors.

Introduction

Musculoskeletal disorders (MSDs), known as cumulative trauma disorders or repetitive stress injuries, are injuries to muscles, nerves, tendons, ligaments, joints, cartilage, and spinal discs (1). Whereas work-related musculoskeletal disorders (WMSDs) are conditions in which the work environment and performance of work contribute significantly to the condition, and/or the condition is made worse or persists longer due to work conditions (2). To the best of our knowledge, we are the first to introduce the term laboratory-related musculoskeletal disorders (LMSDs) and can be defined simply as injuries resulted from working in the laboratory.

Biomedical scientists (BMSs) play an important role in any health care institutions. They work in medical laboratories, where specimens are prepared and processed for the detection, diagnosis and treatment of various diseases. BMSs are professionals that identify pathogenic microorganisms in microbiology, measure chemicals in biochemistry, match blood samples in hematology, help to diagnose cancer samples in histopathology, recognize antigen molecules in immunology, detect gene variants in genetics, and others. In medical laboratory, they are exposed to repetitive activities such as microscopy, microtomy, pipetting, and mostly work on standing positions. With the time, repetitive motion can affect muscles, joints, tendons and nerves (3). The long-term and severe LMSDs can highly affect the productivity of the work and shortness the work duration, therefore this will eventually lead to occupational disability and cause a major health challenge for individuals and healthcare systems around the world (4). About 33% of all sickness absence of health care professionals result from MSDs (5). MSDs are one of the common health problems among all health professionals (6). In addition, about 30% of people worldwide have at least MSDs (7).

Few studies are available for the ergonomic hazard which can lead to LMSDs (8). Ergonomic is simply the balance between the work requirement and the capacity of the worker (9). In addition, MSDs studies on certain professionals such as dentists, nurses and surgeons are well covered, however, data on the prevalence of LMSDs among BMSs is scarce. Hence, this study aimed to evaluate the prevalence of LMSDs among BMSs and assess the risk factors associated with BMSs.

Methods

This is a cross-sectional observational study that was conducted in the year of 2020 at the Sultan Qaboos University Hospital and Sultan Qaboos University. The study is ethically approved from the Medical Research Ethics Committee, College of Medicine and Health Sciences, SQU (SQU-EC/095/2020). Preceding the study, a detailed procedure of the study was explained, a voluntary consent form was also signed by each participant before the study. BMSs with occupational or non-occupational accidents that affected their musculoskeletal system were excluded from the study.

The sample size was calculated using the infinite population sample size formula: $n = \frac{N \times Z^2 \times p \times (1-p)}{d^2 \times (N-1) + Z^2 \times p \times (1-p)}$ where N : Population size (The total number of BMS working at SQU and SQUH) = 170, Z (Standard value with confidence level 95%) = 1.96, d (Permissible error on either side) = 10%, p (Proportion of the characteristic under the study). At the present, there is no available information on the proportion of MSDs among BMS. Therefore, the *P*-value was obtained from a pilot study which was conducted among 20 participants who were fulfilled the research criteria. Those who participated in the pilot study were excluded from the study. The *P*-value was found equal to 63.5%. The sample size was calculated as 97. After sampling, the number was increased 15% to avoid non-response or inappropriately filled questions and that number was randomly selected from the overall population.

The survey included two main sections. The first section was used to obtain data about socio-demographic characteristics such as age, gender, weight, height, nationality, specialty, educational level, nature of the job, and the number of experience years. The second section was from a previously validated Nordic musculoskeletal questionnaire (10). This section was used to assess the prevalence of MSDs among the participants. The first two questions were about the health status of the participants and whether they had any occupational or non-occupational accidents that affected their musculoskeletal system. Therefore, those who answered yes in these two questions were excluded because they did not fulfill the research criteria. The participants were asked in this section if they experienced any pain at different nine anatomical regions of the

body, specifically at: neck, shoulder, upper back, elbows, lower back, wrists/hands, hips/thighs, knees, and feet/ankles in the last year preceding the survey and also in the last seven days before the study. Further, the participants were asked in this section if they visited a physiotherapist or prevented from doing works due to any pain in the same nine regions mentioned above. There was a diagram to show these body sites for clarification to the participants. The section also includes questions about the intensity, frequency, and duration of the pain related to any musculoskeletal injuries.

The data were analyzed using Statistical Package for Social Science (SPSS) version 25 software. Frequencies and percentages were used to represent the categorical data such as age, gender, and specialty. Continuous data were presented as mean and standard deviation. A Chi-square test was performed to measure the significant association between different risk factors and the prevalence of LMSDs among the BMSs. The *P*-value was considered significant if it was equal to or less than 0.05.

Results

A total of 114 BMSs participated in the study. However, 31 BMSs were excluded because they did not fulfil the research criteria. Therefore a total of 83 BMSs were included in this study. The vast majority were females (63.9%) and 75.9% were married. The majority of the BMSs (36.1%) were in the age group of 35-44. Their average body mass index (BMI) was 25.3 (± 4.6). More than half of the participants (53.0%) did regular physical exercises. Almost all the participants are right-handed (94.0%). More than half (73.2%) of the participants used to work while they were standing. A total of 60.2% BMSs were sometimes stressed at work, 50.0% were sometimes working overtime, and 79.0% did not have heavy work at home (Table 1).

The overall prevalence of LMSDs among BMSs in the last 12 months in any parts of the body was 77.1%. Neck complaint was the highest; 50.6% and 25.3% in the last 12 months and 7 days respectively. Followed by shoulders complaint 49.4% and 16.9%, lower back complaint 43.4% and 15.7%, upper back complaint 34.9% and 14.5%, knees complaint 30.1% and 8.4%, ankle or feet 26.5% and 4.8%, wrists or hands 24.1% and 4.8% respectively. Elbows complaint was the lowest in both periods, 9.6% and 2.4% followed by thighs

or hip complaints, 10.8%, and 3.6% respectively (Figure 1). A total of 65.57% of BMSs had irregular symptoms of LMSDs, 54.10% experienced moderate pain due to these symptoms, and 44.26% had symptoms that persisted from hours to days (Figure 2).

There was no significant association between age and marital status and LMSDs among BMSs. However, there was a significant association between the neck and upper back complaints with the female gender (Table 2). In addition, no significant association was found between nationality, designation and degree of qualification with the LMSDs among the participants. None of the complaints were significantly associated with dominant hand, BMI, and regular physical exercises. (Table 3). Heavy work at home was significantly associated with a lower back complaint. There was no significant association between working overtime and stressed at work with LMSDs in this study (Table 4). Regarding the nature of the job, shoulders complaint was significantly associated with microscopy and pipetting. Lower back complaints show a significant association with pipetting (Table 5).

Discussion

In the present study, the overall prevalence of the LMSDs among BMSs was 77.1%. This finding is higher than a similar two studies, one was in Iran, which reported 72.4% MSDs among medical laboratory workers (11). The other one was in India which found that the overall prevalence of MSDs among medical laboratory technicians was 73.3% (8). However, this finding is lower than a recent study in Saudi Arabia which reported 82% MSDs among medical laboratory workers (12).

The most prevalent LMSDs among BMSs were neck followed by shoulders and lower back with 50.6%, 49.4%, 43.4%, respectively. This finding is in line with other study conducted in Metropolitan Washington, which found that neck complaint (61.5%) was the most prevalent symptom. However, their study also found that the wrists/hands (56.4%) symptoms were also high among their participants. This could be because their study only involved cytotechnologists who deal more with microscopes, and frequently use their hands/wrists

to adjust the microscope knob (13). Similar finding also reported that neck (55-60%), followed by upper back (53%), and lower back (57%) were the most prevalent MSDs among cytotechnologist (14). In addition, similar findings were reported among laboratory technicians in which they found that neck was the most prevalent MSDs (3,15,16). Another two studies, which used pathologists as a target health professionals found that neck was the most prevalent MSDs (17,18).

In contrast with our findings, a similar study conducted in India, which included medical laboratory technicians from different departments such as pathology, microbiology haematology, histology, and biochemistry. They found that the lower back complaint (32.5%) was the most prevalent followed by knees (20.7%) (19). Another similar study was conducted in Ethiopia revealed that ankles/feet (21.7%) and knees (20.8%) were the most affected parts (3). In addition, two similar studies in Saudi Arabia and Iran found that lower back was the most prevalent MSDs among clinical laboratory workers (11,12).

The finding of our study could be explained by the fact the BMSs used to work with different procedures and deal mostly with specimens that require the use of microscopes. Therefore, they used to bend their neck for a long time. Further, they used to work with static or repetitive tasks without relaxing their arms. This may explain the reasons for their shoulders complaints. The high prevalence of lower back complaints could be due to nonadjustable chairs with unsupported back posture available in their workstations. Another factor that might be a reason for the lower back complaint is the job nature, in which the BMSs used to work for a long time while they were standing without back support. BMSs in the current study reported that their symptoms of LMSDs occurred irregularly (Figure2). This finding is also consistent with other similar study (13).

Some risk factors such as increasing age of medical laboratory technicians, and worsening of the working conditions as well as standing and sitting for long hours were found to be associated with developing MSDs among the medical laboratory workers (3). The current study found no statistically significant association with age, marital status, nationality, dominant hand, BMI, degree of qualifications, and working overtime. Similar findings were also reported (8). Usually the BMSs work in multitasking jobs in which they move from one

workstation to another. Therefore, they were able to avoid postural stress. So, this might explain the reasons for no significant association between age and the MSDs (8).

The female gender showed a significant association with neck and upper back symptoms among BMSs (Table 1). This finding is in agreement with other similar study (8). In contrast, other study found that the female gender had no association with a neck complaint (11). In general, the prevalence of MSDs was found to be more common among women rather than men (20). A study conducted in Amsterdam also found that female workers are at high risk to develop MSDs in the upper extremity (21). There were several explanations for this gender difference. One of these explanations is the biological differences between females and males. For example, both genders can expose to the same physical work demands, however, they have different maximal physical work capacities. In addition, many workstations or tools are not taking into account the anthropometric differences between the two genders. Furthermore, it is suggested that the female reproductive system and, use of oral contraceptives, pregnancy, and childbirth can explain the high risk of MSDs among females. However, the precise mechanism is still not explored (21).

In the current study, pipetting showed a significant association with shoulders and lower back complaints. This is consistent with other study that found that increasing pipette work was associated with an increase in shoulder complaints among laboratory technicians (22). The significant association between pipetting and lower back complaints can be explained by the fact that BMSs did not use back supports while pipetting. Furthermore, our finding showed a significant association between microscopy and shoulder complaints. This may due to increased stress of the shoulder ligaments while using the microscope for a long time.

The present study found that there was a significant association between the LMSDs specifically the lower back complaints and the heavy work at home. This was in agreement with similar study in which heavy work of home was also significant with MSDs but with wrists/hands (11). However, both studies found that there was no significant association between stress at work and MSDs among medical laboratories.

Several limitations of our study are worth noting. First, due to the COVID-19 pandemic, we were not able to spread the questionnaire face to face to the BMSs. Therefore, we were incapable to provide more explanation about questions for the participants and encourage them to participate. Second, the study was restricted to BMSs from a single hospital and university, even though this hospital serves as a tertiary referral hospital in Oman and the university is the national university of the Sultanate of Oman. Finally, the study did not distinguish if those LMSDs were partially or fully related to the instruments and machines used in different laboratories.

In conclusion, the study highlighted that the prevalence of LMSDs among BMSs is considerably high. The most affected parts were the neck, followed by the shoulder and lower back. The study also found that the prevalence of LMSD was associated with female gender, pipetting, microscopy, and heavy work at home. Good knowledge, attitude, practice, and training of ergonomics may minimize the prevalence of LMSDs among BMS.

Acknowledgments

The authors would like to thank all BMSs who participated in this study.

References

1. Oladeinde BH, Ekejindu IM, Omoregie R, Aguh OD. Awareness and knowledge of ergonomics among medical laboratory scientists in Nigeria. *Ann Med Health Sci Res.* 2015;5(6):423-427. doi:10.4103/2141-9248.177989.
2. Bernard BP, editor. U.S. Department of Health and Human Services, Centers for Disease control and Prevention, National Institute of Occupational Safety and Health. Musculoskeletal disorders and workplace factors: a critical review of epidemiologic evidence for work-related musculoskeletal disorders of the neck, upper extremity, and lower back. July 1997. DHHS (NIOSH) Publication No. 97-141. Available from: <https://www.cdc.gov/niosh/docs/97-141/>.
3. Haile EL, Taye B, Hussen F. Ergonomic workstations and work-related musculoskeletal disorders in the clinical laboratory. *Lab Med* 2012;43:11–9.
4. Dong H, Zhang Q, Liu G, Shao T, Xu Y. Prevalence and associated factors of musculoskeletal disorders among Chinese healthcare professionals working in tertiary hospitals: A cross-sectional study. *BMC Musculoskelet Disord* 2019;20(175):1–8. <https://doi.org/10.1186/s12891-019-2557-5>.
5. Meijssen P, Knibbe HJJ. Work-related musculoskeletal disorders of perioperative personnel in the Netherlands. *AORN J.* 2007;86(2):193–208.
6. Tinubu BM, Mbada CE, Oyeyemi AL, Fabunmi AA. Work-related musculoskeletal disorders among nurses in Ibadan, south-west Nigeria: a cross-sectional survey. *BMC Musculoskelet Disord.* 2010;11:12.
7. Vos T, Abajobir AA, Abate KH, Abbafati C, Abbas KM, Abd-Allah F, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet* 2017;390:1211–59.
8. Maulik S, Iqbal R, De A, Chandra AM. Evaluation of the working posture and prevalence of musculoskeletal symptoms among medical laboratory technicians. *J Back Musculoskelet Rehabil* 2014;27(4):453–61.
9. Smallwood J, Ajayi O. The impact of training on construction ergonomics knowledge and awareness. *Ergonomics* 2007;21(1):23–38.
10. Kuorinka I, Jonsson B, Kilbom A, Vinterberg H, Biering-Sørensen F, Andersson G, et al. Standardised Nordic questionnaires for the analysis of musculoskeletal symptoms. *Appl Ergon.* 1987;18:233–7.
11. Sadeghian F, Kasaeian A, Noroozi P, Vatani J, Taiebi SH. Psychosocial and individual characteristics and musculoskeletal complaints among clinical laboratory workers. *Int J Occup Saf Ergon.* 2014;20:355–61.
12. AlNekhila AF, AlTamimi AM, AlAqeel BY, AlHawary AA, AlFadhel SF, Masuadi EM. Work-related musculoskeletal disorders among clinical laboratory workers. *Avicenna J Med.* 2020;23:10(1):29–34.
13. Kalavar SS, Hunting KL. Musculoskeletal symptoms among cytotechnologists. *Lab Med.* 1996;27:765–9.
14. Thompson, SK, Mason, E. Dukes, S. Ergonomics and cytotechnologists: Reported musculoskeletal discomfort. *Diagn. Cytopathol* 2003;29: 364–7.
15. Maulik S, De A, R Iqbal. Work related musculoskeletal disorders among medical laboratory technicians. in 2012 Southeast Asian Network of Ergonomics Societies Conference (SEANES) 2012. DOI <https://doi.org/10.1109/SEANES.2012.6299585>.
16. Bjorksten MG, Almby B, Jansson ES: Hand and shoulder ailments among laboratory technicians using modern plunger-operated pipettes. *Appl Ergon.* 1994, 25: 88-94.

17. Vahdatpour B, Khosravi S, Rahimi A, Sattari S, Mogtaderi A, Dabiri F, et al. Work-related musculoskeletal disorders among radiologists in Isfahan: a cross-sectional study. *Res J Biol Sci.* 2010; 5(10): 664-9.
18. Fritzsche FR, Ramach C, Soldini D, Caduff R, Tinguely M, Cassoly E, et al. Occupational health risks of pathologists—results from a nationwide online questionnaire in Switzerland. *BMC Public Health.* 2012;12:1054.
19. Maulik S, Iqbal R. Occupational health and musculoskeletal symptoms among Indian medical laboratory technicians. *JOHE.* 2013;2:82–92.
20. Wijnhoven. HAH, De Vet. HCW, Picavet, HSJ. Prevalence of musculoskeletal disorders is systematically higher in woman than in men, *Clinical Journal of Pain* 2006;22(8): 717–724. <https://doi.org/10.1097/01.ajp.0000210912.95664.53>.
21. de Zwart BC, Frings-Dresen MH, Kilbom A. Gender differences in upper extremity musculoskeletal complaints in the working population. *Int Arch Occup Environ Health.* 2001;74(1):21–30.
22. Holm JW, Mortensen OS, Gyntelberg F. Upper limb disorders among biomedical laboratory workers using pipettes. *Cogent Medicine* 2016;3(1):1256849.

Table1: Demographic data of biomedical scientists

Age	N (%)	Gender	N (%)	Marital status	N (%)	Nationality	N (%)
<25	1(1.2%)	Male	30(36.1%)	Single	20(24.1%)	Omani	65(78.3%)
25-34	37(44.6%)	Female	53(63.9%)	Married	63(75.9%)	Non-Omani	18(21.7%)
35-44	30(36.1%)						
45-54	10(12.0%)						
>54	5(6.0%)						
Dominant hand	N (%)	Specialty	N (%)	Designation	N (%)	Qualification	N (%)
Right	78(94.0%)	Biochemistry	19(22.9%)	Superintendent	2(2.4%)	Diploma	7(8.4%)
Left	5(6.0%)	Genetics	7(8.4%)	Chief BMS	9(11.0%)	Degree	54(65.1%)
		Hematology	24(28.90%)	Senior BMS	30(36.6%)	Master	19(22.9%)
		Histopathology	15(18.1%)	Junior BMS	40(48.8%)	PHD	3(3.6%)
		Microbiology & Immunology	11(13.3%)	Researcher	1(1.2%)		
		Anatomy	6(7.2%)				
		Physiology	1(1.2%)				
Nature of the job	N (%)	Working over the time	N (%)	Stressed at work	N (%)	Heavy work at home	N (%)
Microscopy	32(39.0%)	Yes	11(13.4%)	Yes	14(16.9%)	Yes	17(21.0%)
Sitting	42(51.2%)	No	30(36.6%)	No	19(22.9%)	No	64(49.0%)
Standing	60(73.2%)	Sometimes	41(50.0%)	Sometimes	50(60.2%)		
Pipetting	46(56.1%)						
Microtomy	13(15.9%)						
Regular physical exercise	N (%)	Mean height in cm $\pm$SD: 162.46$\pm$8.09 Mean weight in kg $\pm$SD : 67.43$\pm$14.73 Mean BMI in kg/m² $\pm$SD: 25.31$\pm$4.63 Mean total years working $\pm$SD : 11.52$\pm$7.78					
Yes	44(53.0%)						
No	39(47.0%)						

BMI: Body mass index

Table 2: Prevalence and association of LMSDs with age, gender, and nationality of BMSs within 12 months.

Body regions	Risk Factors											
	Age N (%)					Gender N (%)				Marital status N (%)		
	25	25-34	35-44	44-54	>54	P-value	Female	Male	P-value	Married	Single	P-value
Neck	1(100%)	22(59.5%)	16(53.3%)	3(30.0%)	0(0.0%)	0.060	33(62.3%)	9 (30.0%)	0.009	28(44.4%)	14(70.0%)	0.083
Shoulders	0(0.0%)	20(54.1%)	16(53.3%)	4(40.0%)	1(20.0%)	0.468	30(56.6%)	11(36.7%)	0.129	29(46.0%)	12(60.0%)	0.405
Upper back	0(0.0%)	18(48.6%)	8(26.7%)	3(30.0%)	0.0(0.0%)	0.121	24(45.3%)	5(16.7%)	0.017	23(36.5%)	6(30.0%)	0.793
Elbows	0(0.0%)	4(10.8%)	2(6.7%)	2(20.0%)	0(0.0%)	0.693	6(11.3%)	2(6.7%)	0.762	6(9.5%)	2(10.0%)	1.000
Wrists/hands	1(100%)	11(29.7%)	6(20.0%)	2(20.0%)	0(0.0%)	0.219	13(24.5%)	7(23.3%)	1.000	12(19.0%)	8(40.0%)	0.108
Lower back	0(0.0%)	19(51.4%)	13(43.3%)	1(10.0%)	3(60.0%)	0.146	26(49.1%)	10(33.3%)	0.247	26(41.3%)	10(50.0%)	0.669
Hips/ thighs	0(0.0%)	3(8.1%)	4(13.3%)	2(20.0%)	0(0.0%)	0.722	6(11.3%)	3(10.0%)	1.000	7(11.1%)	2(10.0%)	1.000
Knees	0(0.0%)	14(37.8%)	6(20.0%)	3(30.0%)	2(40.0%)	0.530	8(26.7%)	17(32.1%)	0.789	17(27.0%)	8(40.0%)	0.409
Ankles/feet	1(100%)	11(29.7%)	7(23.3%)	3(30.0%)	0(0.0%)	0.288	7(23.3%)	15(28.3%)	0.815	13(20.6%)	9(45.0%)	0.063

Table 3: Prevalence and association of LMSDs with dominant hand, body mass index, and physical exercises of BMSs within 12 months.

Body regions	Risk factors										
	Dominant hand N (%)			BMI N (%)					Physical exercises N (%)		
	Right	Left	<i>P</i> -value	Underweigh t	Normal	Overweight	Obese	<i>P</i> - value	Yes	No	<i>P</i> -value
Neck	41(52.6%)	1(20.0%)	0.342	3(100.0%)	16(47.1%)	17(54.8%)	4(44.4%)	0.333	21(47.7%)	21(53.8%)	0.763
Shoulder	39(50.0%)	2(40.0%)	1.000	1(33.3%)	20(58.8%)	15(48.4%)	3(33.3%)	0.491	20(45.5%)	21(53.8%)	0.587
Upper back	29(37.2%)	0(0.0%)	0.228	3(100.0%)	12(35.3%)	8(25.8%)	4(44.4%)	0.070	13(29.5%)	16(41.0%)	0.387
Elbows	7(9.0%)	1(20.0%)	0.977	0(0.0%)	4(11.8%)	4(12.9%)	0(0.0%)	0.643	4(9.1%)	4(10.3%)	1.000
Wrists &hands	20(25.65)	0(0.0%)	0.447	2(66.7%)	8(23.5%)	9(29.0%)	0(0.0%)	0.105	10(22.7%)	10(25.6%)	0.958
Lower back	35(44.9%)	1(20.0%)	0.534	2(66.7%)	15(44.1%)	11(35.5%)	5(55.6%)	0.573	19(43.2%)	17(43.6%)	1.000
Hips/thighs	9(11.5%)	0(0.0%)	0.950	0(0.0%)	2(5.9%)	6(19.4%)	0(0.0%)	0.186	5(11.4%)	4(10.3%)	1.000
Knees	25(32.1%)	0(0.0%)	0.321	1(33.3%)	10(29.4%)	8(25.8%)	5(55.6%)	0.397	12(27.3%)	13(33.3%)	0.718
Ankles/feet	21(26.9%)	1(20.0%)	1.000	1(33.3%)	7(20.6%)	11(35.6%)	2(22.2%)	0.574	12(27.3%)	10(25.6%)	1.000

BMI: Body mass index

Table 4: Prevalence and association of LMSDs with the working overtime, heavy work at home, and stressed at work among the BMSs within 12 months.

Body regions	Risk factors										
	Working over the time N (%)				Heavy work at home N (%)			Stressed at work N (%)			
	Yes	No	Sometimes	<i>P</i> - value	Yes	No	<i>P</i> - value	Yes	No	Sometimes	<i>P</i> - value
Neck	5(54.5%)	14(46.7%)	21(51.2%)	0.883	9(52.9%)	32(50.0%)	1.000	10(71.4%)	7(36.8%)	25(50.0%)	0.144
Shoulder	5(45.5%)	16(53.5%)	19(46.3%)	0.821	9(52.9%)	31(48.4%)	0.954	8(57.1%)	6(31.6%)	27(54.0%)	0.205
Upper back	5(45.5%)	10(33.3%)	13(31.7%)	0.690	8(47.1%)	20(31.3%)	0.352	4(28.6%)	4(21.1%)	21(42.0%)	0.228
Elbows	1(9.1%)	2(6.7%)	4(9.8%)	0.897	3(17.6%)	4(6.3%)	0.371	2(14.3%)	1(5.3%)	5(10.0%)	0.680
Wrists & hands	2(18.2%)	8(26.7%)	9(22.0%)	0.821	4(23.5%)	14(21.9%)	1.000	2(14.3%)	4(21.1%)	14(28.0%)	0.535
Lower back	6(54.5%)	11(36.7%)	18(43.9%)	0.577	12(70.6%)	23(35.9%)	0.022	8(57.1%)	5(26.3%)	23(46.0%)	0.176
Hips/thighs	1(9.1%)	4(13.3%)	3(7.3%)	0.698	3(17.6%)	5(7.8%)	0.583	1(7.1%)	0(0.0%)	8(16.0%)	0.143
Knees	3(27.3%)	9(30.0%)	12(29.3%)	0.986	8(47.1%)	16(25.0%)	0.496	4(28.6%)	5(26.3%)	16(32.0%)	0.891
Ankles/feet	4(36.4%)	7(23.3%)	10(24.4%)	0.677	6(35.3%)	15(23.4%)	0.141	3(21.4%)	3(18.5%)	16(32.0%)	0.353

Table 5: Prevalence and association of LMSDs with the nature of the job of BMSs within 12 months.

Body regions	Nature of the job N (%)									
	Microscopy	<i>P- value</i>	Sitting	<i>P- value</i>	Standing	<i>P-value</i>	Pipetting	<i>P- value</i>	Microtomy	<i>P-value</i>
Neck	20(62.5%)	0.159	19(45.2%)	0.374	30(50.0%)	0.908	27(58.7)	0.191	7(53.8%)	1.000
Shoulder	21(56.6%)	0.042	21(50.0%)	1.000	29(48.3%)	0.803	28(26.9%)	0.045	5(58.5%)	0.545
Upper back	15(46.9%)	0.132	13(31.0%)	0.532	23(28.3%)	0.504	19(41.3%)	0.299	5(83.5%)	1.000
Elbow	6(18.8%)	0.070	3(7.1%)	0.656	5(8.3%)	0.766	3(6.5%)	0.495	4(30.8%)	*NA
Wrists/Hands	9(28.1%)	0.714	8(19.0)	0.730	16(26.7%)	0.615	12(28.1%)	0.884	5(38.5%)	0.349
Lower back	14(43.8%)	1.000	20(47.6%)	0.736	28(46.7%)	0.561	25(24.3%)	0.054	7(53.8%)	0.629
Hips/Thighs	3(9.4%)	0.993	3(7.1%)	0.433	6(10.0%)	0.647	5(10.9%)	1.000	2(15.4%)	0.944
Knees	9(28.1%)	0.900	11(26.2%)	0.531	20(33.3%)	0.513	16(34.8%)	0.476	5(38.5%)	0.725
Ankles/Feet	10(31.3%)	0.640	9(21.4%)	0.378	17(28.3%)	0.821	15(32.6%)	0.278	3(23.1%)	1.000

NA: Not applicable

Figure legends

Figure 1: Comparison between the prevalence of LMSDs among BMSs in the last 7 days and 12 months preceding the study.

Figure 2: Pain and symptoms characteristics of the LMSDs experienced by BMSs working within 12 months preceding the study.

Intensity of pain

- Mild
- Moderate
- Severe
- Very severe

Duration of symptoms

- Minutes to hours
- Hours to a day
- Days to a week
- weeks to a month
- More than a month

Frequency of Symptoms

- Daily
- Once a week
- Once a month
- Irregular