

Breast and prostate cancer risk: the interplay of polygenic risk, high-impact monogenic variants, and family history




Figure S1: Prevalence of the breast and prostate cancer according to polygenic risk score (PRS) percentiles.




Figure S2: Distributions of the breast and prostate cancer PRS across self-reported ethnic groups in the UK Biobank cohort (Data-Field 21000). Raw PRS was adjusted using first four principal components. Distributions of: a) raw PRS in breast cancer; b) adjusted PRS in breast cancer; c) raw PRS in prostate cancer; d) adjusted PRS in prostate cancer.




Figure S3: PRS among breast and prostate cases versus controls.






Figure S4: Distributions of the breast and prostate cancer PRS across carrier status in the UK Biobank cohort.




(a) breast cancer




(b) prostate cancer


Figure S5: Impact of PRS across individuals with monogenic variants carriers.




(a) PALB2




(b) CHEK2



(c) BRCA1




(d) BRCA2




(e) ATM


Figure S6: Breast cancer odds ratio (OR) among individuals stratified for presence of high-impact variants in each gene and PRS.




(a) CHEK2




(b) ATM



(c) BRCA1




(d) BRCA2




(e) HOXB13

Figure S7: Prostate cancer OR among individuals stratified for presence of high-impact variants in each gene and PRS.



(a) high risk variants (BRCA1, BRCA2)



(b) intermediate risk variants (PALB2, CHEK2, ATM)

Figure S8: Breast cancer OR among individuals stratified for presence of high-impact variants carrier in intermediate and high risk variants and PRS.




Figure S9: Breast cancer OR among individuals stratified for presence of a first-degree family history with breast cancer and PRS.




Figure S10: Prostate cancer OR among individuals stratified for presence of a first-degree family history with prostate cancer and PRS.

Table S1: Characteristics of participants by breast cancer status.

	Cases	Controls
Participants, N	3739	85974
Age*, mean (SD)	52.23 (7.57)	56.83 (8.35)
Family history of breast cancer, n (%)	762 (20.38)	9210 (10.71)

* Age: age at onset for cases, and age at recruitment for controls

Table S2: Characteristics of participants by prostate cancer status.

	Cases	Controls
Participants, N	1332	73111
Age*, mean (SD)	60.35 (4.44)	57.02 (8.68)
Family history of prostate cancer, n (%)	273 (20.5)	5689 (7.78)

* Age: age at onset for cases, and age at recruitment for controls