

# **Brain atrophy progression in Parkinson's disease is shaped by connectivity and local vulnerability**

## ***Supplemental Information***

### **Inclusion criteria (participants with Parkinson's disease)**

- (1) at least two of the following: resting tremor, bradykinesia, and rigidity *or* either asymmetric resting tremor *or* asymmetric bradykinesia
- (2) a diagnosis of Parkinson's disease for 2 years or less at the screening stage
- (3) Hoehn & Yahr stage I or II at the baseline visit
- (4) confirmation of dopamine transporter deficit by single-photon emission computed tomography
- (5) not expected to require Parkinson's disease medications within at least 6 months of the baseline visit
- (6) at least 30 years old at the time of Parkinson's disease diagnosis

## Deformation based morphometry (DBM)

In brief, the raw T1-weighted MRI images were processed through an initial processing during which the image was resampled, corrected for bias field inhomogeneities, affine-registered, and entered into SPM's unified segmentation protocol. Next, the image underwent a more refined processing that includes skull stripping, brain parcellation, local tissue intensity transformation, partial volume estimation, and spatial normalization to a common reference using DARTEL. For each participant at each time point, the DBM pipeline produced a set of processed images including a voxel-wise map of the Jacobian determinants, which was used as a marker of local brain atrophy. A 2-mm FWHM smoothing kernel was then applied on the Jacobian maps and the log of the Jacobian determinant was calculated as the voxel-wise relative deformation value to create DBM maps. DBM processing was carried out with the Computational Anatomy Toolbox (CAT12)<sup>1</sup> implemented in Statistical Parametric Mapping software (SPM12, <https://www.fil.ion.ucl.ac.uk/spm/doc/>).

## Structural and Functional Network Analysis

### *1) Regional brain atrophy progression*

The brain was parcellated into 68, 114, 219, and 448 approximately equally sized cortical regions using the Cammoun atlas.<sup>2</sup> Each analysis was repeated at all four resolutions to ensure that the results were robust regardless of the spatial scale. The average atrophy progression (i.e., mean W-Score difference between the follow-up time point and baseline) was calculated for each cortical region.

### *2) Structural and functional network construction*

Structural and functional connectivity matrices were respectively constructed from diffusion-weighted and resting-state functional MRI scans acquired from 70 healthy adults (age =  $29 \pm 9$  year, 43 men)<sup>3</sup> (<https://doi.org/10.5281/zenodo.2872624>). For the structural connectome, a deterministic streamline tractography approach was used. A binary group-average structural connectivity matrix was computed for each of the four resolution scales using a distance-dependent consensus approach that preserved the connection-length distribution of each participant<sup>4</sup> (<https://www.brainnetworkslab.com/coderesources>). In addition, the mean functional connectivity of all the 70 participants was calculated to obtain the group-average functional connectome.

### 3) *Collective neighborhood atrophy progression*

The group-level structural and functional connectivity matrices were used to compute the collective atrophy progression of the neighborhood of each region, an index of each node's "disease exposure". The collective atrophy progression of structurally connected neighbors ( $D_i$ ) was calculated as the mean atrophy progression of all the regions with a (deterministic) structural connection with a given region  $i$  (node) (see Equation 1). The collective atrophy progression of functional neighbors was calculated in a similar way, but with each neighbor  $j$ 's deformation value being weighted by the strength of the functional connectivity (FC) between node  $i$  and  $j$  (see Equation 2).

$$D_i = \frac{1}{N_i} \sum_{i \neq j, j=1}^{N_i} d_j \quad (1) \qquad D_i = \frac{1}{N_i} \sum_{i \neq j, j=1}^{N_i} d_j \times FC_{ij} \quad (2)$$

In both equations,  $D_i$  is the disease exposure or average deformation progression in the neighbors of region  $i$ ,  $d$  is the region atrophy progression, and  $N_i$  is the number of regions structurally connected to the region  $i$  (i.e., node degree). Therefore, only nodes with a direct structural connection are assumed to contribute to each other's atrophy progression. The deformation in the region under consideration was excluded from equations (1) and (2) i.e. regions did not contribute to their own atrophy progression. For the functional neighborhood atrophy progression (equation 2), only the regions structurally connected to the given region were considered.

### 4) *Correlations*

Pearson's correlations were calculated between the mean atrophy progression of all brain regions and the collective atrophy progression of their structurally connected neighbors and their structurally connected neighbors weighted by their FC.

### 5) *Comparison with a null model*

To investigate whether the relationships between the atrophy progression of a region and the atrophy progression of its structurally and functionally connected neighborhood depends on the network spatial topology, the correlations were tested against a null model preserving the spatial autocorrelation between regions<sup>5</sup> using the netneurotools toolbox (<https://netneurotools.readthedocs.io/en/latest/index.html>). This approach generates a null model by projecting the brain regions onto a sphere and randomly rotating the sphere. To address the loss of data caused by the medial wall rotation, this model assigns the nearest data to a missing parcel (see Markello and

Misic<sup>6</sup>) for a comparison between spatially-constrained null models). Briefly, a surface-based representation of the different resolution of the Cammoun atlas on the FreeSurfer (release v6.0.0; <http://surfer.nmr.mgh.harvard.edu/>) fsaverage surface was created. The spherical projection of the fsaverage surface was then used to define spatial coordinates for each region by selecting the vertex closest to the center of mass of the region. The spatial coordinates were used to create null models by rotating and reassigning region values 10,000 times. The regions centroids were assigned to the closest rotated centroids based on the minimum Euclidean distance. The spatial rotation was performed at the parcel resolution and in one hemisphere before being mirrored in the other. Finally, Pearson's correlation coefficients that were derived from the empirical networks (i.e., observed values) were tested against those derived from the null networks (i.e., permuted values).

### References

1. Gaser C, Dahnke R. CAT-A Computational Anatomy Toolbox for the Analysis of Structural MRI Data. *Human Brain Mapping*. 2016;32(7):336-348.
2. Cammoun L, Gigandet X, Meskaldji D, et al. Mapping the human connectome at multiple scales with diffusion spectrum MRI. *Journal of Neuroscience Methods*. 2012;203(2):386-397.  
doi:10.1016/j.jneumeth.2011.09.031
3. Mišić B, Betzel RF, Nematzadeh A, et al. Cooperative and Competitive Spreading Dynamics on the Human Connectome. *Neuron*. 2015;86(6):1518-1529. doi:10.1016/j.neuron.2015.05.035
4. Betzel RF, Griffa A, Hagmann P, Misic B. Distance-dependent consensus thresholds for generating group-representative structural brain networks. *Network Neuroscience*. 2019;3(2):475-496.  
[http://dx.doi.org/10.1162/netn\\_a\\_00075](http://dx.doi.org/10.1162/netn_a_00075).
5. Vázquez-Rodríguez B, Suárez LE, Markello RD, et al. Gradients of structure–function tethering across neocortex. *Proceedings of the National Academy of Sciences of the United States of America*. 2019;116(42):21219-21227. doi:10.1073/pnas.1903403116
6. Markello R, Misic B. Comparing spatially-constrained null models for parcellated brain maps. *bioRxiv*. 2020:2020.08.13.249797. <http://biorxiv.org/content/early/2020/08/14/2020.08.13.249797.abstract>.

**Table 1.** Regions showing a significant atrophy progression with time (F-Contrast)

F-Value	Cluster Size Number of voxels	Peak xyz [mm]	Overlap with the neuromorphometric atlas	Brain regions			
25.34	24366	57 -4 -16	41.00%	Right Cerebral White Matter			
			14.00%	Left Cerebral White Matter			
			6.00%	Right MTG middle temporal gyrus			
			5.00%	Right AnG angular gyrus			
			4.00%	Right PCu precuneus			
			4.00%	Left Lateral Ventricle			
			3.00%	Right TMP temporal pole			
			3.00%	Right Lateral Ventricle			
			3.00%	Right ITG inferior temporal gyrus			
			3.00%	Unknown			
			2.00%	Right FuG fusiform gyrus			
			2.00%	Right SPL superior parietal lobule			
			2.00%	Right STG superior temporal gyrus			
			2.00%	Right PCgG posterior cingulate gyrus			
			1.00%	Right MOG middle occipital gyrus			
			1.00%	Right IOG inferior occipital gyrus			
			21.33	11517	-52 -46 8	35.00%	Left Cerebral White Matter
						19.00%	Left MTG middle temporal gyrus
						8.00%	Left TMP temporal pole
						6.00%	Left ITG inferior temporal gyrus
6.00%	Left STG superior temporal gyrus						
5.00%	Left AnG angular gyrus						
5.00%	Left PCu precuneus						
3.00%	Left SPL superior parietal lobule						
3.00%	Left MOG middle occipital gyrus						
2.00%	Left FuG fusiform gyrus						
2.00%	Left IOG inferior occipital gyrus						
1.00%	Left PCgG posterior cingulate gyrus						
1.00%	Left SMG supramarginal gyrus						
1.00%	Left Amygdala						
20.6	2396	12 -32 -40	66.00%	Left Ent entorhinal area			
			17.00%	Brain Stem			
16.18	388	-46 -26 44	17.00%	Left Cerebellum White Matter			
			17.00%	Right Cerebellum White Matter			
21.16	286	-14 12 58	37.00%	Left Cerebral White Matter			
			34.00%	Left PoG postcentral gyrus			
			25.00%	Left SMG supramarginal gyrus			
			4.00%	Left PO parietal operculum			
17.4	159	-8 20 -12	54.00%	Left Cerebral White Matter			
			46.00%	Left SFG superior frontal gyrus			
15.74	116	14 16 0	44.00%	Left Cerebral White Matter			
			20.00%	Left SCA subcallosal area			
			19.00%	Left MFC medial frontal cortex			
			14.00%	Left ACgG anterior cingulate gyrus			
			3.00%	Left Accumbens Area			
			97.00%	Right Caudate			
19.74	61	14 -74 -6	3.00%	Right Cerebral White Matter			
			52.00%	Right LiG lingual gyrus			
16.78	88	-40 21 12	48.00%	Right Cerebral White Matter			
			88.00%	Left Cerebral White Matter			
			9.00%	Left TrIFG triangular part of the inferior frontal gyrus			
15.34	59	16 26 16	3.00%	Left OpIFG opercular part of the inferior frontal gyrus			
			100.00%	Right Cerebral White Matter			
15.26	60	8 28 32	98.00%	Right MSFG superior frontal gyrus medial segment			
			2.00%	Right ACgG anterior cingulate gyrus			
15.16	59	4 16 -18	56.00%	Right Cerebral White Matter			
			31.00%	Right SCA subcallosal area			
			14.00%	Right Accumbens Area			
			100.00%	Left CO central operculum			
14.83	13	-46 -20 20	72.00%	Right Hippocampus			
			72.00%	Right Hippocampus			
14.45	79	22 -9 -21	28.00%	Right Amygdala			
			74.00%	Right SMG supramarginal gyrus			
			14.00%	Right Cerebral White Matter			
13.89	99	42 -34 44	7.00%	Right PoG postcentral gyrus			
			5.00%	Right SPL superior parietal lobule			
			100.00%	Right Cerebral White Matter			
			100.00%	Right Cerebral White Matter			
13.7	16	16 33 3	70.00%	Left SMC supplementary motor cortex			
			20.00%	Left SFG superior frontal gyrus			
13.69	18	16 12 46	10.00%	Unknown			
			79.00%	Right PHG parahippocampal gyrus			
			14.00%	Right Cerebral White Matter			
13.67	10	-3 20 64	7.00%	Right FuG fusiform gyrus			
			91.00%	Unknown			
13.66	14	30 -9 -32	9.00%	Left SMC supplementary motor cortex			
			100.00%	Left MCgG middle cingulate gyrus			
13.55	11	2 14 64	100.00%	Left MSFG superior frontal gyrus medial segment			
			100.00%	Right Cerebral White Matter			
13.46	13	-4 9 38	100.00%	Right Cerebral White Matter			
			55.00%	Right TMP temporal pole			
13.29	15	-2 39 44	45.00%	Right Ent entorhinal area			
			100.00%	Right Cerebral White Matter			
13.28	12	12 12 58					
13.09	11	26 4 -30					


**Table 2.** Regions showing a significant atrophy progression after two and four years (T-Contrasts)

T-Value	Cluster Size Number of voxels	Peak xyz [mm]	Overlap with the neuromorphometric atlas	Brain regions
<b>Contrast: Baseline - 2 years</b>				
6.55	487	46 -62 12 45 -62 26	40.00% 31.00% 25.00%	Right AnG angular gyrus Right Cerebral White Matter Right MTG middle temporal gyrus
6.68	285	57 -4 -16 57 -15 -10	53.00% 43.00% 5.00%	Right MTG middle temporal gyrus Right STG superior temporal gyrus Right Cerebral White Matter
6.66	188	14 15 2	94.00%	Right Caudate
5.98	143	-57 -28 -22	62.00% 20.00% 19.00%	Left MTG middle temporal gyrus Left Cerebral White Matter Left ITG inferior temporal gyrus
6.16	69	-52 3 -22	51.00% 36.00%	Left TMP temporal pole Left MTG middle temporal gyrus
6.94	43	-6 -54 38	98.00%	Left PCu precuneus
5.77	27	33 9 -42	100.00%	Right TMP temporal pole
5.69	18	46 12 -33	100.00%	Right TMP temporal pole
5.78	14	-56 -6 -10	100.00%	Left STG superior temporal gyrus
5.73	13	-56 -45 9	100.00%	Left STG superior temporal gyrus
5.65	10	-60 -15 -12	100.00%	Left MTG middle temporal gyrus
<b>Contrast: Baseline - 4 years</b>				
8.53	48093	40 -64 39	22.00% 18.00% 7.00% 5.00% 4.00% 3.00% 3.00% 3.00% 3.00% 2.00% 2.00% 2.00% 2.00% 2.00% 2.00% 2.00% 2.00% 1.00% 1.00% 1.00% 1.00% 1.00% 1.00% 1.00% 1.00% 1.00% 1.00%	Right Cerebral White Matter Left Cerebral White Matter Left MTG middle temporal gyrus Right MTG middle temporal gyrus Right PCu precuneus Right AnG angular gyrus Left ITG inferior temporal gyrus Left TMP temporal pole Right TMP temporal pole Left PCu precuneus Left AnG angular gyrus Left STG superior temporal gyrus Right ITG inferior temporal gyrus Right SPL superior parietal lobule Right FuG fusiform gyrus Left IOG inferior occipital gyrus Right STG superior temporal gyrus Left MOG middle occipital gyrus Left SPL superior parietal lobule Right MOG middle occipital gyrus Right PCgG posterior cingulate gyrus Right SMG supramarginal gyrus
7.63	234	14 -74 -6	58.00% 42.00%	Right LiG lingual gyrus Right Cerebral White Matter
7.42	286	-14 12 58	50.00% 50.00%	Left Cerebral White Matter Left SFG superior frontal gyrus
6.81	131	-40 21 12	94.00% 5.00%	Left Cerebral White Matter Left OplFG opercular part of the inferior frontal gyrus
6.73	239	-9 21 -12	39.00% 21.00% 19.00% 18.00% 2.00%	Left Cerebral White Matter Left SCA subcallosal area Left MFC medial frontal cortex Left ACgG anterior cingulate gyrus Left Accumbens Area
6.14	47	6 38 27	100.00%	Right MSFG superior frontal gyrus medial segment
6.13	296	15 20 -4	58.00% 23.00% 10.00% 9.00%	Right Caudate Right Cerebral White Matter Right Accumbens Area Right SCA subcallosal area
5.76	106	-33 -45 42	63.00% 25.00% 11.00%	Left SPL superior parietal lobule Left SMG supramarginal gyrus Left Cerebral White Matter
5.53	16	58 -3 -36	100.00%	Unknown
5.48	55	20 -84 -12	49.00% 35.00% 16.00%	Right LiG lingual gyrus Right Cerebral White Matter Right OFuG occipital fusiform gyrus
5.46	15	-14 -44 70	60.00% 40.00%	Left PoG postcentral gyrus Left Cerebral White Matter
5.43	14	15 -78 -14	100.00%	Right LiG lingual gyrus
5.42	10	20 -98 -9	60.00% 20.00% 20.00%	Right OCP occipital pole Right Cerebral White Matter Right OFuG occipital fusiform gyrus


**Table 3.** Regions showing a significant atrophy progression after one and two years after including all the subjects with a T1-MRI at baseline and one year ( $N=120$ ), and baseline and two years ( $N=109$ )





T-Value	Cluster Size Number of voxels	Peak xyz [mm]	Overlap with the neuromorphometric atlas	Brain regions
<b>Contrast: Baseline - 1 year</b>				
6.38	58	-63 -32 -4	98.00%	Left MTG middle temporal gyrus
			2.00%	Left Cerebral White Matter
6.37	42	-52 -18 -30	86.00%	Left ITG inferior temporal gyrus
			14.00%	Left Cerebral White Matter
6.24	89	-58 -16 -14	92.00%	Left MTG middle temporal gyrus
			6.00%	Left Cerebral White Matter
			2.00%	Left STG superior temporal gyrus
6.13	20	18 18 4	95.00%	Right Caudate
			5.00%	Right Cerebral White Matter
6.01	40	10 8 10	100.00%	Right Caudate
5.97	53	-56 -48 -4	51.00%	Left MTG middle temporal gyrus
			49.00%	Left Cerebral White Matter
5.94	20	-45 -30 -21	90.00%	Left ITG inferior temporal gyrus
			5.00%	Left Cerebral White Matter
			5.00%	Left FuG fusiform gyrus
5.77	17	-56 -28 -22	59.00%	Left ITG inferior temporal gyrus
			41.00%	Left Cerebral White Matter
5.74	13	-54 -3 -26	62.00%	Left MTG middle temporal gyrus
			38.00%	Left Cerebral White Matter
<b>Contrast: Baseline -2 years</b>				
8.71	4530	-56 -27 -22	34.00%	Left MTG middle temporal gyrus
			19.00%	Left TMP temporal pole
			15.00%	Left Cerebral White Matter
			11.00%	Left STG superior temporal gyrus
			10.00%	Left ITG inferior temporal gyrus
			5.00%	Left AnG angular gyrus
			4.00%	Left FuG fusiform gyrus
7.62	817	57 -3 -18	37.00%	Right TMP temporal pole
			36.00%	Right MTG middle temporal gyrus
			16.00%	Right STG superior temporal gyrus
			8.00%	Right Cerebral White Matter
			2.00%	Right ITG inferior temporal gyrus
7.61	655	15 21 -3	87.00%	Right Caudate
			6.00%	Right Cerebral White Matter
			6.00%	Right Accumbens Area
7.55	823	46 -60 12	46.00%	Right Cerebral White Matter
			27.00%	Right MTG middle temporal gyrus
			23.00%	Right AnG angular gyrus
			3.00%	Right MOG middle occipital gyrus
6.68	129	-50 24 6	74.00%	Left Cerebral White Matter
			13.00%	Left OpIFG opercular part of the inferior frontal gyrus
			13.00%	Left TrIFG triangular part of the inferior frontal gyrus
6.28	22	58 -3 -36	86.00%	Unknown
			14.00%	Right ITG inferior temporal gyrus
6.27	39	26 -68 36	72.00%	Right SPL superior parietal lobule
			13.00%	Right Cerebral White Matter
			8.00%	Right AnG angular gyrus
			8.00%	Right SOG superior occipital gyrus
6.19	14	4 -51 24	64.00%	Right PCgG posterior cingulate gyrus
			36.00%	Right PCu precuneus
6.04	26	-6 -56 38	100.00%	Left PCu precuneus
5.76	12	28 -8 -33	50.00%	Right PHG parahippocampal gyrus
			42.00%	Right FuG fusiform gyrus
			8.00%	Right Cerebral White Matter
5.75	14	44 15 -26	57.00%	Right TMP temporal pole
			43.00%	Right Cerebral White Matter
5.74	36	39 -57 28	67.00%	Right Cerebral White Matter
			33.00%	Right AnG angular gyrus





**Figure 1.** Overlap between the regions showing significant atrophy progression ( $P_{FWE}\text{-value}<0.05$ ) after including the subjects with a T1-MRI at all time points ( $N=74$ , Bl-2Y: red) and all the subjects with a T1-MRI at baseline and one year ( $N=120$ , Bl-1Y: green), and baseline and two years ( $N=109$ , Bl-2Y: blue) using the MNI152-2009c template for visualization purpose (Top row:  $x = 45, y = -62, z = -19$ ; Middle row:  $x = 14, y = 15, z = 7$ ; Bottom row:  $x = 54, y = -6, z = 30$ ). P: posterior; A: anterior; R: right; L: left




**Figure 2.** Relationship between the regional atrophy progression and the atrophy progression in their structurally (SC) and functionally connected (FC) neighbors after two and four years using the Cammoun atlas with 68 (I), 114 (II) and 219 (III) regions.




**II****Atrophy progression after 2 years****Atrophy progression after 4 years**

**III****Atrophy progression after 2 years****Atrophy progression after 4 years**


**Figure 3.** Relationship between the regional atrophy progression after two and four years and the prevalence of endothelial cells and oligodendrocytes using the Cammoun atlas with 68 (I), 114 (II) and 219 (III) regions.








## Atrophy progression after 2 years


III




## Atrophy progression after 4 years






**Figure 4.** No significant correlation between the atrophy progression after two and four years in *de novo* Parkinson's disease and the prevalence of astrocytes (I), microglia (II), oligodendrocyte precursors cells (III), excitatory neurons (IV) and inhibitory neurons (V) in the cortex.



## Atrophy progression after 2 years




## Atrophy progression after 4 years


**V**

**Figure 5.** Gene ontology (GO) enrichment analysis of all the genes related to atrophy progression in Parkinson's disease using the GOrilla platform. The fold enrichment was obtained by calculating the ratio of the genes positively related to the atrophy progression after two (A) and four years (B) over the number of expected genes for each GO term, based on the background genes list. Only results with a  $p\text{-value}_{Bonferroni} < 0.05$  were presented. No significant categories were found for the genes negatively related to the atrophy progression after two and four years. The GO terms from the GOrilla platform positively associated with atrophy progression after two ( $N=26$ ) and four years ( $N=23$ ) respectively overlap with 88% and 74% of the terms from PANTHER.


**A** GO terms: genes related to the atrophy progression after 2 years



**B** GO terms: genes related to the atrophy progression after 4 years



**Figure 6.** Gene ontology (GO) enrichment analysis of all the genes related to atrophy progression in Parkinson's disease using the PANTHER platform. The fold enrichment was obtained by calculating the ratio of the genes positively related to the atrophy progression after two (A) and four years (B) over the number of expected genes for each GO term, based on the background genes list. Only results with a  $p\text{-value}_{Bonferroni} < 0.05$  were presented. No significant categories were found for the genes negatively related to the atrophy progression after two and four years.



**Table 4.** Genes located in PD GWAS loci associated with each GO term related to atrophy progression after two years in *de novo* Parkinson's disease

GO terms	Genes located in PD GWAS loci	Nearest Genes
Regulation of synaptic transmission glutamatergic	None	
Synapse organization	BSN - bassoon presynaptic cytomatrix protein	IP6K2
Regulation of synaptic plasticity	PRRT2 - proline-rich transmembrane protein 2	CD19
Anterograde trans-synaptic signaling	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	EGR3 - early growth response 3	BIN3
Chemical synaptic transmission	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	EGR3 - early growth response 3	BIN3
Trans-synaptic signaling	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	EGR3 - early growth response 3	BIN3
Synaptic signaling	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	EGR3 - early growth response 3	BIN3
Modulation of chemical synaptic transmission	PRRT2 - proline-rich transmembrane protein 2	CD19
Regulation of trans-synaptic signaling	None	
Neuron projection morphogenesis	None	
Cell-cell signaling	FGFR3 - fibroblast growth factor receptor 3	TMEM175, GAK
	EGR3 - early growth response 3	BIN3
Cellular component morphogenesis	SHROOM3 - shroom family member 3	FAM47E, SCARB2, FAM47E-STBD1
Signaling	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	EGR3 - early growth response 3	BIN3
	FGFR3 - fibroblast growth factor receptor 3	TMEM175, GAK
	CAMK2D - calcium/calmodulin-dependent protein kinase ii delta	CAMK2D
Nervous system process	TTC8 - tetratricopeptide repeat domain 8	GALC
System process	TTC8 - tetratricopeptide repeat domain 8	GALC
	CAMK2D - calcium/calmodulin-dependent protein kinase ii delta	CAMK2D

Multicellular organismal process	ABCA1 - atp-binding cassette sub-family a (abc1) member 1	SLC44A1
	TRIM8 - tripartite motif containing 8	GBF1
	FGFR3 - fibroblast growth factor receptor 3	TMEM175, GAK
	TTC8 - tetratricopeptide repeat domain 8	GALC
	CAMK2D - calcium/calmodulin-dependent protein kinase ii delta	CAMK2D
	SHROOM3 - shroom family member 3	FAM47E, SCARB2, FAM47E-STBD1
	SMAD4 - smad family member 4	MEX3C
	TMEM107 - transmembrane protein 107	CHRNB1
Regulation of cell communication	FAM13A - family with sequence similarity 13 member a	SNCA
	ABCA1 - atp-binding cassette sub-family a (abc1) member 1	SLC44A1
	LGALS3 - lectin galactoside-binding soluble 3	GCH1
	UBAP2 - ubiquitin associated protein 2	UBAP2
	TRIM8 - tripartite motif containing 8	GBF1
	FGFR3 - fibroblast growth factor receptor 3	TMEM175, GAK
	PDCD4 - programmed cell death 4 (neoplastic transformation inhibitor)	ADRA2A, SLC44A1
	CAMK2D - calcium/calmodulin-dependent protein kinase ii delta	CAMK2D
	GRK5 - g protein-coupled receptor kinase 5	INPP5F, BAG3
	GPAM - glycerol-3-phosphate acyltransferase mitochondrial	ADRA2A
	SMAD4 - smad family member 4	MEX3C
Regulation of localization	ABCA1 - atp-binding cassette sub-family a (abc1) member 1	SLC44A1
	LGALS3 - lectin galactoside-binding soluble 3	GCH1
	TRIM8 - tripartite motif containing 8	GBF1
	C5orf30 - chromosome 5 open reading frame 30	PAM
	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	CAMK2D - calcium/calmodulin-dependent protein kinase ii delta	CAMK2D
	SEMA4A - sema domain immunoglobulin domain (ig) transmembrane domain (tm) and short cytoplasmic domain (semaphorin) 4a	PMVK, KRTPCAP2, GBAP1
	IPO5 - importin 5	MBNL2
	SMAD4 - smad family member 4	MEX3C
Regulation of signaling	FAM13A - family with sequence similarity 13 member a	SNCA
	ABCA1 - atp-binding cassette sub-family a (abc1) member 1	SLC44A1

	LGALS3 - lectin galactoside-binding soluble 3	GCH1
	UBAP2 - ubiquitin associated protein 2	UBAP2
	TRIM8 - tripartite motif containing 8	GBF1
	FGFR3 - fibroblast growth factor receptor 3	TMEM175, GAK
	PDCD4 - programmed cell death 4 (neoplastic transformation inhibitor)	ADRA2A, SLC44A1
	CAMK2D - calcium/calmodulin-dependent protein kinase ii delta	CAMK2D
	GRK5 - g protein-coupled receptor kinase 5	INPP5F, BAG3
	GPAM - glycerol-3-phosphate acyltransferase mitochondrial	ADRA2A
	SMAD4 - smad family member 4	MEX3C
Signal transduction	FAM13A - family with sequence similarity 13 member a	SNCA
	ABCA1 - atp-binding cassette sub-family a (abc1) member 1	SLC44A1
	TRIM8 - tripartite motif containing 8	GBF1
	SERP1 - stress-associated endoplasmic reticulum protein 1	MED12L
	FGFR3 - fibroblast growth factor receptor 3	TMEM175, GAK
	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	PDCD4 - programmed cell death 4 (neoplastic transformation inhibitor)	ADRA2A, SLC44A1
	CAMK2D - calcium/calmodulin-dependent protein kinase ii delta	CAMK2D
	SEMA4A - sema domain immunoglobulin domain (ig) transmembrane domain (tm) and short cytoplasmic domain (semaphorin) 4a	PMVK, KRTPCAP2, GBAP1
	PITPNM2 - phosphatidylinositol transfer protein membrane-associated 2	HIP1R
	GRK5 - g protein-coupled receptor kinase 5	INPP5F, BAG3
	SMAD4 - smad family member 4	MEX3C
Developmental process	LGALS3 - lectin galactoside-binding soluble 3	GCH1
	EGR3 - early growth response 3	BIN3
	TRIM8 - tripartite motif containing 8	GBF1
	PALLD - palladin cytoskeletal associated protein	CLCN3
	TMEM107 - transmembrane protein 107	CHRNA1
	SMC6 - structural maintenance of chromosomes 6	KCNS3
	ASXL3 - additional sex combs like 3 (drosophila)	ASXL3
	FGFR3 - fibroblast growth factor receptor 3	TMEM175, GAK
	PDCD4 - programmed cell death 4 (neoplastic transformation inhibitor)	ADRA2A, SLC44A1

	SEMA4A - sema domain immunoglobulin domain (ig) transmembrane domain (tm) and short cytoplasmic domain (semaphorin) 4a	PMVK, KRTCAP2, GBAP1
	SHROOM3 - shroom family member 3	FAM47E, SCARB2, FAM47E-STBD1
	SMAD4 - smad family member 4	MEX3C
Regulation of cellular process	FAM13A - family with sequence similarity 13 member a	SNCA
	ABCA1 - atp-binding cassette sub-family a (abc1) member 1	SLC44A1
	LGALS3 - lectin galactoside-binding soluble 3	GCH1
	EGR3 - early growth response 3	BIN3
	TRIM8 - tripartite motif containing 8	GBF1
	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	ZNF681 - zinc finger protein 681	TMEM175, GAK
	CAMK2D - calcium/calmodulin-dependent protein kinase ii delta	CAMK2D
	IPO5 - importin 5	MBNL2
	CELF5 - cugbp elav-like family member 5	SPPL2B
	CCNG2 - cyclin g2	FAM47E, SCARB2, FAM47E-STBD1
	UBTF - upstream binding transcription factor rna polymerase i	UBTF, FAM171A2
	YEATS2 - yeats domain containing 2	MCCC1
	SMC6 - structural maintenance of chromosomes 6	KCNS3
	ASXL3 - additional sex combs like 3 (drosophila)	ASXL3
	ZNF184 - zinc finger protein 184	LOC100131289
	UBAP2 - ubiquitin associated protein 2	UBAP2
	SERP1 - stress-associated endoplasmic reticulum protein 1	MED12L
	FGFR3 - fibroblast growth factor receptor 3	TMEM175, GAK
	C5orf30 - chromosome 5 open reading frame 30	PAM
	PDCD4 - programmed cell death 4 (neoplastic transformation inhibitor)	ADRA2A, SLC44A1
	SEMA4A - sema domain immunoglobulin domain (ig) transmembrane domain (tm) and short cytoplasmic domain (semaphorin) 4a	PMVK, KRTCAP2, GBAP1
	PITPNM2 - phosphatidylinositol transfer protein membrane-associated 2	HIP1R
	GRK5 - g protein-coupled receptor kinase 5	INPP5F, BAG3
	SHROOM3 - shroom family member 3	FAM47E, SCARB2, FAM47E-STBD1
	ZNF26 - zinc finger protein 26	FBRSL1

	GPAM - glycerol-3-phosphate acyltransferase mitochondrial	ADRA2A
	SMAD4 - smad family member 4	MEX3C
Regulation of biological process	ABCA1 - atp-binding cassette sub-family a (abc1) member 1	SLC44A1
	LGALS3 - lectin galactoside-binding soluble 3	GCH1
	IPO5 - importin 5	MBNL2
	CELF5 - cugbp elav-like family member 5	SPPL2B
	YEATS2 - yeats domain containing 2	MCCC1
	SMC6 - structural maintenance of chromosomes 6	KCNS3
	ZNF184 - zinc finger protein 184	LOC100131289
	C5orf30 - chromosome 5 open reading frame 30	PAM
	PITPNM2 - phosphatidylinositol transfer protein membrane-associated 2	HIP1R
	GRK5 - g protein-coupled receptor kinase 5	INPP5F, BAG3
	SHROOM3 - shroom family member 3	FAM47E, SCARB2, FAM47E-STBD1
	ZNF26 - zinc finger protein 26	FBRSL1
	GPAM - glycerol-3-phosphate acyltransferase mitochondrial	ADRA2A
	SMAD4 - smad family member 4	MEX3C
	FAM13A - family with sequence similarity 13 member a	SNCA
	EGR3 - early growth response 3	BIN3
	TRIM8 - tripartite motif containing 8	GBF1
	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	ZNF876P - zinc finger protein 876 pseudogene	TMEM175, GAK
	CAMK2D - calcium/calmodulin-dependent protein kinase ii delta	CAMK2D
	CCNG2 - cyclin g2	FAM47E, SCARB2, FAM47E-STBD1
	UBTF - upstream binding transcription factor rna polymerase i	UBTF, FAM171A2
	ASXL3 - additional sex combs like 3 (drosophila)	ASXL3
	UBAP2 - ubiquitin associated protein 2	UBAP2
	SERP1 - stress-associated endoplasmic reticulum protein 1	MED12L
	FGFR3 - fibroblast growth factor receptor 3	TMEM175, GAK
	PDCD4 - programmed cell death 4 (neoplastic transformation inhibitor)	ADRA2A, SLC44A1
	SEMA4A - sema domain immunoglobulin domain (ig) transmembrane domain (tm) and short cytoplasmic domain (semaphorin) 4a	PMVK, KRTCAP2, GBAP1

**Table 5.** Genes located in PD GWAS loci associated with each GO term related to atrophy progression after four years in *de novo* Parkinson's disease

GO terms	Genes located in PD GWAS loci	Nearest Genes
Regulation of synaptic plasticity	RIMS1 - regulating synaptic membrane exocytosis 1	RIMS1
	MAPT - microtubule-associated protein tau	CRHR1, WNT3
Anterograde trans-synaptic signaling	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	EGR3 - early growth response 3	BIN3
	P2RX2 - purinergic receptor p2x ligand-gated ion channel 2	FBRSL1
	DOC2A - double c2-like domains alpha	SETD1A
Chemical synaptic transmission	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	DOC2A - double c2-like domains alpha	SETD1A
	EGR3 - early growth response 3	BIN3
	P2RX2 - purinergic receptor p2x ligand-gated ion channel 2	FBRSL1
Trans-synaptic signaling	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	DOC2A - double c2-like domains alpha	SETD1A
	EGR3 - early growth response 3	BIN3
	P2RX2 - purinergic receptor p2x ligand-gated ion channel 2	FBRSL1
Synaptic signaling	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	DOC2A - double c2-like domains alpha	SETD1A
	EGR3 - early growth response 3	BIN3
	P2RX2 - purinergic receptor p2x ligand-gated ion channel 2	FBRSL1
Modulation of chemical synaptic transmission	MAPT - microtubule-associated protein tau	CRHR1, WNT3
	RIMS1 - regulating synaptic membrane exocytosis 1	RIMS1
Regulation of trans-synaptic signaling	MAPT - microtubule-associated protein tau	CRHR1, WNT3
	RIMS1 - regulating synaptic membrane exocytosis 1	RIMS1
Cell-cell signaling	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	DOC2A - double c2-like domains alpha	SETD1A
	EGR3 - early growth response 3	BIN3
	FGFR3 - fibroblast growth factor receptor 3	TMEM175, GAK
	MAPT - microtubule-associated protein tau	CRHR1, WNT3

	P2RX2 - purinergic receptor p2x ligand-gated ion channel 2	FBRSL1
Signaling	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	DOC2A - double c2-like domains alpha	SETD1A
	EGR3 - early growth response 3	BIN3
	FGFR3 - fibroblast growth factor receptor 3	TMEM175, GAK
	CAMK2D - calcium/calmodulin-dependent protein kinase ii delta	CAMK2D
	MAPT - microtubule-associated protein tau	CRHR1, WNT3
	P2RX2 - purinergic receptor p2x ligand-gated ion channel 2	FBRSL1
Regulation of plasma membrane bounded cell projection organization	SEMA4A - sema domain immunoglobulin domain (ig) transmembrane domain (tm) and short cytoplasmic domain (semaphorin) 4a	PMCK, KRTCAP2, GBAP1
	RIMS1 - regulating synaptic membrane exocytosis 1	RIMS1
	MAPT - microtubule-associated protein tau	CRHR1, WNT3
	TAPT1 - transmembrane anterior posterior transformation 1	BST1
Regulation of cell projection organization	MAPT - microtubule-associated protein tau	CRHR1, WNT3
	RIMS1 - regulating synaptic membrane exocytosis 1	RIMS1
	SEMA4A - sema domain immunoglobulin domain (ig) transmembrane domain (tm) and short cytoplasmic domain (semaphorin) 4a	PMCK, KRTCAP2, GBAP1
	TAPT1 - transmembrane anterior posterior transformation 1	BST1
Cell communication	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	DOC2A - double c2-like domains alpha	SETD1A
	EGR3 - early growth response 3	BIN3
	FGFR3 - fibroblast growth factor receptor 3	TMEM175, GAK
	MAPT - microtubule-associated protein tau	CRHR1, WNT3
	P2RX2 - purinergic receptor p2x ligand-gated ion channel 2	FBRSL1
Multicellular organismal process	CAMK2D - calcium/calmodulin-dependent protein kinase ii delta	CAMK2D
	ABCA1 - atp-binding cassette sub-family a (abc1) member 1	SLC44A1
	FGFR3 - fibroblast growth factor receptor 3	TMEM175, GAK
	BTRC - beta-transducin repeat containing e3 ubiquitin protein ligase	GBF1
	MAPT - microtubule-associated protein tau	CRHR1, WNT3
	LEFTY1 - left-right determination factor 1	ITPKB
	NMT1 - n-myristoyltransferase 1	UBTF, FAM171A2, CRHR1

	P2RX2 - purinergic receptor p2x ligand-gated ion channel 2	FBRSL1
	RIMS1 - regulating synaptic membrane exocytosis 1	RIMS1
	SHROOM3 - shroom family member 3	FAM47E, SCARB2, FAM47E-STBD1
	RUNDC1 - run domain containing 1	RETREG3
	SMAD4 - smad family member 4	MEX3C
	TRIM8 - tripartite motif containing 8	GBF1
	TTC8 - tetratricopeptide repeat domain 8	GALC
	UBE2B - ubiquitin-conjugating enzyme e2b	C5orf24
	TAPT1 - transmembrane anterior posterior transformation 1	BST1
	TMEM107 - transmembrane protein 107	CHRNA1
<hr/>		
Regulation of localization	ABCA1 - atp-binding cassette sub-family a (abc1) member 1	SLC44A1
	BSN - bassoon presynaptic cytomatrix protein	IP6K2
	CAMK2D - calcium/calmodulin-dependent protein kinase ii delta	CAMK2D
	DOC2A - double c2-like domains alpha	SETD1A
	MAPT - microtubule-associated protein tau	CRHR1, WNT3
	NMT1 - n-myristoyltransferase 1	UBTF, FAM171A2, CRHR1
	LGALS3 - lectin galactoside-binding soluble 3	GCH1
	IPO5 - importin 5	MBNL2
	P2RX2 - purinergic receptor p2x ligand-gated ion channel 2	FBRSL1
	RIMS1 - regulating synaptic membrane exocytosis 1	RIMS1
	SEMA4A - sema domain immunoglobulin domain (ig) transmembrane domain (tm) and short cytoplasmic domain (semaphorin) 4a	PMCK, KRTCAP2, GBAP1
	SMAD4 - smad family member 4	MEX3C
	TRIM8 - tripartite motif containing 8	GBF1
	STX7 - syntaxin 7	RPS12
<hr/>		
Regulation of cell communication	ABCA1 - atp-binding cassette sub-family a (abc1) member 1	SLC44A1
	CAMK2D - calcium/calmodulin-dependent protein kinase ii delta	CAMK2D
	BTRC - beta-transducin repeat containing e3 ubiquitin protein ligase	GBF1
	BRD7 - bromodomain containing 7	NOD2
	FGFR3 - fibroblast growth factor receptor 3	TMEM175, GAK
	FAM13A - family with sequence similarity 13 member a	SNCA

GPAM - glycerol-3-phosphate acyltransferase mitochondrial	ADRA2A
MAPT - microtubule-associated protein tau	CRHR1, WNT3
LGALS3 - lectin galactoside-binding soluble 3	GCH1
LEFTY1 - left-right determination factor 1	ITPKB
GRK5 - g protein-coupled receptor kinase 5	INPP5F, BAG3
P2RX2 - purinergic receptor p2x ligand-gated ion channel 2	FBRSL1
NMT1 - n-myristoyltransferase 1	UBTF, FAM171A2, CRHR1
PYDC1 - pyd (pyrin domain) containing 1	SETD1A
RIMS1 - regulating synaptic membrane exocytosis 1	RIMS1
SMAD4 - smad family member 4	MEX3C
TRIM8 - tripartite motif containing 8	GBF1
SH3RF1 - sh3 domain containing ring finger 1	CLCN3
UBE2B - ubiquitin-conjugating enzyme e2b	C5orf24
UBAP2 - ubiquitin associated protein 2	UBAP2

---