Running Title (40 characters): **Fertility in breast cancer survivors**

Title: Pregnancy, fertility concerns, and fertility preservation procedures in French breast cancer survivors in the FEERIC national study (on behalf of the Seintinelles research network).

Manon Mangiardi-Veltin (1)*, Clara Sebbag (2)*, Christine Rousset-Jablonski (3), Isabelle Ray-Coquard (3), Clémentine Berkach (2), Lucie Laot (2), Yuting Wang (1), Inès Abdennebi (2), Julie Labrosse (2), Camille Sautter (2), Aullène Toussaint (2), Laura Sablone (2), Enora Laas (2), Sarah Khallouch (2), Florence Coussy (1), Pietro Santulli (4), Charles Chapron (4), Angelique Bobrie (5), William Jacot (5), Nadir Sella (6), Elise Dumas (6), Claire Sénéchal-Davin (7), Marc Espie (8), Sylvie Giacchetti (8), Lorraine Maitrot (4), Geneviève Plu-Bureau (4), Charles Coutant (9), Julien Guerin (11), Bernard Asselain (12), Pierre Fumoleau (1), Manuel Rodrigues (1), Christine Decanter (13), Audrey Mailliez (14), Lidia Delrieu (6), Seintinelles research network (15), Fabien Rey (2,6), Anne-Sophie Hamy (1,2,6)

¹ Department of Medical Oncology, Institut Curie, Paris, France; University Paris, Paris France

² Department of Surgical Oncology, Institut Curie, Paris, France; University Paris, Paris France

³ Centre Léon Bérard, Lyon, France; RESEARCH on HealthcARE PERFORMANCE (RESHAPE), University Claude Bernard Lyon 1, Lyon, France.

⁴ Department of Obstetrics and Gynecology, Hôpital Cochin, Paris, France; University Paris, Paris France

⁵ Department of Medical Oncology, Institut du cancer de Montpellier, Montpellier, France

⁶ Residual Tumor & Response to Treatment Laboratory, RT2Lab, Translational Research Department, INSERM, U932 Immunity and Cancer, University Paris, Paris, France

⁷ Department of Oncogenetics, Institut Bergonié, Bordeaux, France

⁸ Sénopole, Hôpital Saint Louis, AP-HP, Paris, France; University Paris, Paris France

⁹ Department of Surgical Oncology, Centre Georges-François Leclerc - Unicancer, Dijon, France.

¹⁰ Clinical Research Department, Centre Georges-François Leclerc - Unicancer, Dijon, France

¹¹ Data Factory, Data Office, Institut Curie, 25 rue d'Ulm, 75005 Paris, France

¹² Department of Biostatistics, Institut Curie, Paris, France

¹³ ART department, Centre hospitalier régional Lille, Lille, France

¹⁴ Department of Medical oncology, Centre Oscar Lambert, Lille, France

¹⁵ Seintinelles research network, 40 Rue Rémy Dumoncel, 75014 Paris, guillemette.jacob@seintinelles.com

* Manon Mangiardi and Clara Sebbag contributed equally to this work

Correspondence: fabien.reyal@curie.fr; Tel.: 00 33 144324087, 00 33 615271980

Capsule: The FEERIC prospective study conducted on breast cancer (BC) survivors confirmed the importance of adequate oncofertility counseling including information on contraception and showed frequent spontaneous pregnancies after BC.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Abstract

Objective

To study fertility concerns and oncofertility practices at time of breast cancer (BC) diagnosis.

Design

The FEERIC study (Fertility, Pregnancy, Contraception after BC in France) is a prospective, multicenter study.

Setting

Web-based collaborative research platform *Seintinelles*.

Patients

517 patients with prior BC diagnosis free from relapse and aged 18 to 43 years at inclusion (from 12th March 2018 to 27th June 2019).

Intervention

Baseline online self-administered questionnaires.

Main Outcome Measure

Fertility preservation procedures at BC diagnosis.

Results

Median age at BC diagnosis was 33.6 years and 424 patients (82.0%) received chemotherapy. Overall, 236 (45.6%) patients were offered specialized oncofertility counseling, 124 (24.0%) underwent one or more FP procedures with material preservation (oocytes n=108, 20.9%; embryos n=31, 6.0%; both oocytes and embryos n=13, 2.5%; ovarian cryopreservation n=6, 1.2%) and 78 patients received gonadotropin-releasing hormone agonists (15.1%). With a median follow-up of 26.7 months after the end of treatments, 133 pregnancies (25.7%) had occurred in 85 patients (16.4%), including 20 unplanned pregnancies (15.0%). Most of the pregnancies were spontaneous (n=113, 87.6%), while 16 (12.4%) required medical interventions. Patients who had an unplanned pregnancy were less likely to have received fertility counseling (p=0.02) and contraceptive counseling (p=0.08) at BC diagnosis.

Conclusion

Most of the patients were not offered proper specialized oncofertility counseling at the time of BC diagnosis. Spontaneous pregnancies after BC were very much more frequent than pregnancies resulting

from the use of cryopreserved gametes. Adequate contraceptive counseling seems as important as information about fertility and might prevent unplanned pregnancies.

Key words (3 to 5): Breast cancer, fertility, preservation, pregnancy, contraception

Introduction

Breast cancer (BC) is the most common **cancer among young women** (1), with approximately 6000 women under the age of 45 years diagnosed in France every year (2). Primary BC treatments include surgery, chemotherapy, radiotherapy and targeted therapy as hormonal therapy and trastuzumab. BC treatments can have a direct (chemotherapy-induced ovarian damage) or indirect (fertility decline due to delayed pregnancy project) impact on fertility (3). In the general context of a trend towards women delaying their first pregnancies in the Western world, increasing numbers of women are being diagnosed with BC before they have had their children (4). Furthermore, BC survival rates are increasing (2), and survivorship issues (including motherhood after cancer) are increasingly being encountered in routine practice. In addition, recent studies have confirmed the long-term safety of pregnancy after BC independently of hormonal receptor status (5,6), *HER2* status (7) or the presence of a *BRCA* mutation (8).

Fertility preservation (FP) procedures have developed rapidly over the last decade and are now part of the supportive care offered to patients in routine practice. Available FP methods (9) include (i) the retrieval and storage of frozen material sampled before the start of BC treatment (mature oocyte or embryo cryopreservation, ovarian-tissue cryopreservation, and immature oocyte cryopreservation after *in vitro* maturation (IVM)); and (ii) ovarian protection through treatment with gonadotropin-releasing hormone agonists (GnRHa). In France, the National Institute of Cancer (InCA) considers improving access to oncofertility services a priority (10). Efforts have been made to guarantee universal access, and all FP costs in the context of cancer are covered by the national social security system, a feature rendering French oncofertility practices unique worldwide. Nevertheless, FP counseling and access remain heterogeneous in France, as many patients receive no FP information or are not referred at the optimal time (11).

Even though it is becoming more frequent with the inflow of reassuring data (12), **pregnancy after BC** remains a rare event (13). In a recent meta-analyses, out of 711 BC survivors who received systemic treatments, 14% had a pregnancy, which is on average 40% lower than in the general population (14).

However, few - if any - data is available to determine whether these low pregnancy rates are due to an actual decrease in fertility, to a lower pregnancy attempt rate due to medical reasons, or to the women's reproductive choices. Spontaneous fertility rates after cancer treatment remain difficult to assess. The pregnancy rates achieved through the use of reproductive techniques after cancer treatments have increased with the development of FP (15–17), but recent data suggest that they remain lower after BC than in women without cancer, or women with other types of cancer (18). Furthermore, registry data show that only a few patients ask for fertility restoration (15).

The **FEERIC national study** was launched in France in March 2018 and was designed to compare fertility, pregnancy and contraception outcomes in young BC survivors and in matched cancer-free women. This online study was performed via the Seintinelles network, a collaborative social network set up to provide a source of volunteers for participation in research studies. Data were collected on twice-yearly self-completed forms. The current study focuses on BC survivors only, and is based on the enrollment's questionnaire. The objective was to analyze patients' attitudes and the information provided regarding fertility at the time of diagnosis, rates of referral to fertility specialists and access to FP procedures, and to describe the reproductive history between BC diagnosis and study inclusion.

Materials and methods

Study design

The FEERIC (Fertility, Pregnancy, Contraception after BC in France) study is a prospective study aiming at assessing the impact of BC treatment on fertility, pregnancy and contraception. The data are collected through self-administered online questionnaires via the Seintinelles* research platform (19). Seintinelles is a collaborative social network created in 2012 to accelerate the recruitment of French volunteers for cancer research studies, by connecting researchers with men and women of various ages, social and medical backgrounds, with or without a history of BC, willing to participate in research studies. The scientific board of the Seintinelles approved the FEERIC project in December 2015, and the ethics board of Sud Ouest Outre Mer II approved the project 5th October, 2017.

Patients

Patients were recruited from 12th March 2018 to 27th June 2019. The inclusion criteria were: female patients with a previous diagnosis of localized, relapse-free BC (invasive or *in situ*), aged 18 to 43 years at the time of diagnosis, and who had completed treatment (surgery and/or chemotherapy, and/or radiotherapy) at the time of enrollment, without any time criteria since the date of the end of treatments. Women on ongoing endocrine therapy or trastuzumab treatment were accepted. The exclusion criteria were previous hysterectomy and/or bilateral oophorectomy and/or bilateral salpingectomy. The controls were women aged from 18 to 43 years, free from BC and other cancers, who had not undergone hysterectomy and/or bilateral oophorectomy and/or bilateral salpingectomy.

Enrollment

Cases were recruited by both (i) the Seintinelles network and (ii) eight breast care/oncofertility and gynecology centers: Institut Curie (Paris and Saint Cloud), Hôpital Saint Louis (Paris), Hôpital Cochin (Paris), Centre Léon Berard (Lyon), Centre Oscar Lambret and Centre Hospitalier Régional Universitaire (Lille), Institut Bergonie (Bordeaux). Patients were asked to participate in the study at follow-up consultations, or through the display of invitation flyers and posters in waiting rooms. In addition to the direct proposal to participate during follow-up consultations, 6 centers sent invitations to

participate through mailing to patients (electronic e-mail addresses when available, or by postal mail else) (Institut Curie, Léon Bérard (Lyon), Centre Georges-François Leclerc (Dijon), Hôpital Saint-Louis (Paris), Institut du Cancer de Montpellier (Montpellier)).

Forms

The forms were designed by workgroups including physicians involved in BC, fertility or contraceptive care, and members of the Seintinelles staff. Volunteers matching the inclusion criteria (11 questions) were sent a link to the survey and were asked to complete a baseline form at inclusion and follow-up forms every six months (a total of six forms). The baseline form contained 181 questions relating to demographic data, reproductive and contraceptive history, pregnancy plans, BC history and treatments, if applicable, data for FP procedures (FPPs) if performed, and reproductive life since the end of treatment. The questions regarding patient satisfaction were rated from 0 (unsatisfied) to 10 (extremely satisfied). In total, 28 questions specifically related to FP concerns. The database was hosted by a specialized health data provider (www.ids-assistance.com) in accordance with European General Data Protection Regulation, and was forwarded to the statistical team for analysis in a safe and anonymized manner. The current study concerns baseline characteristics and attitudes to fertility in BC survivors.

Study endpoints

Fertility preservation procedures (FPPs) was defined as a procedure intending to help patients's ability to have children, and included: (i) FPP with material preservation (either oocyte, embryo, or ovarian tissue cryopreservation); (ii) FPP without frozen material, *i.e.* GnRHa during chemotherapy;

Oocyte or embryo retrieval was achieved either after controlled ovarian stimulation (COS) or without ovarian stimulation (*in vitro* maturation, IVM)). Evolutive pregnancies were defined as pregnancies evolutive after the first trimester of pregnancy. Ectopic pregnancies, miscarriage, elective abortion or abortions for medical reasons were not considered evolutive pregnancies.

For each pregnancy declared, the specific question: "was this pregnancy desired?" was asked. A pregnancy was considered desired when the patient answered "yes" and was considered unplanned if the patient answered "no".

Median time to pregnancy was defined as time from first attempt to the occurrence of pregnancy, and median time to evolutive pregnancy was defined as time from first attempt to the occurrence of an evolutive pregnancy.

Statistical analysis

We anticipated to include 251 patients in the FEERIC study. Thanks to a large involvement of the Seintinelles Network, a total of 517 BC patients replied to the questionnaire.

The study population was described in terms of frequencies for qualitative variables, or medians and associated ranges for quantitative variables. To compare continuous variables among different groups, Wilcoxon-Mann-Whitney test was used for groups including less than 30 patients, and for variables displaying multimodal distributions, otherwise, we used student t-test. Association between categorical variables was assessed with the chi-square test, or with the Fisher's exact test if at least one category included less than three patients. In boxplots, lower and upper bars represent the first and third quartile, respectively, the medium bar is the median, and whiskers extend to 1.5 times the inter-quartile range.

For analysis of the association between clinical variables (age, BMI, profession, study level, pregnancy plans, marital status and chemotherapy) and FPP rates, we performed a univariate analysis, with values of $p \leq 0.05$ considered statistically significant. Analyses were performed with R software, version 3.1.2.

Results

Characteristics of the patients at BC diagnosis

We included 517 BC patients in the study (Table 1, figure S1). Median age at BC diagnosis was 33.6 years and median age at study inclusion was 38.0 years. In total, 455 patients (88.0%) had been educated to university level. Most patients were living with a partner at the time of diagnosis ($n=431$, 83.4% (cohabitation $n=245$, 47.4%; married $n=186$, 36.0%)). Sixty-seven patients (13.0%) were current smokers, and 368 patients (71.2%) declared no other comorbid conditions at the time of inclusion. Median BMI was 22.2 kg/m² and 121 patients (23.4%) were overweight or obese. A history of infertility prior to BC diagnosis was recorded for 77 patients (14.9%) and 210 (40.6%) patients had no children at BC diagnosis. Two hundred seventy-one patients (52.4%) had a family history of BC (first degree: $n=115$ (22.2%); second or third degree: $n=156$ (30.2%)).

Most of the patients ($n=419$, 81.0%) had received oncogenetic counseling and 410 (79.3%) had undergone genetic testing. No constitutional mutation was found in 287 (70.0%) (oncogenetic results available at the time of study inclusion, $n=352$), whereas germline mutations were detected in 61 patients (*BRCA1* $n=39$ (9.5%); *BRCA2* mutation $n=22$ (5.4%); others $n=4$ (1.0%)).

BC treatments

Median time from BC diagnosis to inclusion was 34.9 months (Supplemental table 1). The first-line treatment was surgery in 66.3% of the patients ($n=343$), whereas 33.5% underwent first-line systemic treatment ($n=173$) (Fig. 1A). All but one of the patients underwent surgery for BC. Surgical procedures were mastectomy $n=239$ (46.5%) / lumpectomy $n=275$ (53.5%), axillary node dissection $n=252$ (54.2%) / sentinel node biopsy $n=213$ (45.8%). Chemotherapy was administered to 424 patients (82.0%), mostly with anthracyclines followed by taxanes regimens, 172 (40.6%) in a neoadjuvant setting (including 9 with adjuvant treatment, 2.2%) and 252 (59.4%) in an adjuvant setting. In total, 443 (85.7%) patients received radiotherapy, 126 (24.4%) patients received trastuzumab treatment for *HER2*-positive cancers, and 330 (63.8%) received endocrine therapy (Fig. 1A). Figure 1B displays the different care pathways for the entire cohort.

Pregnancy plans, fertility counseling and oncofertility preservation procedures at BC diagnosis

At the time of BC diagnosis, 218 patients (42.2%) declared no wish of pregnancy, 206 stated they could consider a pregnancy in the future (39.8%), and 67 (13.0%) reported that they were trying to become pregnant (Fig. 2A, supplemental table 2). BC was diagnosed during pregnancy in 26 patients (5.0%), and the outcome of the pregnancy was favorable in most cases (full-term pregnancy, $n=17$ (65.4%); elective abortion, $n=3$ (11.5%); abortion for medical reasons, $n=3$ (11.5%); miscarriage, $n=3$ (11.5%)) (Supplemental table 2).

A health care practitioner inquired if the patient would consider future pregnancy after treatment in 363 patients (70.2%) (Fig. 2B) and 72.4% ($n=364$) patients recalled being informed about chemotherapy-induced ovarian toxicity (Fig. 2C). Among them, the degree of satisfaction of this information was high (median score: 8/10) (Fig. 2D). Specialized oncofertility counseling was offered to 236 patients (45.6%) and 124 women (24.0%) underwent at least one FPP with material preservation (Fig. 2E). The distribution of procedures was as follows: oocyte cryopreservation ($n=108$, 87.1%), embryo cryopreservation ($n=31$, 25.0%), ovarian cryopreservation ($n=6$, 4.8%) (Fig. 2F). In addition, GnRHa were used during treatment in 78 patients (15.1%), representing a total of 181 patients with at least one FPP performed (35.0%) (Fig. 2F). Eighty-eight patients (71.0%) had material preservation after controlled ovarian stimulation, while 36 patients had an *in vitro* maturation procedure (29.0%) (Supplemental table 2). Controlled ovarian stimulation was less frequent for FPPs performed in the before surgery ($n=30$, 37.5%) than for FPPs in after surgery ($n=50$, 62.5%) (Fig. 2G).

Age and year of BC diagnosis ($p<0.001$), study level ($p=0.003$), previous children ($p<0.001$), plans for pregnancy at diagnosis ($p<0.001$) and BC treatment with chemotherapy ($p=0.001$) were significantly associated with the likelihood of undergoing a FPP procedure with material preservation (Supplemental table 3). The reasons for which women did not undergo FPP procedures based on cryopreservation techniques was a lack of such procedures being offered ($n=149$, 42.6%), personal choice of the patient ($n=110$, 31.4%), insufficient time available ($n=50$, 14.3%), oncological concerns ($n=32$, 9.1%), and exclusion on the base of fertility conditions ($n=9$, 2.6%) (Fig. 2H). The personal reasons for not

undergoing FPPs cited by patients included not wanting to have a baby ($n=79$, 59.8%), followed by a desire to focus on BC treatment ($n=27$, 20.5%), constraints ($n=10$, 7.6%) or fears ($n=8$, 6.1%) relating to treatment (Fig. 2I).

Gynecological and reproductive history since BC treatment

Amenorrhea, endocrine therapy, couple situation

Of the 424 patients who received CT, 406 (95.8%) reported amenorrhea during treatment (median duration: 11 months), and the duration of amenorrhea increased with age (Fig. 3A; supplemental table 4). Amenorrhea was reversible in most patients ($n=317$, 74.8%) (Supplemental table 4). Endocrine treatment was stopped prematurely (median: 28 months) in 99 of 330 patients (30.0%), either temporarily ($n=77$, 23.3%) or definitively ($n=22$, 6.7%). The main reasons for stopping endocrine treatment were a desire to become pregnant ($n=51$, 51.5%), adverse side effects ($n=35$, 35.4%), or other reasons ($n=16$, 16.2%). Median time of endocrine therapy intake was significantly shorter in patients who stopped due to side effects than in patients who stopped because of a pregnancy desire (19.8 months *versus* 36.4 months respectively, $p<0.001$)

Between the BC diagnosis and the inclusion of the study, 19.3% of patients ($n=100$) changed marital status (Fig. 3B). The proportion of patients who formed new couples (29/86; 33.7%) was higher than the proportion of patients who were initially in a couple but became single (29/402; 7.2%) (Fig. 3B).

Unplanned pregnancies

During a median follow-up of 26.7 months between the end of BC treatment and inclusion in the study, 18 patients (3.5%) experienced an unplanned pregnancy. Patients who had an unplanned pregnancy were significantly younger at BC diagnosis ($p=0.03$), were more likely to have previous pregnancies ($p=0.01$) or previous children at BC diagnosis ($p=0.02$), were less likely to have received fertility counseling ($p=0.04$) and tended to have less contraceptive counseling ($p=0.08$) (Supp table 5). The satisfaction regarding information on fertility and contraception was significantly lower in patients who

had subsequent unplanned pregnancy than in patients who did not ($p=0.021$ and $p<0.001$ respectively) (Supplemental table 5).

Pregnancies after BC

Overall a total of 133 pregnancies occurred in 85 patients (16.4%), with 20 of them (15.0%) being unplanned). Pregnancy onset was spontaneous in 113 cases (87.6%), and medical intervention was required for 16 pregnancies (ART $n=10$ (13.8%); oocyte donation $n=2$; use of cryopreserved material obtained after controlled ovarian stimulation $n=2$; and ovulation induction $n=2$) (Supplemental table 6). One of the two pregnancies achieved with cryopreserved material ended in a live birth; the other pregnancy was still underway at the time of inclusion in the study. The outcomes of pregnancies were as follows: full-term pregnancy, $n=68$ (51.1%); miscarriage, $n=31$ (23.3%); ongoing pregnancy, $n=20$ (15.0%) ectopic pregnancy, $n=6$ (4.5%); elective abortion, $n=6$ (4.5%); abortion for medical reasons, $n=1$ (0.8%), other $n=1$ (0.8%) (Fig. 3E).

The outcome of the pregnancy was significantly different according to whether the pregnancy was unplanned or desired ($p<0.001$), with notably a higher proportion of elective abortions in unplanned pregnancies than in desired pregnancies (30.0% *versus* 0% respectively) (Fig. 3F). There were very few significant obstetrical complications (twins $n=2$, gestational diabetes $n=3$, pre-eclampsia $n=2$, preterm birth $n=2$, risk of preterm delivery $n=1$, post-partum peritonitis $n=1$, CMV seroconversion $n=1$) and no malformations was reported in the infants. The infants were delivered by the vaginal route ($n=55$; 80.9%) or by cesarean section ($n=13$; 19.1%) (Supplemental table 6).

Pregnancies attempts after BC and time to pregnancy.

Overall, since the end of treatment, 127 patients (24.6%) had attempted to get pregnant; 78 of these patients (61.4%) had at least one pregnancy, 52 patients had at least one live birth (40.9%) and 20 (25.6%) were pregnant at inclusion in the study (five patients were pregnant on inclusion and also had one prior live birth during the follow-up period). For the 109 desired pregnancies, both median time to pregnancy (TTP) and median time to evolutive pregnancy were three months, and 68.6% of the patients

had conceived in the 6 months following first attempt. Median TTP was significantly higher in patients aged 40 and older (median TTP: 15 months), than in younger women (median TTP <30 y.o.: 1 month; 30 to 34 y.o.: 3.5 months; 35 to 39: 3 months, $p=0.02$) (Fig. 3G). At study inclusion, 66 BC patients (12.8%) declared that they were still currently trying to get pregnant.

Discussion

This large study on young BC survivors provides important insights into patients' attitudes and expectations towards fertility and pregnancy, as well as oncofertility practices nationwide in France.

First, one third of women declared that they had not been informed of possible fertility damage before treatment, nor were they asked about future pregnancies plans after BC treatment. Academic societies strongly recommend informing patients about oncofertility (10,16,17), but little was known to date about the prevalence of counseling in French BC survivors. American data have shown that 47 to 68% receive information about FP (18–21). In France, one single-center retrospective study showed that only four of 230 (1.7%) BC patients treated between 2000 and 2010 received information before treatment, with another 49 (21.7%) receiving information after gonadotoxic treatments had been initiated (22). Another French national survey (23) on 104 female cancer patients under the age of 45 years diagnosed in 2002 reported that 31 patients (30.0%) had not been informed about the risk of infertility before treatment. Finally, in the VICAN study (24), which included 427 women aged from 18 to 40 years treated in 2010, 291 women (68.1%) reported having received no FP counseling before cancer treatment.

In line with the lack of information on chemotherapy-induced ovarian damage, almost one third of the patients of our cohort reported not undergoing FPPs because this option was not offered ($n=149$, 28.8%), or because too little time was available ($n=50$, 9.7%). Both these obstacles to FPPs are avoidable and represent actionable ways of improving patient care. On the other hand, a significant subset of women ($n=110$, 21.3%) decided not to go through with FPPs despite receiving counseling. FPPs are costly and the low rates of referral in other studies have been linked to the financial burden (26). However, this bias can be ruled out here, because all French citizens are covered by a universal social security system guaranteeing the full reimbursement of FP fees.

Second, the main technique used was oocyte cryopreservation (87.1%), which was much more frequently used than embryo (25.0%) or cortex cryopreservation (4.8%). This distribution seems to be clinically relevant, as, in addition to yielding similar results to embryo freezing for efficacy (9), oocyte cryopreservation has the key advantage of preserving the fertility of the woman, rather than the fertility

of the couple (3). Recent studies have not found divorce or separation rates to be particularly high following BC diagnosis (27,28), nevertheless the preservation of the woman's fertility (oocyte), rather than that of the couple (embryo), is particularly advantageous for single women or women whose relationships are breaking down. Surprisingly, our results showed that nearly half of patients who had FPP with material preservation in the neoadjuvant setting had ovarian stimulation. This result was unexpected, as ovarian stimulation (OS) is not recommended before breast tumor excision in France (3), and there is only limited evidence concerning the safety of OS in the neoadjuvant setting (20,21). Despite the disappointingly low birth rates previously reported after IVM (12,30), approximatively 30% of the patients had IVM. As this rate is expected to increase due to the widespread use of neoadjuvant chemotherapy in young women, results of such techniques should be monitored closely to assess the efficacy of this procedure with long term follow-up. Of note, only 15.1% of patients received GnRHa treatment together with chemotherapy. This low rate may be accounted for by the study period during which most of the patients were treated, as only 117 patients from the cohort (22.6%) were treated in or after 2017. Indeed, the pre-2017 guidelines considered GnRHa to be an experimental technique (31–33), and GnRHa has been considered a valid option for FP in the National Comprehensive Cancer Network (NCCN) and ASCO guidelines only since 2017 and 2018, respectively (34). Two randomized controlled trials (35,36) have since reported significantly lower rates of post-chemotherapy ovarian failure in patients receiving GnRHa during chemotherapy, for both hormone receptor-positive and -negative BC, leading to an update of NCCN guidelines in 2019 (37). All the patients of our study were treated before 2019, and presumably, this treatment will be increasingly used in routine care as an ovary-protecting agent, in addition to other fertility preservation methods.

Third, with a rather short median time between the end of treatment and inclusion, 16.4% of the patients of the cohort had already experienced at least one pregnancy, most of which were spontaneous, and only 2 pregnancies occurred after use of cryopreserved material. The occurrence of unwanted pregnancies (3.9%) was even more frequent than the use of ART (3.1%). These results highlight the statement that contraception is as important as FP in young breast cancer patients (38). In addition, the quality of the fertility and the contraceptive counseling was significantly associated with a lower likelihood of

experiencing unplanned pregnancies. Properly addressing this topic could spare women the burden of undergoing elective abortions or abortions for medical reasons. Finally, for patients who became pregnant after BC, the median time between pregnancy desire and its occurrence was short (three months). Although we cannot formally exclude the possibility that a subset of women currently attempting to get pregnant may remain genuinely infertile due to chemotherapy-induced ovarian damage, the data provided here are reassuring concerning the likelihood of spontaneous pregnancies after BC.

Our study has several strengths. It reports one of the largest recent study on fertility concerns and attitudes at BC diagnosis based on patient reported outcomes. It also provides unprecedented data on unplanned pregnancies, and time between pregnancy attempt and pregnancy occurrence, and provides relative comparison in the number of pregnancies occurring after PF material reuse *versus* spontaneously in a real-world cohort. It also has limitations. Our data could have overestimated the information rate given in the whole young breast cancers population, as most of the FEERIC study women came from high-level socio-professional backgrounds. A similar bias has been reported in other studies recruiting patients via online networks (18,25). Moreover, study inclusions performed by cancer centers or teaching hospitals might not exactly reflect the field reality. Finally, responders to our study may represent a subset of BC patients with a particular interest in fertility following BC, and underrepresent either patients who experienced relapse or those who already have completed their parental project.

In conclusion, our work provides an overview of the fertility and pregnancy concerns in a selected cohort of young French BC survivors. Information about and access to FPPs could be further improved. Healthcare professionals should be aware of the need to systematically offer fertility counseling to patients before BC treatment. In addition, care pathways should be optimized to facilitate rapid access to FPPs if desired by the patient and feasible. Finally, contraception should also be considered a critical topic at BC diagnosis and during the follow-up, as unplanned pregnancies represent a preventable burden for BC survivors. Finally, the longitudinal follow-up of the FEERIC cohort for three years will make it possible to determine whether pregnancy rates differ between BC survivors and matched controls.

Acknowledgments: The FEERIC study was funded by Institut du Cancer InCA, InCA-SHS, grant No. 2016-124. The authors thank all the study participants from the Seintinelles Network, and Lili Sohn who is the sponsor of the study.

Conflicts of interest: none

REFERENCES

1. International Agency for Research on Cancer. Global Cancer Observatory. (2018) Available at: <http://globocan.iarc.fr/Default.aspx> [Accessed August 5, 2018]
2. Institut National du Cancer. Données globales d'épidémiologie des cancers. (2018) Available at: <http://www.e-cancer.fr/Professionnels-de-sante/Les-chiffres-du-cancer-en-France/Epidemiologie-des-cancers/Donnees-globales> [Accessed February 19, 2018]
3. Agence de Biomédecine, Institut National du Cancer. Conséquences des traitements des cancers et préservation de la fertilité - Etat des connaissances et propositions. (2012)
4. Dabrosin C. An overview of pregnancy and fertility issues in breast cancer patients. *Ann Med* (2015) **47**:673–678. doi:10.3109/07853890.2015.1096953
5. Azim HA, Santoro L, Pavlidis N, Gelber S, Kroman N, Azim H, Peccatori FA. Safety of pregnancy following breast cancer diagnosis: a meta-analysis of 14 studies. *Eur J Cancer Oxf Engl 1990* (2011) **47**:74–83. doi:10.1016/j.ejca.2010.09.007
6. Lambertini M, Kroman N, Ameye L, Cordoba O, Pinto A, Benedetti G, Jensen M-B, Gelber S, Del Grande M, Ignatiadis M, et al. Long-term Safety of Pregnancy Following Breast Cancer According to Estrogen Receptor Status. *J Natl Cancer Inst* (2018) **110**:426–429. doi:10.1093/jnci/djx206
7. Lambertini M, Martel S, Campbell C, Guillaume S, Hilbers FS, Schuehly U, Korde L, Azim HA, Di Cosimo S, Tenglin RC, et al. Pregnancies during and after trastuzumab and/or lapatinib in patients with human epidermal growth factor receptor 2-positive early breast cancer: Analysis from the NeoALTTO (BIG 1-06) and ALTTO (BIG 2-06) trials. *Cancer* (2019) **125**:307–316. doi:10.1002/cncr.31784
8. Lambertini M, Ameye L, Hamy A-S, Zingarello A, Poorvu P, Carrasco E, Grinshpun A, Han S, Rousset-Jablonski C, Ferrari A, et al. Pregnancy After Breast Cancer in Patients With Germline BRCA Mutations. *J Clin Oncol* (2020) **38**:JCO.19.02399. doi:10.1200/JCO.19.02399
9. Donnez J, Dolmans M-M. Fertility Preservation in Women. *N Engl J Med* (2017) **377**:1657–1665. doi:10.1056/NEJMr1614676
10. Institut National du Cancer. Plan cancer 2014-2019. (2014) Available at: <http://www.e-cancer.fr/Plan-cancer/Plan-cancer-2014-2019-priorites-et-objectifs> [Accessed March 21, 2018]
11. Dolmans MM. Recent advances in fertility preservation and counseling for female cancer patients. *Expert Rev Anticancer Ther* (2018) **18**:115–120. doi:10.1080/14737140.2018.1415758
12. Agence de la Biomédecine. Rapport médical et scientifique. (2017) Available at: <https://www.agence-biomedecine.fr/annexes/bilan2017/donnees/procreation/01-amp/synthese.htm> [Accessed September 23, 2018]
13. Cobo A, García-Velasco JA, Coello A, Domingo J, Pellicer A, Remohí J. Oocyte vitrification as an efficient option for elective fertility preservation. *Fertil Steril* (2016) **105**:755–764.e8. doi:10.1016/j.fertnstert.2015.11.027
14. Donnez J, Dolmans MM, Diaz C, Pellicer A. Ovarian cortex transplantation: time to move on from experimental studies to open clinical application. *Fertil Steril* (2015) **104**:1097–1098. doi:10.1016/j.fertnstert.2015.08.005

15. Luke B, Brown MB, Missmer SA, Spector LG, Leach RE, Williams M, Koch L, Smith YR, Stern JE, Ball GD, et al. Assisted reproductive technology use and outcomes among women with a history of cancer. *Hum Reprod Oxf Engl* (2016) **31**:183–189. doi:10.1093/humrep/dev288
16. Oktay K, Harvey BE, Loren AW. Fertility Preservation in Patients With Cancer: ASCO Clinical Practice Guideline Update Summary. *J Oncol Pract* (2018) **14**:381–385. doi:10.1200/JOP.18.00160
17. Martinez F, International Society for Fertility Preservation–ESHRE–ASRM Expert Working Group. Update on fertility preservation from the Barcelona International Society for Fertility Preservation-ESHRE-ASRM 2015 expert meeting: indications, results and future perspectives. *Fertil Steril* (2017) **108**:407–415.e11. doi:10.1016/j.fertnstert.2017.05.024
18. Partridge AH, Gelber S, Peppercorn J, Sampson E, Knudsen K, Laufer M, Rosenberg R, Przypyszny M, Rein A, Winer EP. Web-based survey of fertility issues in young women with breast cancer. *J Clin Oncol Off J Am Soc Clin Oncol* (2004) **22**:4174–4183. doi:10.1200/JCO.2004.01.159
19. Quinn GP, Vadaparampil ST, Lee J-H, Jacobsen PB, Bepler G, Lancaster J, Keefe DL, Albrecht TL. Physician referral for fertility preservation in oncology patients: a national study of practice behaviors. *J Clin Oncol Off J Am Soc Clin Oncol* (2009) **27**:5952–5957. doi:10.1200/JCO.2009.23.0250
20. Letourneau JM, Ebbel EE, Katz PP, Katz A, Ai WZ, Chien AJ, Melisko ME, Cedars MI, Rosen MP. Pretreatment fertility counseling and fertility preservation improve quality of life in reproductive age women with cancer. *Cancer* (2012) **118**:1710–1717. doi:10.1002/cncr.26459
21. Ruddy KJ, Gelber SI, Tamimi RM, Ginsburg ES, Schapira L, Come SE, Borges VF, Meyer ME, Partridge AH. Prospective Study of Fertility Concerns and Preservation Strategies in Young Women With Breast Cancer. *J Clin Oncol* (2014) **32**:1151–1156. doi:10.1200/JCO.2013.52.8877
22. Chirol A, Debled M, Fournier M, Papaxanthos A, Hoppe S, Brouste V, Conri V, Bonnefoi H, Von Théobald P, Mathoulin S, et al. [Oncofertility and breast cancer: Where have we come from, where are we going?]. *Bull Cancer (Paris)* (2015) **102**:489–496. doi:10.1016/j.bulcan.2015.04.004
23. Mancini J, Rey D, Préau M, Malavolti L, Moatti J-P. Infertility induced by cancer treatment: inappropriate or no information provided to majority of French survivors of cancer. *Fertil Steril* (2008) **90**:1616–1625. doi:10.1016/j.fertnstert.2007.08.064
24. Jegaden M, Bouhnik A-D, Préau M, Bendiane M-K, Peretti-Watel P, Mancini J, Courbiere B. Fertility status perception, fertility preservation and desire to have children in cancer survivors: French VICAN survey. *Future Sci OA* (2018) **4**:FSO343. doi:10.4155/fsoa-2018-0018
25. Bethlehem J. Selection Bias in Web Surveys. *Int Stat Rev* (2010) **78**:161–188. doi:10.1111/j.1751-5823.2010.00112.x
26. Jones G, Hughes J, Mahmoodi N, Smith E, Skull J, Ledger W. What factors hinder the decision-making process for women with cancer and contemplating fertility preservation treatment? *Hum Reprod Update* (2017) **23**:433–457. doi:10.1093/humupd/dmx009
27. Laitala VS, Saarto T, Einiö EK, Martikainen P, Silventoinen K. Early-stage breast cancer is not associated with the risk of marital dissolution in a large prospective study of women. *Br J Cancer* (2015) **113**:543–547. doi:10.1038/bjc.2015.216
28. Préau M, Bouhnik A-D, Rey D, Mancini J. Le devenir de la relation de couple.57.

29. Hershlag A, Mullin C, Bristow SL. Is Fertility Preservation Feasible and Safe With Neoadjuvant Therapy for Breast Cancer? *J Glob Oncol* (2018)1–5. doi:10.1200/JGO.17.00213
30. Waks AG, Partridge AH. Fertility Preservation in Patients With Breast Cancer: Necessity, Methods, and Safety. *J Natl Compr Canc Netw* (2016) **14**:355–363. doi:10.6004/jnccn.2016.0038
31. Peccatori FA, Azim HA, Orecchia R, Hoekstra HJ, Pavlidis N, Kesic V, Pentheroudakis G, ESMO Guidelines Working Group. Cancer, pregnancy and fertility: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol Off J Eur Soc Med Oncol* (2013) **24 Suppl 6**:vi160-170. doi:10.1093/annonc/mdt199
32. Chen H, Xiao L, Li J, Cui L, Huang W. Adjuvant gonadotropin-releasing hormone analogues for the prevention of chemotherapy-induced premature ovarian failure in premenopausal women. *Cochrane Database Syst Rev* (2019) **3**:CD008018. doi:10.1002/14651858.CD008018.pub3
33. Chen H, Li J, Cui T, Hu L. Adjuvant gonadotropin-releasing hormone analogues for the prevention of chemotherapy induced premature ovarian failure in premenopausal women. *Cochrane Database Syst Rev* (2011)CD008018. doi:10.1002/14651858.CD008018.pub2
34. NCCN Evidence Blocks, TM. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines) - Breast Cancer. (2016) Available at: NCCN.org
35. Moore HCF, Unger JM, Phillips K-A, Boyle F, Hitre E, Porter D, Francis PA, Goldstein LJ, Gomez HL, Vallejos CS, et al. Goserelin for ovarian protection during breast-cancer adjuvant chemotherapy. *N Engl J Med* (2015) **372**:923–932. doi:10.1056/NEJMoa1413204
36. Del Mastro L, Boni L, Michelotti A, Gamucci T, Olmeo N, Gori S, Giordano M, Garrone O, Pronzato P, Bighin C, et al. Effect of the gonadotropin-releasing hormone analogue triptorelin on the occurrence of chemotherapy-induced early menopause in premenopausal women with breast cancer: a randomized trial. *JAMA* (2011) **306**:269–276. doi:10.1001/jama.2011.991
37. NCCN. National Comprehensive Cancer Network: National Comprehensive Cancer Network Guidelines - Breast Cancer. 2019. (2019) Available at: https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf
38. Loibl S, Schmidt A, Gentilini O, Kaufman B, Kuhl C, Denkert C, von Minckwitz G, Parokonnaya A, Stensheim H, Thomssen C, et al. Breast Cancer Diagnosed During Pregnancy: Adapting Recent Advances in Breast Cancer Care for Pregnant Patients. *JAMA Oncol* (2015) **1**:1145–1153. doi:10.1001/jamaoncol.2015.2413

Figure titles and abbreviations (intended for color reproduction on the Web and in print)

Supplemental figure 1: Patient characteristics of the 517 patients at the time of BC diagnosis. A, Median age at BC diagnosis was 33.6 years. B, Median age at study inclusion was 38.0 years. C, Socio-economic level at diagnosis. D, Marital status at diagnosis. E, Smoking status at diagnosis. F, Presence of comorbidities at diagnosis. G, BMI at diagnosis, median was 22.2 kg/m². H, Number of children at diagnosis. I, Family history of BC. J, Presence of a genetic mutation among 410 patients who had oncogenetic screening. Abbreviations: breast cancer (BC); body mass index (BMI); Breast Cancer gene (*BRCA*)

Figure 1: Breast cancer treatments and care pathway. A, Repartition of BC treatments. B, Chart representing the sequential care pathways. Twelve patients were discarded from the sunburst because of discrepancies between the treatment dates mentioned in the study and the pathway. Abbreviations: human epidermal growth factor receptor 2 (Her2), neoadjuvant chemotherapy (NAC).

Figure 2: Oncofertility counseling, fertility preservation procedures and decision making. A, Repartition of patients' pregnancy plans at diagnosis. B, Barplot of medical inquiry about patient's future pregnancy desire. C, Barplot representing the proportion of women who received an information about chemotherapy-induced (CI) ovarian toxicity. D, Patient's satisfaction towards information on chemotherapy-induced (CI) ovarian toxicity (0-to-10 Numerical Rating Scale (NRS), where 0 is strongly dissatisfied and 10 is fully satisfied), out of 364 patients who received information. E, Sankey diagram representing the association between oncofertility counseling proposal and completion of a fertility preservation procedure (FPP). Thirteen patients had a FPP with material preservation by their own initiative in the absence of counselling being offered by oncology team. F, Repartition of FPP with or without material preservation (MP). A combination of methods was used in 39 patients: mostly oocyte cryopreservation plus GnRHa treatment ($n=15$), followed by the cryopreservation of oocytes and embryos ($n=13$). G, Barplots representing ovarian stimulation performance according to chemotherapy setting. H, Repartition of the reasons for not benefiting from a FPP. Main reason was the absence of offer ($n=149$; 42.6%). I, Repartition of the personal reasons for not undergoing a FPP. Abbreviations: chemotherapy-induced (CI); gonadotropin-releasing hormone agonists (GnRHa); controlled ovarian stimulation (COS); in vitro maturation (IVM); neoadjuvant chemotherapy (NAC); chemotherapy (CT); fertility preservation (FP); fertility preservation procedure (FPP); material preservation (MP).

Figure 3: Occurrence and outcome of pregnancies after BC treatment. A, Association between age and duration of amenorrhea. Median duration of amenorrhea is as follows: <30 y.o.: 8 months; [30-35[: 9 months; [35-40[: 12 months; ≥ 40 : 15 months. B, Sankey diagram of the evolution of marital status between diagnosis and study inclusion. C, Association between the information received on fertility and the occurrence of an unplanned pregnancy. D, Association between the information received on contraception and the occurrence of an unplanned pregnancy. E, Outcomes of the 133 pregnancies achieved after breast cancer in the FEERIC cohort. F, Barplots of pregnancy outcomes depending on the status unplanned ($n=20$) or desired ($n=109$) and the pregnancy occurrence (spontaneous $n=113$ or use of assisted reproductive technology $n=16$) out of the 129 pregnancies for which the unplanned/desired status was known. G, Time from first attempt to pregnancy occurrence in months according to age classes and pregnancy achievement method out of the 102 patients for which the attempt duration was known. Abbreviations: years old (y.o.); assisted reproductive technology (ART); breast cancer (BC)

Table titles and abbreviations

Variable	Class	All
<i>n</i> =		517
Age at BC diagnosis (years)		33.6 [30.0, 37.1]
Age at study inclusion (years)		38.0 [34.0, 41.0]
Age (by class)	<30	38 (7.4)
	[30-35)	132 (25.5)
	[35-40)	215 (41.6)
	>=40	132 (25.5)
Study level	High school or lower	62 (12.0)
	University	455 (88.0)
Profession (class)	Intermediate	221 (42.7)
	Low or unemployed	48 (9.3)
	Superior	248 (48.0)
Marital status (at BC diagnosis)	Couple	431 (83.4)
	<i>Relationship</i>	
	Cohabitation	245 (56.8)
	Married	186 (43.2)
	Single	86 (16.6)
BMI (continuous)		22.2 [20.4, 24.6]
BMI (4 classes)	Underweight	30 (5.8)
	Normal weight	366 (70.8)
	Pre-obesity	84 (16.2)
	Obesity	37 (7.2)
Smoking status	Current	67 (13.0)
	Former	207 (40.0)
	Never	243 (47.0)
Comorbidity	No	368 (71.2)
	Yes	149 (28.8)
Co-medications	No	401 (77.6)
	Yes	116 (22.4)
Infertility history	No	440 (85.1)
	Yes	77 (14.9)
Previous pregnancy	No	150 (29.0)
	Yes	367 (71.0)
Number of children (3 classes)	0	210 (40.6)
	1	119 (23.0)
	More than 1	188 (36.4)
Previous births	No	210 (40.6)
	Yes	307 (59.4)
	<i>Breastfeeding history</i>	
	No	81 (26.4)
	Yes	226 (73.6)
	<i>Breastfeeding duration</i> (months)	4.0 [2.25, 9.00]
Familial history of BC	No	246 (47.6)

	At least 1 first-degree relative	115 (22.2)
	At least 1 second-degree relative	156 (30.2)
Oncogenetic counseling	No	98 (19.0)
	Yes	419 (81.0)
Genetic testing	No	107 (20.7)
	Yes	410 (79.3)
	<i>Oncogenetic results</i>	
	No mutation	287 (70.0)
	BRCA1	39 (9.5)
	BRCA2	22 (5.4)
	Others	4 (1.0)
	Results awaited / not known	58 (14.1)

Table 1: Patient's characteristics

Abbreviations: breast cancer (BC); body mass index (BMI); breast cancer gene (*BRCA*). Missing data: Age at BC diagnosis, n=3.

Variable	Class	All
<i>n</i> =		517
Time from BC diagnosis to inclusion (months)		34.9 [18.7, 64.1]
Primary treatment	Surgery	343 (66.3)
	Neoadjuvant treatment	173 (33.5)
	Chemotherapy	172 (33.3)
	Trastuzumab	43 (8.3)
	Endocrine therapy	2 (0.4)
	Radiotherapy	12 (2.3)
BC surgery (type)	Lumpectomy	275 (53.5)
	Mastectomy	239 (46.5)
Axillary surgery (type)	Axillary node dissection	252 (54.2)
	Sentinel node biopsy	213 (45.8)
Chemotherapy	No	93 (18.0)
	Yes	424 (82.0)
	Chemotherapy regimen	
	Anthra-taxans	370 (87.3)
	Others	54 (12.7)
	Chemotherapy setting	
	NAC	163 (38.4)
	NAC & adjuvant CT	9 (2.1)
	Adjuvant CT	252 (59.4)
Radiotherapy	No	74 (14.3)
	Yes	443 (85.7)
Anti-HER2 therapy	No	391 (75.6)
	Yes	126 (24.4)
Endocrine therapy	No	187 (36.2)
	Yes	330 (63.8)
	<i>At least one intake</i>	317 (96.1)

Supplemental table 1: BC treatments.

All but one patient had surgery. Abbreviations: breast cancer (BC); human epidermal growth factor receptor 2 (Her2). Missing data: Time from BC diagnosis to inclusion, n=3; BC surgery, n=3; Axillary surgery, n=52

Variable	Class	All
n=		517
Pregnancy plans at BC diagnosis	Attempting pregnancy	67 (13.0)
	Future pregnancy desire	206 (39.8)
	No	218 (42.2)
	Pregnant at diagnosis	26 (5.0)
	Outcomes of pregnancy	
	Full-term pregnancy	17 (65.4)
	Elective abortion	3 (11.5)
	Abortion for medical reasons	3 (11.5)
	Miscarriage	3 (11.5)
Future pregnancy inquiry	No	154 (29.8)
(by any health care provider)	Yes	363 (70.2)
Information about chemotherapy induced ovarian toxicity	No	139 (27.6)
<i>out of 503</i>	Yes	364 (72.4)
Quality of information (from 1 to 10) (median)		8 [6 - 10]
Oncofertility counselling offered	No	281 (54.4)
	Yes	236 (45.6)
Oncofertility counselling performed	No	312 (60.3)
	Yes	205 (39.7)
FPP with material preservation	Material preserved	124 (24.0)
	<i>Oocyte cryopreservation</i>	108
	<i>Embryo cryopreservation</i>	31
	<i>Ovarian cortex cryopreservation</i>	6
	Ovarian stimulation	
	No (IVM)	36 (29.0)
	Yes	88 (71.0)
FPP without material preservation	LHRH agonists during CT	78 (15.1)
Any FPP	No	336 (65.0)
	Yes	181 (35.0)
FPP timing	Before surgery	64 (48.9)
	After surgery	67 (51.1)

Supplemental table 2: Fertility status, oncofertility counseling and procedures at BC diagnosis

Nine patients underwent FPPs despite not having chemotherapy. Abbreviations: gonadotropin-releasing hormone agonists (GnRHa); chemotherapy (CT); fertility preservation procedure (FPP); *in vitro* maturation (IVM)

Variable	Class	No	Yes	<i>p</i>
<i>n</i> = 517		393	124	
Age at BC diagnosis		34.0 (4.7)	31.8 (3.9)	<0.001
Age at BC diagnosis (class)	<30	19 (4.8)	19 (15.3)	<0.001
	[30-35[	80 (20.4)	52 (41.9)	
	[35-40[	170 (43.3)	45 (36.3)	
	>=40	124 (31.6)	8 (6.5)	
Year BC diagnosis	<=2010	50 (12.8)	1 (0.8)	<0.001
	2010-2015	195 (50.0)	60 (48.4)	
	>2015	145 (37.2)	63 (50.8)	
Study level	Bachelor or lower	57 (14.5)	5 (4.0)	0.003
	University	336 (85.5)	119 (96.0)	
Profession (class)	Intermediate	178 (45.3)	43 (34.7)	0.105
	Low CSP or unemployed	36 (9.2)	12 (9.7)	
	Superior	179 (45.5)	69 (55.6)	
BMI (4 classes)	Underweight	23 (5.9)	7 (5.6)	0.902
	Normal weight	275 (70.0)	91 (73.4)	
	Pre-obesity	66 (16.8)	18 (14.5)	
	Obesity	29 (7.4)	8 (6.5)	
Smoking status	No	341 (86.8)	109 (87.9)	0.861
	Yes	52 (13.2)	15 (12.1)	
BRCA mutation genes		43 (8.3)	22 (4.3)	
	BRCA1	24 (55.8)	15 (68.2)	0.358
	BRCA2	17 (39.5)	5 (22.7)	
	Others	2 (4.7)	2 (9.1)	
Infertility history	No	334 (85.0)	106 (85.5)	0.99
	Yes	59 (15.0)	18 (14.5)	
Previous children	No	120 (30.5)	90 (72.6)	<0.001
	Yes	273 (69.5)	34 (27.4)	
Marital status (BC diagnosis)	Single	60 (15.3)	26 (21.0)	0.178
	Coupled up	333 (84.7)	98 (79.0)	
Attitude towards pregnancy desire	Attempted pregnancy	44 (11.2)	23 (18.5)	<0.001
	Future pregnancy desire	128 (32.6)	78 (62.9)	
	No	196 (49.9)	22 (17.7)	
	Pregnant at diagnosis	25 (6.4)	1 (0.8)	
Chemotherapy	No	84 (21.4)	9 (7.3)	0.001
	Yes	309 (78.6)	115 (92.7)	

Supplemental table 3: Association between clinical characteristics and the performance of fertility preservation procedures

Abbreviations: breast cancer (BC); body mass index (BMI); breast cancer gene (BRCA). Missing data: year of BC diagnosis, n=3

Variable	Class	All
<i>n</i> =		517
Time from end of treatment to inclusion (months)		35.8 (32.4)
Amenorrhea during CT		424 (82.0)
	No	18 (4.2)
	Yes	406 (95.8)
	<i>Amenorrhea duration (months)</i>	11.0 [6.0, 18.0]
	<i>Amenorrhea reversibility</i>	
	No	107 (25.2)
	Yes	317 (74.8)
Endocrine therapy early discontinuation	Yes	99 (30.0)
	<i>Definitively</i>	22 (6.9)
	<i>Temporarily</i>	77 (24.3)
Marital status (current)	Cohabitation	206 (39.8)
	Married	225 (43.5)
	Single	86 (16.6)

Supplemental table 4: Amenorrhea, endocrine therapy, and marital status at study inclusion
Abbreviations: breast cancer (BC); chemotherapy (CT)

Variable	Class	No unplanned pregnancy	At least one unplanned pregnancy	<i>p</i>
<i>n</i> = 517		499	18	
Age at BC diagnosis		33.5 (4.6)	31.1 (4.1)	0.029
Study level	Bachelor or lower University	58 (11.6) 441 (88.4)	4 (22.2) 14 (77.8)	0.322
Profession (class)	Intermediate Low CSP or unemployed Superior	215 (43.1) 45 (9.0) 239 (47.9)	6 (33.3) 3 (16.7) 9 (50.0)	0.473
Marital status (BC diagnosis)	Cohabitation Married Single	238 (47.7) 181 (36.3) 80 (16.0)	7 (38.9) 5 (27.8) 6 (33.3)	0.153
Marital status (current)	Cohabitation Married Single	197 (39.5) 217 (43.5) 85 (17.0)	9 (50.0) 8 (44.4) 1 (5.6)	0.394
BMI		23.3 (4.8)	23.2 (4.9)	0.896
Smoking status	Current Former Never	66 (13.2) 198 (39.7) 235 (47.1)	1 (5.6) 9 (50.0) 8 (44.4)	0.528
Infertility history	No Yes	426 (85.4) 73 (14.6)	14 (77.8) 4 (22.2)	0.581
Previous pregnancy (at BC diagnosis)	No Yes	150 (30.1) 349 (69.9)	0 (0.0) 18 (100.0)	0.013
Previous children	No Yes	208 (41.7) 291 (58.3)	2 (11.1) 16 (88.9)	0.019
Attitude towards pregnancy desire	Attempted pregnancy Future pregnancy desire No Pregnant at diagnosis	66 (13.2) 197 (39.5) 211 (42.3) 25 (5.0)	1 (5.6) 9 (50.0) 7 (38.9) 1 (5.6)	0.722
Fertility counselling out of 503	No Yes <i>Quality of information on fertility (mean)</i>	130 (26.7) 356 (73.3) 8 [6- 10]	9 (52.9) 8 (47.1) 5.50 [0- 7]	0.036 0.021
Contraceptive counselling	No Yes <i>Quality of information on contraception (mean)</i>	164 (32.9) 335 (67.1) 9 [6 - 10]	10 (55.6) 8 (44.4) 5 [3.25 - 8]	0.081 <0.001
Fertility preservation procedure	No Yes	321 (64.3) 178 (35.7)	15 (83.3) 3 (16.7)	0.159

Supplemental table 5: Factors associated with the occurrence of an unplanned pregnancy.
Abbreviations: breast cancer (BC); body mass index (BMI)

Variable	Class	All
Pregnancies (in 85 patients)		133
Pregnancy planning <i>out of 129</i>	Unplanned (in 18 patients)	20 (15.5)
	Desired	109 (84.5)
Pregnancy onset <i>out of 129</i>	Spontaneous	113 (87.6)
	Medical intervention	16 (12.4)
	<i>ART</i>	10 (62.5)
	<i>Oocyte donation</i>	2 (12.5)
	<i>Cryopreserved material reuse</i>	2 (12.5)
	<i>Ovulation induction</i>	2 (12.5)
Pregnancy outcome	Ectopic pregnancy	6 (4.5)
	Induced abortion	6 (4.5)
	Medical abortion	1 (0.8)
	Miscarriage	31 (23.3)
	Ongoing pregnancy	20 (15.0)
	Other	1 (0.8)
	Full-term pregnancy	68 (51.1)
	<i>Delivery Route</i>	
	Caesarean section	13 (19.1)
	Vaginal birth	55 (80.9)
Obstetrical conditions <i>out of 12</i>	Twins	2 (1.6)
	Pre-eclampsia	2 (1.6)
	Preterm birth	2 (1.6)
	Gestational diabetes	3 (2.4)
	Risk of preterm birth	1 (0.8)
	Post-partum peritonitis	1 (0.8)
	CMV seroconversion	1 (0.8)

Supplemental table 6: Occurrence and outcome of pregnancies after BC treatment.

Abbreviations: assisted reproductive technology (ART). Missing data: Pregnancy planning, n=4, Pregnancy onset, n=4

A

Primary treatment

Chemotherapy

Trastuzumab

Breast surgery type

Chemotherapy regimen

Radiotherapy

Axillary surgery type

Chemotherapy setting

Endocrine therapy

B