


# Figure S1

Absolute number of SARS-CoV-2 positive cases and overall incidence per 1000 pregnancies per month in Denmark between March 1 and October 31, 2020.



## Supporting Information Table S1

Procedural and diagnostic codes used to identify the overall study population in the Danish National Patient Register (DNPR).

First, pregnancies were identified in the DNPR if they had a contact with a primary (A) or secondary (B) diagnosis of any of the following codes from March 1 to October 31, 2020 (N = 84 862).

Code	Description
DZ32*	Confirmed pregnancy diagnoses
DZ34*–DZ35*	Pregnancy classification codes
DZ36*	Prenatal screening codes
DO0*–DO07*	Abortion diagnoses
DO200*– DO209*	Bleeding in pregnancy before gestational week 22 diagnoses
DO80*–DO84*	Delivery diagnoses
UXUD86*	Fetal and placental ultrasound/Doppler procedures

\* indicates that all subdiagnoses/procedures within the overarching code are also included.

Secondly, only pregnancies that could be classified according to the following hierarchy were included in the final register-based pregnancy population. Otherwise, they were excluded.

Hierarchy of classification:

Hierarchy	Classification	Code	Conditions
1	Live birth	DZ38*, DO80*–DO84*	Active personal identification number of the child in the Civil Registration System <sup>a</sup>
2	Stillbirths/IUFD	DP95*, DO365*	Gestational week $\geq 22$ –45
3	Miscarriage	DO00*–DO03*, DO08*, DO365*	Gestational week <22
4	Termination	DO04–DO07*, BKHD*– BKHD5*, KLCH*	Gestational week <23
5	Pregnant at end of follow-up	UXUD86A–UXUD86B, UXUD86N	Gestational age <45 weeks at end of follow-up

<sup>a</sup>Except births with a record of miscarriage before gestational age (GA) 22 or a termination of pregnancy before GA 23.

Criteria for exclusion from the pregnancy population were:

- I. Liveborn child with a missing gestational age (n = 406)

- II. Women without a registration of a birth, miscarriage, termination of pregnancy, or ultrasound during follow-up (n = 2965)
- III. Women without a registration of a birth, miscarriage, or termination of pregnancy AND a pregnancy of more than 45 gestational weeks at the end of follow-up (n = 187).

## Supporting Information Table S2

### Procedure and diagnostic codes used to identify basic characteristics and outcomes in the Danish National Patient Register (DNPR).

Variable	Code
BMI	RDA26* RDA27*
Smoking	RDA25*
Pre_existing_astma	DO995A DJ45*
Hypertension	DO100 DI10*
Diabetes	DO240 DO241 DO245
Parity	DZ340 DZ348A DZ348B
Multiple pregnancy	DO30* (the variable also gets the value 1 if there are twins in the population of children)
Eclampsia	DO14 DO15
Gestational Diabtes	DO244
Thromboembolic_event	DO882 DO223 DO225
Pneumonia	DJ12* DJ18*
Intensive_care	NABB NABE
Death	90
Termination of pregnancy	DO04*-DO07* BKHD4 KLCH03 KLCH00
Miscarriages	DO364 DO00*-DO03*
Ectopic pregnancy	DO00
Total deliveries	D080*-D084*
Vaginal deliveries	DO80* DO81* DO83* DO84* (not DO842)  If codes for emergency_cd OR

	elective_cd are recorded the variable gets the value 0  Restricted to deliveries
Operative vaginal delivery	KMAE00 KMAE03 KMAF00  Restricted to deliveries
Emergency Cesarean section	KMCA10A KMCA10D KMCA10E  Restricted to deliveries
Elective Cesarean section	KMCA10B  Restricted to deliveries
Induction	KMAC00* KMAC96A* BKHD20* BKHD20A* BKHD21*  Restricted to deliveries
Total number of children	Derived variable of live birth + IUFD/stillbirths
Live birth	pnr!=. and DZ38
IUFD / stillborn	DO364 AND DP95* Restricted to children
Neonatal_death	D_status D_doddata D_foddata Restricted to live born children
Apgar score 5 min.	DVA00-DVA10 DVAXX Restricted to live born children
Arterial pH	RDA46* Restricted to live born children
Venous pH	RDA47* Restricted to live born children

\* indicates that all subdiagnoses/procedures within the overarching code are also included.

### Supporting Information Table S3

#### **Definition of pre-existing medical problems and previous pregnancy-related complications in DCOD.**

*Pre-existing medical problems* comprised cardiac, renal, endocrine, psychiatric, hematologic, and autoimmune diseases; cancer; and HIV.

*Previous pregnancy-related complications* included thromboembolic events, eclampsia, habitual abortion, preterm birth, mid-trimester loss, neonatal death, stillbirth, baby with major congenital abnormality, small/large for gestational age infant, infant requiring intensive care, puerperal psychosis, placenta praevia, gestational diabetes, placental abruption, postpartum haemorrhage requiring transfusion, surgical procedure during pregnancy, hyperemesis and dehydration requiring admission, ovarian hyperstimulation syndrome or severe infection.

**Table S4: Additional basic characteristics of SARS-CoV-2 infected women overall and according to severity of infection and all pregnant women in Denmark between March 1 and October 31, 2020.**

		SARS-CoV-2 infected pregnant women identified in DCOD	SARS-CoV-2 infected pregnant women / pregnancies identified in the national registers	Non-infected pregnancies identified in the national registers	SARS-CoV-2 infected vs. non-infected pregnancies		SARS-CoV-2 infected women identified in DCOD		
		N = 418	N = 420	N = 82 262	DCOD SARS-CoV-2 cases OR <sub>D</sub> (95% CI)	Register SARS-CoV-2 cases OR <sub>R</sub> (95% CI)	Women requiring admission to hospital due to COVID-19 symptoms N = 23	Women with uncomplicated SARS-CoV-2 infection* N = 395	Women requiring hospital admission due to COVID-19 symptoms vs. women with uncomplicated SARS-CoV-2 infection* OR <sub>DCOD</sub> (95% CI)
<b>Basic characteristics</b>									
Hypertension		< 3	<5	719 (0.9)	NA	NA	NA	NA	
Diabetes		3 (0.7)	<5	492 (0.6)	1.20 (0.39–3.75)	NA	NA	NA	
Women in employment		314 (79.3)	NA	NA			16 (72.7)	298 (79.7)	0.68 (0.26–1.80)
	Unknown	22 (5.3)					1 (4.3)	21 (5.3)	
<b>COVID-19 characteristics</b>									
Test type	PCR or antigen test	399 (97.3)	NA	NR			23 (100)	376 (97.2)	1.03 (1.01–1.05)
	IgG in serum	11 (2.7)	NA	NR			0 (0)	11 (2.8)	NR
	Unknown	8 (1.9)					0 (0)	8 (2.0)	NR
Administration of anti-virals		< 3	NA	NR			< 3	0 (0)	NA
	Unknown	7 (1.7)					1 (4.3)	6 (1.5)	
<b>Pregnancy characteristics</b>									
Previous pregnancies	Multigravida	275 (66.9)	NA	NA			17 (73.9)	258 (66.5)	1.43 (0.55–3.71)
	Unknown	7 (1.7)						7 (1.8)	
Previous pregnancy related complications among multigravid women <sup>†</sup>		60 (21.9)	NA	NA			3 (17.6)	57 (22.2)	0.75 (0.21–2.71)
	Unknown	1 (0.4)					0 (0)	1 (0.4)	

Data are presented as count (%) or median (interquartile range)

OR<sub>D</sub> was estimated using 2 x 2 contingency tables. OR<sub>R</sub> was estimated with univariate logistic regression.

OR<sub>DCOD</sub> was estimated using logistic regression.

\* Uncomplicated COVID-19 included cases where women did not require admission to hospital due to COVID-19 symptoms. Includes women in home isolation or women admitted for obstetric reason within 14 days of first positive COVID-19 test or symptoms.

† Previous pregnancy-related complications include thromboembolic event, eclampsia, habitual abortion, preterm birth and mid trimester loss, neonatal death, stillbirth, baby with major congenital abnormality, SGA, LGA, infant requiring intensive care, puerperal psychosis, placenta praevia, GDM, placental abruption, PPH requiring transfusion, surgical procedure during pregnancy, hyperemesis or dehydration requiring admission, OHSS or severe infection.

BMI Body mass index; DCOD Danish COVID-19 in pregnancy database; GA Gestational age; NR Not relevant; NA Not available