

Table S1. Pharmacokinetic parameters for metabolite RDV-TP from single-dose experiments from Humeniuk et al. 2020 (main text ref. 15), with different RDV dose size and infusion durations, compared to our pharmacokinetic model prediction (equation (1), main text). Pharmacokinetic parameters are given in terms of mean and %CV, with the exception of the half-life $t_{1/2}$ for which uncertainty is explained via IQR. Model predictions extending beyond the %CV/IQR are italicized.

Dosing	Pharmacokinetic parameter (units)	Median (%CV or IQR)	Model prediction
75 mg infused over 2 hours	C_{max} (μM)	2.5 (16.2)	2.5
	C_{24} (μM)	2.2 (23.3)	1.67
	AUC_{∞} ($\text{h}\cdot\mu\text{M}$)	176 (31.1)	184
	$t_{1/2}$ (h)	42.7 (30.6-47.4)	<i>48.0</i>
150 mg infused over 30 minutes	C_{max} (μM)	6.0 (46.1)	8.79
	C_{24} (μM)	3.7 (40.9)	4.0
	AUC_{∞} ($\text{h}\cdot\mu\text{M}$)	297 (28.3)	381
	$t_{1/2}$ (h)	36.0 (27.3-41.5)	34.4
75 mg infused over 2 hours	C_{max} (μM)	5.9 (37.7)	5.5
	C_{24} (μM)	3.3 (55.7)	2.0
	AUC_{∞} ($\text{h}\cdot\mu\text{M}$)	394 (49.9)	291
	$t_{1/2}$ (h)	49.0 (26.6-69.5)	39.4