Knowledge and use of lactational amenorrhea as a family planning method among adolescent mothers in Uganda: a secondary analysis of Demographic and Health Surveys between 2006 and 2016

Catherine Birabwa^{1,4,*}, Pamela Bakkabulindi^{1,2+}, Solomon T Wafula¹, Peter Waiswa^{1,3}, Lenka Benova⁴

¹School of Public Health, Makerere University College of Health Sciences, Kampala, Uganda.

²Health Support Initiatives, P.O Box 10364, Kampala, Uganda.

³Division of Global Health, Department of Public Health Sciences, Karolinska Institutet, Stockholm, Sweden.

⁴Institute of Tropical Medicine, Antwerp, Belgium.

* Correspondence to

Catherine Birabwa

Cathybirabwa3@gmail.com

+Contributed equally to this paper

ABSTRACT

Objective: To assess the level of knowledge and use of lactational amenorrhea method (LAM) among adolescents in Uganda between 2006 and 2016 using nationally representative data from Demographic and Health Surveys (DHS).

Design: Cross-sectional design involving analysis of three DHS (2006, 2011, and 2016) in Uganda.

Setting: The data was collected in Uganda. The DHS are nationally representative surveys on a wide range of indicators including contraception knowledge and use.

Participants: A total of 8,250 adolescents (15-19 years) and 7,110 young women (20-24 years) were included.

Primary outcome measure: Use of LAM among adolescents and young women with a livebirth within six months before each survey.

Results: In 2016, less than 1% of eligible adolescents correctly used LAM and 56% were passively benefitting from LAM. The median duration of postpartum amenorrhea (PPA) among adolescents in 2016 was 6.9 months, declining from 8.3 months in 2006. Compared to adolescents, eligible young women had higher knowledge of LAM and higher median PPA duration in 2016. The percentage of eligible adolescents who met the LAM criteria irrespective of whether they reported LAM use (protected by LAM) decreased from 76% in 2006 to 57% in 2016. More than 50% of eligible adolescents were aware of LAM in 2016, increasing from 6% in 2006, potentially in part due to change in survey question.

Conclusion: Despite increasing awareness of LAM, reported and correct use of LAM was low among adolescents who could benefit from this method in Uganda, and declining over time. Support for adolescents to harness the benefits of correct LAM use should be increased. Additional research is needed to better understand the dynamics of LAM use in adolescents, including the transition to use of other modern contraceptive methods.

Key words: Lactational Amenorrhea Method, adolescents, Uganda, contraceptive methods, breastfeeding, birth spacing

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Strengths and limitations of this study

- Lactational amenorrhea method has the potential to promote healthy timing and spacing of pregnancies and to reduce repeat adolescent births, yet, its use among Ugandan adolescents has not been examined.
- This study provides useful insights into the behavior of adolescents mothers which can be targeted to improve their reproductive knowledge and wellbeing.
- The study used data from three standardised nationally representative surveys, thus findings are generalizable and comparable over time and across age groups.
- Assessment of knowledge of LAM was affected by a change in the phrasing of the question in 2016, thus providing limited comparability with previous surveys.
- The measurement of LAM use was based on self-report.

INTRODUCTION

Reducing adolescent pregnancy rates is an important public health issue in sub-Saharan Africa (SSA), where high fertility and unmet need for family planning (FP) persist. Estimates show that 21 million adolescents aged 15-19 years in developing countries become pregnant every year (Neal et al., 2012, World Health Organization, 2020). Of these, 2.5 million are younger adolescents 12-15 years and at least 10 million are unintended pregnancies (Neal et al., 2012, World Health Organization, 2020). The prevalence of adolescent first births in SSA is estimated at 50% (Neal et al., 2012). As of 2019, about 218 million women in low- and middle-income countries (LMICs) had an unmet need for FP (Sully et al., 2020). In SSA, 25% of women aged 15-49 years are estimated to have an unmet need for FP (Ahinkorah et al., 2020). In 2019, 43% of adolescents in LMICs were estimated to have an unmet need for FP (Sully et al., 2020). Tackling the unmet need for FP among adolescents is critical especially in LMICs because it has significant implications on their reproductive health and wellbeing (Figaroa et al., 2020). Adolescent childbearing has been associated with multiple adverse health outcomes (Amongin et al., 2020a, Burke et al., 2018), and the literature also shows several challenges of the motherhood transition period for adolescents, which leave them disempowered with feelings of shame and embarrassment (Erfina et al., 2019, Mangeli et al., 2017, Watts et al., 2015). Specifically, concern has been raised about the unmet need for FP among postpartum women (Abraha et al., 2018). The acceptability and correct use of various FP methods among women during this period is affected by various socio-cultural and structural factors (Coomson and Manu, 2019, Joshi et al., 2020).

Lactational Amenorrhea Method (LAM) is a recognized modern contraceptive method where postpartum amenorrheic women depend on the contraceptive effect of breastfeeding within the first six months after childbirth (Van der Wijden and Manion, 2015). LAM provides 98% protection against pregnancy if 1) the woman's menstrual period has not returned since childbirth, 2) the baby is fully or nearly fully breastfed, and 3) the baby is less than six months old (Van der Wijden and Manion, 2015, World Health Organization, 2018). The correct use of LAM provides an effective and affordable contraceptive option for breastfeeding women, while also providing an opportunity for linkage and transition of mothers to other FP methods and services. Furthermore, LAM use does not require replenishment of contraceptive supplies or a healthcare provider after appropriate LAM counselling is given. Therefore, LAM can play an important role in preventing unwanted pregnancies during the postpartum period and consequently maternal deaths (Alege et al., 2016). However, its effectiveness may be undermined by factors or practices that affect the three criteria, especially breastfeeding. Evidence shows low breastfeeding rates (Rahman et al., 2020, Yilmaz et al., 2016) and a higher likelihood of exclusive breastfeeding (EBF) discontinuation before infants reach 6 months among adolescent mothers (Jara-Palacios et al., 2015). A study of Nigerian adolescents found lower rates of early breastfeeding initiation and EBF compared to women 20-24 years (Benova et al., 2020). Education level, mode of delivery, antenatal care attendance, and postnatal breastfeeding counselling are some of the factors influencing breastfeeding practices among adolescents (Benova et al., 2020, Yilmaz et al., 2016).

Uganda has one of the highest total fertility rates (5.4 children per woman), maternal mortality rates of 336/100,000 livebirths (Uganda Bureau of Statistics and ICF, 2017), and the youngest population with 78% under 30 years (Ejang, 2020) in the world. Adolescent pregnancy rate is 25% with about 360,000 teenage pregnancies occurring annually (Ministry of Health, 2016, Uganda Bureau of Statistics and ICF, 2017). Evidence shows that over 40% of pregnancies among Ugandan adolescents are unintended (Nalwadda et al., 2019) and there is an increasing occurrence of repeat adolescent births (<20 years) estimated at 56%, with a prevalence of short-birth intervals of 5.4% (Amongin et al., 2020b). Unmet need for child spacing among women with adolescent births or repeat adolescent births has also been reported (Amongin et al., 2020a, Amongin et al., 2020b). Lack of knowledge and access to general health services and youth-friendly FP care, in particular, are some of the key challenges affecting adolescent health in Uganda (Atuyambe et al., 2015). Evidence on use of LAM in adolescents in LMICs is still scarce. There is a need to better understand the duration, knowledge, and correct use of LAM as a postpartum FP method among adolescents (Figaroa et al., 2020).

The main objective of this study was to assess the knowledge and use of LAM among adolescents in Uganda between 2006 and 2016 using nationally representative data from Demographic and Health Surveys (DHS).

METHODOLOGY

Data

This study is a secondary analysis of the cross-sectional data collected during Uganda DHS in 2006, 2011, and 2016. These are nationally representative household surveys carried out every five years. The DHS employs a stratified two-stage cluster sampling design to select participating households. All women of reproductive age (15-49 years) who are either permanent residents of the selected households or visitors who stayed in the household the night before the survey are eligible to be interviewed. The women interviewed provide, among others, self-reported information on their education, marital status, reproductive history, family planning use, maternal health-seeking for recent live births, and breastfeeding practices. Data were collected by trained data collectors using pre-tested tools.

Study population

Our analysis sample included women of reproductive age (15-49 years) at the time of each survey. The primary study population was adolescent girls aged 15-19 years, who were compared to young women aged 20-24 years. Within this sample, we also included a sub-group of adolescent girls and young women with a live birth in the six months before each survey as they are the population benefitting from LAM.

Definitions

We estimated the following indicators capturing three dimensions of lactational amenorrhea: 1) Knowledge of LAM; 2) Use of LAM – including reported, correct, passive, and overall protection by lactational amenorrhea; and 3) Duration of postpartum amenorrhea (Table 1).

Knowledge of LAM was defined as the respondent being aware of LAM as a method of family planning, and was measured based on respondents' binary (yes, no) answer to "Have you ever heard of lactational amenorrhea as a family planning method?" It should be noted that, in the 2016 survey, this question was followed with a probe that describes LAM further, that is, "Up to six months after childbirth, before the menstrual period has returned, women use a method requiring frequent breastfeeding day and night". This probe was not used in the 2006 and 2011 surveys. We assessed knowledge of LAM among all women and among eligible women who could have been using this method at the time of the survey (those with infants below six months of age at the time of each survey).

Use of LAM was based on the reported current use of lactational amenorrhea as a family planning method at the time of each survey (yes or no). This was measured by asking women whether they were doing anything or using any method to delay or avoid getting pregnant; and if so, what method they were using (multiple responses were allowed, for example, LAM and male condoms). We examined the use of LAM through four indicators:

- 1) Current use of LAM - women who reported using LAM as their method of family planning at the time of the survey. We applied two denominators: all women and eligible women (those with infants below six months at the time of the survey).
- 2) Correct use of LAM – women who reported LAM use and met the three LAM criteria, among eligible women (those with infants below six months at the time of the survey). The three criteria for correct use of LAM were; i) woman has a child <6months of age at the time of the survey, ii) woman's menstrual period had not returned since the most recent birth, and iii) her infant was being exclusively breastfed at the time of the survey. *Exclusive breastfeeding* was defined as infants <6 months of age at the time of the survey who were fed exclusively with breastmilk, that is not given anything other than breastmilk in the 24 hours before the survey, except for oral medications.
- 3) Passive use of LAM - women meeting the three LAM criteria but who did not report LAM use, among eligible women (those with infants <6 months at the time of the survey).
- 4) Protection by LAM – women who met the three LAM criteria, regardless of whether they reported LAM use or not), among all women and eligible women.

Duration of postpartum amenorrhea: Postpartum amenorrhea (PPA) period was defined as the time to the resumption of menstruation period following childbirth. This was captured by calculating the median duration in months using the child recode file, following the standard DHS method (Tsfayi et al., 2008).

Table 1. Definitions of key study variables

Indicator (estimate)	Definition (numerator/ denominator)	Stratification by woman's age group (at time of survey)
1 Knowledge of LAM as FP method among all women (% with 95%CI)	Women who report ever hearing of LAM / All women	15-19.9 20-24.9 All women 15-49
2 Knowledge of LAM among eligible women (% with 95%CI)	Women who report ever hearing of LAM / Women with a child <6 months of age	15-19.9 20-24.9 All women 15-49
3 Current reported use of LAM as FP method among all women (% with 95%CI)	Women who report using LAM as their FP method at time of survey / All women	15-19.9 20-24.9 All women 15-49
4 Current reported use of LAM as FP method among eligible women (% with 95%CI)	Women who report using LAM as their FP method at time of survey / Women with a child <6 months of age	15-19.9 20-24.9 All women 15-49
5 Correct use of LAM among eligible women (% with 95%CI)	Women who report using LAM and meet the three criteria/ Women with a child <6 months of age	15-19.9 20-24.9 All women 15-49
6 Passive use of LAM among eligible women (% with 95%CI)	Women who meet the three LAM criteria but do not report using LAM / Women with a child <6 months of age	15-19.9 20-24.9 All women 15-49
7 Protection by LAM among all women (% with 95%CI)	Women who meet the three LAM criteria regardless of whether they report the use of LAM or not / All women	15-19.9 20-24.9 All women 15-49
8 Protection by LAM among eligible women (% with 95%CI)	Women who meet the three LAM criteria regardless of whether they report the use of LAM or not / Women with a child <6 months of age	15-19.9 20-24.9 All women 15-49
9 Duration of postpartum amenorrhea in months (median +IQR)	Number of months before menstrual period returned after most recent birth in the 36 months before the survey	15-19.9 20-24.9 All women 15-49

LAM=Lactational Amenorrhea Method, FP=family planning

Other variables: We used data on women's socio-demographic characteristics at the time of the surveys, including women's age in years, marital status, education status, place and region of residence, and household wealth quintile. Marital status was categorized as "never in union", currently in union/living with a man, and formerly in union/living with a man. Education was categorized as no education, primary education, secondary education, and higher education. Participants' place of residence was either urban or rural, and household wealth quintiles as produced by the DHS.

Statistical Analysis

Analysis was conducted using STATA 14 (StataCorp, Texas, USA). We utilized the *svyset* command in STATA to adjust for the sampling design used and for non-response. Descriptive statistics including frequencies and percentages were used to summarize categorical data while continuous data (duration of PPA) was expressed in terms of median and inter-quartile range. We computed the unweighted numbers of women for the given variable and the weighted percentages and 95% confidence interval. For all denominators using eligible women (with infants below six months at the time of the survey), we only included women with infants who were alive at the time of the survey due to breastfeeding as a criterion for correct LAM use. In the computation of reported use of LAM at the time of the survey, we assumed that all women who did not report using LAM were indeed not using LAM (missing values were recorded as 0). There was no missing data on other key variables for the study samples included in the analysis.

Ethical approval

Data used for the analysis of this paper were obtained with permission from the Demographic and Health Survey Program (DHS) program of the US Agency for International Development. The DHS

receives government permission, uses informed consent, and assures respondents of confidentiality. We did not require an ethics approval for this study since it involved secondary data analysis.

RESULTS

Socio-demographic characteristics of the study samples

The total number of adolescents included in the study was 8,250 and 7,110 young women (Table 2). Across the decade, primary education was the highest level attained in both adolescents (at an average 65%) and young women ($\approx 55\%$) (Table 3). Approximately 77% of adolescents had never been married as compared to approximately 20% in the young women. Table 3 also shows an increasing trend in the proportion of adolescents (17.7% to 24.3%) or young women (22.0% to 29.9%) living in urban areas.

Table 2. Sample size of women according to age group at the time of each survey and recent birth history, by survey

Category	2006	2011	2016
All female respondents (15-49 years)	8,531	8,674	18,506
Adolescents (15-19 years)	1,948	2,026	4,276
Adolescents with a live birth <6months before the survey	367	376	837
Young women (20-24 years)	1,662	1,666	3,782
Young women with a live birth <6months before the survey	1,291	1,225	2,831

Table 3: Socio-demographic characteristics of the study samples for UDHS 2006, 2011 and 2016

Attribute	Survey year	2006				2011				2016			
	Age group	15-19.9		20-24.9		15-19.9		20-24.9		15-19.9		20-24.9	
	n	1,948		1,662		2,026		1,666		4,276		3,782	
		%	95%CI	%	95%CI	%	95%CI	%	95%CI	%	95%CI	%	95%CI
Education	None	3.5%	2.5-4.8	11.5%	9.6-13.8	2.9%	2.1-4.2	4.9%	3.9-6.2	1.8%	1.4-2.4	3.3%	2.7-4.1
	Primary	66.7%	63.6-69.7	59.2%	56.0-62.4	64.8%	61.7-67.8	54.7%	51.6-57.8	64.7%	62.4-67.0	51.0%	48.5-53.4
	Secondary	28.5%	25.5-31.7	23.2%	20.5-26.2	30.8%	28.1-33.7	32.4%	29.5-35.4	31.7%	29.5-33.9	35.1%	32.8-37.4
	Higher	1.3%	0.9-2.0	6.0%	4.8-7.5	1.5%	0.8-2.6	8.0%	6.5-9.8	1.8%	1.4-2.4	10.7%	9.5-12.0
Wealth	Lowest	15.3%	12.9-17.9	18.2%	15.6-21.0	15.4%	13.2-18.0	16.4%	13.9-19.2	17.9%	16.2-19.8	16.9%	15.4-18.6
	Second	17.5%	15.1-20.2	20.1%	17.7-22.7	16.9%	14.6-19.5	17.8%	15.6-20.3	19.7%	18.0-21.5	18.4%	16.8-20.0
	Middle	17.3%	14.9-20.0	17.7%	15.4-20.2	18.0%	15.6-20.5	20.1%	17.1-23.3	19.1%	17.6-20.7	16.1%	14.6-17.7
	Fourth	20.2%	17.7-22.9	15.4%	13.2-17.8	23.5%	20.6-26.6	15.5%	13.1-18.3	20.0%	18.3-21.9	19.3%	17.6-21.2
	Highest	29.8%	26.0-33.9	28.7%	24.8-32.9	26.2%	22.6-30.2	30.3%	26.8-34.0	23.2%	20.8-25.9	29.3%	26.3-32.5
Marital status	Never married	77.6%	75.0-80.0	21.9%	19.4-24.6	77.3%	74.4-79.8	23.9%	21.3-26.7	77.2%	75.5-78.8	25.5%	23.6-27.5
	Married / cohabiting	19.6%	17.4-22.1	67.1%	64.4-69.8	20.0%	17.5-22.7	67.3%	64.3-70.2	19.9%	18.4-21.5	64.0%	61.9-66.0
	Divorced or widowed or separated	2.8%	2.1-3.7	11.0%	9.3-13.0	2.8%	2.0-3.8	8.7%	7.1-10.6	2.9%	2.3-3.6	10.6%	9.5-11.7
Residence	Urban	17.7%	14.0-22.1	22.0%	17.8-26.8	19.3%	16.3-22.8	25.5%	22.2-29.2	24.3%	22.2-26.5	29.9%	27.4-32.6
	Rural	82.3%	77.9-86.0	78.0%	73.2-82.2	80.7%	77.2-83.7	74.5%	70.8-77.8	75.8%	73.5-77.9	70.1%	67.4-72.6
Region	Central 1	11.8%	9.5-14.5	11.5%	9.4-14.0	9.3%	7.0-12.2	13.6%	10.8-16.9	12.1%	10.2-14.2	15.0%	12.7-17.6
	Central 2	9.3%	8.0-10.9	5.6%	4.4-7.0	11.2%	8.2-15.2	10.4%	9.0-12.0	9.8%	8.3-11.4	11.2%	9.2-13.5
	Kampala	8.9%	7.2-11.1	13.3%	10.8-16.4	9.7%	8.1-11.5	10.0%	8.0-12.4	4.7%	3.8-5.8	6.8%	5.9-7.9
	East central	10.3%	8.3-12.7	8.9%	7.2-10.9	9.9%	8.2-11.8	9.5%	7.7-11.5	9.1%	7.5-11.0	8.9%	7.5-10.5
	Eastern	12.2%	10.1-14.7	14.1%	11.6-16.9	15.5%	13.0-18.4	15.5%	13.3-18.1	20.1%	18.5-21.9	16.8%	15.3-18.5
	North	14.0%	11.6-16.7	14.7%	12.0-17.9	8.8%	7.3-10.7	7.4%	5.7-9.6	11.7%	10.6-13.0	9.8%	8.8-11.0

Karamoja	*	*	*	*	3.2%	1.8-5.6	3.0%	2.0-4.4	1.9%	1.5-2.4	1.7%	1.2-2.3
West Nile	5.7%	4.6-7.0	5.0%	3.6-6.8	6.2%	4.9-7.7	5.5%	4.6-6.7	7.6%	5.9-9.6	6.2%	5.1-7.5
Western	15.3%	13.1-17.9	15.2%	12.9-17.9	14.1%	11.1-17.6	12.7%	10.2-15.8	12.8%	11.1-14.8	13.3%	11.9-14.9
South west	12.6%	10.5-15.0	11.8%	9.5-14.4	12.2%	10.3-14.3	12.4%	10.6-14.4	10.2%	8.9-11.6	10.3%	9.1-11.6
% with live birth in <6months	13.4%	10.9-16.4	27.9%	24.3-31.7	13.2%	10.7-16.3	29.7%	25.9-33.8	14.6%	12.7-16.9	31.5%	28.9-34.3

* not sampled due to insecurity

Knowledge of LAM

Knowledge of LAM as a FP method was analyzed for two categories of women; all women and eligible women (who had a child <6 months at the time of the survey). Between 2006 and 2016, knowledge of LAM among all and eligible adolescent girls increased. Among all adolescents, knowledge of LAM increased from 3.7% in 2006 to 36.9% in 2016; whilst among the eligible adolescents, it increased from 6.0% to 56.7%, respectively (Table 4). Compared to young women, knowledge of LAM among all or eligible women was generally lower among the adolescents. Among eligible young women, knowledge of LAM increased from 9.7% in 2006 to 64.1% in 2016.

Use of LAM

We determined the use of LAM using four broad indicators: current reported use, correct use, passive use of LAM, and protection by LAM. The overall use of LAM over the ten years was low among adolescents, young women and all women of reproductive age. The overall current reported use or correct use of LAM was less than 2% (Table 4). The percentage of eligible adolescents or young women correctly using LAM was lower than the percentage who reported using LAM as their FP method. Among eligible women, the current reported use as of LAM in adolescents was 1.3% in 2016 as compared to 0.7% in young women; whilst the correct use of LAM in 2016 was 0.9% in adolescents compared to 0.5% in young women. Table 4 further shows that across the three surveys, the percentage of eligible women passively using LAM was higher compared to those reporting intentional LAM use (>50% of eligible women in all surveys and age groups). Also, the passive use of LAM decreased among all age groups over the review period. Among adolescent girls, passive LAM use decreased from 76.4% in 2006 to 56.3% in 2016, compared to young women among whom it decreased from 84.9% to 56.9% respectively. Between 2006 and 2016, the percentage of eligible women protected by LAM decreased across all age groups. As of 2016, at least 50% of women eligible for LAM were indeed protected by LAM in all three age groups, more so among all women of reproductive age at 62.0%. The percentage of eligible adolescents protected by LAM decreased from 76.4% in adolescents in 2006 to 57.2% in 2016, as compared to young women where it decreased from 84.9% in 2006 to 57.3% in 2016 (table 4).

Table 1. Knowledge and use of lactational amenorrhea and duration of postpartum amenorrhea, by age group and survey

LAM outcome	Age group	2016			2011			2006		
		N	%	95%CI	N	%	95%CI	N	%	95%CI
1. Knowledge of LAM among all women	15-19.9	4,276	36.9%	35.0-38.9	2,026	7.3%	5.7-9.4	1,948	3.7%	2.7-5.0
	20-24.9	3,782	65.1%	63.3-67.0	1,666	13.4%	11.0-16.1	1,662	6.6%	5.4-8.1
	15-49	18,506	64.0%	62.9-65.1	8,674	13.0%	11.6-14.6	8,531	6.9%	5.8-8.1
2. Knowledge of LAM among eligible women	15-19.9	220	56.7%	49.2-63.8	107	13.5%	7.5-23.2	110	6.0%	3.5-13.8
	20-24.9	461	64.1%	58.7-69.2	230	14.5%	9.5-21.4	216	9.7%	6.2-14.9
	15-49	1,495	68.9%	66.1-71.5	783	15.3%	12.4-18.7	802	7.6%	5.7-10.0
3. Current reported Use of LAM as FP method among all women	15-19.9	1,595	0.4%	0.2-0.8	2,026	**	**	1	**	**
	20-24.9	2,448	1.4%	0.9-2.1	1,666	0.1%	**	6	**	**
	15-49	11,811	1.0%	0.8-1.2	8,674	0.1%	**	27	**	**
4. Current reported Use of LAM as FP method among eligible women	15-19.9	122	2.2%	0.7-6.5	107	**	**	0	**	**
	20-24.9	302	1.0%	0.4-2.9	230	**	**	2	**	**
	15-49	1,026	2.5%	1.6-3.7	783	0.3%	**	3	**	**
5. Correct use of LAM among women reporting LAM use & child is alive	15-19.9	6	35.2%	8.8-75.2	1	**	**	0	**	**
	20-24.9	30	6.0%	1.7-19.3	3	**	**	0	**	**
	15-49	109	15.6%	9.8-24.0	13	**	**	2	**	**
6. Passive use of LAM among eligible women	15-19.9	122	97.5%	89.8-99.3	107	**	**	0	**	**
	20-24.9	302	98.8%	95.9-99.7	230	99.6%	98.6-99.9	2	**	**
	15-49	1,026	97.1%	95.3-98.2	783	82.4%	79.2-85.2	3	**	**
7. Protection by LAM among all women	15-19.9	4,276	2.9%	2.3-3.5	2,026	3.7%	2.8-5.0	1,948	4.2%	3.3-5.3
	20-24.9	3,782	6.9%	6.0-7.9	1,666	11.5%	9.8-13.6	1,662	10.9%	9.4-12.7
	15-49	18,506	4.9%	4.5-5.3	8,674	7.5%	6.8-8.3	8,531	8.0%	7.4-8.7
8. Protection by LAM among eligible women	15-19.9	220	57.2%	49.5-64.6	107	73.5%	62.7-82.1	110	76.4%	66.5-83.99
	20-24.9	461	57.3%	52.5-62.0	230	80.3%	73.8-85.5	216	84.9%	78.8-89.5
	15-49	1,495	62.0%	58.9-64.9	783	82.7%	79.6-85.5	802	86.4%	83.3-88.98

**Percentages not computed due to small sample size

Duration of postpartum amenorrhea

Over the ten-year period, the median duration of postpartum amenorrhea declined among both adolescents and young women who had had a live birth in the 3 years preceding each survey (Figure 1). Among the adolescents, the median duration of PPA decreased from 8.3 months in 2006 to 6.9 months in 2016, while in young women it decreased from 11.2 months to 8.0 months respectively. Across all age groups, the median duration of PPA was lowest among adolescents. The median duration of postpartum amenorrhea in 2016 was 1.1 months shorter among adolescents compared to young women.

Figure 1. Duration of postpartum amenorrhea by age group and survey

DISCUSSION

This study explored the knowledge and use of lactational amenorrhea as a FP method among adolescents in Uganda. The findings indicate that nearly 60% of eligible adolescents in 2016 had knowledge about LAM, and there was an increase in the percentage of adolescents with knowledge of LAM in the examined period. The reported and correct use of LAM among eligible adolescents was very low, and there was a high but declining prevalence of passive LAM use. The results also show a decrease in the median duration of postpartum amenorrhea among adolescents between 2006 and 2016.

We found large increases in knowledge of LAM among all and eligible adolescents, especially between 2011 and 2016. Part of this increase may be attributed to changes in the measurement of LAM knowledge in 2016 where a description of LAM was read out to the respondents by survey enumerators, which was not the case in the years before. While some views have previously considered this as a quality concern resulting in misreporting by women, other analyses have supported the inclusion of the LAM description, as higher proportions of self-reported LAM users who correctly practised LAM were observed in surveys with the description (Fabic and Choi, 2013). Any true increase in levels of knowledge about LAM may be attributed to various efforts that have been implemented over time by the Ugandan government including free primary education, family planning initiatives including the use of community health workers, and promoting the provision of adolescent and youth friendly services (Atuyambe et al., 2015, Nalwadda et al., 2019). The percentage of eligible adolescents who were aware of LAM in this study was higher than what has been reported among women of reproductive age in other SSA countries (Abraha et al., 2018, Alvey, 2015, Ekpenyong et al., 2013). This may be explained by differences in the methods for determining knowledge of LAM. For example, studies by Abraha et al (2018) and Ekpenyong et al (2013) went beyond the DHS approach and included an assessment of respondent's understanding of the three criteria for LAM practice. Regardless, the observed increase in knowledge may result in better attitudes towards and acceptability of LAM as an effective contraceptive method, which would contribute to increased adoption. But this would be dependent on the appropriateness of the information to which adolescents are exposed and their ability to apply it correctly, among other factors.

Findings from this study show very low percentages of eligible adolescents reporting the use of LAM as their primary FP method, and those that were correctly using it. The low prevalence of reported LAM use may partly be explained by methodological limitations posed by the DHS where if a woman reports use of both LAM and other modern methods, the most effective method is recorded and the woman is not recorded as a LAM user (Fabic and Choi, 2013). The use of other FP methods besides LAM during the first 6 months after childbirth has been reported (Afifi, 2007). This may explain the observed high proportion of women that were passively using LAM (meet the LAM criteria but do not report LAM use). The study findings relate to previous studies that also show a low prevalence of LAM use among women of reproductive age in general (Fabic and Choi, 2013) and adolescents in particular (Figaroa et al., 2020). Low correct use of LAM is a result of not fulfilling one or more of the three criteria, which also influence the effectiveness of LAM as a FP method. This would mean that, if new adolescent mothers are not correctly using LAM and they are not using any other FP methods within the first 6 months after childbirth, they would be at higher risk of becoming pregnant again. This would also vary based on factors like sexual activity or access to clinical or social support during this period. In other studies, violation of the exclusive breastfeeding and postpartum amenorrhea criteria have been reported to be the primary factors affecting the correct use of LAM (Van der Wijden and Manion, 2015, Sipsma et al., 2013). This is particularly important in adolescents among whom, early initiation and exclusive breastfeeding have been reported to be low, with a higher likelihood of discontinuation (Benova et al., 2020, Jara-Palacios et al., 2015).

Postpartum amenorrhea is one of the key determinants for correct LAM use and its effectiveness. The study findings showed a decrease in the median duration of PPA among adolescents from 8.3 (2006) to 6.9 (2016). The resumption of menses is a commonly used reported determinant of FP use among postpartum women (Gahungu et al., 2021), yet variably understood as reported by Cooper et al (2019), who found misconceptions about the return of fecundity in Tanzania (Cooper et al., 2019). The decreasing duration of PPA points to the need to strengthen postpartum FP among adolescents, as the risk of becoming pregnant may be higher in this sub-population. The review by Figaroa and colleagues noted wide variation in PPA duration among adolescents in LMICs (Figaroa et al., 2020); while a study in India among women with at least one livebirth reported a lower duration of PPA (5.7 months) than what was found in the present study (Singh et al., 2012). Breastfeeding practices have

also been linked to a shorter total duration of PPA (Singh et al., 2012). According to the authors, the risk of resumption of menses decreases with the increasing duration of breastfeeding. This may explain the decreasing duration of PPA among adolescents and why it was lower compared to young women in this study.

Limitations

This study benefitted from three rounds of highly comparable nationally representative data collected over ten years in Uganda to provide an understanding of patterns of LAM use among adolescents. We constructed several indicators of LAM knowledge and use, with sensitivity to the population under consideration (i.e. denominators). Nonetheless, this study has limitations. First, it was not possible to investigate factors associated with LAM knowledge or use among adolescents because of the small sample size. Such information would help inform strategies to improve the use of LAM in this key sub-population. The findings are also limited by the use of self-reported data on the use of LAM and its criteria especially exclusive breastfeeding and return of menses. Correct/accurate reporting on such variables is rather subjective and is affected by the individual's understanding which is often limited. Also, the scope of knowledge of LAM assessed in the surveys is limited and may not provide a proper indication of the understanding of LAM among adolescents. Further qualitative studies can help elucidate the understanding and considerations for use of LAM in this important population segment.

CONCLUSIONS

There is increasing awareness about LAM but, the reported and correct use of LAM is low among adolescents with a live baby under 6 months of age in Uganda. There is need to improve uptake of contraceptive methods especially in postpartum adolescents through targeted counselling and educational programs. This will go a long way in not only reducing their short-interpregnancy intervals but improving their health and wellbeing including that of their children. Further, additional research is needed to better understand the dynamics of correct LAM use and transition to other FP methods among adolescent mothers.

Acknowledgements

CB and PB would like to thank the Institute of Tropical Medicine in Belgium for the training that contributed to the development of this paper. We would also like to convey our thanks to the Demographic and Health Survey program for granting access to the survey datasets which were used in this study.

Contributions

CB and PB share first authorship and contributed equally to this paper. CB, PB conceptualized the study, conducted data analysis and developed the manuscript. LB provided analysis guidance and supported the conceptualization process and data analysis. CB, PB and LB wrote the first manuscript draft. STW supported data analysis and participated in manuscript review. PW provided technical guidance to the manuscript development process. All authors reviewed the manuscript and approved the final manuscript.

Funding statement

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests

The authors declare no competing interests

Patient consent for publication

Not required.

Patient and public involvement: Patients or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Data availability statement

Data are available in a public, open access repository. Data are available for research purposes on www.dhsprogram.com after registration

References

- ABRAHA, T. H., TEFERRA, A. S., GELAGAY, A. A., WELESAMUEL, T. G., FISSEHA, G. K., AREGAWI, B. G. & BELAY, D. S. 2018. Knowledge and associated factors of lactational amenorrhea as a contraception method among postpartum women in Aksum town, Tigray Region, Ethiopia. *BMC Research Notes*, 11, 641.
- AFIFI, M. 2007. Lactational amenorrhoea method for family planning and women empowerment in Egypt. *Singapore Med J*, 48, 758-62.
- AHINKORAH, B. O., AMEYAW, E. K. & SEIDU, A.-A. 2020. Socio-economic and demographic predictors of unmet need for contraception among young women in sub-Saharan Africa: evidence from cross-sectional surveys. *Reproductive Health*, 17, 163.
- ALEGE, S. G., MATOVU, J. K., SSENSALIRE, S. & NABIWEMBA, E. 2016. Knowledge, sources and use of family planning methods among women aged 15-49 years in Uganda: a cross-sectional study. *Pan African Medical Journal*, 24.
- ALVEY, J. 2015. *Measuring the use of the Lactational Amenorrhea Method as a postpartum family planning method in urban Nigeria: challenges in measurement and recommendations for future MIYCN-FP integrated programs*.
- AMONGIN, D., BENOVA, L., NAKIMULI, A., NAKAFEERO, M., KAHARUZA, F., ATUYAMBE, L. & HANSON, C. 2020a. Trends and determinants of adolescent childbirth in Uganda- analysis of rural and urban women using six demographic and health surveys, 1988–2016. *Reproductive Health*, 17, 74.
- AMONGIN, D., NAKIMULI, A., HANSON, C., NAKAFEERO, M., KAHARUZA, F., ATUYAMBE, L. & BENOVA, L. 2020b. Time trends in and factors associated with repeat adolescent birth in Uganda: Analysis of six demographic and health surveys. *PLOS ONE*, 15, e0231557.
- ATUYAMBE, L. M., KIBIRA, S. P., BUKENYA, J., MUHUMUZA, C., APOLOT, R. R. & MULOGO, E. 2015. Understanding sexual and reproductive health needs of adolescents: evidence from a formative evaluation in Wakiso district, Uganda. *Reproductive health*, 12, 1-10.
- BENOVA, L., SIDDIQI, M., ABEJIRINDE, I.-O. O. & BADEJO, O. 2020. Time trends and determinants of breastfeeding practices among adolescents and young women in Nigeria, 2003–2018. *BMJ Global Health*, 5, e002516.
- BURKE, H. M., SANTO, L. D., BERNHOLC, A., AKOL, A. & CHEN, M. 2018. Correlates of Rapid Repeat Pregnancy Among Adolescents and Young Women in Uganda. *Int Perspect Sex Reprod Health*, 44, 11-18.
- COOMSON, J. I. & MANU, A. 2019. Determinants of modern contraceptive use among postpartum women in two health facilities in urban Ghana: a cross-sectional study. *Contracept Reprod Med*, 4, 17.
- COOPER, C. M., KAVLE, J. A., NYONI, J., DRAKE, M., LEMWAYI, R., MABUGA, L. & PFITZER, A. 2019. Perspectives on maternal, infant, and young child nutrition and family planning: Considerations for rollout of integrated services in Mara and Kagera, Tanzania. *Matern Child Nutr*, 15 Suppl 1, e12735.
- EJANG, M. 2020. Tackling Youth Unemployment with Youth Livelihood Programme (YLP) in Northern Uganda.
- EKPENYONG, C. E., DANIEL, N. E., UWAH, A. F., ETTEBONG, E. O. & IBU, J. O. 2013. Lactational amenorrhoea method of contraception: An in-depth study of awareness, knowledge and practice by breast feeding mothers with unintended pregnancies. *International Journal of Medicine and Medical Sciences*, 5, 6-13.
- ERFINA, E., WIDYAWATI, W., MCKENNA, L., REISENHOFER, S. & ISMAIL, D. 2019. Adolescent mothers' experiences of the transition to motherhood: An integrative review. *International journal of nursing sciences*, 6, 221-228.
- FABIC, M. & CHOI, Y. Measuring use of the lactational amenorrhea method through the Demographic and Health Surveys: data quality and implications. annual meeting of the Population Association of America, New Orleans, 2013.

- FIGAROA, M. N., BELLIZZI, S., DELVAUX, T. & BENOVA, L. 2020. Lactational amenorrhoea among adolescent girls in low-income and middle-income countries: a systematic scoping review. *BMJ global health*, 5, e002492.
- GAHUNGU, J., VAHDANINIA, M. & REGMI, P. R. 2021. The unmet needs for modern family planning methods among postpartum women in Sub-Saharan Africa: a systematic review of the literature. *Reproductive Health*, 18, 1-15.
- JARA-PALACIOS, M. Á., CORNEJO, A. C., PELÁEZ, G. A., VERDESOTO, J. & GALVIS, A. A. 2015. Prevalence and determinants of exclusive breastfeeding among adolescent mothers from Quito, Ecuador: a cross-sectional study. *International breastfeeding journal*, 10, 33-33.
- JOSHI, A. K., TIWARI, D. P., POUDYAL, A., SHRESTHA, N., ACHARYA, U. & DHUNGANA, G. P. 2020. Utilization of Family Planning Methods Among Postpartum Mothers in Kailali District, Nepal. *International journal of women's health*, 12, 487-494.
- MANGELI, M., RAYYANI, M., CHERAGHI, M. A. & TIRGARI, B. 2017. Exploring the Challenges of Adolescent Mothers From Their Life Experiences in the Transition to Motherhood: A Qualitative Study. *Journal of family & reproductive health*, 11, 165-173.
- MINISTRY OF HEALTH, U. 2016. Reproductive, Maternal, Newborn, Child and Adolescent Health Sharpened Plan for Uganda 2016/17 – 2019/20. Kampala, Uganda.
- NALWADDA, G., NAMUTEBI, M. & VOLGSTEN, H. 2019. Health care providers' perceptions of family planning and contraception education for adolescents in Kampala, Uganda—A qualitative study. *Sexual & Reproductive Healthcare*, 21, 15-20.
- NEAL, S., MATTHEWS, Z., FROST, M., FOGSTAD, H., CAMACHO, A. V. & LASKI, L. 2012. Childbearing in adolescents aged 12-15 years in low resource countries: a neglected issue. New estimates from demographic and household surveys in 42 countries. *Acta Obstet Gynecol Scand*, 91, 1114-8.
- RAHMAN, A., NOMANI, D. & TANEERANICHSKUL, S. 2020. Trends and Determinants of EBF among Adolescent Children Born to Adolescent Mothers in Rural Bangladesh. *International Journal of Environmental Research and Public Health*, 17, 9315.
- SINGH, N. S., SINGH, N. S. & NARENDRA, R. 2012. Postpartum amenorrhoea among Manipuri women: a survival analysis. *Journal of health, population, and nutrition*, 30, 93.
- SIPSMA, H. L., BRADLEY, E. H. & CHEN, P. G. 2013. Lactational amenorrhea method as a contraceptive strategy in Niger. *Maternal and child health journal*, 17, 654-660.
- SULLY, E., BIDDLECOM, A., DARROCH, J. E., RILEY, T., ASHFORD, L. S., LINCE-DEROCHÉ, N., FIRESTEIN, L. & MURRO, R. 2020. Adding it up: investing in sexual and reproductive health 2019.
- TESFAYI, G., SHEA, R. & VINOD, M. 2008. Contraceptive Use, Breastfeeding, Amenorrhea and Abstinence During the Postpartum Period: An Analysis of Four Countries. *DHS Analytical Studies* 14.
- UGANDA BUREAU OF STATISTICS & ICF 2017. Uganda Demographic and Health Survey 2016: Key Indicators Report. Kampala, Uganda. In: UBOS & ROCKVILLE (eds.). Maryland, USA: UBOS ICF.
- VAN DER WIJDEN, C. & MANION, C. 2015. Lactational amenorrhoea method for family planning. *Cochrane Database of Systematic Reviews*.
- WATTS, M. C. N. C., LIAMPUTTONG, P. & MCMICHAEL, C. 2015. Early motherhood: a qualitative study exploring the experiences of African Australian teenage mothers in greater Melbourne, Australia. *BMC public health*, 15, 1-11.
- WORLD HEALTH ORGANIZATION. 2018. *Postpartum family planning: essential for ensuring health of women and their babies* [Online]. Available: https://www.who.int/reproductivehealth/topics/family_planning/world-contraception-day-2018/en/ [Accessed].
- WORLD HEALTH ORGANIZATION. 2020. *Adolescent pregnancy* [Online]. Available: <https://www.who.int/news-room/fact-sheets/detail/adolescent-pregnancy> [Accessed].

YILMAZ, E., YILMAZ, Z., ISI, H., GULTEKIN, I. B., TIMUR, H., KARA, F. & KUCUKOZKAN, T. 2016.
Factors associated with breastfeeding initiation and exclusive breastfeeding rates in Turkish adolescent mothers. *Breastfeeding Medicine*, 11, 315-320.