Title: Spectrum of HRCT chest findings in RT-PCR positive asymptomatic COVID-19 patients at a COVID designated hospital in Nepal.

Authorship Details

Dr. Bina Basnet, MD

Nepalese Army Institute of Health Sciences, College of Medicine. Nepal

Email: basnet.bina@naihs.edu.np / binprisha@gmail.com

Dr. Sujit Pant, MD

Nepalese Army Institute of Health Sciences, College of Medicine. Nepal

Email: sujit.pant046@gmail.com

Dr. Sujata Pant, MD

Nepalese Army Institute of Health Sciences, College of Medicine. Nepal

Email: sujata23@gmail.com

Dr. Kalpana Rai, MD

Nepalese Army Institute of Health Sciences, College of Medicine. Nepal

Email: drkalpanarai@gmail.com

Dr. Niraj Basanta Tulachan, MD

Nepalese Army Institute of Health Sciences, College of Medicine. Nepal

Email: nbtulachan@gmail.com

Dr. Bibek Karki, MD

Nepalese Army Institute of Health Sciences, College of Medicine. Nepal

Email: forbibek@gmail.com

Dr. Rajiv Shahi, FRCR

Nepalese Army Institute of Health Sciences, College of Medicine. Nepal

Email: rsshahi@hotmail.com

Dr. Shiva Bahadur Basnet, MD

Nepalese Army Institute of Health Sciences, College of Medicine. Nepal

Email: shiva.basnet@naihs.edu.np

Dr. Bikash Bikram Thapa, MS MSPH

Nepalese Army Institute of Health Sciences, College of Medicine. Nepal

Email: bikashbikram.thapa@naihs.edu.np

Abstract:

Introduction: COVID-19 pandemic is grappling the world with the surge of infection time and again. Clinicians are trying to justify the ethics of public health care. Asymptomatic COVID-19 cases are going undocumented and most of them practice self-isolation. Studies have revealed significant radiological changes among RT-PCR positive asymptomatic COVID-19 cases.

Objective: The aim of this cross-sectional study is to characterized chest CT findings of asymptomatic RT-PCR-positive patients in one of the COVID-designated hospitals in Nepal.

Results: Out of 43, 26 (60.5%) participants had positive Chest CT scan findings consistent with COVID pneumonia. 65% had bilateral and 77% had multifocal lesions. The ground-glass opacities (92%), mixed (ground-glass opacities and consolidation) pattern (30.7%), and consolidation only (34.6%) were common chest CT findings. The median CT score was 3.5 (Interquartile range; 2-6). **Conclusion:** The majority of the RT-PCR positive asymptomatic patient present with CT scan changes of lungs which are important to determine clinical status, prognosis, and long-term sequel in those cohorts.

Key Words: COVI-19; Chest CT; Viral Pneumonia; Nepal

Introduction:

Since the first reports of the Corona Virus Disease-2019 (COVID-19) from Wuhan, city in China, the cases have been reported from all the seven continents and had already caused death in millions. The COVID-19 disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is primarily transmitted through respiratory droplets and infect lungs. [1] Reverse Transcriptase Polymerase reaction (RT-PCR) is the confirmatory laboratory diagnosing test for SARS-CoV2. The RT-PCR is recommended in asymptomatic cases especially in close contacts, screening and early identification of infection in high risk population and locations like healthcare centers, prior to surgical procedures, and prior to receiving immunosuppressive therapy. [2,3]

In a metanalysis the pooled percentage of the asymptomatic infection was 15.6%. [4] Asymptomatic cases are less infectious than symptomatic. [5,6] Despite low infectivity, 20-50% of the asymptomatic cases had specific changes in the chest computed tomography (CT). But the probability of progression of the asymptomatic cases into pneumonia is very less (<1%). [7,8]

CT changes of the asymptomatic patients are important not only to evaluate the chest CT as alternative diagnostic modalities for selected COVID-19 cases but also to evaluate the long term outcomes of COVID-19 related respiratory pathophysiology, which is mostly unknown. [9]

The objective of our study is to characterize the chest CT scan features and clinical outcomes of RT-PCR confirmed asymptomatic COVID-19 patients in COVID-19 designated hospital (Shree Birendra Hospital), Nepal.

Methodology

This is a cross sectional study where a cohort of laboratory proven RT-PCR positive COVID-19 asymptomatic adult (age > 18 years) patients were included and were subjected to the HRCT scan of the chest. All the cases included in this study fulfill the WHO criteria of the “close contacts” of the COVID-19 confirmed cases. The study is conducted between October 2020 and December 2020 with ethical approval from institutional review board and consent of the patient and

HRCT scan were acquired in the Hitachi Multidetector 128 slice CT scanner. The parameters for used for CT acquisition were helical mode volumetric HRCT with Tube voltage 100kVp-120kVp and tube current 80-500mA, and slice thickness of 1.0mm with reconstruction interval, 0.6mm using a sharp reconstruction algorithm. CT images were obtained with the patient in supine position with full inspiration. Intravenous contrast administration was not used. Acquired images were transferred to a separate workstation for further processing. Image was reconstructed in axial, coronal, and sagittal planes to detect the craniocaudal and axial/peripheral distribution of the lung parenchymal involvement.

All images were viewed on both lung (width, 1500 HU; level, -700 HU) and mediastinal (width, 350 HU; level, 40 HU) settings. The chest CT scan was evaluated by two radiologist characterizing the parenchymal involvement on the basis of the (a) characteristic findings and morphology: ground glass opacities, consolidation, linear bands, bronchial wall thickening, nodules and additional findings like pleural effusion and mediastinal lymphadenopathy. (b) Distribution of the involvement: Laterality, craniocaudally distribution, number of the lobes involved, percentage of involvement in each lobe. Then the CT severity score (0-25) was calculated following the semi- quantitative scoring system which depends on the visual

assessment of the each 5 lung lobes (0- 0%; <5%; 2- 5 to 25%; 3- 26 to 50%; 4- 51 to 75%; 5- > 75%) that was initially proposed by Pan et al.[10]

Results

In this cross-sectional study total of 43 RT-PCR positive COVID-19 cases were included. Among the study cohort 26 (60.5%) had positive Chest CT scan findings (Figure 1-3). The positive Chest CT had mostly bilateral (65.3%) and multifocal (77%) lesion. 92% of them had ground glass opacities (Table 1).

The median global CT score of the abnormal CT scan chest was 3.5 (Interquartile range; 2-6). The mean CT score value was significantly higher (5.3 ± 2.6 vs 3.0 ± 1.0) in bilateral lung disease. The ratio of upper to lower lobe involvement was 2:3. The patients with normal CT scan had higher mean cycle threshold (CT) value of RT-PCR test than abnormal chest CT scan group (23.8 vs 21.7;). However, the difference was statistically not significant. One patient developed mild COVID-19 symptoms (cough and headache) during follow up.

Table1: HRCT details of the study cohort:

Particulars	Value
Age	41.42 $\pm$ 11.7 (23-66)
Male: Female	3:2
Positive CT findings	60.5%(26/43)
Lobar involvement	Right- 7 (27%) , Left-2 (8%) and Bilateral -17 (65%)
Number of Focus	Single- 23% (6/26; Multiple – 77%(20/26)
Characteristic Chest HRCT scan findings	Consolidation 34.6%; (9) Ground Glass opacities 92%; (24) GGO and Consolidation 30.7% (8) Atelectatic Band 19% (5) Septal thickening 30.7%; (8)

Figure 1: Multifocal predominantly peripheral consolidative and ground glass opacities in bilateral lung. Interspersed areas of interlobular septal thickening seen in the inferior lingual segment of left upper lobe.

Figure 2: Patchy peripheral ground glass opacity with focal traction bronchiectasis within the affected area

Figure 3: Consolidation (a) Subsegmental (b) multifocal peripheral rounded morphology

Discussion:

Pneumonia is primary manifestation of the COVID-19 disease. WHO has classified symptomatic COVID-19 infection into mild cases, pneumonia, severe pneumonia, and critical disease (sepsis, septic shock and or ARDS). In a large cohort of population based study the 30- 40% were asymptomatic young adults. The virus can be cultured from infected individual as early as six days prior to the developments of symptoms. Whereas it takes at least a week for CT scan to detect changes in lung parenchyma.[11-14] Screening CT chest had positivity rate of 1.6%.[15] In RT-PCR confirmed COVID-19 cases HRCT help in prognostication, evaluating the disease progression and monitoring the response to therapy. [16]

With RT-PCR as reference, the sensitivity, specificity, accuracy of chest CT in indicating COVID-19 infection were 97%, 25% and 68% respectively, the accuracy of which is higher in age more than 60 years. Studies have shown that Bilateral lung (90%) involvement with ground glass opacities (46-50%) and mixed GGO along with consolidation (44-50%), and consolidation only (25%) were the most common pattern in Chest CT scan characteristics of COVID-19 patients. [12,14,16,17] Similar pattern of Ground glass opacities (95%) and consolidation (5%) with predominantly bilateral, sub pleural, and multiple lesion were described among asymptomatic cases.[9] [18] 83% had features of lung consolidation among asymptomatic COVID-19 cases (54%) found in the cruise ship “Diamond Princess”.[19] Parenchymal inflammation, endothelial dysfunction, and cytokine release syndrome are responsible for COVID-lung and final radiological manifestations. The pathophysiological pathway can lead to one or more of the following: acute respiratory distress syndrome (ARDS) with diffuse alveolar

damage (DAD), diffuse thrombotic alveolar microvascular occlusion, and inflammatory mediator-associated airway inflammation.[17,20,21,20]

The median CT severity score in this study was 3.5 with range (range, 1-11). Franccone et al found significant positive correlation ($p < 0.0001$) of CT score between age of patient, inflammatory biomarkers, and with severity of the disease. CT score >18 has hazard ratio of 8.33 (95% CI, 3.19-21.7) for COVID-19 related mortality.[22] The significance of CT score in asymptomatic patients is yet to be evaluated in short and long term follow up.

Based on SARS-COV-1 data of 2003 two third of the survivors suffered from TGF- β -mediated pulmonary fibrosis and SARS-COV-2 is expected to share similar chronic sequel.[23,24]. CT chest is widely available modality to assess and follow the pulmonary changes in COVID-19 patients both symptomatic and asymptomatic. During early 2020 asymptomatic or undocumented cases were responsible for 79% of the documented cases. [25]Probability of having incidental CT findings among asymptomatic cases is high due to ongoing progression of the pandemic. Though it is difficult to explain the temporal phase of CT changes in asymptomatic cases screening CT is recommended for all RT-PCR positive COVID-19 patients for the purpose of characterization of the findings and its long-term sequel.[26] The study was conducted during first COVID-19 pandemic wave in Nepal when the average nation wide case positivity rate was 15.6%. Study with large sample size and follow up scan add more to the scientific evidence on radiological features of asymptomatic COVID-19 cases. Study in a context of higher incidence rate can unfold different data and evidences.

Conclusion:

Chest CT plays an important role in the diagnosis and management of the RT-PCR negative COVID suspected as well as RT-PCR positive asymptomatic COVID-19 cases. The long term sequel of the COVID-Lung is little known to us. The present evidences suggest that the COVID-19 asymptomatic cases should be followed clinically and radiologically in order to evaluate its outcome.

Declarations

Funding

Not applicable

Conflicts of interest/Competing interests

We declare no conflict of interest or competing interests

Ethics approval

Ethical approval for this study was taken from Institutional review board

Consent to participate

Written consent taken

Consent for publication

All authors consent to publication of this paper

Availability of data and material

All data is available for review by contacting Dr. Bikash Thapa

Code availability

Not applicable

Authors' contributions

All author has made substantial contributions to the conception or design of the work; the acquisition, analysis, interpretation of data; and the creation of manuscript. BB agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

References:

1. Meyerowitz EA, Richterman A, Gandhi RT, Sax PE. Transmission of SARS-CoV-2: A Review of Viral, Host, and Environmental Factors. *Ann Intern Med* 2021, 174(1):69-79.
2. Infectious Diseases Society of America Guidelines on the Diagnosis of COVID-19, updated December 23, 2020. [<https://www.idsociety.org/practice-guideline/covid-19-guideline-diagnostics>]. Accessed 19 March 2019.
3. Overview of Testing for SARS-CoV-2 [<https://www.cdc.gov/coronavirus/2019-ncov/hcp/testing-overview.html>]. Accessed 19 March 2019
4. He J, Guo Y, Mao R, Zhang J. Proportion of asymptomatic coronavirus disease 2019: A systematic review and meta-analysis. *J Med Virol* 2021, 93(2):820-830.
5. Madewell ZJ, Yang Y, Longini IM, Jr., Halloran ME, Dean NE. Household Transmission of SARS-CoV-2: A Systematic Review and Meta-analysis. *JAMA Netw Open* 2020, 3(12):e2031756.
6. Buitrago-Garcia D, Egli-Gany D, Counotte MJ, Hossmann S, Imeri H, Ipekci AM *et al.* Occurrence and transmission potential of asymptomatic and presymptomatic SARS-CoV-2 infections: A living systematic review and meta-analysis. *PLoS Med* 2020, 17(9):e1003346.
7. Wang Y, Liu Y, Liu L, Wang X, Luo N, Li L. Clinical Outcomes in 55 Patients With Severe Acute Respiratory Syndrome Coronavirus 2 Who Were Asymptomatic at Hospital Admission in Shenzhen, China. *J Infect Dis* 2020, 221(11):1770-1774.
8. Hu Z, Song C, Xu C, Jin G, Chen Y, Xu X *et al.* Clinical characteristics of 24 asymptomatic infections with COVID-19 screened among close contacts in Nanjing, China. *Sci China Life Sci* 2020, 63(5):706-711.
9. Meng H, Xiong R, He R, Lin W, Hao B, Zhang L *et al.* CT imaging and clinical course of asymptomatic cases with COVID-19 pneumonia at admission in Wuhan, China. *J Infect* 2020, 81(1):e33-e39.

10. Pan F, Ye T, Sun P, Gui S, Liang B, Li L *et al.* Time Course of Lung Changes at Chest CT during Recovery from Coronavirus Disease 2019 (COVID-19). *Radiology* 2020, 295(3):715-721.
11. Oran DP, Topol EJ. Prevalence of Asymptomatic SARS-CoV-2 Infection : A Narrative Review. *Ann Intern Med* 2020, 173(5):362-367.
12. Clinical management of COVID-19: interim guidance [WHO-2019-nCoV-clinical-2020.5-eng%20(1).pdf]. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>. Accessed 19 March 2021.
13. Lavezzo E, Franchin E, Ciavarella C, Cuomo-Dannenburg G, Barzon L, Del Vecchio C *et al.* Suppression of a SARS-CoV-2 outbreak in the Italian municipality of Vo'. *Nature* 2020, 584(7821):425-429.
14. Ramanan RV, Joshi AR, Venkataramanan A, Nambi SP, Badhe R. Incidental chest computed tomography findings in asymptomatic Covid-19 patients. A multicentre Indian perspective. *Indian J Radiol Imaging* 2021, 31(Suppl 1):S45-S52.
15. Achour A, Dkhil O, Saad J, Abdelali M, Zrig A, Hmida B *et al.* Chest CT-scan finding of asymptomatic COVID-19 pneumonia: a prospective 542 patients' single center study. *Pan Afr Med J* 2020, 36:257.
16. Ojha V, Mani A, Pandey NN, Sharma S, Kumar S. CT in coronavirus disease 2019 (COVID-19): a systematic review of chest CT findings in 4410 adult patients. *Eur Radiol* 2020, 30(11):6129-6138.
17. Ai T, Yang Z, Hou H, Zhan C, Chen C, Lv W *et al.* Correlation of Chest CT and RT-PCR Testing for Coronavirus Disease 2019 (COVID-19) in China: A Report of 1014 Cases. *Radiology* 2020, 296(2):E32-E40.
18. Shi H, Han X, Jiang N, Cao Y, Alwalid O, Gu J *et al.* Radiological findings from 81 patients with COVID-19 pneumonia in Wuhan, China: a descriptive study. *Lancet Infect Dis* 2020, 20(4):425-434.
19. Inui S, Fujikawa A, Jitsu M, Kunishima N, Watanabe S, Suzuki Y *et al.* Chest CT findings in cases from the cruise ship “Diamond Princess” with coronavirus disease 2019 (COVID-19). *Radiol Cardiothorac Imaging* 2020, 2(2).
20. Calabrese F, Pezzuto F, Fortarezza F, Hofman P, Kern I, Panizo A *et al.* Pulmonary pathology and COVID-19: lessons from autopsy. The experience of European Pulmonary Pathologists. *Virchows Arch* 2020, 477(3):359-372.
21. Tian S, Xiong Y, Liu H, Niu L, Guo J, Liao M *et al.* Pathological study of the 2019 novel coronavirus disease (COVID-19) through postmortem core biopsies. *Mod Pathol* 2020, 33(6):1007-1014.
22. Marco Francone, Franco Iafrate, Giorgio Maria Masci, Simona Coco, Francesco Cilia, Lucia Manganaro *et al.* Chest CT score in COVID-19 patients: correlation with disease severity and short-term prognosis. *Eur Radiol* 2020, 30:6808–6817
23. Ngai JC, Ko FW, Ng SS, To KW, Tong M, Hui DS. The long-term impact of severe acute respiratory syndrome on pulmonary function, exercise capacity and health status. *Respirology* 2010, 15(3):543-550.
24. Wang F, Kream RM, Stefano GB. Long-Term Respiratory and Neurological Sequelae of COVID-19. *Med Sci Monit* 2020, 26:e928996.
25. Li R, Pei S, Chen B, Song Y, Zhang T, Yang W *et al.* Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV-2). *Science* 2020, 368(6490):489-493.

26. Simpson S, Kay FU, Abbara S, Bhalla S, Chung JH, Chung M *et al.* Radiological Society of North America Expert Consensus Statement on Reporting Chest CT Findings Related to COVID-19. Endorsed by the Society of Thoracic Radiology, the American College of Radiology, and RSNA - Secondary Publication. *J Thorac Imaging* 2020, 35(4):219-227.