

APPENDICES

Right For Me: a pragmatic multi-arm cluster randomised controlled trial of two interventions for increasing shared decision-making about contraceptive methods

APPENDIX 1	Research questions and hypotheses
APPENDIX 2	Study poster and information sheet
APPENDIX 3	Right For Me prompt card
APPENDIX 4	Right For Me decision aids
APPENDIX 5	Right For Me written guidance
APPENDIX 6	Implementation slide deck: Video + Prompt Card
APPENDIX 7	Implementation slide deck: Decision Aids + Training
APPENDIX 8	Strategies for maximising enrolment, retention, and data quality
APPENDIX 9	Outcomes and measures
APPENDIX 10	Impact of patient and public involvement
APPENDIX 11	Protocol refinements and deviations
APPENDIX 12	Participant flow
APPENDIX 13	Supplementary results tables: primary outcome
APPENDIX 14	Supplementary results tables: secondary outcomes
APPENDIX 15	Supplementary results tables: process outcomes

APPENDIX 1 Research questions and hypotheses

Table A1. Research questions and hypotheses


Research Questions	Hypotheses
OBJECTIVE 1 (PRIMARY OUTCOME)	
1. Does implementing the video + prompt card increase the rate of shared decision-making about contraceptive methods compared to usual care?	Implementing the video + prompt card will increase the rate of shared decision-making about contraceptive methods compared to usual care.
2. Does implementing the decision aids + training increase the rate of shared decision-making about contraceptive methods compared to usual care?	Implementing the decision aids + training will increase the rate of shared decision-making about contraceptive methods compared to usual care.
3. Does implementing the video + prompt card and the decision aids + training result in greater increases in the rate of shared decision-making about contraceptive methods compared to usual care than implementing either of the interventions alone?	Implementing the video + prompt card and the decision aids + training will result in greater increases in the rate of shared decision-making about contraceptive methods compared to usual care than implementing the video + prompt card alone or the decision aids + training alone.
4. What patient characteristics and other factors modify the effect of implementing the interventions on the rate of shared decision-making about contraceptive methods?	This heterogeneity of treatment effects analysis is exploratory (i.e., hypothesis generating) and thus no a priori hypotheses have been developed.
OBJECTIVE 2 (SECONDARY OUTCOMES)	
<i>For each of the secondary outcomes:</i>	
5. Does implementing the video + prompt card increase or decrease (as relevant) the [rate/level] of [secondary outcome] compared to usual care?	Implementing the video + prompt card will increase the rate of conversation about contraception, satisfaction with the conversation about contraception, values concordance of intended contraceptive method(s), use of intended contraceptive method(s), adherence to contraceptive method(s) used, and satisfaction with contraceptive method(s) used; and decrease the level of decision regret, the rate of unintended pregnancy ¹ , and the rate of unwelcome pregnancy compared to usual care. Analyses pertaining to intended contraceptive method(s), intention to use a highly effective contraceptive method, contraceptive method(s) used, and use of a highly effective contraceptive method are exploratory and thus no a priori hypotheses have been developed.
6. Does implementing the decision aids + training increase or decrease (as relevant) the [rate/level] of [secondary outcome] compared to usual care?	Implementing the decision aids + training will increase the rate of conversation about contraception, satisfaction with the conversation about contraception, values concordance of intended contraceptive method(s), use of intended contraceptive method(s), adherence to contraceptive method(s) used, and satisfaction with contraceptive method(s) used; and decrease the level of decision regret, the rate of unintended pregnancy ¹ , and the rate of unwelcome pregnancy compared to usual care. Analyses pertaining to intended contraceptive method(s), intention to use a highly effective contraceptive method, contraceptive method(s) used, and use of a highly effective contraceptive method are exploratory and thus no a priori hypotheses have been developed.
7. Does implementing the video + prompt card and the decision aids + training result in greater increases or decreases (as relevant) in the [rate/level] of [secondary outcome] compared to usual care than implementing either of the interventions alone?	Implementing the video + prompt card and the decision aids + training will result in greater increases in the rate of conversation about contraception, satisfaction with the conversation about contraception, values concordance of intended contraceptive method(s), use of intended contraceptive method(s), adherence to contraceptive method(s) used, and satisfaction with contraceptive method(s) used; and greater decreases in the level of decision regret, the rate of unintended pregnancy ¹ , and the rate of unwelcome pregnancy compared to usual care than implementing the video + prompt card alone or the decision aids + training alone. Analyses pertaining to intended contraceptive method(s), intention to use a highly effective contraceptive method, contraceptive method(s) used, and use of a highly effective contraceptive method are exploratory and thus no a priori hypotheses have been developed.
OBJECTIVE 3 (PROCESS OUTCOMES)	
8. Of participants receiving care in a trial arm implementing the video + prompt card, what proportion report having watched the whole video?	Of participants receiving care in a trial arm implementing the video + prompt card, at least 70% will report having watched the whole video.
9. Of participants receiving care in a trial arm implementing the video + prompt card, what proportion report having received the prompt card?	Of participants receiving care in a trial arm implementing the video + prompt card, at least 70% will report having received the prompt card.

- | | |
|---|--|
| <p>10. Of participants receiving care in a trial arm implementing the decision aids + training, what proportion report having used a decision aid together with a health care provider?</p> | <p>Of participants receiving care in a trial arm implementing the decision aids + training, at least 70% will report having used a decision aid together with a health care provider.</p> |
| <p>11. Is the proportion of participants who report having watched the whole video higher among those receiving care in a trial arm implementing the video + prompt card and the decision aids + training than in a trial arm implementing the video + prompt card alone?</p> | <p>The proportion of participants who report having watched the whole video will be higher among those receiving care in a trial arm implementing the video + prompt card and the decision aids + training than in a trial arm implementing the video + prompt card alone.</p> |
| <p>12. Is the proportion of participants who report having received the prompt card higher among those receiving care in a trial arm implementing the video + prompt card and the decision aids + training than in a trial arm implementing the video + prompt card alone?</p> | <p>The proportion of participants who report having received the prompt card will be higher among those receiving care in a trial arm implementing the video + prompt card and the decision aids + training than in a trial arm implementing the video + prompt card alone.</p> |
| <p>13. Is the proportion of participants who report having used a decision aid together with a health care provider higher among those receiving care in a trial arm implementing the video + prompt card and the decision aids + training than in a trial arm implementing the decision aids + training alone?</p> | <p>The proportion of participants who report having used a decision aid together with a health care provider will be higher among those receiving care in a trial arm implementing the video + prompt card and the decision aids + training than in a trial arm implementing the decision aids + training alone.</p> |
| <p>14. What proportion of participants who report having watched the whole video would recommend it to a friend?</p> | <p>A majority of participants who report having watched the whole video would recommend it to a friend.</p> |
| <p>15. What proportion of participants who report having received the prompt card would recommend it to a friend?</p> | <p>A majority of participants who report having received the prompt card would recommend it to a friend.</p> |
| <p>16. What proportion of participants who report having used a decision aid together with a health care provider would recommend it to a friend?</p> | <p>A majority of participants who report having used a decision aid together with a health care provider would recommend it to a friend.</p> |
| <p>17. Is the proportion of participants who would recommend the video to a friend higher among those receiving care in a trial arm implementing the video + prompt card and the decision aids + training than in a trial arm implementing the video + prompt card alone?</p> | <p>The proportion of participants who would recommend the video to a friend will be higher among those receiving care in a trial arm implementing the video + prompt card and the decision aids + training than in a trial arm implementing the video + prompt card alone.</p> |
| <p>18. Is the proportion of participants who would recommend the prompt card to a friend higher among those receiving care in a trial arm implementing the video + prompt card and the decision aids + training than in a trial arm implementing the video + prompt card alone?</p> | <p>The proportion of participants who would recommend the prompt card to a friend will be higher among those receiving care in a trial arm implementing the video + prompt card and the decision aids + training than in a trial arm implementing the video + prompt card alone.</p> |
| <p>19. Is the proportion of participants who would recommend the decision aids to a friend higher among those receiving care in a trial arm implementing the video + prompt card and the decision aids + training than in a trial arm implementing the decision aids + training alone?</p> | <p>The proportion of participants who would recommend the decision aids to a friend will be higher among those receiving care in a trial arm implementing the video + prompt card and the decision aids + training than in a trial arm implementing the decision aids + training alone.</p> |

Notes. ¹Unintended pregnancy was measured based both on pregnancy timing preferences immediately before conception and pregnancy seeking immediately before conception.

APPENDIX 2

Study poster and information sheet



right for me

This clinic is involved in the Right For Me study
Participating involves doing a brief health care survey

You can participate if you:

had a **health care visit** at this clinic today

can read and write **English** or **Spanish**

are **15 to 49** years old

were assigned **female** sex at birth

have **not participated before**

Participants receive a \$10 gift card

To participate, please ask a clinic staff member

You can participate in
this study if you:

had a **health care visit**
at this clinic today

can read and write
English or **Spanish**

are **15 to 49** years old

were assigned
female sex at birth

have **not**
participated before

We will check that you meet these
criteria at the start of the study

Who can I contact?

If you have questions or concerns about this
study, you can contact:

Ruth Manski

Right For Me Project Manager
Dartmouth College
(e) ruth.manski@dartmouth.edu
(t) 603 653 0879
(during business hours)

Rachel Thompson

Right For Me Principal Investigator
Dartmouth College
(e) rachel.thompson@dartmouth.edu
(t) 603 653 0860
(during business hours)

If you have questions, concerns, complaints, or
suggestions about research at Dartmouth, you
can call:


Office of the Committee for the Protection of Human Subjects

Dartmouth College
(t) 603 646 6482
(during business hours)

THE
Dartmouth
INSTITUTE

FOR HEALTH POLICY & CLINICAL PRACTICE

This study is funded through a Patient-Centered Outcomes
Research Institute (PCORI) Award (CDR-1403-12221).



right for me

Study Information

Right For Me is a research study led
by Dartmouth College. Your clinic is
involved in the study.

This handout gives information about the Right For Me study.

If there's something you don't understand, please ask questions.

What is the purpose?

The purpose of this study is to learn new things about people's experiences of health care and how improvements can be made to help people get the health care they want and need.

What is involved?

Participating involves doing a 5-minute survey on a tablet computer before you leave the clinic. The survey asks about your health care today. If relevant, the survey also asks about your experiences and views about birth control.

A small group of people will be invited to do two more 5-minute surveys at home later. The surveys ask more questions about experiences and views about birth control. The first survey will be sent in 4 weeks. The second survey will be sent in 6 months.

Do I have to participate?

No, participating is voluntary. Your choice about whether to participate will not affect the quality of your health care. You can also stop participating at any time. If you stop participating, the information we have already collected will continue to be used.

How many people will participate?

This is a large and important study. Altogether, 16 clinics from New England are involved. We expect that 390 people from your clinic and 5,850 people from other clinics will participate.

Will I be paid?

You will receive a \$10 gift card for Amazon.com for every survey you complete. If you are aged under 20, you will need an email address to receive the gift card. If you are aged 20 or older, you will need either an email address or a mailing address. The gift card will be sent no more than two weeks after you complete each survey.

If the information we collect in the study is used to develop a product sold for a profit, you will not receive a share of the profits.

What are the benefits?

You may enjoy completing the survey and find it valuable to give feedback on your health care. People receiving health care in the future may also benefit from what we learn from the study.

What are the risks?

There are risks involved in most life activities. We think that the risks involved in participating in this study are minimal. You may find some of the survey questions personal. You can skip any questions you don't want to answer and remain in the study.

How will you use the information?

The information we collect will be stored for an indefinite period of time.

We will use the information we collect to learn new things about people's experiences of health care and how improvements can be made. We will share what we learn with participants, researchers, health professionals, people who design health care, and the public.

We may also share the information we collect with other people so that they can confirm what we learned or do new studies on related topics.

How will you protect my privacy?

Protecting your privacy is important to us.

Some of the information we collect could identify you. Information that could identify you will be treated as confidential.

None of the people that provide health care in clinics during the study will be allowed to see information that could identify you.

The people that will be allowed to see information that could identify you are: (1) the study principal investigator, project manager, and data manager, and (2) people who work in information security, information management, and translation at Dartmouth College and elsewhere. It is also possible for a court or government official to order the release of study data including information about you.

If you choose to receive your gift card via email, we will give your email address to a company so they can email it to you.

We will store information that could identify you in password-protected computer files and locked filing cabinets. We will only ever transfer this information using secure methods.

We will remove or change all information that could identify you before we share what we learn with others. We will also remove or change all information that could identify you before we share the information we collect with others.

How can I find out what you learn?

You can find out what we learn from the study by visiting the study website (www.rightforme.org) after January 2018.

APPENDIX 3


Right For Me prompt card

Try asking these questions today

What are my options?

What are the possible pros and cons of those options?

How likely are each of those pros and cons to happen to me?



right for me

APPENDIX 4

Right For Me decision aids

Types of Birth Control Methods

This decision aid is to help you and your health care provider talk about methods of birth control and choose what's right for you. Most people can safely use these methods. Your health care provider can tell you whether these methods are safe for you.



	LONG-ACTING	SHORT-ACTING	BARRIER	NATURAL*	PERMANENT	EMERGENCY
What are they?	Birth control methods that are placed by a health care provider and last between 3 and 10 years	Birth control methods that are used every day, every week, every 4 weeks, or every 13 weeks	Birth control methods that are used every time you have sex	Birth control methods that do not involve any hormones or devices	Birth control methods that involve a procedure to close off the tubes that carry eggs or sperm	Birth control methods that are used after unprotected sex
What are the options?	<ul style="list-style-type: none"> • Implant • Hormonal IUD • Copper IUD 	<ul style="list-style-type: none"> • Injection • Progestin Pill • Combined Pill • Patch • Ring 	<ul style="list-style-type: none"> • Male Condom • Female Condom • Spermicide • Sponge • Cervical Cap • Diaphragm 	<ul style="list-style-type: none"> • Withdrawal Method <p><i>Fertility Awareness Methods:</i></p> <ul style="list-style-type: none"> • Standard Days Method® • TwoDay Method® • Ovulation Method • Symptothermal Method 	<p><i>Female Sterilization:</i></p> <ul style="list-style-type: none"> • By Laparoscopy • By Minilaparotomy • By Hysteroscopy <p><i>Male Sterilization:</i></p> <ul style="list-style-type: none"> • Vasectomy 	<ul style="list-style-type: none"> • Copper IUD • Ulipristal Pill • Progestin Emergency Pill • Combined Pill
Who might choose them?	<p>People who want or are comfortable with:</p> <ul style="list-style-type: none"> • Almost no chance of pregnancy (fewer than 1 in 100 people become pregnant in the first year) • A method they can almost forget about • A procedure to start and stop using the method 	<p>People who want or are comfortable with:</p> <ul style="list-style-type: none"> • Some chance of pregnancy (6 to 9 in 100 people become pregnant in the first year) • A method they need to remember • A hormonal method • A method they can stop without a health care visit 	<p>People who want or are comfortable with:</p> <ul style="list-style-type: none"> • A higher chance of pregnancy (12 to 29 in 100 people become pregnant in the first year) • A method they need to remember • A non-hormonal method • Protection against sexually transmitted infections (STIs) (not all methods offer this) 	<p>People who want or are comfortable with:</p> <ul style="list-style-type: none"> • A higher chance of pregnancy (22 to 24 in 100 people become pregnant in the first year) • A method they need to remember • A non-hormonal method • A method that does not involve birth control devices 	<p>People who want or are comfortable with:</p> <ul style="list-style-type: none"> • Almost no chance of pregnancy (fewer than 1 in 100 people become pregnant in the first year) • A method they can forget about • A procedure • Never becoming pregnant in the future 	<p>People who:</p> <ul style="list-style-type: none"> • Have had unprotected sex and don't want to become pregnant

*The Lactational Amenorrhea Method is another natural birth control method that may be used by some people who are breastfeeding. Your health care provider can tell you about this method.

Long-Acting Reversible Birth Control Methods

This decision aid is to help you and your health care provider talk about methods of birth control and choose what's right for you. Most people can safely use these methods. Your health care provider can tell you whether these methods are safe for you.



	IMPLANT	HORMONAL IUD	COPPER IUD
How is it used?	A health care provider puts a flexible device the size of a matchstick under the skin of your arm	A health care provider puts a small, T-shaped, plastic device in your uterus	A health care provider puts a small, T-shaped plastic and copper device in your uterus
How often?	Every 3 years	Every 3 to 5 years (depends on the brand)	Every 10 years
How does it work?	Releases progestin	Releases progestin	Releases copper
When does it start working?	Immediately or after 7 days (depends on when you get the implant)	Immediately or after 7 days (depends on when you get the IUD)	Immediately
How many people become pregnant in the first year?			
Not always following the instructions:	Fewer than 1 in 100 people ○○○○○○○○○○	Fewer than 1 in 100 people ○○○○○○○○○○	Fewer than 1 in 100 people ○○○○○○○○○○
Always following the instructions:	Fewer than 1 in 100 people ○○○○○○○○○○	Fewer than 1 in 100 people ○○○○○○○○○○	Fewer than 1 in 100 people ○○○○○○○○○○
What are some of the side effects?			
Unscheduled spotting or bleeding?	Possible, may or may not improve over time	Possible, usually improves over time	Possible, usually improves over time
Heavy or prolonged bleeding?	Possible		Possible, usually improves over time
More painful periods?			Possible, usually improves over time
Fewer or no periods?	Possible	Possible	
Other side effects (e.g., nausea, headaches, breast tenderness, mood changes or depression)?	Possible	Possible	
Device comes out?		Possible	Possible
Does it protect against sexually transmitted infections (STIs)?	No	No	No

Short-Acting Reversible Birth Control Methods

This decision aid is to help you and your health care provider talk about methods of birth control and choose what's right for you. Most people can safely use these methods. Your health care provider can tell you whether these methods are safe for you.



	INJECTION	PROGESTIN PILL	COMBINED PILL	PATCH	RING
How is it used?	A health care provider gives you a shot in your arm or buttock	You swallow a small pill	You swallow a small pill	You stick a small, thin patch on your body	You put a flexible, plastic ring in your vagina
How often?	Every 13 weeks	Every day at the same time	Every day at the same time	Every week	Every 4 weeks
How does it work?	Releases progestin	Releases progestin	Releases progestin and estrogen	Releases progestin and estrogen	Releases progestin and estrogen
When does it start working?	Immediately or after 7 days (depends on when you get the first shot)	Immediately or after 2 days (depends on when you start taking the pill)	Immediately or after 7 days (depends on when you start taking the pill)	Immediately or after 7 days (depends on when you start using the patch)	Immediately or after 7 days (depends on when you start using the ring)
How many people become pregnant in the first year?					
Not always following the instructions:	6 in 100 people ●○○○○○○○○○	9 in 100 people ●○○○○○○○○○	9 in 100 people ●○○○○○○○○○	9 in 100 people ●○○○○○○○○○	9 in 100 people ●○○○○○○○○○
Always following the instructions:	Fewer than 1 in 100 people ○○○○○○○○○	Fewer than 1 in 100 people ○○○○○○○○○	Fewer than 1 in 100 people ○○○○○○○○○	Fewer than 1 in 100 people ○○○○○○○○○	Fewer than 1 in 100 people ○○○○○○○○○
What are some of the side effects?					
Unscheduled spotting or bleeding?	Possible	Possible	Possible, usually improves over time	Possible, usually improves over time	Possible, usually improves over time
Heavy or prolonged bleeding?	Possible	Possible			
Fewer or no periods?	Possible	Possible			
Other side effects (e.g., nausea, headaches, breast tenderness)?		Possible	Possible	Possible	Possible
Bone loss?	Possible, usually improves when you stop getting the shot				
Skin irritation?				Possible	
Vaginal irritation or discharge?					Possible
Does it protect against sexually transmitted infections (STIs)?	No	No	No	No	No

Barrier Birth Control Methods

This decision aid is to help you and your health care provider talk about methods of birth control and choose what's right for you. Most people can safely use these methods. Your health care provider can tell you whether these methods are safe for you.



	MALE CONDOM	FEMALE CONDOM	SPERMICIDE	SPONGE	CERVICAL CAP (with spermicide)	DIAPHRAGM (with spermicide)
How is it used?	A thin latex or polyurethane sleeve is put on the erect penis before sex	You put a thin sleeve in your vagina before sex	You put a foam, cream, gel, film, or suppository in your vagina before sex	You put a round foam device (that already contains spermicide) in your vagina before sex	You put a silicone-rubber device in your vagina, with a spermicide, before sex	You put a latex or silicone device in your vagina, with a spermicide, before sex
How often?	Every time you have sex	Every time you have sex	Every time you have sex	Every time you have sex	Every time you have sex	Every time you have sex
How does it work?	Prevents sperm from reaching an egg	Prevents sperm from reaching an egg	Prevents sperm from reaching an egg	Prevents sperm from reaching an egg	Prevents sperm from reaching an egg	Prevents sperm from reaching an egg
When does it start working?	Immediately	Immediately	Depends on the brand	Immediately	Immediately	Immediately
How many people become pregnant in the first year?						
Not always following the instructions:	18 in 100 people ●●○○○○○○○○	21 in 100 people ●●●○○○○○○○	28 in 100 people ●●●○○○○○○○	12 in 100 people (if never given birth) ●●○○○○○○○○ 24 in 100 people (if given birth) ●●●○○○○○○○	14 in 100 people (if never given birth vaginally) ●●○○○○○○○○ 29 in 100 people (if given birth vaginally) ●●●○○○○○○○	12 in 100 people ●●○○○○○○○○
Always following the instructions:	2 in 100 people ●○○○○○○○○○	5 in 100 people ●○○○○○○○○○	18 in 100 people ●●○○○○○○○○	9 in 100 people (if never given birth) ●○○○○○○○○○ 20 in 100 people (if given birth) ●●●○○○○○○○	<i>No information available</i>	6 in 100 people ●○○○○○○○○○
What are some of the side effects?						
Allergic reaction?	Possible	Possible	Possible	Possible	Possible	Possible
Vaginal symptoms (e.g., odor, irritation, infection, vaginitis)?			Possible	Possible	Possible	Possible
Toxic shock syndrome?				Possible	Possible	Possible
Does it protect against sexually transmitted infections (STIs)?	Provides the best protection	Provides some protection	No	No	No	No

Natural Birth Control Methods

This decision aid is to help you and your health care provider talk about methods of birth control and choose what's right for you. Most people can safely use these methods. Your health care provider can tell you whether these methods are safe for you.



	WITHDRAWAL METHOD	FERTILITY AWARENESS METHODS:			
		STANDARD DAYS METHOD®	TWODAY METHOD®	OVULATION METHOD	SYMPTOTHERMAL METHOD
How is it used?	During sex, the penis is removed from your vagina and away from your genitals before ejaculation	You monitor the days of your menstrual cycle. You avoid sex on days 8 through 19 of your cycle.	You monitor your cervical secretions. You avoid sex on days you notice secretions and for one day after.	You monitor your cervical secretions. You use a set of rules to know what days to avoid sex.	You monitor your cervical secretions and your body temperature. You use a set of rules to know what days to avoid sex.
How often?	Every time you have sex	Every day	Every day (checking your secretions two or more times a day)	Every day	Every day
How does it work?	Prevents sperm from reaching an egg	Prevents sperm from reaching an egg when there is a higher chance of pregnancy	Prevents sperm from reaching an egg when there is a higher chance of pregnancy	Prevents sperm from reaching an egg when there is a higher chance of pregnancy	Prevents sperm from reaching an egg when there is a higher chance of pregnancy
When does it start working?	Immediately	You can start monitoring the days of your menstrual cycle anytime. It may take time before it is a 'safe day' to have sex.	You can start monitoring your cervical secretions anytime. It may take time before it is a 'safe day' to have sex.	You can start monitoring your cervical secretions anytime. It may take time before it is a 'safe day' to have sex.	You can start monitoring your cervical secretions and your body temperature anytime. It may take time before it is a 'safe day' to have sex.
How many people become pregnant in the first year?					
Not always following the instructions:	22 in 100 people ●●●○○○○○○○	24 in 100 people ●●●○○○○○○○	24 in 100 people ●●●○○○○○○○	24 in 100 people ●●●○○○○○○○	24 in 100 people ●●●○○○○○○○
Always following the instructions:	4 in 100 people ●○○○○○○○○○	5 in 100 people ●○○○○○○○○○	4 in 100 people ●○○○○○○○○○	3 in 100 people ●○○○○○○○○○	Fewer than 1 in 100 people ○○○○○○○○○
Are there side effects?	No	No	No	No	No
Does it protect against sexually transmitted infections (STIs)?	No	No	No	No	No

Note: The Lactational Amenorrhea Method is another natural birth control method that may be used by some people who are breastfeeding. Your health care provider can tell you about this method.

Permanent Birth Control Methods

This decision aid is to help you and your health care provider talk about methods of birth control and choose what's right for you. Most people can safely use these methods. Your health care provider can tell you whether these methods are safe for you.




	FEMALE STERILIZATION:			MALE STERILIZATION:
	BY LAPAROSCOPY	BY MINILAPAROTOMY	BY HYSTEROSCOPY	VASECTOMY
How is it used?	A health care provider uses instruments inserted through one or two small incisions in your abdomen to be able to see and close your fallopian tubes	A health care provider makes an incision in your abdomen and moves your fallopian tubes up so they can be seen. The provider then closes your tubes.	A health care provider inserts an instrument through your vagina and places small devices in your fallopian tubes. Scar tissue forms around the devices and closes your tubes.	A health care provider makes one or two incisions or a small puncture in the skin of the scrotum and closes the tubes that carry sperm
How often?	Once	Once	Once	Once
How does it work?	Prevents sperm from reaching an egg	Prevents sperm from reaching an egg	Prevents sperm from reaching an egg	Prevents sperm from being released
When does it start working?	Immediately	Immediately	After 3 months, when an X-ray shows that your fallopian tubes are closed	After 2 to 4 months, when a test shows that there are no longer sperm in the semen
How many people become pregnant in the first year?				
Not always following the instructions:	Fewer than 1 in 100 people ○○○○○○○○○○	Fewer than 1 in 100 people ○○○○○○○○○○	Fewer than 1 in 100 people ○○○○○○○○○○	Fewer than 1 in 100 people ○○○○○○○○○○
Always following the instructions:	Fewer than 1 in 100 people ○○○○○○○○○○	Fewer than 1 in 100 people ○○○○○○○○○○	Fewer than 1 in 100 people ○○○○○○○○○○	Fewer than 1 in 100 people ○○○○○○○○○○
What are some of the side effects?				
Abdominal cramps or other pain following the procedure?	Possible	Possible	Possible	Possible
Dizziness, nausea, vomiting, bleeding, or other symptoms following the procedure?	Possible	Possible	Possible	Possible
Minor complication (e.g., infection)?	Possible	Possible	Possible	Possible
Major complication (e.g., injury requiring surgery)?	Possible	Possible	Possible	Possible
Pain that continues for some time?	Possible	Possible	Possible	Possible
Does it protect against sexually transmitted infections (STIs)?	No	No	No	No

Emergency Birth Control Methods

This decision aid is to help you and your health care provider talk about methods of birth control and choose what's right for you. Most people can safely use these methods. Your health care provider can tell you whether these methods are safe for you.



	COPPER IUD	ULIPRISTAL PILL	PROGESTIN EMERGENCY PILL	COMBINED PILL
How is it used?	A health care provider puts a small, T-shaped plastic and copper device in your uterus. You can then use this as regular birth control for up to 10 years.	You swallow one pill	You swallow one pill or you swallow two pills, 12 hours apart (depends on the brand)	You swallow a specific number of combined birth control pills and then repeat 12 hours later. The number of pills needed depends on the brand.
When?	Within 5 days after unprotected sex	As soon as possible within 5 days after unprotected sex	As soon as possible within 5 days after unprotected sex	As soon as possible within 5 days after unprotected sex
How does it work?	Releases copper to prevent a pregnancy from occurring	Releases ulipristal acetate to prevent a pregnancy from occurring	Releases progestin to prevent a pregnancy from occurring	Releases progestin and estrogen to prevent a pregnancy from occurring
How effective is it at preventing pregnancy?	Highly effective – the most effective method	The most effective method after the Copper IUD	As effective as the Ulipristal Pill if taken within 3 days after unprotected sex Less effective than the Ulipristal Pill if taken 3 to 5 days after unprotected sex	The least effective method
What are some of the side effects?				
Temporary bleeding changes (e.g., irregular bleeding or spotting, next period not at the expected time)?		Possible	Possible	Possible
Other temporary side effects (e.g., headaches, nausea or vomiting, breast tenderness, abdominal pain, dizziness, fatigue)?		Possible	Possible	Possible
Non-temporary bleeding changes or other side effects?	Possible (More information on the 'Long-Acting Reversible Birth Control Methods' decision aid)			
Does it protect against sexually transmitted infections (STIs)?	No	No	No	No



right for me

Frequently Asked Questions

What is shared decision-making?

The process of health professionals and patients making health decisions together.

What are the Right For Me Decision Aids?

A set of seven one-page tools that compare birth control methods. They are intended to be used by health professionals during the health care visit. The Decision Aids are available in English and Spanish.

Who developed the Decision Aids?

Researchers at Dartmouth College and their project partners, including patients, health professionals, and other stakeholders. Decision Aid authors are listed in a Supporting Document, along with a conflict of interest disclosure statement. There is a link to the Supporting Document on each Decision Aid.

Where did the information come from?

Reviews of scientific evidence, relevant national and international guidelines, and existing patient resources. Information sources are listed in a Supporting Document. There is a link to the Supporting Document on each Decision Aid.

Where did the questions and format come from?

A survey of over 600 women and health professionals⁴, focus groups with patients in three states, and best practice guidelines for decision aid development.

Who can use the Decision Aids?

Any health professional who provides information or counseling about contraception to patients receiving health care in your clinic.

With which patients can the Decision Aids be used?

Any English- or Spanish-speaking patients receiving health care in your clinic. You may wish to use the Decision Aids with patients who are considering contraception for the first time, already using a contraceptive method, or interested in changing methods. The reading level of each Decision Aid is 8th grade or less.

Are the Decision Aids intended to be used by the patient alone?

No. The decision aids are a framework for a conversation with a health professional. Patients may need medical terminology explained. You may also wish to share further information, such as non-contraceptive benefits, other side effects, usage instructions, mechanism of actions, brand names, or off-label use.

How can I use the Decision Aids with patients?

A demonstration of how to use the Decision Aids is provided in the Decision Aid Training Video. As explained in the Training Video, following three steps may help:

- ① Explain It
- ② Give It
- ③ Use It

How do I know which Decision Aids to use?

You are encouraged to use as many or as few Decision Aids as makes sense for the patient. The 'Types of Birth Control Methods' Decision Aid may be helpful to use with patients without a lot of background knowledge before narrowing down to a more specific Decision Aid.

What if some contraceptive methods are unsafe for a patient?

The U.S. Medical Eligibility Criteria for Contraceptive Use and published updates¹⁻³ provide guidance on who can safely use each method of contraception and may be used to facilitate the provision of individualized information on method safety.

Can I write on the Decision Aids?

Of course. You are welcome to add or change content to suit the individual patient.

What does 'not always following the instructions' mean?

This is the terminology adopted to convey 'typical-use effectiveness' after consultation with patients.

What does 'always following the instructions' mean?

This is the terminology we adopted to convey 'perfect-use effectiveness' after consultation with patients.

Why are some side effects listed and not others?

Side effects routinely cited in the evidence, guidelines, and existing patient resources reviewed were prioritized for inclusion in the Decision Aids. Due to space constraints, the list of side effects is not exhaustive.

Under side effects, what does a blank cell mean?


This signifies that the side effect was not attributed to the contraceptive method in the evidence, guidelines, and existing patient resources reviewed.

References

1. Centers for Disease Control and Prevention. U.S. Selected Practice Recommendations for Contraceptive Use, 2013. *MMWR Recomm Rep*. 2013;62(5):1-60.
<http://www.cdc.gov/reproductivehealth/unintendedpregnancy/usspr.htm>.
2. Centers for Disease Control and Prevention. Update to CDC's U.S. Medical Eligibility Criteria for Contraceptive Use, 2010: Revised Recommendations for the Use of Hormonal Contraception Among Women at High Risk for HIV Infection or Infected with HIV. *MMWR*. 2012;61(24):449-452.
<http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6124a4.htm>.
3. Centers for Disease Control and Prevention. Update to CDC's U.S. Medical Eligibility Criteria for Contraceptive Use, 2010: Revised Recommendations for the Use of Contraceptive Methods During the Postpartum Period. *MMWR*. 2011;60(26):878-883. <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6026a3.htm>.
4. Donnelly KZ, Foster TC, Thompson R. What matters most? The content and concordance of patients' and providers' information priorities for contraceptive decision making. *Contraception*. 2014;90(3):280-287.
doi:10.1016/j.contraception.2014.04.012.

APPENDIX 6

Implementation slide deck: Video + Prompt Card



right for me

Video + Prompt Card

Introduction

This slide deck is intended to support people in your clinic (administrative staff, health professionals, and others) to implement the Right For Me Video + Prompt Card.

About

The Right For Me Video is a three-minute video intended to be viewed by patients immediately before the health care visit. It is viewed on an iPad with headphones. The Video is available in English and Spanish, with and without on-screen captions.

The Right For Me Prompt Card is a small card intended to be taken by patients immediately before the health care visit. The Prompt Card is available in English and Spanish.

Objective

The aim of the Video is to enhance patients' motivation, skills, and self-efficacy to ask health professionals three specific questions:

- (1) What are my options?
- (2) What are the pros and cons of those options?
- (3) How likely are those pros and cons to happen to me?

The aim of the Prompt Card is to remind patients of the three questions presented in the video.

Audience

The target audience for the Video + Prompt Card is any patient waiting to see a health professional in your clinic (for example, while in the waiting room or after the rooming-in process).

Development

The Video + Prompt Card were developed by researchers at Dartmouth College and their project partners, based on previous research and extensive consultation with patients, health professionals, and other stakeholders.

Preview

You can preview the Video + Prompt Card by:

1. Visiting www.rightforme.org
2. Navigating to 'For Clinics' and then 'Video + Prompt Card'
3. Entering the clinic username and password

Your Project Contact will give you the clinic username and password.
Please do not share this information with others outside your clinic.

Implementation

Implementing the Video + Prompt Card is a team effort. The tasks involved are explained below and on the remaining slides.

What?	Who?
1. Supplying the materials	The Right For Me project team
2. Facilitating patients viewing the Video	Administrative staff, health professionals, and/or others
3. Facilitating patients taking the Prompt Card	Administrative staff, health professionals, and/or others
4. Other support tasks	Administrative staff, health professionals, and/or others

1. Supplying the materials

The Right For Me project team

Right For Me Video

We have supplied your clinic with:

- 2 iPads programmed to view the Video
- 2 sets of headphones
- Cleaning wipes



Right For Me Prompt Card

We have supplied your clinic with:

- 3 display stands of English Prompt Cards
- 1 display stand of Spanish Prompt Cards
- Extra refills of Prompt Cards



2. Facilitating patients viewing the Video

For administrative staff, health professionals, and/or others

Facilitating Patients Viewing the Video

We encourage you to facilitate patients viewing the Video in a way that works for your clinic. Things you may wish to think about include:


- Who will hand the iPads and headphones to patients?
- When and where will you hand the iPads and headphones to patients?
- What will you say to patients about the Video?
- Who will collect the iPads and headphones from patients?

The next slides provide some practice guidance that may be helpful.

Guidance for Clinic Staff

Before handing an iPad to a patient, open the cover. You should see four videos displayed on the screen.

If you see that a video is playing, ① tap anywhere on the screen and ② tap 'Done' in the top left corner. Close the cover.



Guidance for Patients

Basic instructions for patients are provided on the iPad cover in English and Spanish.



Guidance for Patients



Patients may choose to watch the Video in English or Spanish, with or without on-screen captions.

Patients should put on the headphones and ① tap the video they want to watch. It will play automatically.



Guidance for Patients

If a patient wishes to exit a video before it has finished playing, they should ① tap anywhere on the screen and ② tap 'Done' in the top left corner.



Guidance for Patients

If a patient wishes to increase or decrease the volume of a video, they should ① press the long up or down button on the right side of the iPad.



3. Facilitating patients taking the Prompt Card

For administrative staff, health professionals, and/or others

Facilitating Patients Taking the Prompt Card

We encourage you to facilitate patients taking the Prompt Card in a way that works for your clinic. Things you may wish to think about include:

- Will you hand the patient a Prompt Card when you give them the iPad?
- Will you put the Prompt Cards out on display and refer patients to them?
- What will you say to patients about the Prompt Card?

Patients may take the Prompt Card home with them to keep.

4. Other support tasks

For administrative staff, health professionals, and/or others

iPad Management

Please ensure the iPads and headphones are stored securely in the clinic when not in use.

Please charge the iPads as often as needed, including every night. Plug the iPad charger into a wall outlet (not a computer).

iPad Management

Please check periodically that the iPads remain locked into the Video application. To do this, press the iPad home button at the bottom of the iPad. When you press this, the screen should not change. If the screen changes, please tell your Project Contact.


Prompt Card Management

Please check periodically to make sure you have enough Prompt Cards. If you are running low on Prompt Cards in English and/or Spanish, please tell your Project Contact. They will ask us to send extra Prompt Cards to your clinic.

Thank you.
Good luck!

APPENDIX 7

Implementation slide deck: Decision Aids + Training



right for me

Decision Aids + Training

Introduction

This slide deck is intended to support people in your clinic (administrative staff, health professionals, and others) to implement the Right For Me Decision Aids + Training.

About

The Right For Me Decision Aids are a set of seven one-page decision aids on birth control methods. They are intended to be used by health professionals during the health care visit. The Decision Aids are available in English and Spanish.

The Right For Me Decision Aid Training consists of a four-minute Video and Frequently Asked Questions intended to be reviewed by health professionals before beginning to use the Decision Aids.

Objective

The aim of the Decision Aids is to support health professionals to facilitate shared decision-making about birth control methods with patients in the health care visit.

The aim of the Training Video and Frequently Asked Questions is to enhance health professionals' motivation, skills, and self-efficacy to use the Decision Aids.

Audience

The target audience for the Decision Aids + Training is any health professional who provides information or counseling about contraception to patients receiving health care in your clinic.

Development

The Decision Aids + Training were developed by researchers at Dartmouth College and their project partners, based on previous research and extensive consultation with patients, health professionals, and other stakeholders.

Implementation

Implementing the Decision Aids + Training is a team effort. The tasks involved are explained below and on the remaining slides.

What?	Who?
1. Supplying the materials	The Right For Me project team
2. Taking the Training	Health professionals
3. Using the Decision Aids	Health professionals
4. Other support tasks	Administrative staff, health professionals, and/or others

1. Supplying the materials

The Right For Me project team

Decision Aids

We have supplied your clinic with:

- Tear-pads of the seven different Decision Aids in English and Spanish (each set is bound with a rubber band)
- 1 desktop or wall-mounted display stand for each exam room

Types of Birth Control Methods

This decision aid is to help you and your health care provider talk about methods of birth control and choose what's right for you. Most people can safely use these methods. Your health care provider can tell you whether these methods are safe for you.

What are they?	LONG-ACTING	SHORT-ACTING	BARRIER	NATURAL	PERMANENT	EMERGENCY
What are the options?	<ul style="list-style-type: none"> • Implant • Hormonal IUD • Copper IUD 	<ul style="list-style-type: none"> • Injection • Progestin Pill • Combined Pill • Patch • Ring 	<ul style="list-style-type: none"> • Male Condom • Female Condom • Spermicide • Sponge • Cervical Cap • Diaphragm 	<ul style="list-style-type: none"> • Withdrawal Method • Lactation Amenorrhea Method • Standard Days Method • TwoDay Method • Ovulation Method • Symptothermal Method 	<ul style="list-style-type: none"> • Female Sterilization • By Laparoscopy • By Hysteroscopy • Vasectomy 	<ul style="list-style-type: none"> • Copper IUD • Ulipristal Pill • Progestin Emergency Pill • Combined Pill
Who might choose them?	<ul style="list-style-type: none"> • People who want or are comfortable with: • Almost no chance of pregnancy • Fewer than 1 in 100 people become pregnant in the first year • A method they can almost forget about • A procedure to start and stop using the method 	<ul style="list-style-type: none"> • People who want or are comfortable with: • Some chance of pregnancy • 18 to 20 people become pregnant in the first year • A method they need to remember • A hormonal method • A method they can stop without a health care visit 	<ul style="list-style-type: none"> • People who want or are comfortable with: • A higher chance of pregnancy • 18 to 20 people become pregnant in the first year • A method they need to remember • A non-hormonal method • Protection against sexually transmitted infections (STIs) not all methods offer this 	<ul style="list-style-type: none"> • People who want or are comfortable with: • A higher chance of pregnancy • 18 to 20 people become pregnant in the first year • A method they need to remember • A non-hormonal method • A method that does not remove birth control devices 	<ul style="list-style-type: none"> • People who want or are comfortable with: • Almost no chance of pregnancy • Fewer than 1 in 100 people become pregnant in the first year • A method they can forget about • A procedure 	<ul style="list-style-type: none"> • People who: • Have had unprotected sex and don't want to become pregnant

*The Lactational Amenorrhea Method is another natural birth control method that may be used by some people who are breastfeeding. Your health care provider can tell you about this method.

For more information, including authors, information sources, and terms of use, see www.rightfor.me/decision © 2016 Trustees of Dartmouth College | Version 1.0

Training

We have supplied your clinic with online access to:

- The Training Video
- The Frequently Asked Questions

2. Taking the Training

For health professionals

Taking the Training

You can take the Training as often as you wish by:

1. Visiting www.rightforme.org
2. Navigating to 'For Clinics' and then 'Decision Aids + Training'
3. Entering the clinic username and password

Your Project Contact will give you the clinic username and password.
Please do not share this information with others outside your clinic.

3. Using the Decision Aids

For health professionals

Using the Decision Aids

All guidance on using the Decision Aids is provided in the Training Video and Frequently Asked Questions.

4. Other support tasks

For administrative staff, health professionals, and/or others

Decision Aid Management

Please place tear-pads of the seven different Decision Aids on display in each exam room.


Please check periodically to make sure that each exam room has enough Decision Aids. If you are running low on Decision Aids in English and/or Spanish, please tell your Project Contact. They will ask us to send extra Decision Aids to your clinic.

Thank you.
Good luck!

Table A2. Strategies for maximising enrolment, retention, and data quality

Goal	Strategies
Maximise enrolment	<ul style="list-style-type: none"> • A study name, branding, and engaging recruitment materials designed with patient input • No requirement for documented informed consent or parental consent to participate • Option to participate in the study without a commitment to complete all three surveys • Option to participate in the study without providing name and contact details • Face-to-face training in approved recruitment and data collection processes for clinic staff • Regular feedback to clinic staff on the number of participants enrolling in the study • Participant compensation of a \$10 gift card for completing the T1 survey
Maximise retention	<ul style="list-style-type: none"> • Multiple completion modes for the T2 and T3 surveys for participants aged 20 years and older • The use of advance prompts and/or reminders for the T2 and T3 surveys (see Figure A1 below) • Provision of an addressed, reply-paid envelope to facilitate the return of paper surveys • Sending and receiving of T2 and T3 surveys from the research institution (rather than clinic) • Participant compensation of a \$10 gift card for completing each of the T2 and T3 surveys
Maximise data completeness and quality	<ul style="list-style-type: none"> • Collection of data via surveys rather than interviews given vulnerability of topic to social desirability bias • Development of surveys and measures by experts in patient-reported outcome and experience measurement in partnership with patients and stakeholders • Online mode of survey completion for the T1 survey to ensure no handling of surveys by clinic staff and thus reinforce the confidentiality of responses • Reassurance in the study information sheet that no health care providers from participating clinics would have access to identified participant-level data • Use of programmed or instructional skips to ensure participants were asked only relevant questions, thereby minimising survey fatigue • Pop-up messages that notified participants of missed questions and invited them to respond to them prior to proceeding to the next page in online surveys • Reminders of clinic name on the tablet computer used to administer the T1 survey • Use of intervention images in items assessing intervention exposure • Participant-specific reminders of the date and clinic of the health care visit and the (previously reported) intended contraceptive method(s) in follow-up surveys • Duplicate data entry for a random subsample of paper surveys to confirm accuracy

Figure A1. Survey invitation and reminder schedule customised to survey mode



APPENDIX 9

Outcomes and measures

Table A3. Outcomes and measures

Outcome	Variable	Coding	Eligible	Excluded
Primary outcome				
Shared decision-making about contraceptive methods	Binary	<ul style="list-style-type: none"> • Optimal (12 out of 12) • Suboptimal (0-11 out of 12) 	Participants who experienced a conversation about contraception in the health care visit	Of the 4,209 participants who were eligible for the study and experienced a conversation about contraception in the health care visit, 21 were excluded due to missing data on shared decision-making about contraceptive methods and a further 108 due to missing data on one or more covariates
	Categorical (post-hoc)	<ul style="list-style-type: none"> • Optimal (12 out of 12) • Moderate (10-11 out of 12) • Low (0-9 out of 12) 		
	Continuous (post-hoc)	Range: 0-12		
Secondary outcomes				
Conversation about contraception	Binary	<ul style="list-style-type: none"> • Conversation • No conversation 	All participants	Of the 5,038 participants who were eligible for the study, 20 were excluded due to missing data on conversation about contraception and a further 153 due to missing data on one or more covariates
			<i>Prespecified subgroup:</i> All participants excluding those not at risk of unintended pregnancy at the time of the health care visit ¹	Of the 5,038 participants who were eligible for the study, 20 were excluded due to missing data on conversation about contraception, 257 due to not meeting subgroup criteria, and a further 142 due to missing data on one or more covariates
			<i>Prespecified subgroup:</i> All participants excluding those not at risk of unintended pregnancy at the time of the health care visit ¹ and those who reported that they did not want or need to talk about contraception	Of the 5,038 participants who were eligible for the study, 20 were excluded due to missing data on conversation about contraception, 687 due to not meeting subgroup criteria, and a further 133 due to missing data on one or more covariates
Satisfaction with conversation about contraception	Binary	<ul style="list-style-type: none"> • Very satisfied • Somewhat satisfied, neither satisfied nor dissatisfied, somewhat dissatisfied, very dissatisfied 	Participants who experienced a conversation about contraception in the index health care visit	Of the 4,209 participants who were eligible for the study and experienced a conversation about contraception in the health care visit, 15 were excluded due to missing data on satisfaction with conversation about contraception and a further 109 due to missing data on one or more covariates
Intended contraceptive method(s)	Binary	<ul style="list-style-type: none"> • ≥1 listed method(s) • 0 listed methods 	Participants who experienced a conversation about contraception in the index health care visit	Of the 4,209 participants who were eligible for the study and experienced a conversation about contraception in the health care visit, 26 were excluded due to missing data on intended contraceptive method(s) and a further 101 due to missing data on one or more covariates
			<i>Prespecified subgroup:</i> Participants who experienced a conversation about contraception in the health care visit excluding those not at risk of unintended pregnancy at the time of the health care visit ¹	Of the 4,209 participants who were eligible for the study and experienced a conversation about contraception in the health care visit, 26 were excluded due to missing data on intended contraceptive method(s), 224 due to not meeting subgroup criteria, and a further 94 due to missing data on one or more covariates.

			<p>Prespecified subgroup: Participants who experienced a conversation about contraception in the health care visit excluding those not at risk of unintended pregnancy at the time of the health care visit¹ and those who reported that they did not want or need to use a contraceptive method</p>	n/a
Intention to use a highly effective contraceptive method	Binary	<ul style="list-style-type: none"> • ≥1 highly effective method(s) • 0 highly effective methods 	<p>Participants who experienced a conversation about contraception in the index health care visit</p> <p>Prespecified subgroup: Participants who experienced a conversation about contraception in the health care visit excluding those not at risk of unintended pregnancy at the time of the health care visit¹</p> <p>Prespecified subgroup: Participants who experienced a conversation about contraception in the health care visit excluding those not at risk of unintended pregnancy at the time of the health care visit¹ and those who reported that they did not want or need to use a contraceptive method</p>	<p>Of the 4,209 participants who were eligible for the study and experienced a conversation about contraception in the health care visit, 26 were excluded due to missing data on intended contraceptive method(s) and a further 101 due to missing data on one or more covariates</p> <p>Of the 4,209 participants who were eligible for the study and experienced a conversation about contraception in the health care visit, 26 were excluded due to missing data on intended contraceptive method(s), 224 due to not meeting subgroup criteria, and a further 94 due to missing data on one or more covariates.</p> <p>Of the 4,209 participants who were eligible for the study and experienced a conversation about contraception in the health care visit, 26 were excluded due to missing data on intended contraceptive method(s), 317 due to not meeting subgroup criteria, and a further 87 due to missing data on one or more covariates.</p>
Values concordance of intended contraceptive method(s)	Binary	<ul style="list-style-type: none"> • Very confident • Confident, somewhat confident, not very confident, not at all confident 	<p>Participants who experienced a conversation about contraception in the health care visit and intended to use ≥1 contraceptive method(s)</p>	<p>T1: Of the 3,772 participants who were eligible for the study, experienced a conversation about contraception in the health care visit, and intended to use ≥1 contraceptive method(s), 5 were excluded due to missing data on values concordance of intended contraceptive method(s) and a further 79 due to missing data on one or more covariates</p> <p>T2,T3: Of the 3,772 participants who were eligible for the study, experienced a conversation about contraception in the health care visit, and intended to use ≥1 contraceptive method(s), 1,301 were excluded due to not giving permission to be contacted for follow-up data collection, 1,213 due to not participating at follow-up, 1 due to missing data on values concordance of intended contraceptive method(s), and a further 5 due to missing data on one or more covariates</p>
Decision regret about intended contraceptive method(s)	Continuous	Range: 0-4	<p>Participants who experienced a conversation about contraception in the index health care visit and intended to use ≥1 contraceptive method(s)</p>	<p>Of the 3,772 participants who were eligible for the study, experienced a conversation about contraception in the health care visit, and intended to use ≥1 contraceptive method(s), 1,301 were excluded due to not giving permission to be contacted for follow-up data collection, 1,213 due to not participating at follow-up, 3 due to missing data on decision regret about intended contraceptive method(s), and a further 5 due to missing data on one or more covariates</p>

Contraceptive method(s) used	Binary	<ul style="list-style-type: none"> • ≥1 listed method(s) • 0 listed methods 	Participants who experienced a conversation about contraception in the index health care visit and intended to use ≥1 contraceptive method(s)	n/a
Use of a highly effective contraceptive method	Binary	<ul style="list-style-type: none"> • ≥1 highly effective method(s) • 0 highly effective methods 	Participants who experienced a conversation about contraception in the index health care visit and intended to use ≥1 contraceptive method(s)	n/a
Use of intended contraceptive method(s)	Binary	<ul style="list-style-type: none"> • Perfect concordance of intended and used method(s) • Imperfect concordance of intended and used method(s) 	Participants who experienced a conversation about contraception in the index health care visit and intended to use ≥1 contraceptive method(s)	Of the 3,772 participants who were eligible for the study, experienced a conversation about contraception in the health care visit, and intended to use ≥1 contraceptive method(s), 1,301 were excluded due to not giving permission to be contacted for follow-up data collection, 1,213 due to not participating at follow-up, 1 due to missing data on use of intended contraceptive method(s), and a further 5 due to missing data on one or more covariates
Adherence to contraceptive method(s) used	Binary	<ul style="list-style-type: none"> • Perfect adherence to ≥1 method • Perfect adherence to 0 methods 	Participants who experienced a conversation about contraception in the index health care visit, intended to use ≥1 contraceptive method(s), and used ≥1 contraceptive method(s)	Of the 1,212 participants who were eligible for the study, experienced a conversation about contraception in the health care visit, intended to use ≥1 contraceptive method(s), and used ≥1 contraceptive method(s), 2 were excluded due to missing data on adherence to contraceptive method(s) used and a further 5 due to missing data on one or more covariates
Satisfaction with contraceptive method(s) used	Binary	<ul style="list-style-type: none"> • Very satisfied • Somewhat satisfied, neither satisfied nor dissatisfied, somewhat dissatisfied, very dissatisfied 	Participants who experienced a conversation about contraception in the index health care visit, intended to use ≥1 contraceptive method(s), and used either none of the listed contraceptive methods or ≥1 contraceptive method(s)	Of the 1,257 participants who were eligible for the study, experienced a conversation about contraception in the health care visit, intended to use ≥1 contraceptive method(s), and used either none of the listed contraceptive methods or ≥1 contraceptive method(s), 3 were excluded due to missing data on satisfaction with contraceptive method(s) used and a further 5 due to missing data on one or more covariates
Unintended pregnancy (timing preferences)	Binary	<ul style="list-style-type: none"> • ≥1 unintended pregnancies • 0 unintended pregnancies 	Participants who experienced a conversation about contraception in the index health care visit and intended to use ≥1 contraceptive method(s)	n/a
Unintended pregnancy (pregnancy seeking)	Binary	<ul style="list-style-type: none"> • ≥1 unintended pregnancies • 0 unintended pregnancies 	Participants who experienced a conversation about contraception in the index health care visit and intended to use ≥1 contraceptive method(s)	n/a
Unwelcome pregnancy	Binary	<ul style="list-style-type: none"> • ≥1 unwelcome pregnancies • 0 unwelcome pregnancies 	Participants who experienced a conversation about contraception in the index health care visit and intended to use ≥1 contraceptive method(s)	n/a
Process outcomes				
Intervention exposure ²	Binary ³	<ul style="list-style-type: none"> • Intended exposure • Unintended exposure 	Participants with data on at least one study outcome ⁴	n/a
Acceptability of interventions ²	Binary	<ul style="list-style-type: none"> • Would recommend to a friend • Would not recommend to a friend 	Participants exposed to the relevant intervention component as intended ⁵	n/a

Notes. Grey cells signify planned analyses that could not be conducted. ¹Participants were considered not at risk of unintended pregnancy if they reported they were pregnant, were trying to get pregnant, were born without ovaries or a uterus, had had their ovaries or uterus removed, had entered menopause, were infertile, or did not plan to have vaginal sex with a person who produces sperm. ²Only

descriptive analyses of these outcomes were conducted (see *Analysis*).³Exposure outcomes were planned as binary variables. Because only descriptive analyses were conducted, categorical data are also presented.⁴Each outcome was measured among all participants and is reported among participants with data on at least one study outcome.⁵Each outcomes was measured among participants with any exposure to the relevant intervention component and is reported among participants exposed to the relevant intervention component as intended.

Table A4. Impact of patient and public involvement

Domain	Illustrative impacts on decisions and materials
Objectives	<ul style="list-style-type: none"> • Decision to include the prompt card in the video intervention strategy • Refinement of conceptualisation of some of the secondary outcomes (e.g., for values concordance of intended contraceptive method(s), to focus on values concordance as perceived by the patient and not by a third party) • Addition of the secondary outcome of unwelcome pregnancy • Planned subgroup analyses for some secondary outcomes
Interventions	<ul style="list-style-type: none"> • Content domains and content of the decision aids • Visual presentation of textual and numerical information in the decision aids • Language used in the decision aids • Content of the script for the (provider) training video • Refinement of the wording of one of the questions advocated by the video and prompt card to “pros and cons” • Creation of a live rather than animated (patient) video that featured an authentic patient story • Content of the script for the (patient) video • Creation of versions of the (patient) video with on-screen captions for patients with hearing impairments
Methods	<ul style="list-style-type: none"> • Creation of a patient-friendly study name (and the specific name, 'Right For Me') • Decision not to create eligibility criteria pertaining to current gender identity • Language used to communicate eligibility criteria to prospective participants • Language used in surveys and in new or adapted patient-reported outcome measures (including question stem and response options for the Measure of Alignment of Choices (MATCH)) • Conceptualisation and design of strategies for enhancing recruitment, retention, and data quality (including materials to provide regular feedback to clinic staff on the number of participants enrolling in the study) • Inclusion of four-week follow-up data collection in addition to six-month follow-up data collection • Coding of variables for heterogeneity of treatment effects analyses
Results	<ul style="list-style-type: none"> • Interpretation of findings

Table A5. Protocol refinements and deviations

Method	Change or addition
Study setting	During the study, we became aware that some clinics used staff members from affiliated clinics that were also participating in the study to temporarily cover staff shortages. Attempts were made by clinics themselves to avoid any contamination arising from this.
Participants	Some participants (n=29) were believed to have enrolled in the study more than once, despite passing eligibility screening. Suspected duplicate participation attempts were retrospectively classified as ineligible by researchers. During the study, after publication of the study protocol, we devised eligibility rules pertaining to the required timing of completion of the T2 and T3 surveys relative to the target completion date. These rules required T2 surveys to be completed within one week before or after the target completion date and T3 surveys to be completed within two weeks before or after the target completion date.
Study outcomes	In our study protocol, we did not specify precisely how the outcomes of intended contraceptive method(s) or contraceptive method(s) used would be coded. Given the diversity in participant responses to items assessing these two outcomes, we elected to code these outcomes as binary variables.
Study time frame	Due to delays, the enrolment of participants and administration of the T1 survey among the pre-implementation cohort began after April 1, 2016 in two clinics, and the implementation of the intervention(s) and administration of the T1 survey among the post-implementation cohort began after July 1, 2016 in one clinic. Enrolment of participants and administration of the T1 survey among the post-implementation cohort continued until December 30, 2016 in one clinic.
Data collection	Due to weekends, public holidays, or other planned staff absences, there were some minor deviations from the planned survey mailing timeframe.

Figure A2. Flowchart of participants who consented to the study by trial arm




Table A6. Intervention effects on shared decision-making about contraceptive methods (continuous scoring method): differences between pre-implementation and post-implementation cohorts and difference-in-differences analyses

	Pre-implementation cohort M (SD)	Post-implementation cohort M (SD)	Difference between pre-implementation [REF] and post-implementation cohorts ALSM ¹ (95% CI)
All participants (n=4,080)			
Video (n=995)	10.51 (2.20)	10.61 (1.96)	-0.02 (-0.26, 0.23)
Decision aids (n=1,031)	10.94 (1.80)	11.16 (1.51)	0.14 (-0.09, 0.37)
Dual (n=1,117)	10.89 (1.87)	10.81 (1.82)	-0.00 (-0.24, 0.23)
Usual care (n=937)	10.98 (1.57)	10.69 (1.95)	-0.29 (-0.54, -0.05)
	Difference-in-differences ALSM ¹ (95% CI)	P-value unadjusted for multiple comparisons	P-value adjusted for multiple comparisons
All participants (n=4,080)			
Video vs. Usual care [REF]	0.28 (-0.07, 0.63)	0.12	0.49
Decision aids vs. Usual care [REF]	0.44 (0.10, 0.77)	0.01	0.09
Dual vs. Video [REF]	0.01 (-0.33, 0.35)	0.95	1.00
Dual vs. Decision aids [REF]	-0.15 (-0.47, 0.18)	0.38	0.86

Notes. ALSM = Adjusted Least Squares Mean, CI = Confidence Interval, M = Mean, REF = Reference group, SD = Standard Deviation.

¹Analyses adjusted for age, health insurance coverage, health literacy, educational attainment, ethnicity, race, number of pregnancies, number of births, number of abortions, number of miscarriages, previous contraceptive method(s), and survey language.

Table A7. Intervention effects on shared decision-making about contraceptive methods (categorical scoring method): differences between pre-implementation and post-implementation cohorts and difference-in-differences analyses

	Pre-implementation cohort			Post-implementation cohort			Difference between pre-implementation [REF] and post-implementation cohorts
	% low	% moderate	% optimal	% low	% moderate	% optimal	ACOR ¹ (95% CI)
All participants (n=4,080)							
Video (n=995)	31.8	12.5	55.7	30.9	13.0	56.1	0.92 (0.70, 1.22)
Decision aids (n=1,031)	22.7	11.6	65.7	18.5	12.3	69.2	1.11 (0.85, 1.45)
Dual (n=1,117)	23.5	12.2	64.3	25.4	15.2	59.5	0.90 (0.69, 1.18)
Usual care (n=937)	23.7	12.7	63.5	28.1	15.5	56.4	0.75 (0.57, 0.99)
	Difference-in-differences ACOR ¹ (95% CI)			P-value unadjusted for multiple comparisons			P-value adjusted for multiple comparisons
All participants (n=4,080)							
Video vs. Usual care [REF]	1.23 (0.83, 1.83)			0.30			0.78
Decision aids vs. Usual care [REF]	1.48 (1.01, 2.19)			0.05			0.26
Dual vs. Video [REF]	0.98 (0.67, 1.44)			0.91			1.00
Dual vs. Decision aids [REF]	0.81 (0.56, 1.19)			0.28			0.76

Notes. ACOR = Adjusted Cumulative Odds Ratio, CI = Confidence Interval, REF = Reference group. ¹Analyses adjusted for age, health insurance coverage, health literacy, educational attainment, ethnicity, race, number of pregnancies, number of births, number of abortions, number of miscarriages, previous contraceptive method(s), and survey language.

Table A8. P-values for three-way interactions to assess heterogeneity of treatment effects on shared decision-making about contraceptive methods

Variable ¹	Variable levels	P-value for three-way interaction between arm, cohort, and variable ²
Age	15 to 24 years, 25 to 49 years	0.66
Health insurance coverage	None, Public only, Other	0.06
Health literacy	Limited, Adequate	<0.0001
Educational attainment	High school graduate or less, Some college or more	0.58
Ethnicity and race	White alone and not of Hispanic, Latino, or Spanish origin, Of Hispanic, Latino, or Spanish origin and/or another race(s)	0.61
Number of pregnancies	None, One or more	0.39
Number of births	None, One or more	0.22
Number of abortions	None, One or more	0.77
Number of miscarriages	None, One or more	0.44
Previous contraceptive method(s)	None, Less effective, Highly effective	0.85

Notes. ¹For the variables of age, health insurance coverage, health literacy, educational attainment, ethnicity and race, and previous contraceptive method(s), analyses adjusted for age, health insurance coverage, health literacy, educational attainment, ethnicity, race, number of pregnancies, number of births, number of abortions, number of miscarriages, previous contraceptive method(s), and survey language (except the variable itself). For the variables of number of pregnancies, number of births, number of abortions, and number of miscarriages, analyses adjusted for age, health insurance coverage, health literacy, educational attainment, ethnicity, race, previous contraceptive method(s), and survey language. ²P-values were compared to a critical value of .005 to account for the consequences of multiple comparisons.

Table A9. Intervention effects on secondary outcomes: differences between pre-implementation and post-implementation cohorts

	Pre-implementation cohort %	Post-implementation cohort %	Difference between pre-implementation [REF] and post-implementation cohorts AOR ¹ (95% CI)
CONVERSATION ABOUT CONTRACEPTION (n=4,865)			
Video (n=1,202)	82.5	83.5	0.86 (0.60, 1.24)
Decision aids (n=1,116)	89.4	94.5	1.58 (0.97, 2.55)
Dual (n=1,342)	81.0	84.4	1.35 (0.94, 1.93)
Usual care (n=1,205)	76.5	78.6	1.41 (0.99, 2.03)
Prespecified subgroup excluding those not at risk of unintended pregnancy (n=4,619)			
Video (n=1,173)	84.4	85.6	0.86 (0.58, 1.26)
Decision aids (n=1,097)	91.2	96.0	1.69 (0.98, 2.92)
Dual (n=1,212)	92.5	92.1	1.05 (0.64, 1.72)
Usual care (n=1,137)	81.0	83.3	1.43 (0.97, 2.12)
Prespecified subgroup excluding those not at risk of unintended pregnancy and those who reported that they did not want or need to talk about contraception (n=4,198)			
Video (n=1,017)	98.0	98.5	1.18 (0.42, 3.33)
Decision aids (n=1,038)	98.8	99.7	3.26 (0.61, 17.35)
Dual (n=1,148)	96.6	97.7	1.90 (0.86, 4.21)
Usual care (n=995)	92.9	95.1	1.80 (0.92, 3.51)
SATISFACTION WITH CONVERSATION ABOUT CONTRACEPTION (n=4,085)			
Video (n=1,000)	85.9	89.2	1.19 (0.78, 1.83)
Decision aids (n=1,031)	93.5	94.8	1.21 (0.70, 2.08)
Dual (n=1,118)	90.4	89.8	1.02 (0.65, 1.59)
Usual care (n=936)	90.9	91.4	1.09 (0.66, 1.79)
INTENDED CONTRACEPTIVE METHOD(S) (n=4,082)			
Video (n=998)	90.6	91.4	1.11 (0.65, 1.89)
Decision aids (n=1031)	91.8	91.9	0.86 (0.51, 1.46)
Dual (n=1117)	92.4	88.0	0.67 (0.39, 1.16)
Usual care (n=936)	88.6	89.2	1.13 (0.67, 1.90)
Prespecified subgroup excluding those not at risk of unintended pregnancy (n=3,865)			
Video (n=955)	95.1	95.4	0.99 (0.48, 2.05)
Decision aids (n=980)	96.7	96.6	0.73 (0.34, 1.60)
Dual (n=1,045)	97.2	94.7	0.54 (0.23, 1.24)
Usual care (n=885)	94.0	94.2	1.55 (0.75, 3.18)
INTENTION TO USE A HIGHLY EFFECTIVE CONTRACEPTIVE METHOD (n=4,082)			
Video (n=998)	26.2	27.3	1.01 (0.68, 1.51)
Decision aids (n=1,031)	30.4	31.1	1.10 (0.77, 1.56)
Dual (n=1,117)	39.5	31.4	0.74 (0.52, 1.05)
Usual care (n=936)	37.5	38.1	1.04 (0.71, 1.53)
Prespecified subgroup excluding those not at risk of unintended pregnancy (n=3,865)			
Video (n=955)	27.5	28.5	0.95 (0.63, 1.43)
Decision aids (n=980)	32.1	32.7	1.14 (0.79, 1.63)
Dual (n=1,045)	41.6	33.8	0.72 (0.50, 1.04)
Usual care (n=885)	39.7	40.3	1.07 (0.72, 1.60)
Prespecified subgroup excluding those not at risk of unintended pregnancy and those who reported that they did not want or need to use a contraceptive method (n=3,779)			
Video (n=938)	27.9	29.0	0.99 (0.65, 1.50)

Decision aids (n=961)	32.6	33.4	1.17 (0.81, 1.69)
Dual (n=1,027)	42.2	34.5	0.71 (0.49, 1.04)
Usual care (n=853)	41.3	41.8	1.05 (0.69, 1.57)
VALUES CONCORDANCE OF INTENDED CONTRACEPTIVE METHOD(S)			
Immediately after the health care visit (n=3,688)			
Video (n=910)	54.8	53.8	0.91 (0.67, 1.22)
Decision aids (n=947)	66.8	67.0	0.98 (0.73, 1.30)
Dual (n=998)	63.8	64.9	1.05 (0.79, 1.40)
Usual care (n=833)	62.3	59.3	0.88 (0.64, 1.19)
Four weeks after the health care visit (n=1,252)			
Video (n=264)	43.4	44.8	0.91 (0.53, 1.59)
Decision aids (n=232)	54.0	46.2	0.71 (0.41, 1.23)
Dual (n=336)	57.5	52.2	0.88 (0.55, 1.43)
Usual care (n=262)	44.8	49.7	1.27 (0.74, 2.17)
Six months after the health care visit (n=1,252)			
Video (n=238)	50.6	44.7	0.75 (0.42, 1.33)
Decision aids (n=215)	47.1	35.9	0.58 (0.32, 1.03)
Dual (n=275)	50.0	49.7	1.06 (0.62, 1.83)
Usual care (n=242)	48.2	49.7	1.13 (0.65, 1.96)
USE OF INTENDED CONTRACEPTIVE METHOD(S)			
Four weeks after the health care visit (n=1,252)			
Video (n=264)	57.8	64.1	1.21 (0.70, 2.10)
Decision aids (n=232)	56.0	56.8	1.03 (0.60, 1.78)
Dual (n=337)	61.7	61.3	1.00 (0.62, 1.63)
Usual care (n=262)	58.6	63.4	1.24 (0.72, 2.12)
Six months after the health care visit (n=1,252)			
Video (n=237)	59.2	56.5	0.84 (0.47, 1.48)
Decision aids (n=215)	59.8	46.9	0.59 (0.33, 1.05)
Dual (n=275)	57.5	61.0	1.22 (0.71, 2.10)
Usual care (n=242)	50.6	55.3	1.32 (0.76, 2.27)
ADHERENCE TO CONTRACEPTIVE METHOD(S) USED			
Four weeks after the health care visit (n=1,205)			
Video (n=255)	74.4	77.4	1.19 (0.61, 2.32)
Decision aids (n=219)	89.2	77.0	0.43 (0.19, 0.97)
Dual (n=316)	84.5	83.6	1.13 (0.57, 2.24)
Usual care (n=250)	83.5	84.8	1.00 (0.47, 2.13)
Six months after the health care visit (n=1,205)			
Video (n=225)	69.5	75.0	1.52 (0.77, 2.99)
Decision aids (n=198)	94.0	80.9	0.29 (0.10, 0.84)
Dual (n=259)	88.3	79.7	0.58 (0.26, 1.32)
Usual care (n=230)	87.7	84.6	0.69 (0.30, 1.60)
SATISFACTION WITH CONTRACEPTIVE METHOD(S) USED			
Four weeks after the health care visit (n=1,249)			
Video (n=264)	56.6	70.2	1.58 (0.90, 2.78)
Decision aids (n=231)	62.0	62.6	1.09 (0.62, 1.90)
Dual (n=336)	67.3	69.0	1.15 (0.70, 1.91)
Usual care (n=262)	64.4	73.1	1.55 (0.88, 2.74)
Six months after the health care visit (n=1,249)			
Video (n=237)	65.8	112 (69.6)	1.13 (0.62, 2.06)
Decision aids (n=214)	69.0	81 (63.8)	0.82 (0.45, 1.49)

Dual (n=275)	88.8	138 (70.8)	0.32 (0.15, 0.72)
Usual care (n=240)	70.7	112 (70.9)	1.04 (0.57, 1.89)
	Pre-implementation cohort M (SD)	Post-implementation cohort M (SD)	Difference between pre-implementation [REF] and post-implementation cohorts ALSM ¹ (95% CI)
DECISION REGRET ABOUT INTENDED CONTRACEPTIVE METHOD(S)			
Four weeks after the health care visit (n=1,250)			
Video (n=260)	0.82 (0.83)	0.61 (0.62)	-0.16 (-0.35, 0.02)
Decision aids (n=232)	0.59 (0.64)	0.61 (0.62)	0.00 (-0.19, 0.18)
Dual (n=337)	0.56 (0.63)	0.56 (0.67)	-0.03 (-0.19, 0.13)
Usual care (n=262)	0.65 (0.60)	0.56 (0.62)	-0.08 (-0.26, 0.10)
Six months after the health care visit (n=1,250)			
Video (n=237)	0.66 (0.70)	0.68 (0.77)	0.04 (-0.15, 0.23)
Decision aids (n=215)	0.70 (0.92)	0.92 (0.90)	0.21 (0.02, 0.40)
Dual (n=275)	0.66 (0.77)	0.70 (0.78)	0.01 (-0.17, 0.20)
Usual care (n=241)	0.65 (0.77)	0.65 (0.69)	0.00 (-0.18, 0.19)

Notes. ALSM = Adjusted Least Squares Mean, AOR = Adjusted Odds Ratio, CI = Confidence Interval, M = Mean, REF = Reference group, SD = Standard Deviation. ¹For the outcome of conversation about contraception, analyses adjusted for age, health insurance coverage, health literacy, educational attainment, ethnicity, race, number of pregnancies, number of births, number of abortions, number of miscarriages, and survey language. For the outcomes of satisfaction with conversation about contraception, intention to use one or more contraceptive method(s), intention to use highly effective contraceptive method, and values concordance immediately after the health care visit, analyses adjusted for age, health insurance coverage, health literacy, educational attainment, ethnicity, race, number of pregnancies, number of births, number of abortions, number of miscarriages, previous contraceptive method(s), and survey language. For the outcomes of values concordance four weeks and six months after the health care visit, use of intended contraceptive method(s), adherence to contraceptive method(s) used, satisfaction with contraceptive method(s), and decision regret, analyses adjusted for age, health insurance coverage, health literacy, educational attainment, ethnicity, race, number of pregnancies, number of births, number of abortions, number of miscarriages, and previous contraceptive method(s).

Table A10. Intervention effects on secondary outcomes: difference-in-differences analyses

	Difference-in-differences AOR ¹ (95% CI)	P-value unadjusted for multiple comparisons	P-value adjusted for multiple comparisons
CONVERSATION ABOUT CONTRACEPTION (n=4,865)			
Video vs. Usual care [REF]	0.61 (0.37, 1.02)	0.06	0.31
Decision aids vs. Usual care [REF]	1.12 (0.61, 2.03)	0.72	0.99
Dual vs. Video [REF]	1.56 (0.94, 2.60)	0.09	0.40
Dual vs. Decision aids [REF]	0.86 (0.47, 1.56)	0.61	0.97
Prespecified subgroup excluding those not at risk of unintended pregnancy (n=4,619)			
Video vs. Usual care [REF]	0.60 (0.34, 1.04)	0.07	0.34
Decision aids vs. Usual care [REF]	1.18 (0.60, 2.31)	0.63	0.97
Dual vs. Video [REF]	1.22 (0.65, 2.29)	0.53	0.94
Dual vs. Decision aids [REF]	0.62 (0.30, 1.29)	0.20	0.65
Prespecified subgroup excluding those not at risk of unintended pregnancy and those who reported that they did not want or need to talk about contraception (n=4,198)			
Video vs. Usual care [REF]	0.66 (0.19, 2.25)	0.50	0.93
Decision aids vs. Usual care [REF]	1.81 (0.30, 10.94)	0.52	0.94
Dual vs. Video [REF]	1.60 (0.43, 5.92)	0.48	0.92
Dual vs. Decision aids [REF]	0.58 (0.09, 3.70)	0.57	0.95
SATISFACTION WITH CONVERSATION ABOUT CONTRACEPTION (n=4,085)			
Video vs. Usual care [REF]	1.10 (0.57, 2.12)	0.78	0.99
Decision aids vs. Usual care [REF]	1.11 (0.53, 2.33)	0.78	0.99
Dual vs. Video [REF]	0.85 (0.46, 1.59)	0.62	0.97
Dual vs. Decision aids [REF]	0.84 (0.42, 1.70)	0.63	0.97
INTENDED CONTRACEPTIVE METHOD(S) (n=4,082)			
Video vs. Usual care [REF]	0.98 (0.46, 2.09)	0.96	1.00
Decision aids vs. Usual care [REF]	0.77 (0.36, 1.62)	0.49	0.92
Dual vs. Video [REF]	0.61 (0.28, 1.31)	0.20	0.65
Dual vs. Decision aids [REF]	0.78 (0.37, 1.66)	0.52	0.94
Prespecified subgroup excluding those not at risk of unintended pregnancy (n=3,865)			
Video vs. Usual care [REF]	0.64 (0.23, 1.78)	0.39	0.87
Decision aids vs. Usual care [REF]	0.47 (0.16, 1.38)	0.17	0.60
Dual vs. Video [REF]	0.54 (0.18, 1.64)	0.28	0.76
Dual vs. Decision aids [REF]	0.73 (0.24, 2.28)	0.59	0.96
INTENTION TO USE A HIGHLY EFFECTIVE CONTRACEPTIVE METHOD (n=4,082)			
Video vs. Usual care [REF]	0.97 (0.56, 1.70)	0.92	1.00
Decision aids vs. Usual care [REF]	1.05 (0.62, 1.77)	0.85	1.00
Dual vs. Video [REF]	0.73 (0.42, 1.25)	0.25	0.72
Dual vs. Decision aids [REF]	0.67 (0.41, 1.11)	0.12	0.50
Prespecified subgroup excluding those not at risk of unintended pregnancy (n=3,865)			
Video vs. Usual care [REF]	0.88 (0.50, 1.57)	0.67	0.98
Decision aids vs. Usual care [REF]	1.06 (0.62, 1.82)	0.83	1.00
Dual vs. Video [REF]	0.76 (0.44, 1.32)	0.32	0.81
Dual vs. Decision aids [REF]	0.63 (0.38, 1.06)	0.08	0.39
Prespecified subgroup excluding those not at risk of unintended pregnancy and those who reported that they did not want or need to use a contraceptive method (n=3,779)			
Video vs. Usual care [REF]	0.95 (0.53, 1.70)	0.85	1.00
Decision aids vs. Usual care [REF]	1.12 (0.64, 1.94)	0.70	0.98
Dual vs. Video [REF]	0.72 (0.41, 1.26)	0.25	0.73
Dual vs. Decision aids [REF]	0.61 (0.36, 1.03)	0.07	0.34

VALUES CONCORDANCE OF INTENDED CONTRACEPTIVE METHOD(S)

Immediately after the health care visit (n=3,688)

Video vs. Usual care [REF]	1.04 (0.67, 1.59)	0.87	1.00
Decision aids vs. Usual care [REF]	1.12 (0.73, 1.70)	0.61	0.97
Dual vs. Video [REF]	1.15 (0.76, 1.75)	0.50	0.93
Dual vs. Decision aids [REF]	1.07 (0.71, 1.61)	0.74	0.99

Four weeks after the health care visit (n=1,252)

Video vs. Usual care [REF]	0.72 (0.33, 1.56)	0.40	0.87
Decision aids vs. Usual care [REF]	0.56 (0.26, 1.20)	0.14	0.53
Dual vs. Video [REF]	0.97 (0.47, 2.02)	0.93	1.00
Dual vs. Decision aids [REF]	1.25 (0.60, 2.58)	0.55	0.95

Six months after the health care visit (n=1,252)

Video vs. Usual care [REF]	0.66 (0.30, 1.47)	0.31	0.80
Decision aids vs. Usual care [REF]	0.51 (0.23, 1.14)	0.10	0.45
Dual vs. Video [REF]	1.42 (0.65, 3.12)	0.38	0.86
Dual vs. Decision aids [REF]	1.84 (0.83, 4.07)	0.13	0.51

USE OF INTENDED CONTRACEPTIVE METHOD(S)

Four weeks after the health care visit (n=1,252)

Video vs. Usual care [REF]	0.98 (0.45, 2.11)	0.96	1.00
Decision aids vs. Usual care [REF]	0.83 (0.39, 1.79)	0.64	0.97
Dual vs. Video [REF]	0.83 (0.40, 1.72)	0.61	0.97
Dual vs. Decision aids [REF]	0.97 (0.47, 2.01)	0.94	1.00

Six months after the health care visit (n=1,252)

Video vs. Usual care [REF]	0.63 (0.29, 1.40)	0.26	0.74
Decision aids vs. Usual care [REF]	0.45 (0.20, 0.99)	0.05	0.27
Dual vs. Video [REF]	1.46 (0.66, 3.21)	0.35	0.83
Dual vs. Decision aids [REF]	2.07 (0.94, 4.55)	0.07	0.35

ADHERENCE TO CONTRACEPTIVE METHOD(S) USED

Four weeks after the health care visit (n=1,205)

Video vs. Usual care [REF]	1.20 (0.44, 3.29)	0.73	0.99
Decision aids vs. Usual care [REF]	0.43 (0.14, 1.31)	0.14	0.53
Dual vs. Video [REF]	0.95 (0.37, 2.46)	0.91	1.00
Dual vs. Decision aids [REF]	2.64 (0.92, 7.62)	0.07	0.36

Six months after the health care visit (n=1,205)

Video vs. Usual care [REF]	2.20 (0.75, 6.46)	0.15	0.56
Decision aids vs. Usual care [REF]	0.42 (0.11, 1.63)	0.21	0.67
Dual vs. Video [REF]	0.38 (0.13, 1.11)	0.08	0.37
Dual vs. Decision aids [REF]	1.98 (0.52, 7.50)	0.31	0.80

SATISFACTION WITH CONTRACEPTIVE METHOD(S) USED

Four weeks after the health care visit (n=1,249)

Video vs. Usual care [REF]	1.02 (0.46, 2.27)	0.96	1.00
Decision aids vs. Usual care [REF]	0.70 (0.31, 1.55)	0.38	0.86
Dual vs. Video [REF]	0.73 (0.34, 1.55)	0.41	0.88
Dual vs. Decision aids [REF]	1.06 (0.50, 2.26)	0.87	1.00

Six months after the health care visit (n=1,249)

Video vs. Usual care [REF]	1.09 (0.46, 2.55)	0.85	1.00
Decision aids vs. Usual care [REF]	0.79 (0.33, 1.85)	0.58	0.96
Dual vs. Video [REF]	0.29 (0.11, 0.76)	0.01	0.10
Dual vs. Decision aids [REF]	0.40 (0.15, 1.06)	0.06	0.33

	Difference-in-differences ALSM ¹ (95% CI)	P-value unadjusted for multiple comparisons	P-value adjusted for multiple comparisons
DECISION REGRET ABOUT INTENDED CONTRACEPTIVE METHOD(S)			
Four weeks after the health care visit (n=1,250)			
Video vs. Usual care [REF]	-0.08 (-0.34, 0.18)	0.53	0.94
Decision aids vs. Usual care [REF]	0.08 (-0.18, 0.34)	0.55	0.95
Dual vs. Video [REF]	0.14 (-0.11, 0.38)	0.28	0.76
Dual vs. Decision aids [REF]	-0.02 (-0.27, 0.22)	0.84	1.00
Six months after the health care visit (n=1,250)			
Video vs. Usual care [REF]	0.04 (-0.23, 0.31)	0.78	0.99
Decision aids vs. Usual care [REF]	0.21 (-0.06, 0.48)	0.13	0.52
Dual vs. Video [REF]	-0.03 (-0.29, 0.23)	0.82	1.00
Dual vs. Decision aids [REF]	-0.20 (-0.47, 0.07)	0.14	0.54

Notes. ALSM = Adjusted Least Squares Mean, AOR = Adjusted Odds Ratio, CI = Confidence Interval, REF = Reference group. ¹For the outcome of conversation about contraception, analyses adjusted for age, health insurance coverage, health literacy, educational attainment, ethnicity, race, number of pregnancies, number of births, number of abortions, number of miscarriages, and survey language. For the outcomes of satisfaction with conversation about contraception, intention to use one or more contraceptive method(s), intention to use highly effective contraceptive method, and values concordance immediately after the health care visit, analyses adjusted for age, health insurance coverage, health literacy, educational attainment, ethnicity, race, number of pregnancies, number of births, number of abortions, number of miscarriages, previous contraceptive method(s), and survey language. For the outcomes of values concordance four weeks and six months after the health care visit, use of intended contraceptive method(s), adherence to contraceptive method(s) used, satisfaction with contraceptive method(s), and decision regret, analyses adjusted for age, health insurance coverage, health literacy, educational attainment, ethnicity, race, number of pregnancies, number of births, number of abortions, number of miscarriages, and previous contraceptive method(s).

Table A11. Patient-reported intervention exposure among pre-implementation and post-implementation cohorts

	Video	Decision aids	Dual	Usual care
PRE-IMPLEMENTATION COHORT (n=1683)	(n=370)	(n=470)	(n=443)	(n=400)
Video				
Watched whole video before visit	6 (1.6)	2 (0.4%)	13 (3.0)	4 (1.0)
Watch whole video during or after visit	5 (1.4)	7 (1.5%)	6 (1.4)	1 (0.3)
Watched some of video	4 (1.1)	1 (0.2%)	7 (1.6)	2 (0.5)
Watched none of video	351 (95.9)	458 (97.9%)	413 (94.1)	392 (98.2)
<i>Missing</i>	4	2	4	1
Decision aids				
Looked at decision aid(s) during visit	61 (16.7)	86 (18.3)	59 (13.5)	39 (9.8)
Looked at decision aid(s) before or after visit	72 (19.7)	129 (27.5)	101 (23.2)	71 (17.9)
Did not look at decision aid(s)	233 (63.7)	254 (54.2)	276 (63.3)	287 (72.3)
<i>Missing</i>	4	1	7	3
Video and decision aids				
Watched whole video before visit + looked at decision aid(s) during visit	3 (0.8)	1 (0.2)	4 (0.9)	1 (0.3)
Some other exposure to the video and decision aid(s)	131 (35.8)	216 (46.2)	162 (37.2)	111 (28.0)
Watched none of video + did not look at decision aid(s)	232 (63.4)	251 (53.6)	270 (61.9)	285 (71.8)
<i>Missing</i>	4	2	7	3
POST-IMPLEMENTATION COHORT (n=3335)	(n=875)	(n=671)	(n=945)	(n=844)
Video				
Watched whole video while waiting	160 (18.8)	36 (5.6)	162 (17.7)	40 (5.0)
Watched some of video while waiting	18 (2.1)	14 (2.2)	14 (1.5)	18 (2.3)
Watched none of video while waiting	674 (79.1)	593 (92.2)	737 (80.7)	737 (92.7)
<i>Not sure or missing</i>	23	28	32	49
Prompt card				
Received prompt card while waiting	156 (19.3)	140 (23.9)	238 (27.1)	74 (9.5)
Did not receive prompt card while waiting	652 (80.7)	446 (76.1)	640 (72.9)	702 (90.5)
<i>Not sure or missing</i>	67	85	67	68
Video and prompt card				
Watched whole video while waiting + received prompt card while waiting	75 (9.4)	23 (4.0)	74 (8.6)	16 (2.1)
Watched whole video while waiting + did not receive prompt card while waiting	74 (9.3)	7 (1.2)	75 (8.7)	16 (2.1)
Watched some of video while waiting + received prompt card while waiting	7 (0.9)	7 (1.2)	4 (0.5)	4 (0.5)
Watched some of video while waiting + did not receive prompt card while waiting	7 (0.9)	5 (0.9)	8 (0.9)	14 (1.9)
Watched none of video while waiting + received prompt card while waiting	69 (8.7)	101 (17.6)	149 (17.3)	48 (6.4)
Watched none of video while waiting + did not receive prompt card while waiting	564 (70.9)	431 (75.1)	552 (64.0)	651 (86.9)
<i>Not sure or missing</i>	79	97	83	95
Decision aids				
Used decision aid(s) with provider	71 (9.0)	194 (31.5)	147 (16.9)	94 (12.0)
Did not use decision aid(s) with provider but were given	146 (18.5)	158 (25.6)	194 (22.3)	143 (18.2)
Did not use decision aid(s) with provider and were not given	571 (72.5)	264 (42.9)	528 (60.8)	548 (69.8)
<i>Not sure or missing</i>	87	55	76	59
Video, prompt card, and decision aids				
Watched whole video while waiting + received prompt card while waiting + used decision aid(s) with provider	26 (3.5)	17 (3.2)	41 (5.0)	9 (1.3)
Some other exposure to the video, prompt card, and decision aid(s)	268 (36.3)	294 (54.9)	382 (47.0)	223 (31.3)
Watched none of video while waiting + did not receive prompt card while waiting + did not use decision aid(s) with provider and were not given	444 (60.2)	225 (42.0)	389 (47.9)	481 (67.5)

Notes. Grey cells signify intended exposure. Information on variable coding is available on request.

Table A12. Acceptability of interventions among patients exposed as intended in the post-implementation cohort

	Would recommend to a friend		Would not recommend to a friend	
	Freq.	% (95% CI)	Freq.	% (95% CI)
Watched whole video while waiting (n=322)	305	94.7 (91.7, 96.9)	17	5.3 (3.1, 8.3)
Received prompt card while waiting (n=394)	360	91.4 (88.2, 94.0)	34	8.6 (6.0, 11.8)
Used decision aid(s) with provider (n=341)	336	98.5 (96.6, 99.5)	5	1.5 (0.5, 3.4)