Tracking SARS-CoV-2 in rivers as a tool for epidemiological surveillance

María Noel Maidana-Kulesza^{1,#}, Hugo Ramiro Poma^{1,#}, Diego Gastón Sanguino-Jorquera¹,
Sarita Isabel Reyes¹, María del Milagro Said-Adamo^{1,2}, Martín Mainardi Remis^{1,3}, Dolores
Gutiérrez-Cacciabue^{1,3}, Héctor Antonio Cristóbal^{1,2}, Mercedes Cecilia Cruz¹, Mónica
Aparicio González¹, Verónica Beatriz Rajal^{1,3,4,*}

¹ Laboratorio de Aguas y Suelos, Instituto de Investigaciones para la Industria Química
(INIQUI), Universidad Nacional de Salta (UNSa) – Consejo Nacional de Investigaciones
Científicas y Técnicas (CONICET), Av. Bolivia 5150, Salta, 4400, Argentina

² Facultad de Ciencias Naturales, UNSa, Av. Bolivia 5150, Salta, 4400, Argentina

³ Facultad de Ingeniería, UNSa, Av. Bolivia 5150, Salta, 4400, Argentina

⁴ Singapore Centre for Environmental Life Science Engineering (SCELSE), Nanyang
Technological University, Singapore.

Shared first authorship

* Corresponding author: vbrajal@gmail.com; UNSa, Av. Bolivia 5150, Salta, 4400,
Argentina

21 Abstract

22 The aim of this work was to evaluate if rivers could be used for SARS-CoV-2 surveillance
 23 to support health authorities. Five sampling points from three rivers (AR-1 and AR-2 in
 24 Arenales River, MR-1 and MR-2 in Mojotoro River, and CR in La Caldera River) from the
 25 Province of Salta (Argentina), two of them receiving the discharges of the wastewater
 26 plants (WWTP) of the city of Salta, were monitored from July to December 2020 during the
 27 first wave of COVID-19. Fifteen water samples from each point (75 samples in total) were
 28 collected and characterized physico-chemically and microbiologically and SARS-CoV-2
 29 was quantitatively detected by RT-qPCR. In addition, two targets linked to human
 30 contributions, human polyomavirus (HPyV) and RNase P, were quantified and used to
 31 normalize the SARS-CoV-2 concentration, which was ultimately compared to the active
 32 reported COVID-19 cases. Statistical analyses allowed us to verify the correlation between
 33 SARS-CoV-2 and the concentration of fecal indicator bacteria (FIB), as well as to find
 34 similarities and differences between sampling points. La Caldera River showed the best
 35 water quality, and FIB were within acceptable limits for recreational activities. Although
 36 Mojotoro River receives the discharge of the northern WWTP of the city, it did not affect
 37 the water quality. Instead, the Arenales River presented the poorest water quality and the
 38 river at AR-2 was negatively affected by the discharges of the southern WWTP, which
 39 contributed to the significant increase of fecal contamination. SARS-CoV-2 was only found
 40 in about half of the samples and in low concentrations in La Caldera and Mojotoro Rivers,
 41 while it was high and persistent in the Arenales River. None of the two human tracers was
 42 detected in CR, only HPyV was found in MR-1, MR-2 and AR-1, and both were quantified
 43 in AR-2. The experimental and the normalized (using the two tracers) viral concentrations
 44 strongly correlated with the curve of active reported COVID-19 cases; thus, the Arenales
 45 River at AR-2 reflected the epidemiological situation of the city. This is, to the best of our

46 knowledge, the first study that showed the dynamic of SARS-CoV-2 concentration in an
47 urban river highly impacted by wastewater and proved that can be used for SARS-CoV-2
48 surveillance to support health authorities.

49

50 **Keywords:** SARS-CoV-2; water quality; human tracers; normalization; rivers impacted by
51 wastewater; epidemiological surveillance

52

1. Introduction

The *Coronaviridae* family includes a broad spectrum of viruses that cause infections in various vertebrate animals including humans. Human Coronavirus are generally responsible for common colds. However, severe acute respiratory syndrome coronavirus 1 (SARS-CoV1) and Middle East respiratory syndrome coronavirus (MERS-CoV) were responsible for global epidemics, with high rates of morbidity and mortality, in 2003 and 2012, respectively (Peiris et al, 2003; Zaki et al, 2012). On 31 December 2019, the World Health Organization was informed about pneumonia cases of unknown origin in Wuhan, China. On 7 January 2020 the new virus, called 2019-nCoV at that moment, was identified as the cause and on 11 February 2020, it was named as the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) by the International Committee on Virus Taxonomy (Gorbalenya et al., 2020). The virus spread rapidly among the population in the globe and the pandemic was declared in mid-March 2020. Since then, it has infected more than 176 million people and 3.8 million confirmed deaths by June 16th, 2021 (WHO, 2020).

SARS-CoV-2 is an enveloped single-stranded positive-sense RNA virus with an approximate diameter of 80 to 120 nm. The main transmission pathway is person-to-person, mainly airborne by aerosols, droplets, and sputum, formed during breathing, sneezing, coughing, or talking (Zhang et al., 2020a). Another way of transmission described initially was by contact with contaminated fomites; however, this hypothesis lost strength and is yet under discussion (Harrison et al., 2020; Kampf et al., 2020; Mukhra et al., 2020).

Although the 2019-nCoV infection (COVID-19) is mainly a respiratory disease, SARS-CoV-2 also replicates in the enterocytes from ileum and colon (Zhang et al., 2020b). Thus, a fraction of infected people, symptomatic or asymptomatic, excrete the virus in feces (Xiao et al., 2020b; Zhang et al., 2020c; Zhang et al., 2020d) and urine (Guan et al., 2020; Jeong

et al., 2020) in variable concentrations. Furthermore, in one study the virus was isolated from urine (Sun et al., 2020) and in another from stool samples and propagated in Vero E6 cells (Xiao et al., 2020a). The fact that viral particles remain infective after being excreted through fecal matter, supports the hypothesis of a possible fecal-oral (Amirian, 2020; Jiang et al., 2020; Park et al., 2020; Wölfel et al., 2020) or a fecal-respiratory (Xiao et al., 2020a) route of transmission, although they have not been fully established yet.

The viral circulation in the population of a city, or a region, is reflected in the viral loads in their wastewater (Medema et al., 2020; Randazzo et al., 2020a). In fact, the so-called wastewater-based epidemiology has been applied in many developed countries (Ahmed et al., 2020; Lodder et al., 2020; Betancourt et al., 2021; Gonzalez et al., 2020; Albastaki et al., 2021; Saththasivam et al., 2021) to help authorities to better understand the situation and to make decisions regarding lockdowns. In most places, surveillance has been done in wastewater treatment plants, analyzing therefore, samples that represent the population of a city or part of it (Ahmed et al., 2020; Gonzalez et al., 2020; Kumar et al., 2020; Randazzo et al., 2020b; Rimoldi et al., 2020).

Rivers, as well as other water bodies, usually receive the discharges of treated wastewater (although sometimes with insufficient treatment) and, on some occasions, also (illegal) discharges of raw sewage. In this way, surface water can be impacted by wastewater. At the same time many of those aquatic environments are used as source water, for the irrigation of vegetables or for recreational purposes (Chávez-Díaz et al., 2019; Gutiérrez-Cacciabue et al., 2014; Poma et al., 2012). Previous studies about the microbiological quality of surface water in Salta, northwest of Argentina, demonstrated that fecal contamination was considerable in some rivers (Chavez-Díaz et al., 2020; Cruz et al., 2012; Gutierrez-Cacciabue et al., 2014), and multiple pathogens, including bacteria, enteric virus, and parasites, were found (Pisano et al., 2018; Poma et al., 2012; Prez et al.,

2020), representing a risk for the population in contact with those waters (Poma et al., 2019).

The aim of this work was to evaluate if rivers could be used for SARS-CoV-2 surveillance to support health authorities. Three rivers from the province of Salta (northwest of Argentina), two of them receiving the discharges of the two wastewater plants of the city of Salta (main city in the province, 700,000 inhabitants), were monitored from July to December 2020 during the first wave of COVID-19. Water samples were physico-chemically and microbiologically characterized, and SARS-CoV-2 was quantitatively detected. In addition, two targets linked to human contributions were quantified and used to normalize the SARS-CoV-2 concentration, which was ultimately compared to the reported COVID-19 cases.

2. Materials and methods

2.1. Sampling sites

Three rivers from the province of Salta, northwest of Argentina, were selected for the study: Arenales, Mojotoro, and La Caldera Rivers (Figure 1). The first two are impacted by the discharges of the two wastewater treatments plants (WWTP) of the city of Salta (capital of the province, with 700,000 inhabitants) and vicinities, while the third one is used as source water upstream and as a recreational ambient downstream.

The Arenales River (AR) is the main river that crosses the city of Salta, from west to southeast, that belongs to the Juramento-Salado watershed in the province of Salta, northwest of Argentina. While passing across the city, the river receives stormwater, industrial effluents, and illegal raw sewage (Poma et al., 2012; Lomniczi et al., 2007). Two sampling points were selected (Figure 1 a): AR-1 (Figure 1 c) and AR-2 (Figure 1 d)

located 0.7 km upstream and 0.7 km downstream the southern wastewater treatment plant (WWTP-S) and the municipal landfill, respectively. Despite the chemical and microbiological contamination of this river (Lomniczi et al., 2007; Poma et al., 2012; Prez et al., 2020), it is used for domestic irrigation of vegetables and for recreational purposes.

Figure 1. Geographical representation of the sampling sites (a) in Salta city and vicinities at the Province of Salta, northwest of Argentina (b). Three rivers were monitored: Arenales River before (AR-1) and after (AR-2) the discharge of the southern wastewater treatment plant (WWTP-S); Mojotoro River before (MR-1) and after (MR-2) the discharge of the northern wastewater treatment plant (WWTP-N), and La Caldera River (CR). An old wastewater stabilization pond (WWSP), out of use, located between CR and MR-1 is also

indicated. Shaded pink area corresponds to the coverage of WWTP-N, while shaded orange area corresponds to the coverage of WWTP-S. Illustrative photographs of the sampling points are shown: AR-1 (c), AR-2 (d), MR-1 (e), MR-2 (f), and CR (g).

La Caldera River (CR) is located 13 km north from the city of Salta. It is used as source water and for recreational activities. Indeed, it is the most visited river in Salta city's vicinities because of its landscapes and accessibility. One sampling site (CR) in a recreational area was selected for this study (Figure 1 e).

The Mojotoro River (MR) is originated from the convergence of La Caldera and Vaqueros Rivers (Figure 1 a). It represents a natural border between Salta city and vicinities, going from urban to semi-rural areas, and it is mainly used for recreation and gravel extraction. Two sampling points, 10 km away from each other, were selected for monitoring: MR-1 (Figures 1 a, e) and MR-2 (Figures 1 a, f), located 5 km before and 5 km after the discharge of the northern wastewater treatment plant (WWTP-N) (Figure 1 a), respectively. There is also an old, out of use, wastewater stabilization pond (WWSP) located 3.5 km after CR and 0.7 km before MR-1 (Figure 1 a).

2.2. Sampling and characterization of water samples

A total of 15 monitoring campaigns were carried out weekly initially and then every other week, from the 13 July to 28 December 2020. Water samples were characterized physico-chemical and microbiologically. Physico-chemical variables: pH, temperature (°C), turbidity (NTU), and conductivity (mS/cm), were measured in situ using a multiparametric probe U10 Horiba (Japan). For microbiological characterization, one-liter water samples were collected in sterile bottles and refrigerated until fecal indicator bacteria (FIB) were analyzed in the laboratory. Also 20-liter water samples were collected in clean and rinsed plastic

carboys at each point and brought to the laboratory for concentration and molecular analysis.

Total (TC) and thermotolerant (TTC) coliforms were determined by the multiple-tube fermentation method, culturing them on MacConkey broth (Britania, Argentina) at 37 and 44.5 °C for 48 h, respectively (Eaton et al., 2005). *Escherichia coli* (EC) and enterococci (EN) were enumerated through the Membrane Filter technique (Eaton et al., 2005). *E. coli* was cultured in modified mTEC Agar (Fluka, USA) at 35 °C for 2 h and 44.5 °C for 22 h (Method 1603, USEPA 2002a), and enterococci in mE Agar (Difco, USA) at 41 °C for 48 h and confirmation in Esculin-Iron Agar at 41 °C for 20 min (Method 1106.1, USEPA 2002b). Results for TC and TTC were expressed as Most Probable Number (MPN) per 100 mL, while for EC and EN they were expressed as Colony Forming Unit (CFU) per 100 mL.

2.3. Water concentration and nucleic acid extraction

Twenty-liter water samples were concentrated down to 80 mL by ultrafiltration using a disposable hemodialysis hollow fiber cartridge (FX Classix 100, Fresenius Medical Care, Germany), following a previously optimized and validated protocol for the detection of enteric viruses (Poma et al, 2012; Rajal et al, 2007). A fraction of the final concentrated water sample (140 µL) was employed for nucleic acid extraction using a commercial kit (Puro Virus RNA, Productos Bio-Lógicos, Argentina), producing 50 µL of extract which was used for the quantitative detection of SARS-CoV-2, Polyomavirus (HPyV) and Ribonuclease P (RNase P).

2.4. Quantitative detection of SARS-CoV-2

The detection of SARS-CoV-2 was performed on the RNA extracts from each concentrated water sample by RT-qPCR based on the detection of a region of the N gene (N1 system) validated by the US Centers for Disease Control and Prevention (CDC, 2020).

The determination and quantification of the gene N was performed in a StepOne Plus real time PCR system (Applied Biosystem). The final 20 μ L reaction volume contained 5 μ L of TaqMan Fast virus 1-Step Master Mix 4X (Applied Biosystems), primers and probe solutions, water (to complete 15 μ L), and 5 μ L of template. The final concentrations of forward and reverse primers were 400 nM each while it was 250 nM for the probe. The RT-qPCR was initiated with reverse transcription at 50 °C for 5 minutes followed by a step of 95 °C for 20 seconds and 45 cycles of 95 °C for 15 seconds and 59 °C for 1 minute.

Dilutions of the RNA extract were performed when needed to overcome inhibition. Reactions were performed in duplicate, simultaneously with a non-template control (MiliQ water) and DNA plasmid (nCoV-ALL-Control Plasmid, Eurofins) as positive control. A protocol for rapid digestion with EcoR1 (Promega, USA) was used to linearize the plasmid, following the manufacturer's instructions.

The standard curve for SARS-CoV-2 quantification ($y = -3.4700 x + 43.1616$; $R^2 = 0.9820$; amplification efficiency: 94%; dynamic range: 8) was performed with eight 10-fold serial dilutions, in triplicate, of the DNA plasmid that contains the interest target.

2.5. Normalization with Human Polyomaviruses (HPyV) and Ribonuclease P (RNase P) as human input indicators

The viral concentration was expected to show variations along time. These variations could be due to changing viral discharges into the rivers but also due to fluctuations in the flow rate, i.e. because of stormwater. Thus, to accurately assess the evolution of SARS-CoV-2 concentration, without the impact of external factors, two human indicators (HPyV and RNaseP) were used as normalizers.

Human JC (JCPyV) and BK (BKPyV) polyomaviruses were quantitatively detected by qPCR using the oligonucleotides and reaction conditions described by Barrios et al. (2018). On

the other hand, a fragment of the gene encoding for the human Ribonuclease P (RNase P) was quantitatively detected by qPCR using the system described by the US Centers for Disease Control and Prevention (CDC, 2020).

The final volume of each qPCR reaction contained 15 μ L of reaction mix containing SensiFAST Probe HiRox kit 2X (Bioline) and specific primers and probes to reach the final concentrations established (a summary of all the oligonucleotides used for all the targets, including those for the detection of SARS-CoV-2 is shown in Table 1) and 5 μ L of template. Cycling parameters were 95 °C for 5 minutes for polymerase activation and 45 cycles of 95 °C for 10 seconds for denaturation and 60 °C for 50 seconds for annealing/extension.

Standard curves for HPyV ($y = -3.656 x + 39.317$; $R^2 = 0.996$; amplification efficiency: 88%; dynamic range 8) and RNase P ($-3.71 x + 42.8280$; $R^2 = 0.9991$; dynamic range 7) markers, were built using 10-fold dilutions of the corresponding plasmid DNA, containing the target sequences, in triplicate.

Table 1. Oligonucleotides: forward (F) and reverse (R) primers and Taq-Man® probes (P) used for the detection of SARS-CoV-2 N gene (CDC, 2020), HPyV (Barrios et al., 2018) and RNase P gene (CDC, 2020).

Target	Oligonucleotide	Sequence 5' - 3'	Final concentration (nM)
N1 system of SARS-CoV-2 gene N	2019-nCoV_N1-F	GACCCCAAAATCAGCGAAAT	400
	2019-nCoV_N1-R	TCTGGTTACTGCCAGTTGAATCTG	400
	2019-nCoV_N1-P	FAM-ACCCCGCATTACGTTTGGTGGACC-BHQ1	250
HPyV JC-BK	S1-JC-BK (F)	CTGCTGCTGCCACAGGATT	500
	AS2-JC-BK (R)	CCTCTACAGTAGCAAGGGATGCA	500
	P-JC-BK	FAM-AGCAGCAGCCTCYCCAGCAGCAATTCAGC-BHQ	250

RNase P	RP-F	AGATTTGGACCTGCGAGCG	500
	RP-R	GAGCGGCTGTCTCCACAAGT	500
	RP-P	HEX-TTCTGACCTGAAGGCTCTGCGCG– BHQ-1	250

226

227 All the extracted nucleic acids from the concentrated water samples were analyzed for
 228 HPyV and RNaseP, by duplicate, using positive (plasmid) and negative (ultrapure water)
 229 controls. These results were used for the normalization of the SARS-CoV-2
 230 concentrations; three alternative normalizations were assessed, as follows:

231 Alternative 1: $CN1_{CoV} = \frac{C_{CoV}}{C_{HPyV}}$ (1)

232

233 Alternative 2: $CN2_{CoV} = \frac{C_{CoV}}{C_{RNaseP}}$ (2)

234

235 Alternative 3: $CN3_{CoV} = \frac{C_{CoV}}{\sqrt{\frac{C_{HPyV}}{C_{a,HPyV}} \times \frac{C_{RNaseP}}{C_{a,RNaseP}}}}$ (3)

236 Where C (gc/mL) is the concentration and subindexes CoV (as for SARS-CoV-2), $HPyV$
 237 and $RNaseP$ indicate the targets; C_a (gc/mL) is the average concentration and C_N (gc/mL)
 238 is the normalized concentration. The super index (i) ($i = 1, 2, 3$) indicates the normalization
 239 alternative.

240 2.6. COVID-19 cases

241 Reported COVID-19 cases for the city of Salta were compiled and the prior two-weeks
 242 accumulated amount were assigned to each monitoring date to compare the
 243 epidemiological curve to the evolution of the viral concentration in the Arenales River. The
 244 reason for choosing this river was that it crosses the city receiving the impact of various
 245 human activities (domestic and industrial) and also, it receives the discharges of the main

wastewater treatment plant (WWTP-S) (covers 85% of the population), which is outdated and not sufficient for treating the volume of a fast-growing city.

2.7. Statistical analyses

Physico-chemical variables (temperature, pH, conductivity, and turbidity) measured along the monitoring campaigns were compiled to show descriptive statistics: mean and standard deviation, median and range of variability (minimum – maximum).

To assess data variability during the monitoring campaign within each river and regarding the impact due to SARS-CoV-2, different statistical tests were run.

Most analyses were carried out using non-parametric tests, due to the non-normal distribution of the data ($p < 0.001$). Only the correlation between concentrations of HPyV and RNase P was carried out with a parametric test (Pearson) as both datasets showed normal distribution. Normality was tested by the Shapiro–Wilks W-test (Shapiro and Francia, 1972).

Kruskal-Wallis test (non-parametric) was performed with the physico-chemical and microbiological variables measured to compare all the sampling points. Different letters were assigned when there were significant differences within the variables between the sampling points with p -values < 0.05 .

Correlations between all the physico-chemical variables, and fecal indicator bacteria and SARS-CoV-2 concentrations measured for all the sampling points were assessed by Spearman test (non-parametric). This test was also applied to all the same variables for each sampling point. Correlations were considered significant when the probabilities (p) were lower than 0.05 and they were arbitrary considered weak when coefficients were lower than 0.40, moderate when were between 0.40 and 0.75, and strong when were higher than 0.75.

A Cluster Analysis (CA) and a Linear Discriminant Analysis (DA) were applied using nine variables (TC, TTC, EC, EN, SARS-CoV-2, T, conductivity, pH and turbidity) to assess the behavior of the data in a multivariate sense. The CA aimed to search similarities between sampling points. This multivariate statistical test was performed by the Ward Method. Euclidean distances were calculated for each case. The DA was applied to distinguish between sampling points. The goal is to discriminate two or more groups of objects by maximizing their differences (Di Renzo et al., 2016). All the variables were standardized before the multivariate performance, to eliminate the effect of scale.

Statistical analyses were run with the InfoStat software (Di Rienzo et al., 2016) and RStudio version 4.0.0 (Ihaka and Gentleman, 1996; R Development Core Team, 2005; RStudio Team, 2020; Wickham, 2009).

Finally, the parametric Pearson's test was performed to evaluate if there was correlation between the concentrations of HPyV and RNaseP. In addition, Spearman test was performed to assess for correlation between the evolution of the concentration of SARS-CoV-2 in the Arenales River, at AR-1 and AR-2, and the number of active reported cases in Salta city. Then the test was applied for the normalized SARS-CoV-2 concentration at AR-2 and the number of cases.

3. Results

A total of 15 sampling events were carried out from July to December of 2020 in three rivers, monitoring a total of five points: AR-1 and AR-2 at the Arenales River, CR at La Caldera River, and MR-1 and MR-2 at Mojotoro River.

3.1. Physico-chemical characterization of water samples

Four physico-chemical variables (temperature, pH, conductivity, and turbidity) were measured in water (Table 2). Water temperature (7.2 - 23.8 °C) was directly affected by environmental temperature (5.0 - 25.0 °C) (July to middle September is winter and middle September to middle December is spring in the South Hemisphere), as expected. Although the maximum temperatures were similar for all the rivers, the minimum values were higher for AR-1 and AR-2 (Arenales River is in the middle of the city) compared to those from CR, MR-1 and MR-2 which are in the vicinities of Salta, in open space and more exposed to the wind and elements.

Table 2. Physico-chemical variables measured in surface water, along 15 sampling events, in the Arenales River at AR-1 and AR-2 (before and after the discharges of the southern wastewater treatment plant, WWTP-S, respectively; distance between AR-1 and AR-2 is 1.4 km and WWTP-S is separated 0.7 km from each), in La Caldera River (CR), and in the Mojotoro River at MR-1 (after an old wastewater stabilization pond and before the discharges of the northern wastewater treatment plant, WWTP-N) and MR-2 (after the discharges of the WWTP-N; distance between MR-1 and MR-2 is 10 km and WWTP-N is 5 km away from each). K-W: Bonferroni Post Hoc test from Kruskal Wallis. Medians with different letters are significantly different ($p < 0.05$).

Variables	Sampling points	Mean $\pm$ SD	Median	Range (min-max)	K-W
Temperature (°C)	AR-1	17.41 $\pm$ 3.99	16.30	12.10 - 23.50	AB
	AR-2	19.23 $\pm$ 3.19	18.40	14.70 - 23.80	A
	CR	15.83 $\pm$ 4.73	14.30	7.20 - 23.80	C
	MR-1	15.69 $\pm$ 4.18	15.00	8.00 - 22.50	A
	MR-2	16.28 $\pm$ 4.88	16.00	7.80 - 23.80	BC

pH	AR-1	8.07 ± 0.27	8.12	7.64 - 8.54	E
	AR-2	7.90 ± 0.12	7.91	7.70 - 8.09	E
	CR	8.45 ± 0.28	8.46	7.89 - 8.82	D
	MR-1	7.96 ± 0.40	7.94	6.99 - 8.90	D
	MR-2	8.24 ± 0.32	8.24	7.32 - 8.63	D
Conductivity	AR-1	0.46 ± 0.07	0.48	0.31 - 0.54	G
(mS/cm)	AR-2	0.58 ± 0.07	0.59	0.41 - 0.74	G
	CR	0.21 ± 0.03	0.20	0.17 - 0.27	F
	MR-1	0.24 ± 0.01	0.24	0.21 - 0.27	F
	MR-2	0.25 ± 0.02	0.24	0.20 - 0.28	F
Turbidity	AR-1	17.07 ± 32.51	8.00	4.00 - 134.00	H
(NTU)	AR-2	88.27 ± 28.44	80.00	55.00 - 159.00	J
	CR	16.07 ± 13.78	13.00	5.00 - 56.00	HI
	MR-1	11.47 ± 5.57	10.00	5.00 - 26.00	HI
	MR-2	16.27 ± 5.36	16.00	5.00 - 26.00	I

310

311 Kruskal-Wallis test was performed with physico-chemical variables measured in the five
312 sampling points analyzed. The sampling points CR, MR-1 and MR-2 were similar except
313 for the temperature that was different in MR-1. The two monitoring points at the Arenales
314 River (AR-1 and AR-2; distance was 1.4 km, the WWTP-S was 0.7 km away from each)
315 only showed significant differences in the turbidity, which was much higher in AR-2,
316 probably due to the discharges of the WWTP-S, also reported in a previous work (Poma et
317 al, 2012).

318

319

3.2. Bacterial characterization

The highest concentrations of all FIBs were found in the Arenales River, followed by Mojotoro River and La Caldera River (Figure 2). In general, variability in bacteria concentrations (within two orders of magnitude) was lower in AR-1 and AR-2 than in the other sampling points (Figure 2). Conversely, the concentrations of total and thermotolerant coliforms were highly variable for MR-1, MR-2, and CR.

Bacteria concentrations in Arenales River exceeded 10^4 CFU or MPN per 100 mL and furthermore, they surpassed the acceptable limits for *E. coli* and enterococci (235 CFU/100 mL and 71 CFU/100 mL, respectively, USEPA 2012) in all the samples analyzed. The point AR-2, 700 m downstream from AR-1, was highly impacted by fecal contamination of untreated or insufficiently treated wastewater effluents from the WWTP-S ($p < 0.05$ for all the FBI) (Table 3).

Regarding Mojotoro River, bacteria concentrations were similar in both sampling sites (Table 3) showing that the effluents of the WWTP-N have no impact on the microbial quality of water. However, the concentration of *E. coli* and enterococci exceeded the recreational limits in most of the samples (67 and 80% for MR-1, respectively, and 53 and 33% for MR-2, respectively). In contrast, La Caldera River showed the best water quality, appropriate for recreational activities. In fact, all the samples analyzed (15 in total) were within the accepted values for *E. coli*; however, one of them exceeded the limit for enterococci.

Figure 2. Violin plots representing indicator bacteria determined in 15 monitoring events in five different sampling points (75 samples in total): AR-1 and AR-2 in the Arenales River (before and after the discharges of the southern wastewater treatment plant, respectively), CR in La Caldera River, and MR-1 and MR-2 in the Mojotoro River (before and after the discharges of the northern wastewater treatment plant, respectively). The concentrations of total (TC) and thermotolerant coliforms (TTC) are expressed as the Most Probable

Number per 100 mL, while those of *E. coli* (EC) and *Enterococcus* sp. (EN) are in Colony Forming Units (CFU) per 100 mL. Violin shapes show the data kernel probability with the boxplots embedded. Boxplots show the interquartile range (IQR) in boxes divided by median values (horizontal lines) with whiskers depicting ± 1.5 IQR and outliers as points. The segmented horizontal lines represent the limit values accepted by legislation for recreational water (USEPA, 2012).

Table 3. Bonferroni Post Hoc test from Kruskal Wallis for fecal indicator bacteria (total and thermotolerant coliforms, *E. coli*, and enterococci) determined in five sampling points: CR in La Caldera River, MR-1 and MR-2 in Mojotoro River, and AR-1 and AR-2 in Arenales River. Medians with different letters for each variable are significantly different ($p < 0.05$).

	Total coliforms	Thermotolerant coliforms	<i>E. coli</i>	Enterococci
CR	A	A	A	A
MR-1	A	A	B	B
MR-2	A	A	B	AB
AR-1	B	B	C	C
AR-2	B	C	C	C

3.3. Quantification of SARS-CoV-2, HPyV and RNase P

All the samples (75 in total) collected during the 15 sampling events (from July to December 2020, during the first wave of COVID-19) at five different points in three rivers, were concentrated, the nucleic acids were extracted, and then analyzed for SARS-CoV-2, HPyV and RNase P (Figure 3). In La Caldera River, from the three targets evaluated only SARS-CoV-2 was found in around half of the samples (53.3% samples were non-detects)

and in those cases the viral concentrations were lower than 10^5 gc/L. The same occurred in the Mojotoro River where 7 and 6 out of 15 (46.6 and 40.0%, respectively) samples were non detects at MR-1 and MR-2, respectively. However, HPyV was also detected in that river in both sampling points, in 14 and in 10 (out of 15) samples at MR-1 and MR-2, respectively. The HPyV was detected in all the samples in Arenales River, in both sampling points, showing the impact of human contamination. This effect was also seen through the detection of RNase P that was only found in AR-2 samples. While SARS-CoV-2 was not detected during the first five monitoring campaigns in AR-1, it was quantified in all samples collected at AR-2. The two targets selected for tracing human contamination showed a similar behavior (coefficient was 0.7228; $p = 0.0021$; positive strong correlation) along time. However, the concentrations of HPyV were around one order of magnitude higher than those of RNase P, which may be the reason for RNase P not being found consistently in other sites (average ratio HPyV/RNaseP was 15.8 ± 6.4 , without considering one outlier on 27 July, where the ratio was 299.3).

Figure 3. Evolution of SARS-CoV-2 (C_{CoV} , red solid line), human polyomavirus (C_{HPyV} , green solid line), and RNase P ($C_{RNase P}$, blue solid line) concentrations, determined in five sampling points: AR-1 and AR-2 in Arenales River, MR-1 and MR-2 in Mojotoro River, and CR in La Caldera River, from July to December 2020 (15 sampling events). Dashed lines represent the limit of detection (LOD) for SARS-CoV-2 (red), HPyV (green) and RNase P (blue). Non detects were arbitrarily represented as LOD/2. Values plotted are the average

of duplicates and error bars correspond to the standard deviation (some bars are too small to be seen). Concentrations are expressed in gene copies per liter (gc/L).

3.4. Relationship between water quality and SARS-CoV-2 concentration

Spearman correlation was evaluated using all the physico-chemical, FIBs and SARS-CoV-2 concentration data, including all the sampling points (Supplementary Information, Table S1). Temperature was the only variable that did not show any correlation. Positive strong correlations were found between pH and COND, TURB and FIBs, and among FIBs. Regarding CoV, also positive strong correlations were confirmed with pH and COND, while they were moderate with TURB and all FIBs (Supplementary Information, Table 1).

When Spearman correlation was performed for each sampling point instead, positive strong or moderate correlations were found ($p < 0.05$) between COND and pH (and with temperature for MR-2 only) and also between TC, TTC and EC for all the sampling points (Table 4). In all cases, except for MR-2, there were positive moderate correlations ($p < 0.05$) between TURB and at least two FIBs. Only for AR-1 and AR-2 positive moderate correlations ($p < 0.05$) were found between CoV and pH, CoV and TURB, and EN and another FIB (Table 4).

Table 4. Coefficients and probabilities (below and above the main diagonal, respectively) from Spearman correlation test using the physico-chemical variables, fecal indicator bacteria and SARS-CoV-2 concentration (CoV) for each sampling point (SP): AR-1 and AR-2 in Arenales River; CR in La Caldera River; MR-1 and MR-2 in Mojotoro River. Only significant correlations ($p < 0.05$) are shown and those that are strong are indicated in bold. Physico-chemical variables: temperature (T), pH, conductivity (COND), turbidity (TURB). Fecal indicator bacteria: total coliforms (TC), thermotolerant coliforms (TTC), *E. coli* (EC), enterococci (EN).

SP		T	pH	COND	TURB	TC	TTC	EC	EN	CoV
AR-1	T									
	pH			1.8x10 ⁻⁴						4.6x10 ⁻³
	COND		1							4.6x10 ⁻³
	TURB							0.01		
	TC						0.04	0.04		
	TTC					0.53				
	EC				0.65	0.54			0.03	
	EN							0.57		
	CoV		0.69	0.69						
AR-2	T								0.01	
	pH			1.8x10 ⁻⁴						2.3x10 ⁻³
	COND		1							2.3x10 ⁻³
	TURB								0.05	
	TC								0.02	
	TTC							3.7x10 ⁻³		
	EC						0.70			
	EN	0.66			0.51	0.59				
	CoV		0.81	0.81						
CR	T									
	pH			1.8x10 ⁻⁴						
	COND		1							
	TURB					0.01	0.04			
	TC				0.63		1.1x10 ⁻⁶	0.02		
	TTC				0.55	0.92		4.7x10 ⁻³

415

416 Cluster analysis including all the variables measured for a total of 75 water samples from
 417 five sampling points was performed and the Euclidean distances were determined
 418 (cophenetic correlation was 0.906) (Figure 4). The most similar sampling points were MR-

419 1 and MR-2 (Euclidean distance was 0.69), verifying once again that the Mojotoro River
 420 was not negatively impacted by the discharges of the WWTP-N. Conversely, the most
 421 different sampling points were CR and AR-2 (Euclidean distance was 6.99), in agreement
 422 with other observations, that CR was the point with the best water quality and AR-2 the
 423 most impacted one by the discharges of the WWTP-S.

424

425 **Figure 4.** Cluster analysis performed with nine variables (temperature, pH, conductivity,
 426 turbidity, and concentrations of total and thermotolerant coliforms, *E. coli*, enterococci and
 427 SARS-CoV-2) measured in a total of 75 water samples collected in five sampling points:
 428 AR-1 and AR-2 in the Arenales River, MR-1 and MR-2 in the Mojotoro River, and CR in La
 429 Caldera River, from July to December 2020 (15 sampling events).

430

431 The discriminant analysis performed with nine the variables measured in 75 water samples
 432 along 15 sampling events showed two main groups (Figure 5). At the right side there were
 433 those samples from the Arenales River, where AR-1 was also discriminated from AR-2,
 434 due to the high microbial contamination. The second group, at the left side, included the

435 samples from Mojotoro River, with MR-1 and MR-2 mingled together, slightly separated
436 from those from Caldera River, which was the one with best microbial water quality.

437

438 **Figure 5.** Plots of the first two axes (with eigenvalues higher than 1, explaining 97.7% of
439 the accumulated variance) obtained with a linear discriminant analysis conducted with nine
440 variables: temperature, pH, conductivity, turbidity, and concentrations of total coliforms,
441 thermotolerant coliforms, *E. coli*, enterococci, and SARS-CoV-2. A total of 75 water
442 samples were collected in five sampling points: AR-1 (red circles) and AR-2 (blue circles)
443 in the Arenales River, MR-1 (gray circles) and MR-2 (pink circles) in the Mojotoro River,
444 and CR (green circles) in La Caldera River, from July to December 2020 (15 sampling
445 events).

446

447 3.5. Comparison between SARS-CoV-2 concentration and reported COVID-19
448 cases

The total of COVID-19 reported cases in the city of Salta, accumulated for 14 days before each monitoring campaign, were assigned to that sampling date. The evolution of the number of cases in Salta city was confronted with the SARS-CoV-2 concentration found in the Arenales River (Figure 6). This was not performed for the Mojotoro and La Caldera Rivers as SARS-CoV-2 was detected in around half of the samples, following no pattern, in low concentration, while the number of cases were constantly increasing.

Figure 6. Evolution of SARS-CoV-2 concentration (C_{CoV}) in the Arenales River at two sampling points: AR-1 and AR-2 before and after the southern wastewater treatment plant, respectively, and of the number of COVID-19 active cases accumulated for 14 days before sampling. LOD: limit of detection for SARS-CoV-2 in surface water. Non-detects were arbitrarily represented as LOD/2.

The viral concentration curve anticipated the epidemiological one. The ratio between the viral concentration and the number of active cases (14 days before) was variable, being 32 gc/mL/case the average for AR-1 and 1099 gc/mL/case the average for AR-2. The latest was a consequence of a higher fecal content in AR-2 due to the WWTP discharges in the river and was highly influenced by the extreme values. In fact, at the beginning and towards the end of the epidemiological curve the ratio was around 281 gc/mL/case while it was 2034 gc/mL/case during the two months (August-September) when the viral dissemination was higher.

Spearman correlation coefficient indicated a positive association between the number of cases and the viral concentrations in the Arenales River, at AR-1 ($\rho = 0.89$; $p = 0.00001$) and at AR-2 ($\rho = 0.89$; $p = 0.00084$).

3.6. Normalization of the SARS-CoV-2 concentration

The experimental concentration of SARS-CoV-2, although is calculated based on the detection of the gene of interest in water samples, is subject to fluctuations due to variations in the flow rate, i.e. from precipitations or from discharges of the wastewater plant. One strategy to compensate for that, thus, to help the data to get closer to the real situation, is to normalize the data, with some variable that accounts for the human input in this case. Two targets, T antigen for HPyV and RNase P, linked to human contamination were selected for that. Therefore, the concentration of SARS-CoV-2 in the Arenales River was normalized following three different alternatives, including the concentration of HPyV (Alternative 1), the concentration of RNase P (Alternative 2) and a combination of both (Alternative 3) (Figure 7).

Figure 7. Concentration of SARS-CoV-2 experimentally determined (C_{CoV}) at AR-2 in Arenales River and the normalized concentration using three different alternatives: (a) Alternative 1 ($CN1_{CoV}$), using the concentration of human polyomavirus (HPyV), according to equation 1, (b) Alternative 2 ($CN2_{CoV}$), using the concentration of RNase P, according to equation 2, and (c) Alternative 3 ($CN3_{CoV}$), using the concentration of HPyV and RNase P, according to equation 3.

Regarding the three alternatives for normalization of the SARS-CoV-2 concentration evaluated, the first one was as the ratio with the concentration of human polyomavirus (equation 1), in which case the shape of the curve was not modified. However, the scale of the resulting number was too small, and the sense of the viral concentration magnitude was lost, and the units were (gc of SARS-CoV-2/gc of HPyV). The latest also happened when the ratio of SARS-CoV-2 was calculated with RNase P (equation 2), being the units (gc of SARS-CoV-2/gc of RNase P); however, the shape of the resulting curve changed a bit due to the fluctuations of RNase P concentration. Instead, the third way of performing the normalization was using the two normalizers (HPyV and RNase P), according to equation 3. In this case, the shape of the experimental SARS-CoV-2 concentration remained constant, and the normalized values kept the sense of the concentration magnitude as well as the proper units. Thus, the normalized concentration of SARS-CoV-2, obtained using both HPyV and RNase P, was selected and compared to the number of reported COVID-19 cases in the city (Figure 8). A strong positive correlation was found by Spearman test between the number of COVID-19 cases and the normalized (alternative 3) viral concentrations at AR-2 ($\rho = 0.93$; $p = 0.00054$).

Figure 8. Evolution of the normalized concentration of SARS-CoV-2 (gc/L) (CN3CoV, according to equation 3) in the Arenales River at AR-2 and the accumulated reported COVID-19 cases, from the 12 July to 27 December 2020 during the first wave of COVID-19 in the city of Salta, Argentina.

4. Discussion

A dataset including experimental values for physico-chemical (four variables) and microbiological (four fecal indicator bacteria and SARS-CoV-2 concentration) variables from a total of 75 water samples from five sampling points was used to performed various bi- and multi-variable analyses. These allowed us to compare among sampling points located in the same or different rivers, to evaluate correlations, and to find the most similar and different points.

The results from the monitoring from July to December 2020, allowed to verify that from the three rivers analyzed La Caldera River has the best water quality, with low and

sporadic fecal input, adequate for recreational activities, which was also described in previous work (Chávez Díaz et al, 2020; Gutiérrez Cacciabue et al, 2014). The Mojotero and the Arenales Rivers receive the impact of the discharges of the two wastewater treatment plants of the city, WWTP-N and WWTP-S, respectively. The WWTP-N is new and has capacity for treating efficiently the flow rate from the northern part of the city, thus, the discharges do not affect the water quality of the river at least between MR-1 and MR-2. Instead, the WWTP-S is an outdated treatment plant without enough capacity for a fast-growing city. The discharges of the WWTP-S significantly impacted the Arenales River, verified at AR-2, although the water quality was already poor at AR-1, 700 m upstream, due to domestic and industrial activities and to sporadic illegal discharges of raw sewage (Poma et al., 2012). The latest was evidenced by the high concentrations of fecal bacteria indicators.

Environmental surveillance as a tool to know about the circulation of viruses and other microorganisms causing disease in the population has been proposed for decades already. In fact, rivers that were impacted by wastewater has been monitored in multiple studies, like the analyses of circulation of human polio virus as part of the effort for the eradication (Delogu et al., 2018; Iaconelli et al, 2017; WHO, 2003) and of Hepatitis A virus after vaccination campaigns (Blanco Fernandez et al, 2012a), or of evolutionary relationships between norovirus (Blanco Fernandez, 2012b).

In the case of environmental surveillance of SARS-CoV-2 almost all the efforts around the world were focused on monitoring wastewater (Medema et al., 2020; Randazzo et al., 2020b; Wu et al., 2020), most of the times analyzing the inlets (and sometimes also the outlets) of wastewater treatment plants (Gonzalez et al., 2020; Nasser et al., 2021), but also at maintenance holes that are representative of defined areas of a city, or regions or neighborhoods (Larson et al., 2020). In some cases, the efficiency of wastewater

treatment plants to remove SARS-CoV-2 was evaluated, as in the studies conducted by Wurtzer et al. (2020) in France and by Rimoldi et al. (2020) in Italy.

Wastewater samples are useful to learn about the true viral circulation in a population of geographical interest, especially to estimate the extension of virus dissemination due to asymptomatic people, that seems to play an important role in the COVID-19 pandemic (He et al., 2020). In addition, these samples present the advantages that they are anonymous and non-invasive, they do not depend on the population will (to be tested) under analysis and they can be used by the respective health authorities to make smart decisions regarding the pandemic's management (Thompson et al., 2020).

While monitoring wastewater is a useful tool to evaluate the situation in developed countries this may not be the case in threshold or developing countries, where the sewage systems are insufficient or non-existent, or when the wastewater treatment is inefficient. In most of those places, rivers or other water bodies receive the discharges of untreated (sometimes illegally) or insufficiently treated wastewater, impacting the water quality and thus, the use of the resource (Rimoldi et al., 2020). In such scenarios, surface water like natural or artificial ponds (Iglesias et al., 2020) or rivers could also be used as tools for assessing the epidemiological situation of the population. The first attempt in that direction was done by Guerrero-Latorre and coworkers (2020) that evaluated the presence of SARS-CoV-2 in a context of low sanitation countries. They sampled once in natural streams located in three different regions of Quito, Ecuador. Although they quantified SARS-CoV-2 and human adenoviruses, as well as the number of reported cases, the study is limited due to the small number of samples analyzed. Issues like the presence, persistence, and potential infectivity of SARS-CoV-2 detected in the water environment have been under the discussion for the past year (Carducci et al., 2020; La Rosa et al., 2020).

According to the results of this and previous works, the Arenales River can properly play that role as it does reflect the pathogens circulating in the population. In fact, previous studies performed in the Arenales River revealed the circulation of multiples pathogens including bacteria, parasites, and various enteric viruses (Poma et al, 2012; Blanco Fernandez et al., 2012a; Blanco Fernandez et al., 2012b; Pisano et al., 2018; Prez et al., 2020). Furthermore, the microbiological risk for the population of being in contact with those water through recreational activities was estimated for enterovirus and norovirus (Poma et al, 2019).

The representation of the viral concentration in river water allows to see the variation along time. It is like a succession of instant photos (or moments) showing the situation in that specific point under analysis. Although it allows to understand the magnitude of the viral circulation in the area, there are some limitations that should be considered, especially if the intention is to estimate the number of infected people. A water body like a river will show flow rate variations due to the input of stormwater, illegal raw sewage, and industrial effluents, among others. In other words, the viral concentration will depend on all those human or natural contributions; thus, those flow rates or dilution factors should be considered if the absolute number of virus were to be calculated. On the other hand, there could be some discussion about the influence of population dynamic in the area of impact. One possibility to account for population dynamics and for other human and non-human inputs is to normalize the viral concentration using some other target or surrogate (Medema et al, 2020b). Some chemical compounds, like fecal sterols like coprostanol (Daughton, 2012; Chen et al, 2014), other viruses common in human urinary tract, like human polyomavirus (HPyV) (McQuaig et al., 2009) or pepper mild mottle virus (PMMoV) (D'Aoust et al, 2020), or bacterial fecal indicators like human Bacteroidales (D'Aoust et al, 2020), just to mention some, have been suggested for normalization. The advantage of

using any of them is that, as they are excreted by all the population all the time, then cultural or seasonal effects could be neglected.

In addition to the detection of SARS-CoV-2, two other targets linked to human contamination were quantified in this work. One of them was the gene that encodes for human RNase P, which was used in many studies as an endogenous internal control (Arnold et al., 2020; Boddicker et al., 2007; Dahdouh, et al., 2020), and it has been extensively used as a sample quality control in clinical tests in the current COVID-19 pandemic. The second one was the marker for human polyomaviruses, which has been proposed as a viral indicator of human fecal contamination and it is known to be stable in the environment. In addition, these viruses are excreted by feces and urine fluids (McQuaig et al., 2009; Bofill-Mas et al., 2000), with the species JC and BK reported as the most frequent and prevalent HPyV detected in sewage samples of this geographical region (Barrios et al., 2018).

Neither RNase P nor HPyV were detected in La Caldera River, showing again that fecal human contamination was just sporadic. In the case of Mojotoro River, RNase P was not detected and HPyV was present only in some of the samples, proving that those waters were only slightly impacted by wastewater. Conversely, in the case of the Arenales River, highly impacted by wastewater discharges, HPyV was quantified at both sampling points and the concentration was persistent and steady along the monitoring period. This was also the case for RNase P, which was only detected at AR-2.

Regarding normalization, although most of the times it has been done with only one target as reference, calculating the ratio between the gene of interest and the normalizer (McQuaig et al., 2009), it is highly recommended to use multiple genes for a more accurate normalization. For averaging of the control genes, it is recommended to use the geometric instead of the arithmetic mean, as the former controls are better for possible

outlying values and abundance differences between the different genes (Vandesompele et al., 2002).

In this case the concentration of SARS-CoV-2 from AR-2 was normalized in three different ways. The first one was as the ratio with the concentration of human polyomavirus, in which case the shape of the curve was not modified, the scale of results became too small, and the sense of the viral concentration magnitude was lost, plus the units were gc of SARS-CoV-2 per gc of HPyV. A similar situation resulted when the ratio of SARS-CoV-2 was calculated with RNase P (gc of SARS-CoV-2/gc of RNase P), although the curved was slightly modified due to the fluctuations of RNase P concentration. Instead, the third way of normalizing, using both HPyV and RNase P according to equation 3, seemed more appropriate as the shape of the original concentration curved was not changed but mostly because the normalized values kept the sense of the magnitude.

In the case of the Arenales River the flow rate fluctuations were not measured; however, the difference in flow rate between the sampling points AR-1 and AR-2 (separated 700 m, no additional inputs verified) are only due to the discharges of the WWTP-S. What is interesting is that this discharge, far from producing a “dilution effect”, decreasing the viral concentration, has an “addition effect” increasing it due to the huge fecal contribution. This effect observed in the concentration of SARS-CoV-2 was also reflected in the concentration of the bacterial indicators measured. The magnitude of the discharge modified substantially, in a short distance, the microbiological and physico-chemical characteristics of the water river, differentiating this sampling point from all the others that were analyzed and minimizing any other effect related to the population dynamic or to seasonal effects. Finally, strong correlations were found between both the experimental and the normalized SARS-CoV-2 concentrations and the curve of active COVID-19 cases in the city of Salta. In this way, the water from the Arenales River, at AR-2, reflected the

population situation regarding viral circulation of COVID-19 cases, therefore, it could be well used as a surveillance tool.

This is, to the best of our knowledge, the first study that showed the dynamic of SARS-CoV-2 concentration in an urban river highly impacted by wastewater and proved that it can be used for SARS-CoV-2 surveillance to support health authorities' decisions.

5. Conclusions

- From the three rivers analyzed La Caldera River had the best water quality, and indicator bacteria were within acceptable limits for recreational activities.
- Although Mojotoro River receives the discharges of an old wastewater stabilization pond before MR-1 and the discharges of the northern wastewater treatment plant of the city before MR-2, they did not have a negative effect on the water quality.
- The Arenales River presented the poorest water quality. In addition to the contamination existing at AR-1 the discharges of the southern wastewater treatment plant contributed to the significant increase of fecal contamination at AR-2.
- SARS-CoV-2 was found in about half of the samples analyzed in low concentrations in La Caldera and Mojotoro Rivers. Instead, the virus concentration in the Arenales River was persistently high.
- Two human tracers, human polyomavirus (HPyV) and RNase P, were analyzed. None of them was detected in La Caldera River, only HPyV was found in Mojotoro River and at AR-1 in Arenales River, and both were quantified at AR-2 (Arenales River).
- The Arenales River, at AR-2, was highly impacted by wastewater and the concentration of SARS-CoV-2 was normalized using both human tracers. The experimental and the normalized concentrations correlated with the curve of

accumulated reported COVID-19 cases in the city. Thus, the Arenales River at AR-2 reflects the epidemiological situation of the city.

Acknowledgements

This research was funded by Project COVID-19 233-785, from Fondo para la Investigación Científica y Tecnológica (FONCyT), Agencia Nacional de Promoción de la Investigación, el Desarrollo Tecnológico y la Innovación, Argentina. María Noel Maidana Kulesza, Diego Gastón Sanguino-Jorquera, Sarita Reyes, María del Milagro Said-Adamo, and Martín Mainardi Remis are recipients of doctoral fellowships from CONICET.

References

- Albastaki, A., Naji, M., Lootah, R., Almeheiri, R., Almulla, H., Almarri, I., Alreyami, A., Aden, A., Alghafri, R. (2021) First confirmed detection of SARS-COV-2 in untreated municipal and aircraft wastewater in Dubai, UAE: The use of wastewater-based epidemiology as an early warning tool to monitor the prevalence of COVID-19. *Science of The Total Environment* 760, 143350.
- Amirian, E. S. (2020) Potential fecal transmission of SARS-CoV-2: current evidence and implications for public health. *International Journal of Infectious Diseases* 95, 363-370.
- Arnold, M. T., Temte, J. L., Barlow, S. K., Bell, C. J., Goss, M. D., Temte, E. G., Checovich, M. M., Reisdorf, E., Scott, S., Guenther, K., Wedig, M., Shult, P., Uzicanin, A. (2020) Comparison of participant-collected nasal and staff-collected oropharyngeal specimens for human ribonuclease P detection with RT-PCR during a community-based study. *PloS One* 15(10), e0239000.

697 Boddicker, J. D., Rota, P. A., Kreman, T., Wangeman, A., Lowe, L., Hummel, K. B.,
698 Thompson, R., Bellini, W. J., Pentella, M., DesJardin, L. E. (2007) Real-time reverse
699 transcription-PCR assay for detection of mumps virus RNA in clinical specimens.
700 Journal of Clinical Microbiology 45(9), 2902-2908.

701 Barrios, M. E., Fernández, M. D. B., Cammarata, R. V., Torres, C., Mbayed, V. A. (2018)
702 Viral tools for detection of fecal contamination and microbial source tracking in
703 wastewater from food industries and domestic sewage. Journal of Virological Methods
704 262, 79-88.

705 Betancourt, W. Q., Schmitz, B. W., Innes, G. K., Prasek, S. M., Brown, K. M. P., Stark, E.
706 R., Foster, A. R., Sprissler, R. S., Harris, D. T., Sherchan, S. P., Gerba, C. P., Pepper,
707 I. L. (2021) COVID-19 containment on a college campus via wastewater-based
708 epidemiology, targeted clinical testing and an intervention. Science of the Total
709 Environment 779, 146408.

710 Blanco Fernández, M. D., Torres, C., Riviello-López, G., Poma, H. R., Rajal, V. B., Nates,
711 S., Cisterna, D. M., Campos, R. H., Mbayed, V. A. (2012a) Analysis of the circulation of
712 hepatitis A virus in Argentina since vaccine introduction. Clinical Microbiology and
713 Infection 18(12), E548-E551.

714 Blanco Fernández, M. D., Torres, C., Poma, H. R., Riviello-López, G., Martínez, L. C.,
715 Cisterna, D. M., Rajal, V. B., Nates, S., Mbayed, V. A. (2012b) Environmental
716 surveillance of norovirus in Argentina revealed distinct viral diversity patterns,
717 seasonality and spatio-temporal diffusion processes. Science of the Total Environment
718 437, 262-269.

719 Bofill-Mas, S., Pina, S., Girones, R. (2000) Documenting the epidemiologic patterns of
720 polyomaviruses in human populations by studying their presence in urban sewage.
721 Applied and Environmental Microbiology 66(1), 238-245.

722 Carducci, A., Federigi, I., Liu, D, Thompson, J. R., Veran M. (2020) Making Waves:
723 Coronavirus detection, presence and persistence in the water environment: State of
724 the art and knowledge needs for public health. Water Research 179 (2020) 115907

725 CDC (2020) 2019-Novel Coronavirus (2019-nCoV) Real-Time rRT-PCR Panel Primers
726 and Probes. Centers for Disease Control and Prevention, U.S.
727 <https://www.cdc.gov/coronavirus/2019-ncov/lab/rt-pcrpanel-primer-probes.html> (Last
728 accessed 16th June, 2020).

729 Chávez-Díaz, L. V., Gutiérrez-Cacciabue, D., Poma, H. R., Rajal, V. B. (2020) Sediments
730 quality must be considered when evaluating freshwater aquatic environments used for
731 recreational activities. International Journal of Hygiene and Environmental Health 223,
732 159-170.

733 Chen, C., Kostakis, C., Gerber, J. P., Tschärke, B. J., Irvine, R. J., White, J. M. (2014)
734 Towards finding a population biomarker for wastewater epidemiology studies. Science
735 of the Total Environment 487, 621-628.

736 Cruz, M. C., Gutiérrez Cacciabue, D., Gil, J. F., Gamboni, O., Vicente, M. S., Wuertz, S.,
737 Gonzo, E. E., Rajal, V. B (2012) The impact of point source pollution on shallow
738 groundwater used for human consumption in a threshold country. Journal of
739 Environmental Monitoring 14, 2338-2349.

740 Di Rienzo, J. A., Casanoves, F., Balzarini, M. G., Gonzalez, L., Tablada, M., Robledo, C.
741 W. (2016) InfoStat version 2016. Grupo InfoStat, FCA, Universidad Nacional de
742 Córdoba, Argentina. URL <http://www.infostat.com.ar>.

743 Dahdouh, E., Lázaro-Perona, F., Romero-Gómez, M. P., Mingorance, J., García-
744 Rodríguez, J. (2021) Ct values from SARS-CoV-2 diagnostic PCR assays should not
745 be used as direct estimates of viral load. *Journal of Infection* 82(3), 414-451.

746 D'Aoust, P. M., Mercier, É., Montpetit, D., Jia, J. J., Alexandrov, I., Tariq Baig, A., Mayne,
747 J., Zhang, X., Alain, T., Servos, M. R., MacKenzie, M., Figeys, D., MacKenzie, A. E.,
748 Graber, T. E., Delatolla, R. (2020) Quantitative analysis of SARS-CoV-2 RNA from
749 wastewater solids in communities with low COVID-19 incidence and prevalence. *Water*
750 *Research* 188, 116560.

751 Daughton, C. G. (2012) Using biomarkers in sewage to monitor community-wide human
752 health: isoprostanes as conceptual prototype. *Science of the Total Environment* 424,
753 16-38.

754 Delogu, R., Battistone, A., Buttinelli, G., Fiore, S., Fontana, S., Amato, C., Cristiano, K.,
755 Gamper, S., Simeoni, J., Frate, R., Pellegrinelli, L., Binda, S., Veronesi, L., Zoni, R.,
756 Castiglia, P., Cossu, A., Triassi, M., Pennino, F., Germinario, C., Balena, V., Cicala, A.,
757 Mercurio, P., Fiore, L., Pini, C., Stefanelli, P. (2018) Poliovirus and other enteroviruses
758 from environmental surveillance in Italy, 2009–2015. *Food and environmental*
759 *virology* 10(4), 333-342.

760 Eaton, A. D., Clesceri, L. S., Rice, E. W., Greenberg, A. E., Franson, M. A. H. (2005)
761 *Standard Methods for the Examination of Water and Wastewater*. 21st Edition. APHA.
762 American Public Health Association.

763 Gonzalez, R., Curtis, K., Bivins, A., Bibby, K., Weir, M. H., Yetka, K., Thompson, H.,
764 Keeling, D., Mitchell, J., Gonzalez, D. (2020) COVID-19 surveillance in Southeastern
765 Virginia using wastewater-based epidemiology. *Water Research* 186, 116296.

766 Gorbalenya, A. E., Baker, S. C., Baric, R., Groot, R. J. D., Drosten, C., Gulyaeva, A. A.,
767 Haagmans, B. L., Lauber, C., Leontovich, A. M., Neuman, B. W., Penzar, D., Perlman,
768 S., Poon, L., Samborskiy, D., Sidorov, I. A., Solá-Gurpegui, I., Ziebuhr, J. (2020)
769 Severe acute respiratory syndrome-related coronavirus: The species and its viruses—a
770 statement of the Coronavirus Study Group. *Nature Microbiology*, DOI:
771 10.1038/s41564-020-0695-z.

772 Guan, W. J., Ni, Z. Y., Hu, Y., Liang, W. H., Ou, C. Q., He, J. X., Liu, L., Shan, H. Lei, C-L,
773 Hui, D., Du, B., Li, L. J., Zeng, G., Yuen, K. Y., Chen, R. C., Tang, C. L., Wang, T.,
774 Chen, P. Y., Xiang, J., Li, S. Y., Wang, J. L., Liang, Z., Peng, Y., Wei, L., Liu, Y., Hu,
775 Y., Peng, P., Wang, J., Liu, J., Zhong, C., Li, G., Zheng, Z., Qiu, S., Luo, J., Ye, C.,
776 Zhu, S., Zhong, N. (2020) Clinical characteristics of 2019 novel coronavirus infection
777 in China. *New England Journal of Medicine* 382(18), 1708-1720.

778 Guerrero-Latorre, L., Ballesteros, I., Villacrés-Granda, I., Granda, M. G., Freire-Paspuel,
779 B., Ríos-Touma, B. (2020) SARS-CoV-2 in river water: Implications in low sanitation
780 countries. *Science of the Total environment* 743, 140832.

781 Gutiérrez Cacciabue, D., Teich, I., Poma, H. R., Cruz, M. C., Balzarini, M., Rajal, V. B.
782 (2014) Strategies to optimize monitoring schemes of recreational waters using a
783 multivariate approach. *Environmental Monitoring and Assessment* 186(12), 8359-8380.

784 Harrison, A. G., Lin, T., Wang, P. (2020) Mechanisms of SARS-CoV-2 transmission and
785 pathogenesis. *Trends in Immunology* 41(12), 1100-1115.

786 He, D., Zhao, S., Lin, Q., Zhuang, Z., Cao, P., Wang, M. H., Yang, L. (2020) The relative
787 transmissibility of asymptomatic COVID-19 infections among close contacts.
788 *International Journal of Infectious Diseases* 94, 145-147.

789 Iaconelli, M., Muscillo, M., Della Libera, S., Fratini, M., Meucci, L., De Ceglia, M., Giacosa,
790 D., La Rosa, G. (2017) One-year surveillance of human enteric viruses in raw and
791 treated wastewaters, downstream river waters, and drinking waters. Food and
792 Environmental Virology 9(1), 79-88.

793 Iglesias, N. G., Gebhard, L. G., Carballeda, J. M., Aiello, I., Recalde, E., Terny, G.,
794 Ambrosolio, S., L'Arco, G., Konfino, J., Brardinelli, J. I. (2020) SARS-CoV-2
795 surveillance in untreated wastewater: first detection in a low-resource community in
796 Buenos Aires, Argentina. MedRxiv preprint;
797 <https://doi.org/10.1101/2020.10.21.20215434>.

798 Ihaka, R., Gentleman, R. (1996) R: a language for data analysis and graphics. Journal of
799 Computational and Graphical Statistics 5(3), 299-314.

800 Jarrous, N., Altman, S. (2001) Human ribonuclease P. Methods in Enzymology 342, 93-
801 100.

802 Jeong, H. W., Kim, S. M., Kim, H. S., Kim, Y. I., Kim, J. H., Cho, J. Y., Kim, S. H., Kang,
803 H., Kim, S. G., Park, S. J., Kim, E. H., Choi, Y. K. (2020) Viable SARS-CoV-2 in
804 various specimens from COVID-19 patients. Clinical Microbiology and Infection 26(11),
805 1520-1524.

806 Jiang, X., Luo, M., Zou, Z., Wang, X., Chen, C., Qiu, J. (2020) Asymptomatic SARS-CoV-2
807 infected case with viral detection positive in stool but negative in nasopharyngeal
808 samples lasts for 42 days. Journal of Medical Virology 92(10), 1807-1809.

809 Kampf, G., Todt, D., Pfaender, S., Steinmann, E. (2020) Persistence of coronaviruses on
810 inanimate surfaces and their inactivation with biocidal agents. Journal of Hospital
811 Infection 104(3), 246-251.

812 Kumar, M., Patel, A. K., Shah, A. V., Raval, J., Rajpara, N., Joshi, M., Joshi, C. G. (2020)
813 First proof of the capability of wastewater surveillance for COVID-19 in India through
814 detection of genetic material of SARS-CoV-2. Science of The Total Environment 746,
815 141326.

816 La Rosa, G., Bonadonna, L., Lucentini, L., Kenmoe, S., Suffredini, E. (2020) Coronavirus
817 in water environments: Occurrence, persistence and concentration methods - A
818 scoping review. Water Research 179 (2020) 115899.

819 Larson, R. C., Berman, O., Nourinejad, M. (2020). Sampling manholes to home in on
820 SARS-CoV-2 infections. PLoS ONE 15(10): e0240007.

821 Lomniczi, I., Boemo, A., Musso, H. (2007) Location and characterization of pollution sites
822 by principal component analysis of trace contaminants in a slightly polluted seasonal
823 river: A case study of the Arenales River (Salta, Argentina). Water SA 33(4).

824 McQuaig, S. M., Scott, T. M., Lukasik, J. O., Paul, J. H., Harwood, V. J. (2009)
825 Quantification of human polyomaviruses JC virus and BK virus by TaqMan quantitative
826 PCR and comparison to other water quality indicators in water and fecal samples.
827 Applied and Environmental Microbiology 75(11), 3379-3388.

828 Medema, G., Heijnen, L., Elsinga, G., Italiaander, R., Brouwer, A. (2020a) Presence of
829 SARS-Coronavirus-2 RNA in sewage and correlation with reported COVID-19
830 prevalence in the early stage of the epidemic in the Netherlands. Environmental
831 Science & Technology Letters 7(7), 511-516.

832 Medema, G., Been, F., Heijnen, L., Petersen, S. (2020b) Implementation of environmental
833 surveillance for SARS-CoV-2 virus to support public health decisions: Opportunities
834 and challenges. Current Opinion in Environmental Science & Health 17, 49-71.

835 Mukhra, R., Krishan, K., Kanchan, T. (2020) Possible modes of transmission of Novel
836 coronavirus SARS-CoV-2: a review. *Acta Bio-Medica* 91(3), e2020036.
837 <https://doi.org/10.23750/abm.v91i3.10039>

838 Nasser, S., Yavarian, J., Norouzi Baghani, A., Mokhtari Azad, T., Nejati, A., Nabizadeh,
839 R., Hadi, M., Shafiei Jandaghi, N. Z., Vakili, B., Azam Vaghefi, S. K., Baghban, M.,
840 Yousefi, S., Nazmara, S., Alimohammadi, M. (2021) The presence of SARS-CoV-2 in
841 raw and treated wastewater in 3 cities of Iran: Tehran, Qom and Anzali during
842 coronavirus disease 2019 (COVID-19) outbreak. *Journal of Environmental Health*
843 *Science and Engineering*. <https://doi.org/10.1007/s40201-021-00629-6>.

844 Park, S. K., Lee, C. W., Park, D. I., Woo, H. Y., Cheong, H. S., Shin, H. C., Ahn, K., Kwon,
845 M. J., Joo, E. J. (2020) Detection of SARS-CoV-2 in Fecal Samples from Patients with
846 Asymptomatic and Mild COVID-19 in Korea. *Clinical Gastroenterology and Hepatology*.
847 <https://doi.org/10.1016/j.cgh.2020.06.005>.

848 Peiris, J. S. M., Lai, S. T., Poon, L. L. M., Guan, Y., Yam, L. Y. C., Lim, W., Nicholls, J.,
849 Yee, W. K. S., Yan, W. W., Cheung, M. T., Cheng, V. C. C., Chan, K. H. Tsang, D. N.
850 C., Yung, R. W. H., Ng, T. K., Yuen, K. Y. (2003) Coronavirus as a possible cause of
851 severe acute respiratory syndrome. *The Lancet* 361(9366), 1319-1325.

852 Pisano, M. B., Lugo, B., Poma, H. R., Cristóbal, H. A., Raskovsky, V., Martínez Wassaf, M.
853 G., Rajal, V. B., Ré, V. E. (2018) Environmental hepatitis E virus detection supported
854 by serological evidence in the northwest of Argentina. *Transactions of the Royal*
855 *Society of Tropical Medicine and Hygiene* 112(4), 181-187.

856 Poma, H. R., Cacciabue, D. G., Garcé, B., Gonzo, E. E., Rajal, V. B. (2012) Towards a
857 rational strategy for monitoring of microbiological quality of ambient waters. *Science of*
858 *the Total Environment* 433, 98-109.

859 Poma, H. R., Kundu, A., Wuertz, S., Rajal, V. R. (2019) Data fitting approach more critical
860 than exposure scenarios and treatment of censored data for quantitative microbial risk
861 assessment. *Water Research* 154, 45-53.

862 Prez, V. E., Poma, H. R., Giordano, G. G., Victoria, M., Nates, S. V., Rajal, V. B., Barril, P.
863 A. (2020) Rotavirus contamination of surface waters from the northwest of Argentina.
864 *Journal of Water and Health* 18(3), 409-415.

865 Randazzo, W., Cuevas-Ferrando, E., Sanjuán, R., Domingo-Calap, P., Sánchez, G.
866 (2020b) Metropolitan wastewater analysis for COVID-19 epidemiological surveillance.
867 *International Journal of Hygiene and Environmental Health* 230, 113621.

868 Randazzo, W., Truchado, P., Cuevas-Ferrando, E., Simón, P., Allende, A., Sánchez, G.
869 (2020a) SARS-CoV-2 RNA in wastewater anticipated COVID-19 occurrence in a low
870 prevalence area. *Water Research* 181, 115942.

871 R Development Core Team (2005) R: A Language and Environment for Statistical
872 Computing. Vienna, Austria: R Foundation for Statistical Computing ([http://www.R-](http://www.R-project.org)
873 [project.org](http://www.R-project.org)).

874 Rimoldi, S. G., Stefani, F., Gigantiello, A., Polesello, S., Comandatore, F., Mileto, D.,
875 Maresca, M., Longobardi, C., Mancon, A., Romeri, F., Pagani, C., Cappelli, F.,
876 Roscioli, C., Moja, L., Gismondo, M. R., Salerno, F. (2020) Presence and infectivity of
877 SARS-CoV-2 virus in wastewaters and rivers. *Science of the Total Environment*, 744,
878 140911.

879 RStudio Team (2020). RStudio: Integrated Development for R. RStudio, PBC, Boston, MA
880 URL <http://www.rstudio.com/>.

881 Saththasivam, J., El-Malah, S. S., Gomez, T. A., Jabbar, K. A., Remanan, R.,
882 Krishnankutty, A. K., Ogunbiyi, O., Rasool, K., Ashhab, S., Rashkeeb, S., Bensaad, M.,
883 Ahmed, A. A., Mohamoud, Y. A., Malek, J. A., Abu-Raddad, L. J., Jeremijenko, A.,
884 Abu-Halaweh, H. A., Lawler, J., Mahmoud, K. A. (2021) COVID-19 (SARS-CoV-2)
885 outbreak monitoring using wastewater-based epidemiology in Qatar. Science of the
886 Total Environment 774, 145608.

887 Shapiro, S.S., Francia, R. S. (1972) An Approximate Analysis of Variance Test for
888 Normality. Journal of the American Statistical Association 67(337), 215-216.

889 Sun, J., Zhu, A., Li, H., Zheng, K., Zhuang, Z., Chen, Z., Shi, Y., Zhang, Z., Chen, S. B.,
890 Liu, X., Dai, J., Li, X., Huang, S., Luo, L., Wen, L., Zhuo, J., Li, Y., Wang, Y., Zhang, L.,
891 Zhang, Y., Li, F., Feng, L., Chen, X., Zhong, N., Yang, Z., Huang, J., Zhao, J., Li, Y. M.
892 (2020) Isolation of infectious SARS-CoV-2 from urine of a COVID-19 patient. Emerging
893 Microbes & Infections 9(1), 991-993.

894 USEPA, United States Environmental Protection Agency, 2002. Method 1106.1:
895 Enterococci in Water by Membrane Filtration Using Membrane-Enterococcus-Esculin
896 Iron Agar (mE-EIA). EPA 600-4-85-076.

897 USEPA, United States Environmental Protection Agency, 2014. Method 1603: *Escherichia*
898 *coli* (*E. coli*) in Water by Membrane Filtration Using Modified membrane-
899 Thermotolerant *Escherichia coli* Agar (Modified mTEC). EPA-82 1-R-14-010.

900 USEPA, United States Environmental Protection Agency. 2012. Recreational Water
901 Quality Criteria. Office of Water, United States Environmental Protection Agency
902 (USEPA), Washington, DC. Online: [https://www.epa.gov/sites/production/.../rec-](https://www.epa.gov/sites/production/.../rec-factsheet-2012.pdf)
903 [factsheet-2012.pdf](https://www.epa.gov/sites/production/.../rec-factsheet-2012.pdf).

904 Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A.,
905 Speleman, F. (2002) Accurate normalization of real-time quantitative RT-PCR data by
906 geometric averaging of multiple internal control genes. *Genome Biology* 3(7), 1-12.

907 WHO (2003) Guidelines for environmental surveillance of poliovirus circulation (No.
908 WHO/V&B/03.03). World Health Organization.

909 WHO (2020) World Health Organization. <https://covid19.who.int/> (Last accessed 16th June,
910 2021).

911 Wickham, H. (2009) *ggplot2: Elegant Graphics for Data Analysis* Springer-Verlag New
912 York.

913 Wölfel, R., Corman, V. M., Guggemos, W., Seilmaier, M., Zange, S., Müller, M. A.,
914 Niemeyer, D., Jones, T. C., Vollmar, P., Rothe, C., Hoelscher, M., Bleicker, T., Brünink,
915 S., Schneider, J., Ehmann, R., Zwirgmaier, K., Drosten, C., Wendtner, C. (2020)
916 Virological assessment of hospitalized patients with COVID-2019. *Nature* 581, 465-
917 469.

918 Wu, F., Xiao, A., Zhang, J., Moniz, K., Endo, N., Armas, F., Bushman, M., Chai, P. R.,
919 Duvalle, C., Erickson, T. B., Foppe, K., Ghaeli, N., Gu, X., Hanage, W. P., Huang, K.
920 H., Lee, W. L., Matus, M., McElroy, K. A., Rhode, S. F., Wuertz, S., Thompson, J., Alm,
921 E. J. (2020) Wastewater Surveillance of SARS-CoV-2 across 40 U.S. states. *MedRxiv*
922 preprint doi: <https://doi.org/10.1101/2021.03.10.21253235>

923 Wurtzer, S., Marechal, V., Mouchel, J. M., Moulin, L. (2020) Time Course Quantitative
924 Detection of SARS-CoV-2 in Parisian Wastewaters Correlates with COVID-19
925 Confirmed Cases. *medRxiv*. DOI:10.1101/2020.04.12.20062679.

926 Xiao, F., Sun, J., Xu, Y., Li, F., Huang, X., Li, H., Zhao, Jingxian., Huang, J. Zhao, J.
927 (2020a) Infectious SARS-CoV-2 in feces of patient with severe COVID-19. *Emerging*
928 *Infectious Diseases* 26(8), 1920-1922.

929 Xiao, F., Tang, M., Zheng, X., Liu, Y., Li, X., Shan, H. (2020b) Evidence for gastrointestinal
930 infection of SARS-CoV-2. *Gastroenterology* 158(6), 1831-1833.

931 Zaki, A. M., van Boheemen, S., Bestebroer, T. M., Osterhaus, A. D. M. E., Fouchier, R. A.
932 M. (2012) Isolation of a Novel Coronavirus from a Man with Pneumonia in Saudi
933 Arabia. *New England Journal of Medicine* 367, 1814-1820.

934 Zhang, H., Kang, Z., Gong, H., Xu, D., Wang, J., Li, Zhixiu, Li, Zifu, Cui, X., Xiao, J., Zhan,
935 J., Meng, T., Zhou, W., Liu, J. Xu, H. (2020b) Digestive system is a potential route of
936 COVID-19: an analysis of single-cell coexpression pattern of key proteins in viral entry
937 process. *Gut*, 69(6), 1010-1018.

938 Zhang, J., Wang, S., Xue, Y. (2020c) Fecal specimen diagnosis 2019 novel coronavirus–
939 infected pneumonia. *Journal of Medical Virology* 92(6), 680-682.

940 Zhang, R., Li, Y., Zhang, A. L., Wang, Y., Molina, M. J. (2020a) Identifying airborne
941 transmission as the dominant route for the spread of COVID-19. *Proceedings of the*
942 *National Academy of Sciences* 117(26), 14857-14863.

943 Zhang, Y., Chen, C., Zhu, S., Shu, C., Wang, D., Song, J., Song, Y., Zhen, W., Feng, Z.,
944 Wu, G., Xu, J., Xu, W. (2020d) Isolation of 2019-nCoV from a stool specimen of a
945 laboratory-confirmed case of the coronavirus disease 2019 (COVID-19). *China CDC*
946 *Weekly*, 2(8), 123-124.