Gestational diabetes mellitus in women born small or premature:

Systematic review and meta-analysis

Corresponding author

Yasushi Tsujimoto, MD, MPH

Visiting researcher, Department of Health Promotion and Human Behavior,

Kyoto University Graduate School of Medicine / School of Public Health, Yoshida Konoe-cho,

Sakyo-ku, Kyoto 606-8501, Japan

Email: yssh0108@yahoo.co.jp

Authors and affiliations

Yasushi Tsujimoto, MD, MPH^{1,2}, Yuki Kataoka, MD, DrPH^{2,3}, Masahiro Banno, MD, PhD^{2,4},

Shunsuke Taito PT, PhD^{2,5}, Masayo Kokubo MD⁶, Yuko Masuzawa, CNM, PhD^{7,8}, Yoshiko

Yamamoto, MD, PhD^{8,9}

1. Department of Health Promotion and Human Behavior, Kyoto University Graduate School of Medicine / School of Public Health, Yoshida Konoe cho, Sakyo-ku, Kyoto, Japan.

2. Systematic Review Peer Support Group, Koraibashi, Chuo-ku, Osaka, Japan

3. Department of Internal Medicine, Kyoto Min-Iren Asukai Hosipital, Kyoto, Japan

4. Department of Psychiatry, Seichiryo Hospital, Tsurumai 4-16-27, Showa-ku, Nagoya, Aichi, Japan

5. Division of Rehabilitation, Department of Clinical Practice and Support, Hiroshima University Hospital, Kasumi 1-2-3, Minami-ku, Hiroshima, Japan

- 24 6. Department of Neonatology, Nagano Children's Hospital, Toyoshina 3100, Azumino,
25 Nagano, Japan
- 26 7. Tokyo Healthcare University, Chiba Faculty of Nursing, Division of Nursing, Kaijinchonishi
27 1-1042-2, Funabashi, Chiba, Japan
- 28 8. Cochrane Japan, Akashi Cho 10-1, Chuo-ku, Tokyo, Japan
- 29 9. Department of Health Policy, National Center for Child Health and Development, 2-10-1
30 Okura, Setagaya-ku, Tokyo, Japan

31

32

33 **Word count:** 2540 words

34

35

What is already known on this subject?

Approximately 10–15% of infants are born small or premature worldwide, and these children are at high risk of diseases, such as type 1 and 2 diabetes mellitus, hypertension, obesity, and kidney disease, in adulthood. A narrative review reported in 2007 that women born with LBW are at risk of gestational diabetes mellitus, but included a small number of studies. Several subsequent studies have been published since then, but there is no quantitative summary of the relevant evidence to date.

What this study adds?

This is the first systematic review and meta-analysis of observational studies that provides a comprehensive summary of evidence on the association between birth size or premature birth and future GDM risk including previously unpublished data and a large sample size. LBW, preterm birth, and SGA status may be prognostic factors for GDM.

ABSTRACT

Background: Women born preterm or with low birthweight (LBW) have an increased future risk of gestational diabetes mellitus (GDM) during pregnancy; however, a quantitative summary of evidence is lacking.

In this study, we aimed to investigate whether being born preterm, or with LBW or small for gestational age (SGA) are associated with GDM risk.

Methods: We searched the MEDLINE, Embase, and CINAHL databases and study registries, including ClinicalTrials.gov and ICTRP, from launch until 29 October 2020 for observational studies examining the association between birth weight or gestational age and GDM were eligible. We pooled the odds ratios and 95% confidence intervals using the DerSimonian and Laird random-effects model.

Results: Eighteen studies were included (N = 827,382). The meta-analysis showed that being born preterm, with LBW or SGA was associated with increased risk of GDM (pooled odds ratio = 1.84; 95% confidence interval: 1.54 to 2.20; $I^2 = 78.3\%$; $\tau^2 = 0.07$). Given a GDM prevalence of 2.0%, 10%, and 20%, the absolute risk differences were 1.6%, 7.0%, and 11.5%, respectively. The certainty of evidence was low due to serious concerns of risk of bias and publication bias.

Conclusion: Women born prematurely, with LBW or SGA status, may be at increased risk for GDM. However, whether this should be considered in clinical decision-making depends on the prevalence of GDM.

INTRODUCTION

Gestational diabetes mellitus (GDM) is a common pregnancy complication, with prevalence estimates being 1–36%, depending on the population studied and diagnostic criteria employed¹. GDM is defined as preconceptionally unconfirmed glucose intolerance identified in the second or third trimester of pregnancy². Adverse perinatal outcomes associated with uncontrolled diabetes in pregnancy include spontaneous abortion, foetal anomalies, preeclampsia, stillbirth, macrosomia, neonatal hypoglycaemia, and neonatal hyperbilirubinemia, among others³. Women with a history of GDM are at a higher risk of type 2 diabetes than their counterparts^{4,5}.

Low birth weight (LBW) and preterm birth are the leading causes of neonatal death and childhood-onset morbidity⁶. Approximately 10–15% of infants are born small or premature worldwide^{6,7}. Children who survive are at a higher risk of diseases, such as type 1 and 2 diabetes mellitus, hypertension, obesity, and kidney disease, in adulthood⁸. The exact mechanism underlying these risks remains unclear; the Barker hypothesis proposes that pregnancy may activate biological vulnerability in utero⁹.

A narrative review reported in 2007 that women born with LBW are at risk of GDM¹⁰, but included a small number of studies, and additional research has been published subsequently^{11,12}. There is no quantitative summary of the relevant evidence to date. We performed the first systematic review and meta-analysis of observational studies examining the association between preterm birth, with LBW, or with SGA status and the future risk of GDM.

METHODS

We followed the Meta-analysis of Observational Studies in Epidemiology (MOOSE) guidelines (Supplementary Table S1) in the reporting of this study; the study methodology adhered to the

Cochrane Handbook^{13, 14}. Evidence certainty assessment was based on the Grading of Recommendations, Assessment, Development and Evaluation (GRADE) criteria for prognostic factors¹⁵. The protocol was prospectively registered with PROSPERO (CRD42020142004).

Searches

We searched databases such as MEDLINE, Embase, and CINAHL and study registries including ClinicalTrials.gov and ICTRP from launch until 29 October 2020. Qualified authors (YT and YK) developed the search strategy (Supplementary Table S2). No language or publication status restrictions were imposed. Reference lists of shortlisted studies were searched manually for additional potentially eligible titles.

Study selection

Studies were eligible for inclusion if they were observational cohort or case-control studies. Case reports or series were excluded from the present review. We included studies that involved pregnant women regardless of study setting. The exposures of interest were the infancy parameters of presently pregnant women and were defined as follows: LBW, birth weight <2500 g⁷; small for gestational age (SGA), birth weight <10th percentile for the given gestational age, stratified by sex, using the average weight of gestational age¹⁶; and preterm birth, gestational age of <37 weeks¹⁷. When data on both birth weight and gestational age were reported, we extracted data on birth weight in preference. The comparator group comprised women who were not born small or born at full term.

The outcome of interest was GDM, as defined by the International Association of Diabetes Pregnancy Study Groups (IADPSG), World Health Organization (WHO), American

Diabetes Association or Endocrine Society, or International Classification of Diseases 11th revision (ICD-11) or earlier¹⁸⁻²². If studies used other definitions, they were included in the present review; however, we removed them to assess the robustness of the pooled estimates. For studies that reported LBW, preterm birth, or SGA as a risk factor in pregnant women without reporting the association with GDM, we contacted study authors to acquire estimates of such associations, where available. These additional estimates were included in the present analysis, provided they were measures of an association between at least one of the exposure factors and the outcome of interest.

Two investigators independently screened article titles and abstracts to shortlist relevant studies; subsequently, the same sets of authors assessed the full text for study eligibility. In cases where data were incomplete and precluded study eligibility assessment, we contacted study authors with requests for clarification. Multiple publications were assessed together; the record with the most complete data was included in the present review.

Data extraction and quality assessment

Two investigators independently extracted data from all included studies, using a pilot-tested, uniform data extraction sheet. Any discrepancies between reviewers were resolved through consensus between two reviewers or arbitration by a third reviewer, as required. For studies that compared three or more exposure groups, we contacted study authors to obtain data comparing two groups of interest. In cases where this approach was unsuitable, we extracted the relevant data, as reported, and performed subgroup comparisons between the two groups subdivided by specific thresholds (i.e., birth weight 2500 g, <10th percentile, and gestational age 37 weeks for LBW, SGA, and preterm birth, respectively), as this approach may have resulted in conservative

effect estimates. The same authors who performed data extraction also independently assessed the risk of bias in each study, using the Quality In Prognosis Studies (QUIPS) tool²³. We prospectively identified the following candidate confounders: age, obesity, smoking status, socioeconomic status, diabetes mellitus before the index gestation, and family history of diabetes^{24, 25}.

Data synthesis and analysis

We obtained pooled and adjusted ORs with 95% CI estimates of GDM for the exposure and control groups using the DerSimonian and Laird random-effects method. We calculated the absolute risk difference for GDM between the exposure and control groups in low- (control group: GDM risk was assumed to be 2.0%), moderate- (10%), and high- (20%) prevalence groups, using the pooled odds ratios (OR) and 95% confidence intervals (CIs). This assumption was made based on a previous report and our clinical expertise²⁶.

Publication bias was assessed qualitatively by visual inspection of the funnel plot and quantitatively by Egger's test²⁷. Where asymmetry was observed in the funnel plot, we investigated the likely source of this asymmetry using the contour-enhanced funnel plot. We evaluated between-study heterogeneity visually, using forest plots, and quantitatively, using I^2 and τ^2 statistics. We used the Cochrane chi-square test to calculate I^2 and τ^2 statistics. We performed a pre-specified subgroup analysis based on types of exposure (preterm birth, LBW, or SGA). In pre-specified sensitivity analyses, we used crude ORs instead of adjusted ORs and excluded studies using non-standard definitions of GDM. Some studies assessed the risk of GDM among women born with a weight >4000 g (macrosomia); these studies were excluded

from post-hoc sensitivity analysis, as a previous review has shown a U-shaped association between mother's birth weight and GDM risk¹⁰.

All analyses were performed using STATA 14.2 (StataCorp LP, Texas) and RevMan 5.4 (Cochrane Collaboration, UK). Two-sided *p*-values <0.05 were considered indicative of statistical significance.

RESULTS

Figure 1 presents the flow of studies through the present review selection process. After screening 15,281 records, 59 records representing 44 studies were assessed for eligibility based on the full text. Finally, 18 studies including 827,382 participants were included in the qualitative synthesis; 15 studies including 825,622 participants were included in quantitative synthesis. Supplementary Table S3 lists all excluded studies with reasons for exclusion. We did not find any ongoing or unpublished studies by searching study registries. By contacting authors, we obtained unpublished data from two studies^{12, 28}.

Supplementary Table S4 shows the detail characteristics of the included studies. Nine studies (810,197 participants) used population-based samples, 2 (6,915 participants) were multicentre studies, 6 (9,439 participants) were single-centre studies, and 1 (831 participants) did not specify the study setting. Supplementary Table S5 shows the details of the inclusion and exclusion criteria of the included studies. All studies were conducted in high-income countries, mostly between the late 1990s and early 2010s. The studies included participants of non-Hispanic White, Hispanic, African, Asian, or Indian descent. Two studies (28,722 participants) only included women about to deliver their first child. Two studies (140,714 participants) compared pregnant women born preterm and at full term, 9 (216,439 participants) compared

women born with and without LBW, and 4 (468,469 participants) compared women born with and without SGA status. The remaining 3 studies (1,760 participants) only compared the mean birth weight of women with or without GDM. Figure S1 presents a summary of study quality assessment using the QUIPS tool²³. The overall quality of the included studies was moderate to low, mainly due to uncontrolled confounders.

Prematurity and size at birth and the risk of gestational diabetes mellitus

The median GDM rate in the control groups of the included cohort studies was 2.9% (range: 0.5% to 22%). Figure 2 presents a forest plot summarising the studies that assessed the association between preterm birth or size at birth with GDM. Premature birth, LBW, and SGA status were associated with a higher GDM risk (pooled OR, 1.84; 95% CI: 1.54 to 2.20; $I^2 = 78.3\%$; $\tau^2 = 0.07$). Supplementary Table S5 summarises the absolute risk difference in pregnant women born with LBW, SGA status, or born preterm in the low- (2.0% risk of GDM in the control group), medium- (10%), and high- (20%) GDM prevalence groups. The absolute risk increases were 1.6% (95% CI: 1.0 to 2.1%), 7.0% (95% CI: 4.6 to 9.6%), and 11.5% (95% CI: 7.8 to 15.5%) in low-, moderate-, and high-prevalence settings, respectively. The certainty of evidence was low due to serious concerns of risk of bias and publication bias.

Figure 3 presents study estimates in a funnel plot. The plot appeared asymmetrical, and Egger's test for funnel plot asymmetry was statistically significant (p -value = 0.030). Supplementary Figure S2 shows the contour-enhanced funnel plot, which suggests the existence of some missing studies on the left-hand side of the plot; these studies would have yielded statistically non-significant findings.

Data on the birth weight of mothers with or without GDM, obtained from three studies excluded from the meta-analysis, are presented in Supplementary Table S5. These studies

consistently reported that mothers with GDM were born with lower birth weights than those without GDM.

Subgroup and sensitivity analyses

There was substantial between-study heterogeneity ($I^2 = 78.3\%$). Figure 2 presents the results of subgroup analyses for the types of exposure (LBW, preterm birth, or SGA). Although all types of exposure were associated with GDM, there was significant heterogeneity due to the type of exposure (p for interaction = 0.004). The results of additional sensitivity analyses are presented in Supplementary Figure S3, Figure S4, and Figure S5.

DISCUSSION

Main Findings

We found that women born small or premature may have future risk of GDM. However, the evidence certainty was low, and the presented findings may be overestimated, as we observed some evidence of publication bias. These findings were approximately consistent across the subgroups, including different populations, exposures, and studies of varied methodological quality; these findings were robust in sensitivity analyses.

Our finding that the mother's size at birth or premature birth may affect GDM risk was consistent with that of a previous narrative review¹⁰. The strength of this association was similar to that observed in women with a family history of diabetes mellitus, an established risk factor for GDM²⁹. However, the importance of the risk factor in clinical decision depends on the absolute risk difference. Our findings suggested that careful review of the mother's birth status may indicate her risk of GDM and guide pregnancy management in moderate to high prevalence

settings. The mother's preterm birth status and size at birth are not currently considered risk factors for GDM in any of the major guidelines or risk models^{30,31}. Our findings may help further refine these guidelines and models or to develop new ones.

The certainty of evidence for the association between premature birth or SGA status and GDM was low due to the high risk of publication bias, as shown by funnel-plot analysis. The contour-enhanced funnel plot suggests that studies with non-significant findings may not have been published. Although we did not identify any ongoing or unpublished studies, this did not eliminate the risk of publication bias, as observational studies are less likely to be registered than clinical trials³². Thus, the reported estimates may be overestimates. The studies included in this review tended not to adjust for confounders, such as smoking, obesity, socioeconomic status, and family history of diabetes. Future studies should adjust for these factors.

The main result of this review was subject to substantial between-study heterogeneity, as shown by the I^2 statistic¹³. This heterogeneity may be due to the different types of exposure (LBW, SGA, or preterm birth) considered in this study. However, as all exposure types were associated with increased GDM risk, the high I^2 statistic may be due to the large number of participants and narrow CIs of the primary studies³³. Given these findings, we did not assign a low rating to the inconsistency domain of the GRADE criteria¹⁵.

The underlying mechanism of the association between preterm birth or SGA status and subsequent GDM may be gestational malnutrition due to maternal malnutrition or placental insufficiency³⁴. Findings from animal studies have suggested that malnutrition in utero is associated with reduced β -cell counts, pancreas weight, and pancreatic insulin content³⁵. According to the Barker foetal origin hypothesis, these foetal programming events may affect the future risk of disease⁹. A review of epidemiological studies has suggested that LBW and preterm

birth are associated with the risk of type 2 diabetes in adulthood; a similar mechanism is possible for GDM³⁶.

Strengths and Limitations

A key strength of this review is that it is the first to provide a comprehensive summary of evidence on the association between birth size or premature birth and future GDM risk. This study followed the methodological recommendations presented in the Cochrane Handbook, MOOSE guidelines, and GRADE criteria¹³⁻¹⁵. Moreover, this study included previously unpublished data and a large sample size.

Nevertheless, this study has some limitations. First, the included studies were old and may not represent the current clinical practice. The definition of GDM proposed by the IADPSG in 2010 has resulted in an increase in GDM prevalence³⁷. For example, the prevalence of GDM in the United States increased from 4.6% in 2006 to 8.2% in 2016³⁸. The median prevalence of GDM in the control groups of the included studies was 2.9%. However, empirical evidence suggests that relative effect measures are, on average, consistent across different settings; in the present study, we estimated absolute risk differences separately for low-, moderate-, and high-prevalence settings³⁹. Second, 5 of 15 studies divided birth size and preterm birth categories into three or more comparative groups, which could not be combined into two comparison groups of interest. This lack of data required methodological adjustments, as described previously. Lastly, this review only assessed certainty in estimates of association between prognostic factors and an outcome. Future studies are required to determine whether these factors can help risk-stratify pregnant women and improve the clinical management of GDM.

Conclusions

LBW, preterm birth, and SGA status may be prognostic factors for GDM. Clinicians should consider the prevalence of GDM in their setting when considering maternal preterm birth or size at birth in clinical decision-making. Due to the high likelihood of publication bias, the true association between the exposures and outcome of interest may be weaker than that reported herein. Future studies based on up-to-date diagnostic criteria, examining the dose–response relationship between exposure severity and outcome, and comparing low- and middle-income countries, are required to improve the certainty of evidence.

ACKNOWLEDGMENTS

We wish to thank Prof. Anne Monique Nuyt, Dr. Ariane Boivin, Dr. Prabha H. Andraweera, and Dr. Shalem Leemaqz for kindly providing additional study data that were not included in the original reports. We are also grateful to Dr. Kyosuke Kamijo and Dr. Ayako Shibata for contributing a clinical perspective. We would like to thank Editage (www.editage.com) for English language editing.

COMPETING INTERESTS

The authors have no conflicts of interest to declare.

FUNDING

The English editing fee was supported by the Systematic Review Peer Support Group (not-for-profit organisation). The funders played no role in the design or conduct of the study; collection,

management, analysis, or interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

CONTRIBUTION TO AUTHORSHIP

YT is the guarantor of the review. YT had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: YT, YK, MB, ST, MK, YY

Acquisition, analysis, or interpretation of data: All authors

Drafting of the manuscript: YT, YK, MB, ST

Critical revision of the manuscript for important intellectual content: MK, YM, YY

Statistical analysis: YT

Administrative, technical, or material support: YT, YK, MB, ST

Study supervision: MK, YM, YY

REFERENCES

1. Buchanan TA, Xiang AH, Page KA. Gestational diabetes mellitus: risks and management during and after pregnancy. *Nat Rev Endocrinol*. 2012;8(11):639-49.
2. American Diabetes Association. 2. Classification and diagnosis of diabetes: standards of medical care in diabetes—2019. *Diabetes care*. 2019;42(Supplement 1):S13-S28.
3. American Diabetes Association. 14. Management of diabetes in pregnancy: standards of medical care in diabetes—2019. *Diabetes Care*. 2019;42(Supplement 1):S165-S72.
4. Kim C, Newton KM, Knopp RH. Gestational diabetes and the incidence of type 2 diabetes: a systematic review. *Diabetes Care*. 2002;25(10):1862-8.
5. Bellamy L, Casas J-P, Hingorani AD, Williams D. Type 2 diabetes mellitus after gestational diabetes: a systematic review and meta-analysis. *The Lancet*. 2009;373(9677):1773-9.
6. Chawanpaiboon S, Vogel JP, Moller A-B, Lumbiganon P, Petzold M, Hogan D, et al. Global, regional, and national estimates of levels of preterm birth in 2014: a systematic review and modelling analysis. *The Lancet Global Health*. 2019;7(1):e37-e46.
7. United Nations Children's Fund (UNICEF) WHO. UNICEF-WHO Low birthweight estimates: Levels and trends 2000–2015. Geneva: World Health Organization; 2019.
8. Belbasis L, Savvidou MD, Kanu C, Evangelou E, Tzoulaki I. Birth weight in relation to health and disease in later life: an umbrella review of systematic reviews and meta-analyses. *BMC Med*. 2016;14(1):147.
9. Barker D. J. The fetal and infant origins of adult disease. *Bmj*. 1990;301(6761):1111.
10. Pettitt DJ, Jovanovic L. Low birth weight as a risk factor for gestational diabetes, diabetes, and impaired glucose tolerance during pregnancy. *Diabetes Care*. 2007;30(Supplement 2):S147-S9.

- 336 11. Crusell M, Damm P, Hansen T, Pedersen O, Glumer C, Vaag A, et al. Ponderal index at
337 birth associates with later risk of gestational diabetes mellitus. Archives of gynecology and
338 obstetrics. 2017;296(2):249-56.
- 339 12. Andraweera PH, Dekker G, Leemaqz S, McCowan L, Myers J, Kenny L, et al. Effect of
340 Birth Weight and Early Pregnancy BMI on Risk for Pregnancy Complications. Obesity.
341 2019;27(2):237-44.
- 342 13. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al. Cochrane
343 Handbook for Systematic Reviews of Interventions version 6.0
344 : Cochrane; 2019 [updated July 2019. Available from: www.training.cochrane.org/handbook.
345 .
- 346 14. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, et al. Meta-
347 analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of
348 Observational Studies in Epidemiology (MOOSE) group. Jama. 2000;283(15):2008-12.
- 349 15. Foroutan F, Guyatt G, Zuk V, Vandvik PO, Alba AC, Mustafa R, et al. GRADE
350 Guidelines 28: Use of GRADE for the assessment of evidence about prognostic factors: rating
351 certainty in identification of groups of patients with different absolute risks. J Clin Epidemiol.
352 2020;121:62-70.
- 353 16. Sinha SK. Essential neonatal medicine / Sunil Sinha, Lawrence Miall, Luke Jardine.
354 Jardine L, Levene MIEnm, Miall L, editors. Chichester, West Sussex ; Hoboken, NJ: Wiley-
355 Blackwell; 2012.
- 356 17. ACOG Committee Opinion No 579: Definition of term pregnancy. Obstet Gynecol.
357 2013;122(5):1139-40.

18. World Health Organization. ICD-11 : international statistical classification of diseases and related health problems : eleventh revision Geneva: World Health Organization; 2019 [updated 2019. Available from: <https://icd.who.int/browse11/l-m/en>.
19. American Diabetes Association. 2. Classification and Diagnosis of Diabetes. Diabetes Care. 2016;39 Suppl 1:S13-22.
20. World Health Organization. Diagnostic criteria and classification of hyperglycaemia first detected in pregnancy. World Health Organization; 2013.
21. Blumer I, Hadar E, Hadden DR, Jovanović L, Mestman JH, Murad MH, et al. Diabetes and pregnancy: an endocrine society clinical practice guideline. J Clin Endocrinol Metab. 2013;98(11):4227-49.
22. International Association of D, Pregnancy Study Groups Consensus P, Metzger BE, Gabbe SG, Persson B, Buchanan TA, et al. International association of diabetes and pregnancy study groups recommendations on the diagnosis and classification of hyperglycemia in pregnancy. Diabetes Care. 2010;33(3):676-82.
23. Hayden JA, van der Windt DA, Cartwright JL, Côté P, Bombardier C. Assessing bias in studies of prognostic factors. Ann Intern Med. 2013;158(4):280-6.
24. Beauregard JL, Drews-Botsch C, Sales JM, Flanders WD, Kramer MR. Does Socioeconomic Status Modify the Association Between Preterm Birth and Children's Early Cognitive Ability and Kindergarten Academic Achievement in the United States? Am J Epidemiol. 2018;187(8):1704-13.
25. Gestational Hypertension and Preeclampsia: ACOG Practice Bulletin, Number 222. Obstetrics & Gynecology. 2020;135(6):e237-e60.

26. Zhu Y, Zhang C. Prevalence of Gestational Diabetes and Risk of Progression to Type 2 Diabetes: a Global Perspective. *Current diabetes reports*. 2016;16(1):7-.
27. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *Bmj*. 1997;315(7109):629-34.
28. Boivin A, Luo Z-C, Audibert F, Mâsse B, Lefebvre F, Tessier R, et al. Pregnancy complications among women born preterm. *CMAJ*. 2012;184(16):1777-84.
29. Solomon CG, Willett WC, Carey VJ, Rich-Edwards J, Hunter DJ, Colditz GA, et al. A prospective study of pregravid determinants of gestational diabetes mellitus. *Jama*. 1997;278(13):1078-83.
30. National Collaborating Centre for Ws, Children's H. National Institute for Health and Care Excellence: Clinical Guidelines. *Diabetes in Pregnancy: Management of Diabetes and Its Complications from Preconception to the Postnatal Period*. London: National Institute for Health and Care Excellence (UK) Copyright © 2015 National Collaborating Centre for Women's and Children's Health.; 2015.
31. Artzi NS, Shilo S, Hadar E, Rossman H, Barbash-Hazan S, Ben-Haroush A, et al. Prediction of gestational diabetes based on nationwide electronic health records. *Nat Med*. 2020;26(1):71-6.
32. Boccia S, Rothman KJ, Panic N, Flacco ME, Rosso A, Pastorino R, et al. Registration practices for observational studies on ClinicalTrials.gov indicated low adherence. *J Clin Epidemiol*. 2016;70:176-82.
33. Rucker G, Schwarzer G, Carpenter JR, Schumacher M. Undue reliance on I^2 in assessing heterogeneity may mislead. *BMC Med Res Methodol*. 2008;8:79.

34. Plows JF, Stanley JL, Baker PN, Reynolds CM, Vickers MH. The Pathophysiology of Gestational Diabetes Mellitus. *Int J Mol Sci.* 2018;19(11).
35. McMillen IC, Robinson JS. Developmental origins of the metabolic syndrome: prediction, plasticity, and programming. *Physiol Rev.* 2005;85(2):571-633.
36. Whincup PH, Kaye SJ, Owen CG, Huxley R, Cook DG, Anazawa S, et al. Birth weight and risk of type 2 diabetes: a systematic review. *Jama.* 2008;300(24):2886-97.
37. Cundy T, Ackermann E, Ryan EA. Gestational diabetes: new criteria may triple the prevalence but effect on outcomes is unclear. *Bmj.* 2014;348:g1567.
38. Zhou TAO, Sun D, Li X, Heianza Y, Nisa H, Hu G, et al. Prevalence and Trends in Gestational Diabetes Mellitus among Women in the United States, 2006–2016. *Diabetes.* 2018;67(Supplement 1):121-OR.
39. Deeks JJ. Issues in the selection of a summary statistic for meta-analysis of clinical trials with binary outcomes. *Stat Med.* 2002;21(11):1575-600.

FIGURE LEGENDS

Figure 1. PRISMA flow diagram of study eligibility

Duplicate studies are displayed as a single study.

Figure 2. Risk of gestational diabetes among women born preterm, with low birth weight, or small-for-gestational-age status

Effect size (ES, represented as adjusted odds ratios); CI, confidence interval. ES was determined using the random-effects model weighted by the inverse of the variance estimate. Squares represent ES, with marker size reflecting the statistical weight of the study, obtained using

random-effects meta-analysis; horizontal lines represent 95% CIs; diamonds represent the subgroup and overall odds ratios and 95% CIs for gestational diabetes.

Figure 3. Funnel plot for the evaluation of publication bias

The solid vertical line represents the summary estimate of the association between preterm birth, low birth weight, and small for gestational age status and gestational diabetes (using random-effects meta-analysis). A significant publication bias was detected ($p = 0.030$ for Egger's test). The funnel plot shows asymmetry, which indicates publication bias.

Funnel plot with pseudo 95% confidence limits