

Table 1a. Characteristics of included primary studies

Study ID	Setting	Country	Population/environment	Method
Alamar I 2020	Hospital	USA	1 mother with fever, mild chills, fatigue, dysgeusia, and anosmia beginning 1 day before presentation, with positive COVID-19 test.	Case study
Anand P 2020	Hospital	India	69 COVID19 + mothers, 65 infants (1 abortion, 4 stillbirths). 51 mothers had symptoms and of these 45 had mild symptoms.	Cohort study
Ayed A 2020	Hospital	Kuwait	185 pregnant women with PCR confirmed SARS-CoV-2 infection (median age 31; median gestation at diagnosis 29 weeks). 88% had mild symptoms, (fever & cough the most common presenting symptoms). During the study period 40 (21.6%) gave live birth, 3 (1.6%) had a miscarriage, 1 (0.54%) had intrauterine fetal death, unrelated to COVID19. 2 (1.1%) patients developed severe pneumonia & required intensive care. Most neonates asymptomatic; 2 (5%) of them tested positive on day 5.	Retrospective study, medical records
Bachani S 2020	Hospital	India	348 pregnant women tested for SARS-CoV-2, 57 women (16.3%) confirmed positive. 3 maternal deaths were associated with comorbidities. 5 neonates tested positive for SARS-CoV-2	Retrospective study
Bandyopadhyay T 2020	Hospital	India	1 mother, mild grade fever, close contact of positive case (husband)	Case study

Barbero P 2020	Hospital	Spain	91 women with COVID-19 symptoms (fever (37.8 C), dry cough, shortness of breath or dyspnea, chills & myalgia, headache, coryza & new onset of loss of taste or smell) & diagnosed with SARS-CoV-2 infection during pregnancy/postpartum (<40 days after giving birth). 46.2% rate of hospitalization.	Retrospective cohort study
Birindwa EK 2020	Hospital	DR Congo	1 woman positive for SARS-Cov-2.	Case report
Bordbar A 2020	Hospital	Iran	1 women, gestational diabetes, no COVID-19 symptoms; tested 2 days after giving birth.	Case report
Chaudhary S 2020	Hospital	Pakistan	26 women, RT-PCR COVID-19 positive; 10 asymptomatic, 16 symptomatic; all mild disease, 1 pneumonia.	Case series
Cojocar L 2020	Hospital	US	1989 women were screened for SARS-CoV-2, from which 86 (0.04%) tested positive. 34 delivered during study period (3 excluded from analysis). 5 women in analysis admitted to ICU.	Quality improvement project, prospective
Demirjian A 2020	Hospital	UK	1 woman, fever & respiratory symptoms, tested for COVID on day 6 of admission.	Case report
Farsi Z 2020	Hospital	Iran	1 woman, pregnant with triplets, symptoms of cough, fever, myalgia	Case report
Fenizia B 2020	Hospital	Italy	31 laboratory-confirmed COVID19+ women, also radiological assessment, signs & symptoms; 4 severe cases requiring ICU admission	Cohort study, multicentre
Ferraiolo A 2020	Hospital	Italy	1 woman, no symptoms, breech presentation	Case report
Gale C 2020	Hospital; National Registry	UK	Infants with a diagnosis of SARS-CoV-2 within 28 days	Prospective cohort study
Gao W 2020	Hospital	China	1 asymptomatic woman	Case report

Gao X 2020	Hospital	China	14 pregnant women with lab-confirmed COVID-19	Retrospective observational study
Grimminck K 2020	Hospital	The Netherlands	1 pregnant woman, oropharyngeal swab RT-PCR positive for COVID-19, immunosuppressed (lupus)	Case report
He Z 2020	Hospital	China	22 newborns born to pregnant women with COVID-19	Retrospective cohort study
Hinojosa-Velasco A 2020	Hospital	Mexico	1 symptomatic woman (fever, cough, sore throat), patchy chest radiograph, testing positive for COVID-19	Case report
Hsu AL 2020	Hospital	USA	1 pregnant woman, mild symptoms	Case report
Hu X 2020	Hospital	China	6 preterm infants born to COVID-19 positive mothers	Retrospective cohort study
Khan MA 2020	Hospital	Pakistan	66 pregnant women testing COVID-19 positive, most asymptomatic	Cohort study
Kulkarni R 2020	Hospital	India	Pregnant woman, admitted with fever & body ache, in active labor	Case report
Liu W 2020	Hospital	China	Neonates born to women with: 1. confirmed COVID-19 (symptoms & positive RT-PCR), 2. suspected COVID-19 (symptoms with negative RT-PCR but chest CT pneumonia), 3. control cases (with or without symptoms, confirmed as influenza)	Retrospective analysis of cases
Luo Q 2020	Hospital	China	23 pregnant women, 14 with confirmed COVID-19, 9 with suspected COVID-19 (chest-CT pneumonia but negative PCR and serology).	Cohort study of breast feeding
Lv Y 2020	Hospital	China	1 pregnant woman, fever & cough, SARS-CoV-2 PCR positive	Case report

Maraschini A 2020	Hospital	Italy	146 pregnant women with confirmed COVID-19, 142 confirmed by RT-PCR, 4 confirmed by chest X-ray; 99 no COVID pneumonia, 47 with COVID pneumonia	Cohort study
Marin Gabriel MA 2020	Hospital	Spain	Pregnant women, third trimester, RT-PCR or serology COVID-19 positive: 222 RT-PCR positive; 19 RT-PCR negative but serology positive at delivery.	Multi-centre study, retrospective chart review plus 1 month follow-up
Masmejan S 2020	Hospital	Switzerland	13 pregnant women, positive RT-PCR or positive serology (IgG) for SARS-CoV-2. 11 mild symptoms or asymptomatic, 2 critical fever, severe symptoms	Retrospective case series
Mohakud NK 2020	Hospital	India	Symptomatic pregnant woman, taking treatment for hypothyroidism, at 32+ weeks tested positive for SARS-CoV-2.	Case report
Moreno SC 2020	Hospital maternity unit	USA	Symptomatic pregnant women diagnosed positive for COVID-19 via PCR in the third trimester, & all neonates with complete COVID-19 testing & delivery data.	Retrospective observational study
Nayak AH 2020	Tertiary care hospital maternity unit	India	Pregnant women attending for obstetric care.	Cohort study
Ogamba I 2020	Four acute care hospitals	USA	Pregnant women 18 years or older with a diagnosis of COVID-19 (testing was performed on admission to the labor & delivery unit, outpatient setting, and/or inpatient hospitalization)	Retrospective cohort study
Olivini N 2020	COVID-19 ward of children's hospital	Italy	5 women who tested positive after screening due to close contact with an infected maternity services HCW, & their neonates.	Retrospective case series
Oncel MY 2020	Hospital maternity units, multicentre	Turkey	125 pregnant women & their neonates, tested if symptomatic or had had close contact with family members with COVID-19.	Cohort study

Palalioglu RM 2020	Hospital maternity unit	Turkey	Woman aged 42 years, 37 weeks pregnant, with diet-regulated gestational diabetes mellitus, tested SARS-CoV-2 positive, asymptomatic. Admitted two weeks later & C-section performed due to prelabour rupture of membrane.	Case report
Parsa Y 2020	Hospital maternity unit	Iran	41-yr-old pregnant woman with signs & symptoms of acute respiratory illness presented with labor pain & vaginal leak at 37 weeks gestation, tested positive for COVID-19 using RT-PCR, emergency C-section.	Case report
Pereira A 2020	Hospital maternity unit	Spain	22 pregnant women with SARS-CoV-2 (11 symptomatic), median age 34 years, median gestational age 38+5 weeks.	Case series
Pessoa FS 2020	Hospital maternity unit	Brazil	34 year old woman 33+ weeks pregnant, RT-PCR positive, admitted with flu-like symptoms including fever, dry cough & fatigue from talking; chest computed tomography showed attenuations with ground glass opacification and bilateral consolidations.	Case report
Pissarra S 2020	Hospital maternity unit	Portugal	10 SARS-CoV-2 positive pregnant women (7 were symptomatic) giving birth at the hospital.	Case series
Popofsky S 2020	Telephone survey	USA	Women with positive SARS-CoV-2 PCR & their neonates, born at one of 3 hospitals in New York State.	Observational longitudinal cohort
Pulinx B 2020	Hospital obstetric department.	Belgium	30 year old woman 22 weeks pregnant with dichorionic diamniotic twins, recently diagnosed with gestational diabetes mellitus, tested positive for SARS-CoV-2 by RT-PCR.	Case report
Rebello CM 2020	Hospital maternity unit	China	32-year-old woman pregnant at 34 weeks, with gestational diabetes; symptoms of anosmia, ageusia, runny nose, dry cough, headache, myalgia, no fever, diagnosed with SARS-CoV-2.	Case report

Rivera-Hernandez P 2020	Hospital	USA	38-year-old pregnant woman (32/33 weeks) with obesity, poorly controlled type 2 diabetes, asthma & 3-day history of dyspnea & malaise, admitted to ICU.	Case report
Rubio Lorente AM 2020	Hospital	Spain	2 pregnant women, 1: 20w pregnant, fever and cough, positive SARS-CoV-2 PCR test, negative serology. 2: 12w pregnant, delayed abortion, low fever, serology positive, PCR-negative for SARS-CoV-2	Analysis of amniotic fluid of 2 cases
Sajjan GR 2020	Hospital maternity unit	India	21yr old primigravida at 32 weeks gestational age, diagnosed with PPROM	Case report
Schwartz DA 2020	10 hospitals in different cities in Iran	Iran	19 neonates testing positive for SARS-CoV-2	Cohort study
Shende P 2020	Hospital maternity unit	India	1 woman, asymptomatic, 8 weeks pregnant, tested positive for SARS-CoV-2 by PCR. Fetal demise at 13 weeks followed by dilation & curettage termination procedure.	Case report
Singh MV 2020	Hospital maternity unit	India	1 asymptomatic pregnant woman with positive SARS-CoV-2 test, & her neonate	Case report
Sisman J 2020	Hospital maternity unit	USA	1 pregnant woman with obesity, diabetes mellitus & late latent syphilis, & her neonate.	Case report
Smithgall MC 2020	Hospital	USA	3rd trimester placentas from 51 SARS-CoV-2-positive and 25 SARS-CoV-2-negative women.	Cohort study (placentas)
Stonoga E 2020	Hospital	Brazil	Woman at 27 weeks' gestation with COVID-19 symptoms: dyspnea, dry cough, high temperature (38.5°C), anosmia, nausea, vomiting, & diarrhea had developed 2 days before hospitalization.	Case report

Tang F 2020	Hospital	China	Neonates born during the pandemic period 2 February to 31 March 2020 in Wuhan, whose mother was confirmed with COVID-19 in the 3rd trimester, or with +ve SARS-CoV-2-IgM and SARS-CoV-2-IgG and/or +ve RT-PCR for SARS-CoV-2 within 14 days after birth. All included neonates had excluded diagnoses of adenovirus, enterovirus, influenza A, influenza B, parainfluenza, chlamydia pneumonia and mycoplasma pneumonia.	Observational cohort
Tang J 2020	Hospital	China	Two mothers with symptomatic COVID-19 in the second trimester and their neonates.	Case report (2 cases)
Vendola N 2020	Hospital maternity unit	Italy	2 pregnant women with positive IGG on peripheral blood test.	Case report (2 cases)
Vinuela MC 2020	Hospital maternity unit	Spain	Systematic screening in asymptomatic women admitted for spontaneous delivery: first 100 consecutive participants; 9 tested positive for SARS-CoV-2 by PCR; 13 (including 7 of the PCR-positive women) had antibodies to SARS-CoV-2.	Observational study, asymptomatic screening
Vivanti AJ 2020	Hospital	France	Pregnant woman admitted at 35+ weeks with COVID-19-like symptoms and subsequently tested & treated accordingly.	Case report
Von Kohorn I 2020	Hospital	USA	Pregnant woman 34 weeks with vaginal bleeding & cramping, found to have thrombocytopenia, transaminitis, & hyperuricemia; infant delivered by C-section at 34 weeks (due to hemolysis, elevated liver enzymes, low platelet syndrome, suspicion for systemic COVID-19, & history of prior cesarean section)	Case report
Vouga M 2020	International web registry	International	Women with symptoms for COVID-19 included in the study. Study conducted from 24 March to 26 July 2020.	Cohort study
Woodworth KR 2020	Public health records	USA	4,442 women with known pregnancy outcomes with laboratory-confirmed SARS-CoV-2 infection reported during 29 March to 14 October 2020. 4,495 live births.	Longitudinal surveillance

Yang R 2020	Maternal and Child Health Information System, Wuhan.	China	All pregnant women with singleton live birth recorded by the system between 13 January and 18 March 2020.	Cohort study.
Zaigham M 2020	Hospital	Sweden	Case report of a pregnant woman admitted with COVID-19 symptoms & then diagnosis; subsequent investigations in the neonate.	Case report
Zhang L 2020	Hospital	China	18 patients with COVID-19 during late pregnancy, none critically ill. Epidemiological characteristics, clinical manifestations, laboratory tests, chest CT & pregnancy outcomes reported.	Case series (18 cases).
Zhang P 2020	Case series of pathology of placental samples	USA	364 consecutive women attending to give birth including 74 positive & 290 negative for SARS-CoV-2 by NP swab PCR. Placental pathology & clinical characteristics reported.	Case series (placenta analysis).
Zheng T 2020	Hospital	China	Two pregnant women, admitted due to symptoms suggestive of COVID-19. (The only reported COVID-19 cases among pregnant women in this city during 20 January 2020, to 9 April 2020). Laboratory, imaging & SARS-CoV-2 nucleic acid tests were performed on the 2 women & their neonates	Case report (2 cases).

Abbreviations: NP: nasopharyngeal

Table 1b. Results of included primary studies.

Study ID	Patient numbers	Symptoms in mothers, including timing	Mode of delivery of babies	Sample sources: mother	Sample sources: neonate/ infant	Potential vertical tr, samples PCR-positive n/d for SARS-CoV-2 RNA unless otherwise stated	Live culture positivity	Other results
Alamar I 2020	1 mother & neonate dyad	Mother at 35+6 gestational age presented with vaginal bleeding & contractions; subjective fever, mild chills, fatigue, dysgeusia, anosmia beginning 1 day before presentation.	Urgent C-section due to bleeding	NP swab	NP swab; placental tissue biopsy	1/1	not attempted	Neonate positive at 24 hrs, 48 hrs & DOL 7. Mother & neonate remained asymptomatic during the 14 day follow-up period. Placenta ISH for SARS-CoV-2 RNA revealed strong signal in the villous syncytiotrophoblast; but no signal in villous stromal cells, Hofbauer cells, or villous endothelium

Anand P 2020	69 mothers (39 tested positive)	Among 7 positive testing women: 1. critically ill & died 3 days postpartum, symptoms & timing NR; 2. fever 13 days pre-delivery, negative test day prior to delivery, time to diagnosis NR; 3. symptoms days 4 postpartum, time to diagnosis NR; 4. fever before delivery led to testing positive, time to diagnosis NR; 5. tested day 4 postpartum, symptoms NR; 6. symptoms day 12 postpartum; 7. day 10 symptoms so tested, time to diagnosis NR.	Elective C-section: 9/69; emergency C-section: 17/69. Remainder vaginal.	NP swab	NP/OP swab within 24h	7/65; no deaths	not attempted	2 maternal deaths attributed to COVID-19. All COVID-19 positive neonates remained asymptomatic to final follow-up at day 29.
Ayed A 2020	185 positive testing mothers	88% of the patients had mild symptoms. Fever 58%, cough 51%. Time to diagnosis NR; tested on admission to maternity services.	17/41: emergency C-section; remainder vaginal	NP swab	NP swab	0/41 tested positive within 24h; 2/41 neonates tested positive on day 5; no deaths	not attempted	Cases assessed cases between 15 March and 31 May 2020; follow-up to 15 June 2020. At follow-up 98.8% of women had been discharged, median hospital stay 15 days.

Bachani S 2020	348 pregnant women and 56 neonates	Most women that tested positive experienced mild infection (45/57; 78.9%), with one or two spikes of low-grade fever, cough, and/or diarrhoea that resolved in 2–3 days. Three (5.2%) women had moderate symptoms (fever and breathlessness). Timing of symptoms in relation to testing and/or delivery NR.	Spontaneous: 42; Induced labour: 15; Elective C-section: 9; Emergency C-section: 17	NP swab	NP swab		not attempted	57/348 pregnant women positive; 3 deaths (comorbidities, 2 classed as COVID-19-related complications; 1 classed as COVID-19-related). 5/56 neonates positive; 2 tested positive within 24h; 2 tested positive on day 4; 1 tested positive on day 14 no deaths
Bandyopadhyay T 2020	1 mother & neonate dyad	Mild grade fever, pharyngeal swab taken that day and found positive for SARS-CoV-2 by RT-PCR; admitted to hospital for isolation then became asymptomatic one day later. Delivery 1 month after symptoms. Negative by PCR before delivery.	Vaginal	NP swab	NP swab	1/1	not attempted	Qualitative IgG serum antibody test positive; qualitative IgM serum antibody test negative On day 2 & day 3 of life, neonate tested negative. Mother asymptomatic after admission; tested negative 2 days before birth.
Barbero P 2020	91 women	Symptoms preceded & led to testing. Timing in relation to delivery NR. 40/91 developed pneumonia, and 4/91 required ICU admission	C-section: 11/23; remainder vaginal	NP swab or chest X-ray indicated	NP swab	0/23 babies of women with active infection	not attempted	No suggestive symptoms in neonates on clinical follow-up; 1 neonate tested positive on day 8.

Birindwa EK 2020	1 mother & neonate dyad	3 weeks before admission, woman complained of fever, not responding to acetaminophen; fever persisted; swab test done 2 weeks later confirming SARS-CoV-2 positive by RT-PCR. She was admitted. 4 days later a rapid antigen test was negative for SARS-CoV-2. Preterm labour commenced and C-section was performed.	C-section	NP swab	NP swab	1/1, neonate tested positive at birth, on day 3; died on day 5 due to sepsis	not attempted	Inflamed placenta
Bordbar A 2020	1 mother & neonate dyad	No fever, cough, dyspnea or GI symptoms	C-section	NP swab	NP swab; day 2, chest X-ray	1/1; RT-PCR test on day 2, no symptoms; no symptoms at follow-up on day 14	not attempted	
Chaudhary S 2020	26 pregnant women; 14 deliveries during study period	16/26 positive tested women were symptomatic: 14/16 fever, 11/16 cough; 1 severe pneumonia leading to death. Timing of symptoms NR.	3/14 vaginal, 11/14 C-section	NP swab	NP swab	0/14	not attempted	

Cojocar L 2020	31 neonates	5/31 admitted to ICU. Other symptoms, and timing of symptoms, NR.	8/31 C-section; 23/31 vaginal	Unclear, probably NP swab.	OP and NP swab at 24h and 48h	0/31	not attempted	Study on transmission and maternal bonding defined as rooming-in, skin to skin contact & breastfeeding. None of the neonates who bonded with their mothers tested positive.
Demirjian A 2020	1 mother & neonate dyad	Fever and respiratory symptoms on admission; condition deteriorated to severe, leading to C-section 7 days after admission.	C-section	Respiratory samples (day 6) and sputum (day 8)	NP swab, at birth and day 3, stool and blood samples day 3	1/1; RT-PCR negative at birth, RT-PCR respiratory sample positive on day 3, stool and blood samples negative	not attempted	Neonate no symptoms up to day 5, then developed fever (38.0°C), coryza, and mild tachypnea lasting 2 days. No respiratory support required. No symptoms at discharge on day 18. Mother on ventilation for 17 days. Discharged at 31 days. Genomic viral sequencing of SARS-CoV-2 virus isolated from samples from the mother's & the neonate's respiratory tract secretions. Genetic sequences were identical across the entire genome apart from a single-nucleotide difference between the baby and mother's genomes.
Farsi Z 2020	1 mother & 3 neonates	Woman had hypothyroidism, gestational hypertension (HTN), and gestational diabetes mellitus; also recent symptoms of cough, fever and myalgia 3 days before delivery.	Emergency C-section	PCR, test type and sample not specified	PCR, test type and sample not specified	1/3; baby1: negative (day 3), ; baby2: negative day 3, PCR positive day 6, 23, PCR negative day	Not attempted	Neonate1: died from respiratory and gastrointestinal bleeding; Neonate2: healthy at follow-up; Neonate3: died day 16 of abdominal distension and enterocolitis

						30, healthy at 37 day follow-up; baby3: PCR negative day 3,6		
Fenzia B 2020	31 mother & neonate dyads	4/31 women had severe COVID-19 (defined by the need of urgent delivery for the deterioration of maternal conditions or by ICU/sub-intensive care admission). Radiological confirmation of interstitial pneumonia was obtained on admission or antepartum for all 4 severe cases and in 10 (32%) of the mild cases.	25/31 vaginal; 6/31 caesarean	NP swab, placenta & umbilical cord biopsy, umbilical cord blood; amniotic fluid.	NP swab	2/31, all babies healthy	not attempted	2/31 maternal plasma samples positive (both had severe clinical outcome). SARS-CoV-2 in vaginal swab, placental tissue and cord plasma from 1 mother (severe outcome). SARS-CoV-2 in placental tissue from subject 1 mother. SARS-CoV-2 in one milk specimen only (severe clinical outcome). None of the tested 6 amniotic fluids, nor the 12 umbilical cords, resulted positive. Placentae from SARS-CoV-2 infected patients displayed a generalized immune activation profile compared to the uninfected
Ferraiolo A 2020	1 mother & neonate dyad	Hospitalized in order to undergo C-section. No symptoms on admission of fever/ respiratory/ GI symptoms, myalgia, malaise, ageusia/ anosmia	elective caesarean	NP swab placental biopsy and placental swab	NP swab at 0h and 24h	0/1, healthy, discharged at 21 days	not attempted	Placental swabs COVID-19 positive, some placental inflammation on biopsy

Gale C 2020	66 SARS-CoV-2 positive neonates (<12 HOL)	maternal symptoms NR	not reported	not specified	NP swab? - not specified for all	2/unclear	not attempted	17 babies born to mothers with confirmed infection within 7 days before or after birth. Only 2 babies considered vertical transmission due to positive nasopharyngeal swab
Gao W 2020	1 mother & neonate dyad	no fever or cough; thoracic CT showed no abnormality	emergency C-section	NP swab,	NP swab	0/1; day 4 PCR test negative	not attempted	positive serum IgM and IgG antibody (colloidal gold method) were weak positive and strong positive
Gao X 2020	Unclear, possibly 14 women	11/14 women had fever, 6/14 had cough; all 14 had abnormalities evident by thoracic CT. 8/14 were given oxygen via nasal cannula; none was treated with respirator or mechanical ventilation.	2 vaginal, 12 Caesarean section	OP or NP swab	nasal or oropharyngeal swab	0/14	not attempted	All breast milk samples negative for viral RNA, 3 breast milk samples positive for SARS-CoV-2 antibodies
Grimminck K 2020	1 mother & neonate dyad	Cough developed prior to admission	vaginal, induced labour	OP swab vaginal sample, urinary catheter sample, placenta	OP swab	0/1	not attempted	None of the samples taken from vagina, placenta, catheter or baby were positive

He Z 2020	22 neonates	NR	not reported	not reported	Throat swab, amniotic fluid, umbilical cord blood, faeces, urine, blood.	0/22	not attempted	None of the samples taken from neonates tested positive for SARS-CoV-2. 3 newborns had elevated IgM Ab in umbilical cord & fetal blood; 12/22 had IgG positive for SARS-CoV-2; no deaths; 17/22 had hyperbilirubinaemia; report potential kidney damage in newborns
Hinojosa-Velasco A 2020	1 mother & neonate dyad	Fever, coughing, odynophagia, sore throat, headache, diarrhoea, rhinorhea, 2 days before admission	emergency C-section	NP and OP swab, breast milk and stool sample	nasopharyngeal and oropharyngeal swabs, during delivery, RT-PCR; stool sample	1/1	not attempted	Neonate considered to have a severe case of COVID-19 based on severity of symptoms, placed on oxygen. Day 4 milk & stool samples positive for SARS-CoV-2 RNA. 13 days after delivery, the infant's NP & OP swabs and stool samples were negative; maternal samples remained positive.
Hsu AL 2020	1 mother & neonate dyad	Myalgia 2 days prior to hospitalisation, no fever throughout hospitalisation	vaginal	placenta, RT-PCR and histology with immunohistochemical staining	not stated; RT-PCR	0/1, SARS-CoV-2 RT-PCR test negative at 24h for baby (sample source not stated)	not attempted	IHC using SARS-CoV-2 nucleocapsid-specific monoclonal antibody demonstrated SARS-CoV-2 antigens throughout the placenta, under the umbilical cord, and at the central and peripheral placenta disc in chorionic villi endothelial cells, and rarely in CK7-expressing trophoblasts.

Hu X 2020	6 mothers & 6 preterm neonates	In the 6 mothers, COVID-19 experienced 9, 10, 20, 13, 2, 2 days before delivery. All mild except one termed uncomplicated. Fever in 3/6, cough in 2/6.	5 C-section; 1 vaginal	Unclear	Throat & anal swabs, blood; gastric aspiration before feeding, urine and stool samples.	0/6	not attempted	All samples of amniotic fluid, cord blood and breast milk negative. All neonate samples negative.
Khan MA 2020	66 mothers; 67 infants (2 twins)	Most of the 66 SARS-CoV-2-positive women were asymptomatic. One woman was admitted with respiratory failure.	20 spontaneous vaginal delivery; 7 elective C-section; 40 emergency C-section	Not reported	NP swab	0/67	not attempted	
Kulkarni R 2020	1 mother & neonate dyad	Fever and myalgia one day prior to admission in active labour. Mother tested negative on admission and on day 5, serology negative on day 2, but serology positive on day 10.	Vaginal delivery, ventouse	NP swab	NP aspirate, cord stump, and placenta at birth	1/1;		All neonate samples positive at 12h. Neonate admitted with fever, suspected sepsis. Negative on day 14, discharged on day 21.

Liu W 2020	48 neonates	Fever (10/15, 67%) and cough (6/15, 40%) among mothers with COVID-19 diagnosis.	42 C-section; 6 vaginal (no differences between the 3 groups of women)	NP swab, CT scan	amniotic fluid, placental swab, gastric lavage fluid at birth; neonatal serum, throat swab, and feces	0/48.		15 neonates from COVID-positive women; 17 neonates from suspected COVID women; 16 neonates as control group. All samples from amniotic fluid, placental swab, gastric lavage fluid at birth; neonatal serum, throat swab, and feces negative.
Luo Q 2020	23 mother & neonate dyads	Fever (71%) and cough, 8/14 confirmed COVID-19 cases asymptomatic	C-section: 17; Vaginal: 6	Throat swabs, breast milk, blood.	Throat swabs at delivery, serum at 1 month	0/23	not attempted	All throat swabs negative, all antibody tests at 1 month negative (8 infants). None of the women required ventilation. All breast milk samples negative for viral RNA; IgM antibody present in milk in 4 confirmed cases, IgG antibody negative in all samples
Lv Y 2020	1 mother & neonate dyad	Recurrent fever and cough over 2 weeks prior to admission	C- section	not stated, PCR test	Amniotic fluid, umbilical cord blood, placenta, and neonatal gastric fluid, pharyngeal and anal swab	0/1	not attempted	All neonatal samples negative.

Maraschini A 2020	146 women, 143 singleton babies, 3 twins; 2 stillbirths, 147 live births	41/146 (28.1%) of the COVID-19-positive women asymptomatic. 70/146 had fever, 68 had cough, 52 fatigue. Onset of clinical symptoms occurred in 9.5% of the cases on the day of delivery, and in 90.5% before it, the median value being 8 days (range 1-52 days).	Vaginal: 98; Elective C-section: 12; Emergency C-section (maternal or foetal indication): 25; Emergency C-section (COVID-19): 11	NP swab	not stated, RT-PCR	9/149	not attempted	9 infants tested positive for SARS-CoV-2, testing either at birth or up to 9 days after. Of 5 newborns with positive test within 24h, 4 were delivered vaginally, 1 by pre-labor C-section; none developed serious illness. 11 women received ventilation, no ECMO, no deaths. Two stillbirths at 30 and 35 weeks pregnancy. 23 infants admitted to NICU, 18 premature, no deaths
Marin Gabriel MA 2020	242 mothers, 248 neonates	Cough (33%); fever (30%). Odynophagia (2%) and chest pain (<2%) were uncommon. 7 women required ICU. One mother died due to a massive thromboembolic event.	Vaginal: 179; Caeserean : 63	NP and/or OP swab, blood	NP and/or OP swab	11/230 positive (test at 18h).	not attempted	2 additional cases at second test (48h) 222/248 neonates did not need respiratory support, no deaths. No cases of pneumonia or pneumothorax. No additional cases at 1 month follow-up, 40% breast-fed exclusively at follow-up.
Masmejan S 2020	13 mother & neonate dyads	5 asymptomatic, 6 mild, 1 severe and 1 critical.	Vaginal: 9; Forceps: 2; C-section: 2	NP swab, placental swab.	NP swab, cord blood	0/13; all samples (cord blood, placenta, neonate nasopharyngeal swab) negative	not attempted	1 critical case amongst the women, requiring ventilation. No deaths; placental samples negative.

Mohakud NK 2020	1 mother & neonate dyad	Presented with four-day history of low-grade fever, malaise, and breathing difficulty.	C-section	Not reported	Tracheal aspirate swab at 12 hours of life	1/1	Not attempted.	Woman developed breathing difficulty, decreased fetal movements, edema, and visual disturbance and was admitted to hospital & diagnosed with HELLP syndrome with hypothyroidism and moderate COVID-19 pneumonia. Neonate chest X-ray normal; neonate required ventilation at birth, subsequently experienced a seizure but recovered & was discharged healthy.
Moreno SC 2020	19 women, 21 neonates (including two sets of dichorionic diamniotic twins)	Women included in the study if symptomatic and with positive PCR for SARS-CoV-2. Cough 19/19; fever 7/19; shortness of breath 5/19.	12 spontaneous vaginal; 7 C-section	NP swab	NP swab within first 24 hours of life.	0/21	Not attempted.	All 19 women presented with cough. Abnormal radiologic chest X-ray findings in 13/19 women. 8/19 births were premature deliveries. Among the neonates, no invasive mechanical ventilation was required. No neonatal sepsis or neonatal mortality was observed.
Nayak AH 2020	977 pregnant women: 141 with positive SARS-CoV-2 test; 836 with negative test. 131	97% of the women were asymptomatic or had mild symptoms like fever or cough not requiring any oxygen therapy. (unclear if 97% is of all women in the study, or of all positive-testing women).	481 vaginal (66/141 SARS-CoV-2 positive mothers); 443 C-section (67/141 SARS-CoV-2 positive	Unclear, probably NP swab.	Swab within first 24 hours of life	3/131	Not attempted.	Of 131 neonates tested, 3 tested positive on first swab within 24 hours of birth; all tested negative on day 5.

	neonates tested.		mothers); 10 instrumental (1/141 SARS-CoV-2 positive mothers)					
Ogamba I 2020	40 women, 25 deliveries within the time frame of the study; 23 neonates	30/40 SARS-CoV-2-positive women reported <= one symptom due to COVID-19; 10/40 were asymptomatic. Symptoms included loss of smell & taste, nausea, vomiting, body aches, chills, headache, fever, dry cough, shortness of breath, abdominal pain, difficulty breathing, chest pain.	17/25 vaginal; 8/25 C-section	NP or nasal swab	NPI or nasal swab	0/20	Not attempted.	
Olivini N 2020	5 mother & neonate dyads	3/5 women reported symptoms: 1 low grade fever, 2 anosmia, 1 dysgeusia, 1 musculoskeletal pain. Onset of symptoms occurred at day 1, 5 and 19 after childbirth.	2 vaginal, 3 elective C-section	rhino-pharyngeal swabs	rhino-pharyngeal and rectal swabs	0/1 neonate tested within 24h; 5/5 tested positive, at day 2, 16, 12, 6 and 4	Not attempted.	Neonates were asymptomatic or paucisymptomatic, no fever or respiratory symptoms. Three women reported symptoms: 1 low grade fever, 2 anosmia, 1 dysgeusia, 1 musculoskeletal pain. Onset of symptoms occurred at day 1, 5 and 19 after childbirth. Single samples of breastmilk from each of two of the SARS-CoV-2 positive mothers tested negative by RT-PCR.

Oncel MY 2020	125 pregnant women; NP swabs available for 120 newborns (not possible for 5 [asymptomatic] newborns)	85/125 SARS-CoV-2 positive women had at least one COVID-19 symptom. (40/125 had had close contact.) 8 were admitted to ICU for mechanical ventilation, 6/these 8 died.	89 C-section, 36 other (mode not reported)	Not reported	NP/pharyngeal; subsequently, deep tracheal aspirates	4/120; no NP sample tested positive within 24h; 1 positive on day 2, 2 positive on day 4; 1 deep tracheal aspirate tested positive on day 1	Not attempted.	Additional COVID-19 tests done in some (rationale not recorded): placenta tissue (n: 5), amnion fluid (n: 4), deep tracheal aspirate (n: 9), serum (n: 3), stool (n: 2), and breast milk (n: 6), which were all negative for SARS-CoV-2 except only one positive deep tracheal aspirate. 8/125 mothers were admitted to ICU for mechanical ventilation; 6 died.
Palalioglu RM 2020	1 mother and neonate dyad	asymptomatic	C-section	NP swab	NP swab	0/1	Not attempted.	CT performed after C-section showed ground glass opacities in both lungs of the mother despite no COVID-19 symptoms. Baby was admitted to the ICU after developing feeding intolerance and vomiting 6 h after being breastfed, later discharged healthy. Also amniotic fluid, cord blood and placenta, postoperative breast milk were negative by RT-PCR.

Parsa Y 2020	1 mother & neonate dyad	Presented with signs and symptoms of acute respiratory illness including shortness of breath and cough.	C-section	NP swab	Not reported	1/1 24 hours after delivery	Not attempted.	Mother aged 41 years, with opium addiction, with signs and symptoms of acute respiratory illness presenting with labor pain & vaginal leak at 37 weeks of gestation. CT-scan revealed ground-glass opacities. The RT-PCR results of the amniotic fluid and neonate (less than 24 hours after birth) were positive for COVID-19. The newborn suffered vomiting during the first 24 hours after birth, due to opium withdrawal syndrome. She was admitted to the NICU and received supportive care without need for respiratory support.
Pereira A 2020	22 mother & neonate dyads	6/22 mild symptoms, 5/22 pneumonia, 10/22 no symptoms	4 C-section, 18 vaginal (4 of which instrumental)	Not reported.	NP swab taken within first two hours of life.	0/22	Not attempted.	Two born preterm needed admission to the NICU. During follow-up period, there were no major complications, and no neonates were infected during breastfeeding.
Pessoa FS 2020	1 mother & neonate dyad	Admitted with flu-like symptoms including fever, dry cough & fatigue; symptom onset 7 days prior to admission, progressed to dyspnoea	C-section	Not reported	Swab at 6 HOL, source not reported.	1/1	Not attempted.	

Pissarra S 2020	10 mother & neonate dyads	7/10 symptomatic; 1 mother admitted to ICU; days from beginning of symptoms to delivery ranged from 0-15. No details of symptoms or severity	6 vaginal, 4 C-section (obstetrical indication)	NP swab or oropharyngeal swab	NPI swab and bronchial secretions	0/10	Not attempted.	All neonates separated from mothers at birth and transferred to NICU. All remained symptom free and tested negative for SARS CoV2 at birth & at 48 h of life.
Popofsky S 2020	160 neonates	59 mothers (36.9%) were symptomatic during perinatal hospitalization, with fever (>37.7°C), cough, shortness of breath, or a combination of these. 25 mothers (15.6%) had been symptomatic before the perinatal admission but were no longer symptomatic during hospitalization. 76 (48%) were never symptomatic. Mean day of onset of symptoms 14.	122 vaginal; 38 C-section	NP swab	NP swab	1/ 160 newborns (positive test was on day of life 5, after a negative test at birth)	Not attempted.	Mother of infant testing positive was symptomatic at the time of birth & had close contact with the newborn whilst symptomatic.
Pulinx B 2020	1 mother & two twin neonates	Presented to ED at 22w with rhinitis and fever (39.2 °C)	Vaginal	NP swab	placental and amniotic fluid	2/2 (placental and amniotic fluid)	Not attempted.	Both fetuses died prepartum. Placental histological examinations showed chronic intervillitis and extensive intervillous fibrin depositions with ischemic necrosis of the surrounding villi. Viral localization in the placental syncytiotrophoblast cells was confirmed by immunohistochemistry. No other cause of fetal demise was identified.

Rebello CM 2020	1 mother & neonate dyad	Anosmia, ageusia, runny nose, dry cough, headache & myalgia, no fever (10 days before delivery); diagnosed with COVID-19 eight days before delivery	Vaginal	NP swab	umbilical cord blood and skin swabs	1/1 (cord blood and neonate skin swabs)	Not attempted.	
Rivera-Hernandez P 2020	1 mother & neonate dyad	Presented at 33weeks pregnant, 3-day history of malaise & dyspnoea; ICU admission & ventilation; delivery on day 4 of admission	C-section	NP swab	NPI swab	1/1	Not attempted.	No postdelivery contact of neonate with mother or father; father tested negative. Chest radiography in neonate showed mild ground glass appearance & increased pulmonary vascularity bilaterally, consistent with respiratory distress syndrome. Neonate treated in NICU.
Rubio Lorente AM 2020	2 pregnant women	Case 1, symptomatic; Case 2, fever 2 weeks before, resolved prior to amniocentesis	n/a	serology & NP swab; amniocentesis	amniotic fluid	0/2	not attempted	Amniotic fluid negative for SARS-CoV-2 by RT-PCR for both cases
Sajjan GR 2020	1 mother & neonate dyad	Asymptomatic of respiratory disease	C-section	not reported	not reported	1/1, IgG	Not attempted	Neonate Covid Ig G positive, Neonatal lower limb gangrene considered to result from in utero transmission of SARS-CoV-2 infection. At 15 days, neonate was stable.

Schwartz DA 2020	19 neonates who tested positive and their mothers.	Predelivery symptoms: 10 respiratory symptoms (2 severe), 4 asymptomatic, 5 no information reported.	11 births by C-section; 5 vaginal; 3 mode of delivery not reported	Not reported	NP, endotracheal or OP swab		Not attempted	5/19 mothers negative despite positive neonate. 3 infants readmitted later due to signs of possible COVID-19. One of the included 19 neonates was one of triplets, the other two neonates had died of respiratory distress.
Shende P 2020	1 mother & first trimester fetus	Asymptomatic, tested positive 6 weeks previously.	Termination at 13 weeks	Throat swab	Products of conception, serum, placenta, amniotic fluid	1t	Not attempted	Products of conception negative for IgM. Placenta showed infection. Viremia persisted in the placenta six weeks after the mother was tested positive.
Singh MV 2020	1 mother & neonate dyad	Asymptomatic	C-section	NP swab	NP swab	1/1	Not attempted	C-section performed at 38 weeks due to positive SARS-CoV-2 test. Neonate healthy. NP swab turned negative on day 10 of life.
Sisman J 2020	1 mother and infant dyad	3 days pre-delivery admitted with fever and diarrhoea (admitted due to possible preterm labour)	vaginal	NP swab	NP swab	1/1 at 24 and 48 HOL		Premature rupture of membranes so labour induced. Hyperbilirubinemia in the neonate. Neonatal fever & respiratory distress at day 2, resolved by day 5. Chest radiograph normal. Discharged as healthy at day 21.

Smithgall MC 2020	76 pregnant women	26/51 SARS-CoV-2-positive women were asymptomatic; 32 had cough, 27 had fever, 14 had myalgia, 6 had sore throat, 6 had fatigue. 4 suffered severe disease.	51/76 vaginal; 25/76 C-section	placentas	NR	0/51	Not attempted	No adverse perinatal outcomes; no specific histomorphologic changes in placentas; no evidence of direct viral involvement identified by ISH and IHC. Evidence of maternal–fetal vascular malperfusion was identified: SARS-CoV-2 positive women's placentas significantly more likely to show villous agglutination & subchorionic thrombi than SARS-CoV-2-negative women's placentas.
Stonoga E 2020	1 mother & fetus dyad	Dyspnea, dry cough, high temperature (38.5°C), anosmia, nausea, vomiting, and diarrhea had developed 2 days before hospitalization.	C-section	NP swab tested by RT-PCR	Amniotic fluid (before amniotic membranes ruptured), umbilical cord blood, placental membranes, & cotyledon fragments	0/1	Not attempted	Fetal death. All fetal samples tested negative. Multifocal chronic histiocytic intervillitis in the placenta, potentially relevant to cause of death of the fetus, but other causes not excluded.

Tang F 2020	20 mother & neonate dyads	Unclear. However, the mothers of two neonates who tested positive for RT-PCR at 1–2 days after birth also tested positive & exhibited typical symptoms such as fever, cough, one day pre delivery	5 vaginal, 15 C-section	Unclear	Throat swabs, serum, amniotic fluid, umbilical blood (and radiology)	3/20	Not attempted	No neonate had a fever. Radiological examination showed that 10 neonates, eight neonates, and two neonates were diagnosed with pneumonia, increased lung markings, and no abnormality in the lungs, respectively. All amniotic fluid & umbilical cord blood tested negative.
Tang J 2020	2 mother & neonate dyads	Case 1: fever & respiratory symptoms 13 weeks prior to delivery, symptoms resolved; admission to hospital 13 weeks later with premature rupture of membranes, no respiratory symptoms, but serum antibodies positive for SARS-CoV-2. Case 2: mild respiratory symptoms predelivery.	1 vaginal birth, 1 birth by C-section	Throat swabs, serum	Throat swabs at birth, day 3, day 7. Serum on day 7.	0/2 by PCR; 2/2 positive by IgG, 1/2 positive by IgM.	Not attempted	Mothers: throat swabs negative by PCR, 2/2 were positive for serum IgM, 1/2 positive for IgG antibodies. The two mothers did not develop serious complications; outcomes in the neonates were good. Neonate serum IgM antibody to SARS-COV-2 was negative and IgG was positive on the 7th day after birth.
Vendola N 2020	2 mother & neonate dyads	Both cases asymptomatic	1 vaginal, 1 C-section	NP swab, serum	NP swab, serum, umbilical cord blood, peripheral blood soon after delivery.	0/2 for SARS-CoV-2 RNA in throat swabs; 2/2 for IgG in neonatal serum	Not attempted	

Vinuela MC 2020	100 asymptomatic pregnant woman & their neonates	All asymptomatic	94 vaginal, 6 C-section	NP swab	Unclear	0/100	Not attempted	No fetal transmission was observed (details NR). Maternal & neonatal prognosis excellent. All Cts from maternal swabs were >35.
Vivanti AJ 2020	1 mother & neonate dyad	Fever (38.6 °C) and severe cough	C-section	Blood, NP swab, vaginal swab, amniotic fluid during C-section	NP and rectal swabs at 1 hour, 3 and 18 days. Cerebrospinal fluid sample at 3 days.	1/1	Not attempted	Mother admitted at 35+2 weeks of gestation with fever (38.6 °C), severe cough, abundant expectoration for 2 days before hospitalisation. Neonatal blood culture was negative for bacteria or fungi. All neonatal swab samples tested positive by PCR. CSF sample negative. Placenta histology some positive Ab staining for SARS-CoV-2
Von Kohorn I 2020	1 mother & neonate dyad	Maternal gestational diet-controlled diabetes. Presented with cough, vaginal bleeding at 34 weeks gestation. COVID-19 positive 14h prior to delivery. Mother had a cough for 1 week prior to NP swab & test.	C-section	NP swab	Cord blood at birth. NP swabs at 24 hours and 49 HOL. Neonatal blood & urine over first 7 days of life.	1/1 (several consecutive samples tested, some were positive by PCR with variable Ct values)	Not attempted	Infant asymptomatic with normal laboratory studies but some NP swabs tested positive for SARS-CoV-2 RNA. Placenta histology showed no direct evidence of virus.

Vouga M 2020	1,033 pregnant women: 926 tested positive, 107 tested negative. 384 neonates tested at birth.	Symptoms included fever, anosmia, cough, sore throat, dyspnoea, myalgia, fatigue, headache, nausea. Symptom frequency reported in the cohort, symptom severity not reported. Some asymptomatic	Among 926 SARS-CoV-2 positive women: 469 vaginal, 256 C-section, 6 unknown; among 107 SARS-CoV-2 negative women: 44 vaginal; 22 C-section, 1 unknown.	NP swab	NR	11/384	Not attempted	No difference in obstetrical and neonatal outcomes were observed between positive- and negative-testing women.
Woodworth KR 2020	4,442 positive pregnant mothers, 4,495 live births.	Symptom status was known for 2,691 (60.6%) women (including women with symptoms reported on COVID-19 form), 376 (14.0%) of whom were reported to be asymptomatic. Symptoms and severity not reported	2,589 vaginal, 1,331 C-section. Among 3,920 women testing positive for SARS-CoV-2: 2,589 vaginal, 1,331 C-section.	NR	NR	16/610	Not attempted	% positivity was 4.3% (14 of 328) among infants born to women with documented infection ≤14 days before delivery and 0% (0 of 84) among those born to women with documented infection >14 days before delivery. Molecular testing results available for 610 infants. Other perinatal outcomes also reported.

Yang R 2020	11,078 live singleton birth mother & neonate dyads:	Those with fever and cough or having abnormal CT scan were tested for SARS-CoV-2. Severity of symptoms not reported.	5006/11,078 vaginal, 6,072 C-section (Among 65 SARS-CoV-2 positive women, 13 vaginal, 52 C-section)	Pharyngeal swab	Unclear, probably pharyngeal swab as for mother.	0/38	Not attempted	65/11,078 mothers tested positive during pregnancy. 38 /58 newborns of positive testing mothers were tested for SARS-CoV-2 after birth: 0 positive. OR for perinatal conditions & SARS-CoV-2 positivity also reported.
Zaigham M 2020	1 mother & neonate dyad	34 weeks pregnant. Fever, dry cough, abdominal pain. Treated in postpartum ward (no ICU)	C-section	NP & throat swab	NP swab 48 hours after delivery	1/1	Not attempted	High viral load in placenta with massive perivillous fibrin deposition, acute intervillitis in areas with strong positivity for SARS-CoV-2 & chorangiosis in the areas less affected by infection and inflammation. Whole genome sequencing of isolates from the mother & placenta revealed a single variant of the virus.
Zhang L 2020		35 to 41 weeks pregnant. Most positive pregnant women had fever, cough, sore throat, fatigue, chest tightness or shortness of breath, diarrhea, and runny nose. Clinical typing: 1 case "mild", 16 cases "ordinary", 1 case "severe"	1 vaginal, 17 C-section	Throat swab	Sample source NR	0/18	Not attempted	5/18 neonates diagnosed with bacterial pneumonia, successfully treated.

Zhang P 2020	364 pregnant women	Not reported	137/364 vaginal, 227/364 C-section; among SARS-CoV-2 positive women 54/74 vaginal, 20/74 C-section.	NP swab	Placental tissue	2/53 by in situ hybridization in placenta	Not attempted	74/364 women tested positive. 2/53 placental samples positive from PCR positive mothers. 0/10 neonatal placentas positive from negative mothers. No histopathological features within the placentas specific to maternal SARS-CoV-2 infection.
Zheng T 2020	2 x mother + neonate dyads	Ppt 1: 36weeks pregnant, fever & asthenia at admission, CT showed bilateral lung infection, classified as "severe COVID-19"; Ppt 2: 39weeks, admitted due to abnormal chest CT; no fever or cough, CT showed bilateral lung infection.	C-section (both)	Throat swab	Throat swab, anal swab, urine, & blood; sample timing unclear	0/2	Not attempted	

Table 1c. Summary of findings from reviews

Study	Fulfills systematic review methods (Y/N/U); Risk of bias tool used	Stated research question	Databases and search date	No. included studies (No. participants)	Main results	Authors' key conclusions
Akhtar H 2020	Yes; RoB: Newcastle Ottawa Scale	Review of association of SARS-CoV-2 infections with pregnancy, foetal, and neonatal outcomes	PubMed, Scopus, Medline, Cochrane database, Google Scholar from 1 December 2019 to 22 May 2020	22 (156 pregnant women with COVID-19; 108 neonates)	Most common maternal/foetal complications included intrauterine/foetal distress (14%) and premature rupture of membranes (8%). Neonatal clinical manifestations of COVID-19: shortness of breath (6%), gastrointestinal symptoms (4%), fever (3%).	COVID-19 infection in pregnancy leads to increased risk in pregnancy complications such as preterm birth, PPRM; may rarely lead to maternal death. No evidence to support vertical transmission of SARS-CoV-2. Limited evidence on impact of COVID-19 on newborns.
Al Qahtani MA 2020	No	Summarize research regarding pregnancy and vertical transmission of COVID-19	Pubmed from 1 January 2020 to March 20 2020	9 studies (74 pregnant women, 72 positive for COVID-19, 2 negative but symptomatic; 75 neonates)	All except one neonate tested negative; one tested positive for throat swab, but negative for cord blood and placenta.	Only one infant from a positive mother tested positive for COVID-19 throat swab. No difference between pregnant and nonpregnant patients regarding clinical manifestations.
Amaral W 2020	Yes; RoB: GRADE	To investigate pregnant women infected with COVID-19 in terms of signs and symptoms, type of delivery, comorbidities, maternal and neonatal outcomes, and the possibility of vertical transmission.	Embase and PubMed search on 31 October 2020.	70 studies (1457 pregnant women)	Maternal signs and symptoms: fever, cough, nausea. Maternal and fetal outcomes: premature birth (n = 64), maternal death (n = 15), intrauterine fetal death or neonatal death (n = 16), cases of intrauterine fetal distress (n = 28), miscarriage (n = 7), decreased fetal movements (n = 19), severe neonatal asphyxia (n = 5). 39 newborns positive for SARS-CoV-2. SARS-CoV-2 RNA detected in placenta (n = 13) and breast milk (n = 6).	COVID-19 during pregnancy can result in maternal, fetal, and neonatal complications. SARS-CoV-2 viral exposure of neonates during pregnancy and delivery cannot be ruled out. Need for long-term follow-up.

Bellos I 2020	Yes; RoB: Quality appraisal by domains, tool not stated	Evaluate maternal and neonatal outcomes in COVID-19 pregnancies, factors associated with perinatal viral transmission.	Medline, Scopus, CENTRAL, Web of Science and Google Scholar databases up to 3 June 2020.	60 studies	Maternal symptoms: Fever, cough, shortness of breath, 15% asymptomatic. Severe disease in 7-11%. 2 maternal deaths. Neonatal symptoms: fever, shortness of breath, vomiting. 20% asymptomatic. No difference in neonatal transmission rate between women with and without severe disease (OR: 1.94, 95% CI: 0.50–7.60). Preterm birth in 16-30%. Stillbirth, 3 cases, neonatal deaths, 2 cases. Vertical transmission, 4 cases.	Maternal and neonatal clinical course typically mild, low mortality rates. Risk of vertical transmission low, not linked to maternal disease severity..
Bwire GM 2020	Unclear; RoB: No quality assessment, as majority studies case reports.	Determine likelihood of vertical transmission in COVID-19 exposed infants; whether exposed negative infants had SARS-CoV-2 antibodies	PubMed/MEDLINE, Google Scholar. Websites of WHO, CDC; Google search of specific journals; Research Square, medRxiv. From 1 December 2019 to 18 May 2020	33 studies (205 infants born to COVID-19 positive mothers)	6.3% (13/205; 95% CI: 3.0%–9.7%) infants tested positive for COVID-19 virus at birth. SARS-CoV-2 IgG/IgM in 90% COVID-19-negative infants (10/11; 95% CI: 73.9%–107.9%) (6 studies). Median antibody levels IgG 75.49 AU/ml (range, 7.25–140.32 AU/ml), IgM 3.79 AU/ml (range, 0.16–45.83 AU/ml)	Low possibility of vertical transmission of COVID-19, SARS-CoV-2 antibodies in vertically exposed but negative infants.
Caparros-Gonzalez RA 2020	No	Guide clinicians on management of pregnant women re maternal-fetal-neonatal SARS-CoV-2 infections and breastfeeding during COVID-19 pandemic	Web of Science, PubMed, Scopus, Dialnet, CUIDEN, Scielo, Virtual Health Library from 25 August 2020 to 10 November 2020.	49 studies (329 pregnant women and 331 neonates)	Samples positive for SARS-CoV-2 RNA: 15 placental swabs positive on fetal side of the placenta; 7 breastmilk; 1 umbilical cord; 1 amniotic fluid	Evidence supports potential of congenital, intrapartum, and postnatal maternal-fetal-neonatal SARS-CoV-2 infections. Mothers should follow recommendations e.g. wearing facemask and hand washing before and after breastfeeding.
Cavalcante de Melo 2020	Yes; RoB: Newcastle Ottawa Scale	Analyze whether COVID-19 in pregnant women is related to premature birth and birth weight; summarize diagnostic results of neonates born to mothers with COVID-19 to investigate possibility of vertical transmission.	PubMed, Scopus, LILACS, Web of Science, Google Scholar, Preprints, bioRxiv, and medRxiv. Search date 4 May 2020	8 studies (279 women, 60 diagnosed with COVID-19)	No significant association between COVID-19 and preterm delivery (OR = 2.25; 95%CI: 0.96, 5.31; p = 0.06; I ² = 0%). No significant relationship between birth weight and COVID-19 (MD = -124.16; 95%CI: -260.54, 12.22; p = 0.07; I ² = 0%).	COVID-19 in pregnant women may not be associated with occurrence of preterm deliveries or birth weight of newborns. Level of evidence low. A few reports suggest vertical transmission possible, but evidence uncertain.

Centeno-Tablante E 2020	Yes; RoB: Newcastle Ottawa Scale.	Summarise information on breast milk and breastfeeding to inform guidance during COVID-19 pandemic.	MEDLINE (PubMed), the WHO COVID-19 database, Cochrane Library, Web of Science Core Collection, Embase. COVID-19 subset of WHO International Clinical Trials Registry Platform. Search date 7 July 2020	37 studies (77 breast-feeding mothers)	19/77 children confirmed COVID-19 cases (positive RT-PCR). 9/68 breast milk samples from mothers with COVID-19 were positive for SARS-CoV-2 RNA; of the exposed infants, 4 were positive, 2 were negative for COVID-19.	No evidence of SARS-CoV-2 transmission through breast milk. Long-term follow-up studies needed
Chi J 2020	Yes; RoB: Case report/case series (Murad et al.)	Summarise clinical features and maternal–infant outcomes of pregnant women infected with COVID-19 and their infants, including possibility of vertical transmission.	PubMed, Embase, Medline, MedRxiv, CNKI, and the Chinese Medical Journal Full Text Database. Search up to 18 April 2020.	20 studies (230 women with COVID-19, 156 newborns)	Pregnant women: 34.6% obstetric complications, 59% fever, 40.7% lymphopenia, 5.19% received mechanical ventilation, 7 critically ill. Deaths: 1 maternal, 2 neonatal. Newborns: 24.74% premature, 5 SARS-CoV-2 positive (throat swab), all positives delivered by C-section. 8 negative newborns with elevated SARS-CoV-2 IgM and IgG. PCR tests of vaginal secretions, breast milk, amniotic fluid, placental blood, placental tissues all negative.	Most pregnant patients mildly ill. Mortality of pregnant women with COVID-19 lower than for COVID-19 patients overall. C-section more common than vaginal delivery for pregnant women with COVID-19. Main newborn adverse event: premature delivery. Vertical transmission rate 3.91%.
Della Gatta AN 2020	Yes; RoB: Case report/case series (Murad et al.)	Review clinical outcomes for pregnant patients with COVID-19.	PubMed, CINAHL, Scopus. From 14 March 2020 to 16 March 2020	6 studies (51 pregnant women)	48 pregnant women, 46 cesarean (96%), 2 vaginal delivery. 1 stillbirth and 1 neonatal death. Median gestational age 36.5 weeks (interquartile range, 35–38), 15 preterm birth (39%); indications for cesarean delivery not clearly described.	Vertical transmission of SARS-CoV-2 excluded thus far, outcomes for mothers and neonates generally good; high rate of preterm delivery by cesarean, mostly elective, reason for concern. Reasonable to question whether cesarean delivery for pregnant patients with COVID-19 warranted.
Deniz M 2020	No	Review evidence on vertical transmission of SARS CoV-2.	PubMed, Global Health (OVID), SCOPUS from 1 February 2020 to 1 June 2020	50 studies (714 pregnant women with COVID-19, 606 neonates)	17 newborns testing positive for SARS CoV-2 by RT-PCR. 3 neonates with elevated SARS-CoV-2 IgG and IgM. Reporting samples testing positive: 8 placental tissues, 3 breast milk, 1 amniotic fluid	Possible vertical transmission of SARS CoV-2 has been observed in some studies. More RT-PCR tests on amniotic fluid, placenta, breast milk and cord blood are required.

Dhir SK 2020	Yes; RoB: Newcastle Ottawa Scale	Synthesize literature on various modes of transmission (congenital, intrapartum, and postpartum), clinical features and outcomes of SARS-CoV-2 infection in neonates.	PubMed, EMBASE, Web of Science until 9 June 2020.	86 studies (1992 pregnant women, of which 1125 gave birth to 1141 neonates)	281 (25%) neonates were preterm, preferred delivery mode caesarean section (66%). 41 case reports of 43 mother-baby dyads reporting 16 preterm births, 9 low birth weight. 58 neonates reported with SARS-CoV-2 infection, 29 (50%) symptomatic (23 required ICU). 70% with respiratory symptoms. No neonate mortality.	Limited low-quality evidence suggests risk of SARS-CoV-2 infections in neonates is extremely low. Most COVID-positive neonates symptomatic and required intensive care. Postpartum acquisition the commonest mode of infection in neonates, a few cases of congenital infection.
Di Toro F 2020	Yes; RoB: Joanna Briggs Institute	Assess the impact of COVID-19 on maternal and neonatal outcomes.	PubMed, EMBASE, MedRxiv, Scholar, Scopus, Web of Science database up to 8th May 2020.	24 studies (1100 pregnancies).	Pooled prevalence of pneumonia 89% (95%CI 70–100), prevalence of women admitted to ICU 8% (95%CI 1–20). 3 stillbirths, 5 maternal deaths. Pooled prevalence of 85% (95%CI 72–94) for caesarean deliveries. 3 neonatal deaths. Prevalence of COVID-19-related admission to neonatal ICU 2% (95%CI 0–6). 19/444 neonates positive for SARS-CoV-2 RNA at birth. 1 case of elevated levels of IgM and IgG, but negative swab.	Adverse outcomes e.g. ICU admission or patient death can occur, but clinical course of COVID-19 in most women not severe, infection does not significantly influence the pregnancy. High caesarean delivery rate reported, but no clinical evidence to support this. In most cases disease does not threaten the mother, vertical transmission not clearly demonstrated. COVID-19 should not be considered indication for elective caesarean section.

Diriba K 2020	Yes; RoB: Joanna Briggs Institute	Assess the effect of coronavirus infection (SARS-CoV-2, MERS-CoV, and SARS-CoV) during pregnancy and its possibility of vertical maternal–fetal transmission.	PubMed, Web of Science, Embase, Google Scholar, Cochrane Library until the end of April 2020	39 studies (1316 pregnant women)	Most common clinical features: fever, cough, myalgia (prevalence 30 to 97%); Abnormal lab findings: lymphocytopenia, CRP (55–100%). Pneumonia most diagnosed clinical symptom of COVID-19 and non-COVID-19 infection (prevalence 71-89%). CT imaging: Bilateral pneumonia (57.9%), ground-glass opacity (65.8%). Most common treatment: hydroxychloroquine (79.7%), ribavirin (65.2%), oxygen therapy (78.8%). Maternal outcome: rate of preterm birth < 37 weeks 14.3%, preeclampsia 5.9%, miscarriage 14.5%, premature rupture of membranes 9.2%, fetal growth restriction 2.8%. COVID-19 pregnant women: 56.9% delivered by cesarean, 31.3% admitted to ICU, 2.7% deaths. Perinatal outcomes: fetal distress 26.5%, neonatal asphyxia 1.4%, apgar score < 7 at 5 min 1.2%, neonates admitted to ICU 11.3%, death 2.2%. No reports of in utero transmission of CoV.	Coronavirus infection more likely to affect pregnant women. Respiratory infectious diseases demonstrated increased risk of adverse maternal obstetrical complications than general population. No studies reported transmission of CoV from the mother to the fetus in utero, may be due to very low expression of angiotensin-converting enzyme-2 in early maternal–fetal interface cells.
Dube R 2020	No	Review literature to determine effects of confirmed cases of COVID-19 in pregnant women by estimation of mother to child transmission, perinatal outcome and possible teratogenicity.	PubMed, Embase, LitCovid, Google Scholar, EBSCO MEDLINE, CENTRAL, CINAHL, MedRXiv, BioRXiv, Scopus between 1 November 2019 and 10 August 2020	Mother to child transmission: 72 studies (1408 neonates); Perinatal outcomes: 60 studies (1318 fetuses)	Positive tests for SARS-CoV-2 RNA: 3.67% of neonates (NP swab) cord blood 7.1%, placenta 11.7%, amniotic fluid 6.8%, faecal and rectal swabs 9.6%. Rate of preterm labour 26.4% (OR=1.45, 95% CI 1.03 to 2.03), caesarean delivery 59.9% (OR=1.54, 95% CI 1.17 to 2.03). Most common neonatal symptom breathing difficulty (1.79%). Stillbirth rate (COVID-19 positive mothers) 9.9 per 1000. Congenital transmission rate 9/1408 (0.63%).	Chances of mother to child transmission of the SARS-CoV-2 virus is low. Perinatal outcome for the foetus is favourable. Increased chances of caesarean but not preterm delivery. Stillbirth and neonatal death rates low. No reported congenital anomalies.

El-Wahab EWA 2020	No	Summarize SARS-CoV-2 transmission and provide scientific support for the prevention and control of COVID-19.	ISI Web of Knowledge, PubMed, Medline, ScienceDirect, EMBASE, EBSCO, ProQuest, Google Scholar, WHO, CDC, Mayo Clinic, bioRxiv, medRxiv, SSRN, Qeios from 28 December 2019 to 31 July 2020	302 studies	Broad review on various modes of transmission. Regarding vertical transmission, report generally on possibility of mother-to-child transmission and detection of SARS-Co-V-2 in breastmilk.	Very broad review of all types of transmission, including possible mother-to-child and breastmilk.
Figueiro-Filho EA 2020	No	Identify the most significant studies reporting on COVID-19 during pregnancy and provide an overview of SARS-CoV-2 infection in pregnant women and perinatal outcomes.	PubMed, EMBASE, Cochrane Library, Google Scholar until July 20, 2020.	8 studies (10,966 COVID-19 pregnant women)	Pregnant women not more affected by respiratory complications of COVID-19, compared to general population. Down-regulation of ACE2 receptors induced by SARS-CoV-2 cell entry might have been detrimental in subjects with pre-existing ACE2 deficiency associated with pregnancy, which may explain the worse perinatal outcomes.	Maternal characteristics, clinical symptoms, maternal and neonatal outcomes of 10,996 cases of COVID-19 and pregnancy are not worse or different from the general population. Pregnant women are not more affected by the respiratory complications of COVID-19, compared with non-pregnant available data.
Gao Y 2020	Yes; RoB: Institute of Health Economics	Review the clinical features and outcomes of pregnant women with COVID-19.	PubMed, Web of Science, EMBASE, MEDLINE from 1 January 2020 to 16 April 2020.	14 studies (236 COVID-19 pregnant women)	Positive CT findings (71%; 95% CI, 0.49–0.93), caesarean section (65%; 95% CI, 0.42–0.87), fever (51%; 95% CI, 0.35–0.67), lymphopenia (49%; 95% CI, 0.29–0.70), coexisting disorders (33%; 95% CI, 0.21–0.44), cough (31%; 95% CI, 0.23–0.39), fetal distress (29%; 95% CI, 0.08–0.49), preterm labor (23%; 95% CI, 0.14–0.32), severe case or death (12%; 95% CI, 0.03–0.20). Pregnant women with COVID-19 lower incidences of fever and cough.	Incidences of fever, cough and positive CT findings in pregnant women with COVID-19 are less than those in the normal population with COVID-19, rate of preterm labor is higher among pregnant women with COVID-19. No evidence of vertical transmission.
Goh XL 2020	Unclear; RoB: Newcastle Ottawa Scale	To determine a more precise risk of vertical transmission, either intrauterine or during delivery.	PubMed, Medline, Embase, China National Knowledge Infrastructure until 23 May 2020.	17 studies (402 women with COVID-19, 405 newborns)	9/330 newborns tested positive for SARS-CoV-2. Pooled incidence of vertical transmission 16 per 1000 newborns (95% CI 3.40 to 73.11).	Risk of vertical transmission low. Quality of the evidence moderate.

Han Y 2020	No	Assess perinatal outcomes of COVID-19 infections during pregnancy and the possibility of vertical transmission.	MEDLINE, PubMed, Web of Science, Cochrane library, China National Knowledge Infrastructure (CNKI), WANFANG DATA, VIP, SinoMed, Clinical trials.gov from 1 December 2019 to 10 June 2020.	36 studies (1103 patients)	Most common symptoms: fever (64.78%), cough (59.81%), dyspnea (23.86%). 88.73% had typical COVID-19 signs on chest CT or X-ray. Intubation in 35.87%, 4.95% mothers admitted to ICU, maternal death rate <0.01%, premature delivery 25.32%. Rate of the birth weight <2,500 g 30.65%, and that of NICU admission 24.41%. Positive nasopharynx swabs or sputum from newborns was <0.01%.	Pregnant patients with COVID-19 most commonly presented with fever, cough, dyspnea, most had positive imaging. Risk of intubation and admission to ICU were high. Risk of premature delivery was higher, leading to a high risk of NICU admission and low neonatal birthweight. Vertical transmission unlikely.
Hessami K 2020	No	Review evidence on maternal, fetal and neonatal mortality cases associated with COVID-19.	PubMed, Scopus, Google Scholar, Web of Science up to 20 July 2020	10 studies (37 maternal and 12 perinatal mortality cases)	All maternal deaths in women with previous co-morbidities, of which the most common were obesity, diabetes, asthma and advanced maternal age. Causes of mortality: Acute respiratory distress syndrome (ARDS) and severe pneumonia, one thromboembolism. Fetal and neonatal mortalities: result severity of maternal infection or the prematurity. No evidence of vertical transmission or positive COVID-19 among neonates.	Current available evidence suggests maternal mortality mostly happened among women with previous co-morbidities and neonatal mortality seems to be a result of prematurity rather than infection.
Huntley BJF 2020	Yes; RoB: Case report/case series (Murad et al.)	Review frequency of maternal and neonatal complications, as well as maternal disease severity, in SARS-CoV-2 pregnancies	MEDLINE, Ovid, ClinicalTrials.gov, MedRxiv, Scopus up to 29 April 2020.	13 studies (538 SARS-CoV-2 pregnancies, 435 deliveries)	Maternal ICU admission 3.0% (8/263, 95% CI 1.6–5.9), maternal critical disease 1.4% (3/209, 95% CI 0.5–4.1). No maternal deaths. Preterm birth rate 20.1% (57/284, 95% CI 15.8–25.1), cesarean delivery 84.7% (332/392, 95% CI 80.8–87.9), no vertical transmission, neonatal death rate 0.3% (1/313, 95% CI 0.1–1.8).	Early pandemic data indicates low rates of maternal and neonatal mortality and no vertical transmission with SARS-CoV-2. Preterm birth rate of 20% and cesarean delivery rate >80% seems related to geographic practice patterns.
Juan J 2020	Yes; RoB: Joanna Briggs Institute	Review literature on pregnancies affected by COVID-19 to evaluate the effects of COVID-19 on maternal, perinatal and neonatal outcomes.	PubMed, EMBASE, Cochrane library, China National Knowledge Infrastructure Database, Wan Fang Data up to 20 April 2020	19 studies (266 pregnant women with COVID-19)	Most common symptoms: fever, cough, dyspnea, fatigue. Rate of severe pneumonia low. Most had positive CT chest. 177 cases delivered, majority by Caesarean section. GA at delivery ranged from 28 to 41 weeks. Over one-third of neonates transferred to NICU. One case each of neonatal asphyxia and death. 113 neonates tested for SARS-CoV-2 were negative. 2 maternal deaths.	The clinical characteristics of pregnant women with COVID-19 are similar to those of nonpregnant adults with COVID-19. Vertical transmission unclear.

Khalil A 2020	Yes; RoB: Newcastle Ottawa Scale	Review SARS-CoV-2 infection and pregnancy, summarising evidence on the clinical features, laboratory and radiological findings, as well as the pregnancy and neonatal outcomes.	Medline, Embase, Clinicaltrials.gov, Cochrane Library up to 8 June 2020.	86 studies (2567 COVID-19 pregnancies).	Delivery by caesarean section 48.3%. Most commonly reported symptoms: fever (63.3%), cough (71.4%), dyspnoea (34.4%). Laboratory abnormalities: raised CRP or procalcitonin (54.0%), lymphopenia (34.2%), elevated transaminases (16.0%). Preterm birth before 37 weeks' gestation 21.8%, medically-indicated (18.4%). Maternal ICU 7.0%, intubation 3.4%. Maternal mortality ~1%. Maternal ICU admission higher in cohorts with higher rates of co-morbidities (beta=0.007, p<0.05) and maternal age over 35 years (beta=0.007, p<0.01). Maternal mortality higher in cohorts with higher rates of antiviral drug use (beta=0.03, p<0.001), likely due to confounding. Neonatal NP swab positive in 1.4%.	The risk of preterm birth and caesarean delivery was increased. Maternal morbidity is similar to that of women of reproductive age. Vertical transmission probably occurs, but in small proportion of cases.
Kotlyar AM 2020	Yes; RoB: Newcastle Ottawa Scale	Review current literature to determine estimates of vertical transmission of COVID-19	PubMed, EMBASE, medRxiv, bioRxiv up to 28 May 2020.	68 studies (979 neonates)	27/936 SARS-CoV-2 positive neonates from mothers with COVID-19, pooled proportion of 3.2% (95% confidence interval, 2.2-4.3) for vertical transmission. SARS-CoV-2 positive samples: neonatal cord blood 2.9% (1/34), placenta samples 7.7% (2/26), amniotic fluid 0% (0/51), urine 0% (0/17), fecal or rectal swabs 9.7% (3/31). Neonatal serology positive in 3 of 82 samples (3.7%) (IgM).	Vertical transmission of SARS-Co-V-2 is possible and seems to occur in a minority of cases of maternal COVID19 infection in the third trimester. Rates of infection are similar to those of other pathogens that cause congenital infections. No data on rates of vertical transmission in early pregnancy and potential risk for consequent fetal morbidity and mortality.
Mahyuddin AP 2020	No	Assess tools used to confirm maternal-fetal infection and known protective mechanisms of the placental barrier that prevent transplacental pathogen migration.	No search strategy reported. No search date reported.	40 studies	Lack of consensus on diagnostic strategy for congenital infection for COVID-19 pregnancies. Samples positive for SARS-CoV-2: vaginal secretions (22.5%), amniotic fluid (35%), breast milk (22.5%). Neonatal COVID-19 reports: 8 studies. Histology: sparse viral particles, vascular malperfusion, placenta inflammation	Provides overview biology of maternal-fetal interface, mechanisms for vertical viral transmission. Summarises diagnostics in pregnancy to assess mother-to-child transmission.

Novoa RH 2020	Yes; RoB: Newcastle Ottawa Scale	Describe maternal clinical characteristics, maternal and perinatal outcomes in COVID-19-positive pregnant women.	MEDLINE, EMBASE, Cochrane Library and LILACS; China National Knowledge Infrastructure Database (CNKI), Chinese Science, Technology Periodical Database (VIP) and Wan Fang Data from 1 December 2019 to 27 April 2020	37 studies (322 infected pregnant women)	Maternal comorbidity: obesity (24.2%). 42 (28.4%) asymptomatic at admission. Symptoms: Cough (n = 148,59.7%), fever (n = 147,59.3%). Fever (OR: 0.13% CI 0.05- 0.36) and cough (0.26% CI 0.11-0.59) lower in pregnant women with COVID-19 than non-pregnant women with COVID-19. Cesarean: 99 (50.8%), vaginal delivery: 64 (32.8%). Main adverse obstetric outcome: premature birth (n = 37,18.9%). ICU: 30 (10.3%), deaths: 1 (0.3%). SARS-CoV-2 absent in breast milk, amniotic fluid, placenta, umbilical cord blood.	Pregnant women with COVID-19 less symptomatic than general population. Usual symptoms are fever and cough. Severe morbidity and mortality similar to those reported for non-pregnant women. Main adverse obstetric outcome is preterm birth. Vertical transmission unlikely in third trimester.
Pettirosso E 2020	No	Describe understanding of COVID-19 illness in pregnant women, obstetric outcomes and identify gaps in knowledge.	Medline Ovid, EMBASE, WHO COVID-19 research database, Cochrane COVID-19 in pregnancy spreadsheet up to 23 May 2020.	60 studies (1287 SARS-CoV-2 positive pregnant cases)	8 maternal deaths, 6 neonatal deaths, 7 stillbirths, 5 miscarriages reported. 19 neonates SARS-CoV-2 positive, confirmed by RT-PCR of NP swabs.	SARS-CoV-2 infection in pregnancy often asymptomatic. Severe and critical disease similar to general population. Vertical transmission possible; unclear whether SARS-CoV-2 positive neonates infected in utero, intrapartum or postpartum.
Raschetti R 2020	Yes; RoB: Mayo EBP tool	Clarify transmission route, clinical features, outcomes of COVID-19 infections.	PubMed, The Cochrane Library, Web of Science, BioXRiv, MedXRiv from 1 December 2019 to 30 August 2020	74 studies (176 cases of neonatal SARS-CoV-2 infections)	70% infections due to environmental transmission, 30% vertical. 55% of infected neonates developed COVID-19; symptoms: fever (44%), gastrointestinal (36%), respiratory (52%), neurological (18%), lung imaging abnormal (64%). A lack of mother–neonate separation from birth associated with late SARS-CoV-2 infection (OR 4.94 (95% CI: 1.98–13.08), p = 0.0002; adjusted OR 6.6 (95% CI: 2.6–16), p < 0.0001), breastfeeding not associated (OR 0.35 (95% CI: 0.09–1.18), p = 0.10; adjusted OR 2.2 (95% CI: 0.7–6.5), p = 0.148).	Neonatal infections mainly occur postnatally through environmental exposure, 30% of infections may be vertical transmission. 50% infected neonates develop clinical COVID-19, febrile features similar to older patients, favorable outcomes. Mother-neonate rooming-in associated with higher incidence of SARS-CoV-2 infections after 72 h of life.

Rodrigues C 2020	No	Review knowledge on impact of COVID-19 on pregnancy, describe outcome of published cases of pregnant women diagnosed with COVID-19.	PubMed, Scopus, Web of Science, MedRxiv up to 26th June 2020	161 studies (3,985 pregnant women with COVID-19)	2,059 cases with pregnancy outcomes: 42 abortions, 21 stillbirths, 2,015 live births. Preterm birth: 23%. Maternal ICU: 6%, deaths: 28. Neonatal deaths: 10. From the 10/163 samples (amniotic fluid, placenta, cord blood) positive for SARS-CoV-2. 61 newborns positive for SARS-CoV-2. 4/92 breast milk samples positive.	Evidence suggests vertical transmission is possible, limited number of reported cases with intrapartum samples. Information, counseling and adequate monitoring essential to prevent and manage adverse effects of SARS-CoV-2 in pregnancy.
Romeo G 2020	No	Teview case reports of MERS-CoV, SARS-CoV, SARS-CoV-2, during pregnancy and summarise clinical presentation, course of illness, pregnancy and neonatal outcomes.	MEDLINE, ClinicalTrials.gov up to 23 April 2020.	46 studies. MERS-CoV: 8 (12 cases), SARS-CoV: 7 (17 cases), SARS-CoV-2: 31 (98 cases)	Case fatality for SARS-CoV-2 infection in pregnant women 1%. Mother-to-child transmission of MERS-CoV or SARS-CoV not observed. SARS-CoV-2 RNA detected in 7 newborns; IgM, 1 newborn. Amniotic fluid: 1 positive for SARS-CoV-2 RNA.	Limited case reports suggest possible vertical transmission of SARS-CoV-2. AR, RR, clinical significance, route of transmission unclear. Consistent data collection on timing of exposure, symptom onset, clinical presentation, course of illness, pregnancy and neonatal outcomes, and laboratory results among pregnant women with SARS-CoV-2 required.
Salem D 2020	No	Review evidence on outcomes of COVID-19 infections during pregnancy.	Medline, Google Scholar from January to August 2020	Not reported	COVID-19-positive pregnant women usually asymptomatic or mild-to-moderately symptomatic, similar to non-pregnant women. Most common clinical outcome: pneumonia. Unclear if SARS-CoV-2 infection increases risk of maternal, fetal, neonatal complications. Increased risks of complications in pregnant women with co-morbidities. Vertical transmission is possible.	Should monitor pregnant women before and after delivery, and their infants, during pandemic.
Sampieri CL 2020	Yes; RoB: GRADE	Review studies on SARS-CoV-2 in clinical samples of amniotic fluid, placenta or membranes, umbilical cord blood, and human milk, from women with clinical or confirmed diagnosis of COVID-19.	PubMed from 27 March 2020 to 21 May 2020	17 studies (143 clinical samples: 38 amniotic fluid; 34 placentas/ membranes; 39 umbilical cord blood; 32 milk)	9/143 samples positive for SARS-CoV-2 RNA (1 amniotic fluid before rupturing the membranes; 6 placenta/ membranes, possibility of contamination by maternal blood in 3; 2 human milk).	No studies demonstrate SARS-CoV-2 detection with viral isolation and evaluation of infective capacity of viral particles, in clinical samples of amniotic fluid, placenta/ membranes, umbilical cord blood, human milk, from women with confirmed or clinical diagnosis of COVID-19. Vertical transmission cannot be ruled out.

Sheth S 2020	No	Investigate outcomes in COVID-19 positive neonates and incidence of vertical transmission of the virus.	Pubmed, GoogleScholar from 15 November 2019 to 18 June 2020	39 studies (326 COVID-19 positive mothers with neonatal outcomes).	23 neonates COVID-19 positive. Male neonates affected more (79%). 3% neonates acquired infection through suspected vertical transmission. Strict perinatal infection prevention measures can reduce horizontal transmission. Neonates asymptomatic or mildly symptomatic. No neonatal deaths.	Prognosis of COVID-19 positive neonates is good with no mortality, mild symptoms. Minimal vertical transmission. Possibility of vertical transmission very low.
Shrestha R 2020	No	Summarise evidence on foetal and neonatal outcomes of pregnant women with confirmed COVID-19.	PubMed, MedRxiv from December 2019 to April 2020	21 studies (230 pregnant women)	Fever and cough the most common symptoms in pregnant women. Cesarean section: 162 (70.4%); 68 (29.6%) preterm deliveries. 8/161 newborns positive for SARS-CoV-2. coronavirus infection. 2 miscarriages, 2 still births, 1 neonatal death.	Outcome of pregnancy with COVID-19 in late trimester appears favourable. Preterm delivery and cesarean section higher among infected pregnant women. No conclusive evidence of vertical transmission.
Thomas P 2020	Yes; RoB: CLARITY, McMaster; Joanna Briggs Institute	Summarize evidence on vertical transmission of COVID-19 infection in the third trimester and its effects on neonate.	OVID MEDLINE, EMBASE, Cochrane Central Register of Controlled Trial (CENTRAL) from January 2020 to May 2020	18 studies (157 mothers, 160 neonates).	5/81 (6%) neonates positive for SARS-CoV-2. Earliest test was at 16 h after birth; 1 neonate initially negative at birth, positive when re-tested, suggesting SARS-CoV-2 likely hospital-acquired rather than vertically transmitted. 13 (8%) neonates had complications or symptoms.	Rapid descriptive review; early clinical evidence suggests vertical transmission of SARS-CoV-2 from mother to neonate/newborn did not occur.
Tripella G 2020	Yes; RoB: Joanna Briggs Institute	Review the impact of COVID-19 on pregnant women and neonates.	MEDLINE, EMBASE from 1 December 2019 to 18 April 2020	37 studies (275 pregnant women with COVID-19, 248 neonates)	Most women mild to moderate symptoms; ICU: 10; deaths: 1. 2 stillbirths; prematurity 28%. 16 neonates positive for SARS-CoV-2, 9 born from mothers infected during pregnancy. All infected neonates recovered. RT-PCR for SARS-CoV-2 negative on amniotic fluid, vaginal/cervical fluids, placenta tissue, breast milk samples.	SARS-CoV-2 infection in pregnant women associated with mild or moderate disease in most cases, low morbidity and mortality rate. Outcomes of neonates mainly favorable, although neonates at risk should be closely monitored.

Turan O 2020	Yes; RoB: NIH tools	Summarize clinical characteristics and outcomes among pregnant women hospitalized with COVID-19.	PubMed, Ovid Medline, Web of Science, China Academic Literature Database up to 29 May 2020.	63 studies (637 women with confirmed SARS-CoV-2)	Majority (76.5%) had mild disease. Maternal fatality, 1.6%; stillbirth, 1.4%; neonatal fatality, 1.0%. Older age, obesity, diabetes mellitus, raised serum D-dimer and IL-6 predictive of poor outcomes. Pre-term births: 33.7% of which 50% were iatrogenic in women with mild COVID-19 and no complications. Most had cesarean without clear indication. 8 (2.0%) neonates positive NP test, chest infection within 48h.	Advanced gestation, maternal age, obesity, diabetes mellitus, combination of elevated D-dimer and IL-6 predictive of poor pregnancy outcomes in COVID-19. Tate of iatrogenic preterm birth and cesarean delivery high; vertical transmission possible but not proven.
Walker KF 2020	No	Estimate the risk of neonate becoming infected with SARS-CoV-2 by mode of delivery, type of infant feeding and mother-infant interaction.	MEDLINE, Embase, Maternity and Infant Care Database from September 2019 to June 2020	49 studies (655 women, 666 neonates)	28/666 (4%) neonates tested positive postnatally. Babies born vaginally, 8/292 (2.7%) tested positive, compared with 20/374 (5.3%) born by Caesarean. Information on feeding and baby separation lacking. 7/148 (4.7%) breastfed babies tested positive compared with 3/56 (5.3%) for formula fed. 4/107 (3.7%) babies nursed with mother tested positive, compared with 6/46 (13%) who were isolated.	Neonatal COVID-19 infection is uncommon, rarely symptomatic, and the rate of infection is no greater when the baby is born vaginally, breastfed or remains with the mother.
Yoon SH 2020	No	Evaluate clinical manifestations and outcomes of neonates born to women who had COVID-19 during pregnancy.	Search databases PubMed, Embase up to 15 April 2020	28 studies (223 pregnant women, 201 infants)	4 newborns born to mothers with COVID-19 tested SARS-CoV-2 positive within 48h. Breast milk, placenta, amniotic fluids, cord blood, maternal vaginal secretions all negative. Fetal death: 2; premature birth 48/185. low birth weight (< 2,500 g): 8.3%; small for gestational age: 15.6%. Birth asphyxia: 1/8%; respiratory distress syndrome 6.4%.	Evidence suggests COVID-19 during pregnancy rarely affects fetal and neonatal mortality, but can be associated with adverse neonatal morbidities. Vertical transmission has not been observed in most cases.

Table 4. Findings of studies reporting placenta analysis (n=25)

Study ID	Study design (n)	Vertical transmission	Placenta SARS-CoV-2 test result (RT-PCR)	Histopathology and Biopsy	Whole genome sequencing
Alamar I 2020	Case report (n=1)	1/1	NR	Placenta ISH positive for SARS-CoV-2. Placental histopathology showed no inflammation	Not attempted
Birindwa EK 2020	Case report (n=1)	1/1	NR	Inflammation; thrombotic vasculopathy in the placenta	Not attempted
Fenizia B 2020	Cohort (n=31)	2/31	Positive: placental tissue in 2 cases	Generalised immune activation profile in placentae from infected compared to uninfected mothers.	Not attempted
Ferraiolo A 2020	Case report (n=1)	0/1	Placental swabs positive	Some placental inflammation on biopsy	Not attempted
Grimminck K 2020	Case report (n=1)	0/1	Placental sample negative	Not attempted	Not attempted
Kulkarni R 2020	Case report (n=1)	1/1	Placental sample positive	Not attempted	Not attempted
Hsu AL 2020	Case report (n=1)	0/1	NR	IHC with SARS-CoV-2 nucleocapsid-specific monoclonal antibody detected SARS-CoV-2 antigens throughout the placenta, under the umbilical cord, and at the central and peripheral placenta disc in chorionic villi endothelial cells, and rarely in CK7-expressing trophoblasts.	Not attempted
Liu W 2020	Retrospective case analysis (n=48)	0/48	Placental swabs negative	Not attempted	Not attempted

Lv Y 2020	Case report (n=1)	0/1	Placental swabs negative	Not attempted	Not attempted
Masmejan S 2020	Retrospective case series (n=13)	0/13	Placental swabs negative	Not attempted	Not attempted
Ogamba I 2020	Retrospective cohort (n=25)	0/20	NR	Placental pathology did not detect COVID-19 pathology	Not attempted
Oncel MY 2020	Cohort (n=125)	4/120	Placental tissue negative (n=5)	Not attempted	Not attempted
Palalioglu RM 2020	Case report (n=1)	0/1	Placental tissue negative	Not attempted	Not attempted
Pulinx B 2020	Case report (n=1 mother, 2 neonates)	2/2 (placental and amniotic fluid)	Placental swabs positive	Chronic intervillitis and extensive intervillous fibrin depositions with ischemic necrosis of the surrounding villi. Viral localization in the placental syncytiotrophoblast cells confirmed by IHC.	Not attempted
Rebello CM 2020	Case report (n=1)	1/1	Umbilical cord blood sample positive (placenta tissue not tested)	Not attempted	Not attempted
Shende P 2020	Case report (n=1)	1/1 in amniotic fluid, also in placental cell supernatant	placental cell supernatant positive	Immunofluorescence with mAb anti-spike protein S1 and S2 on placental sections shows diffuse viral presence in cytoplasm of syncytiotrophoblasts and some cytotrophoblasts cells	Not attempted
Sisman J 2020	Case report (n=1)	1/1	NR	Placenta histopathology indicated positive for SARS-CoV-2	Not attempted

Smithgall MC 2020	Cohort (n=76 placentas)	0/51	NR	No histomorphologic changes in placenta; no viral detection by ISH and IHC. Evidence of maternal–fetal vascular malperfusion was identified, with placentas from SARS-CoV-2-positive women being significantly more likely to show villous agglutination (P = 0.003) and subchorionic thrombi (P = 0.026) than placentas from SARS-CoV-2-negative women; however no evidence for viral cause.	Not attempted
Stonoga E 2020	Case report (n=1)	0/1	Placenta & umbilical cord blood tested positive by PCR.	Not attempted	Not attempted
Tang J 2020	Case report (n=2)	0/2	NR	Placental tissue showed no microscopic abnormality	Not attempted
Vivanti AJ 2020	Case report (n=1)	1/1	placenta was positive for both SARS-CoV-2 genes (E and S); Log viral load: 11.5	Inflammation of placenta; high IHC positive Ab staining for SARS-CoV-2 cytoplasm of peri-villous trophoblastic cells.	Not attempted
Von Kohorn I 2020	Case report (n=1)	1/1	NR	Placenta histology - no direct evidence of virus	Not attempted
Zaigham M 2020	Case report (n=1)	1/1	NR	high viral load in placenta with massive perivillous fibrin deposition, acute intervillitis in areas with strong positivity for SARS-CoV-2 and chorangiomas in the areas less affected by infection and inflammation	Whole genome sequencing of isolates from the mother & placenta: Presence of

					single variant of the virus
Zhang P 2020	Case series (n= 364 placenta samples)	2/53	NR	SARS-CoV-2 detected by ISH in placenta of positive cases (2/ 53 placental samples from PCR positive mothers). No histopathological features	Not attempted
Zheng T 2020	Case report (n=2)	0/2	NR	Both placentas were examined and found to be normal	Not attempted

Table 5. Ct and RNA concentration results from included studies.

Study	PCR cycle methods	Ct counts	RNA concentration / viral load	Live culture attempted (positivity rate)
Alamar I 2020	NR	NR	NR	Not attempted
Anand P 2020	First screen using E gene, then if +ve, two genes, RdRp & ORF-1b-nsp14b. The detection limit was 35 for E, RDRp & ORF.	35? Unclear	NR	Not attempted
Ayed A 2020	NR	NR	NR	Not attempted
Bachani S 2020	E, RDRp & ORF-1bns14b. The threshold value for infectivity \leq 35 cycles for E, RDRp & ORF genes.	35	Lowest Cts observed for neonate 3: Ct = 15.82, DOL4; & neonate 4: Ct = 15.33, DOL1.	Not attempted
Bandyopadhyay T 2020	NR	NR	NR	Not attempted
Barbero P 2020	NR	NR	NR	Not attempted

Birindwa EK 2020	NR	NR	NR	Not attempted
Bordbar A 2020	NR	NR	NR	Not attempted
Chaudhary S 2020	NR	NR	NR	Not attempted
Cojocar L 2020	NR	NR	NR	Not attempted
Demirjian A 2020	NR	maternal NP DOL1 Ct = 34.9; DOL3 = 39.4	NR	Not attempted
Farsi Z 2020	NR	NR	NR	Not attempted
Fenzia B 2020	RdRP, N & E genes simultaneously amplified & tested. Ct-value <40 defined as positive test result according to manufacturer's instruction.	<40 for two newborns (remainder 40+ judged negative)	NR	Not attempted
Ferraiolo A 2020	3 target genes: RdRP, E & N, & a process control.	NR	NR	Not attempted
Gale C 2020	NR	NR	NR	Not attempted
Gao W 2020	NR	NR	NR	Not attempted

Gao X 2020	RT-PCR assays for SARS-CoV-2 ORF1ab/N genes	NR	NR	Not attempted
Grimminck K 2020	NR	NR	NR	Not attempted
He Z 2020	NR (state following WHO guidelines)	NR	NR	Not attempted
Hinojosa-Velasc o A 2020	real-time RT-PCR, ORF1ab & N gene targets.	23	reported as "high" based on Ct of 23	Not attempted
Hsu AL 2020	NR	NR	NR	Not attempted
Hu X 2020	NR	NR	NR	Not attempted
Khan MA 2020	NR	NR	NR	Not attempted
Kulkarni R 2020	NR	NR	NR	Not attempted
Liu W 2020	NR	NR	NR	Not attempted
Luo Q 2020	ORF1ab gene & N gene	NR	NR	Not attempted

Lv Y 2020	NR	NR	NR	Not attempted
Maraschini A 2020	NR	NR	NR	Not attempted
Marin Gabriel M 2020	NR	NR	NR	Not attempted
Masmejan S 2020	NR	NR	NR	Not attempted
Mohakud NK 2020	NR	NR	NR	Not attempted
Moreno SC 2020	NR (state methods according to WHO guidelines)	NR	NR	Not attempted
Nayak AH 2020	NR	NR	NR	Not attempted
Ogamba I 2020	NR	NR	NR	Not attempted
Olivini N 2020	NR	NR	NR	Not attempted
Oncel MY 2020	NR (state methods according to WHO guidelines)	NR	NR	Not attempted

Pace RM 2020	NR	NR	NR	Not attempted
Palalioglu RM 2020	NR	NR	NR	Not attempted
Parsa Y 2020	NR	NR	NR	Not attempted
Pereira A 2020	NR	NR	NR	Not attempted
Pessoa FS 2020	NR	NR	NR	Not attempted
Pissarra S 2020	NR	NR	NR	Not attempted
Popofsky S 2020	NR	NR	NR	Not attempted
Pulinx B 2020	rRT-PCR based on the CDC oligonucleotide primers & probes for detection of N gene	at birth placental tissue samples Ct = 33 & Ct = 30; amniotic fluid Ct = 23; maternal blood Ct = 35)	NR	Not attempted
Rebello CM 2020	NR	NR	NR	Not attempted

Rivera-Hernandez P 2020	NR	NR	NR	Not attempted
Rubio Lorente AMR 2020	NR	NR	NR	Not attempted
Sajjan GR 2020	NR	NR	NR	Not attempted
Schwartz DA 2020	NR	NR	NR	Not attempted
Shende P 2020	RT-PCR done as per the ICMR/WHO Guidelines for RdRP & E genes	Ct for E gene 26.3, Ct for RdRp gene 25.4 (amniotic fluid)	NR	Not attempted
Singh MV 2020	NR	NR	NR	Not attempted
Sisman J 2020	NR	NR	NR	Not attempted
Smithgall MC 2020	NR	NR	NR	Not attempted
Stonoga E 2020	N and ORF1ab; Ct value considered +ve if both viral genes are <38.	umbilical cord blood N gene Ct = 30.3, ORF1ab Ct = 31.9; placenta N gene Ct = 25.5, ORF1ab Ct = 24.5	NR	Not attempted

Tang F 2020	two sets of primers to respectively amplify ORF1ab & N gene	NR	NR	Not attempted
Tang J 2020	NR	NR	NR	Not attempted
Vendola N 2020	NR	NR	NR	Not attempted
Vinuela MC 2020	RT-PCR detection of N & ORF 1a1b genes; Ct for N gene recorded	Ranged from 36 to 41; median 40.	NR	Not attempted
Vivanti AJ 2020	RT-PCR detection of E & S gene of SARS-CoV-2; full details of PCR methods provided	40	Viral load (log) mother: NP: 4.22; vaginal: 0.63; placenta: 11.15; amniotic: 2.09; blood: 4.87. Neonate: blood: 1.15; NP (DOL1): 2.21; rectal: 4.71	Not attempted
Von Kohorn I 2020	Two RT-PCRs: Test1: E & N2 gene; Test 2: ORF1 & E gene	For positive tests DOL2-7: range from 17.7 (DOL7) to 44.1 (DOL2)	NR	Not attempted
Vouga M 2020	NR	NR	NR	Not attempted
Woodworth KR 2020	NR	NR	NR	Not attempted

Yang R 2020	NR	NR	NR	Not attempted
Zaigham M 2020	RT-PCR for envelope gene, details of PCR methods provided	Maternal: NP Ct = 26.7; blood Ct = 31.7. Infant: NP DOL 2 to 16: Ct range 18.1 (DOL5) to 36.1(DOL12). Placenta: Day of delivery Ct = 13.6	NR	Not attempted
Zhang L 2020	NR	NR	NR	Not attempted
Zhang P 2020	NR	NR	NR	Not attempted
Zheng T 2020	NR	NR	NR	Not attempted