TITLE PAGE

Vaccination willingness for COVID-19 among health care workers in Switzerland

Kathrin Zürcher ^{1*}, Catrina Mugglin ^{1 2 3*}, Matthias Egger ¹, Sandro Müller ⁴, Michael Fluri ⁵, Laurence Bolick ⁶, Rein Jan Piso ⁶, Matthias Hoffmann ^{6*}, Lukas Fenner ^{1 2 * §}

1 Institute of Social and Preventive Medicine, University of Bern, Switzerland

2 Kantonsärztlicher Dienst, Gesundheitsamt, Kanton Solothurn, Switzerland

3 Department of Infectious Diseases, Inselspital, Bern University Hospital, University of Bern, Switzerland

4 Amt für soziale Sicherheit, Kanton Solothurn, Switzerland

5 Hausarztpraxis Weissenstein, Langendorf, Switzerland

6 Department of Internal Medicine, Infectious Diseases and Hospital Epidemiology, Cantonal Hospital Olten, Switzerland

*contributed equally

Correspondence:

Prof. Lukas Fenner, MD MSc
Institute of Social and Preventive Medicine, University of Bern (ISPM)
Mittelstrasse 43, 3012 Bern, Switzerland
Phone: +41 31 631 35 18
E-Mail: lukas.fenner@ispm.unibe.ch

Key words: COVID-19, Vaccine, vaccine hesitancy, health care workers

Running head: Vaccine willingness in HCWs

Word count, inserts and additional files:

Abstract 243 words, main text 2,744 words

Inserts: 2 tables, 2 figures, 42 references

Supplements: 3 tables, 1 figures

ABSTRACT

Aims of the study: Vaccination is regarded as the most promising response to the COVID-19 pandemic. We assessed opinions towards COVID-19 vaccination, willingness to be vaccinated, and reasons for vaccination hesitancy among health care workers (HCWs).

Methods: We conducted a cross-sectional, web-based survey among 3,793 HCWs in December 2020 in the Canton of Solothurn, Switzerland, before the start of the national COVID-19 vaccination campaign.

Results: Median age was 43 years (interquartile range [IQR] 31-53), 2,841 were female (74.9%). 1,511 HCWs (39.8%) reported willingness to accept vaccination, while 1,114 (29.4%) were unsure, and 1,168 (30.8%) would decline vaccination. Among medical doctors, 76.1% were willing, while only 27.8% of nurses expressed willingness. Among 1,168 HCWs who would decline vaccination, 1,073 (91.9%) expressed concerns about vaccine safety and side effects. The willingness of HCWs to be vaccinated was associated with older age (adjusted odds ratio [aOR] 1.97, 95%CI 1.71-2.27) and having been vaccinated for influenza this year (aOR 2.70, 95%CI 2.20-3.31). HCWs who reported a lack of confidence in government were less likely to be willing to be vaccinated (aOR 0.58, 95%CI 0.40-0.84), and women were less willing to be vaccinated than men (OR 0.33 (0.28-0.38)).

Conclusion: Less than half of HCWs reported willingness to be vaccinated before the campaign start, but proportions varied greatly depending on the profession and workplace. Strategies with clear and objective messages that particularly address the concerns of HCWs are needed if their willingness to be vaccinated is to be increased.

INTRODUCTION

The worldwide spread of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infections and coronavirus disease 2019 (COVID-19) is a major public health threat (1). Globally, SARS-CoV-2 infection has been confirmed in over 131,837,512 people and as of 7 April, 2021 2,862,664 have died from COVID-19 (2). In Switzerland, 612,575 COVID-19 cases have been confirmed and more than 9,772 people have died from COVID-19 (3).

The pandemic has stretched the health care system in Switzerland to its limits and burdened the economy with temporary closures of restaurants and stores and large public and private venues (4). In the absence of effective treatments and a safe and effective vaccine, nonpharmaceutical interventions (NPIs) were implemented to mitigate the pandemic. Measures taken include using personal protective equipment (PPE) such as face masks in public spaces, keeping distance between individuals, and rigorous hand hygiene. Many pharmaceutical companies and research laboratories have been working on vaccines (5-7). Effective vaccination is key to controlling the COVID-19 pandemic, but global vaccine distribution is challenging (8). By the end of 2020, several vaccines had demonstrated efficacy in phase 3 trials (7), and by the beginning of January 2021, two vaccines had been approved by the Swiss Agency for Therapeutic Products (Swissmedic) (9, 10).

The delivery of the COVID-19 vaccines started on 4 January 2021, making it essential to identify and address widespread vaccine uptake barriers. Scepticism about these new vaccines against COVID-19 presents one such challenge to vaccine uptake. Health care workers (HCWs), who face an increased risk of infection with SARS-CoV-2 and can transmit the virus among themselves and to highly

vulnerable patients (11-15) are an important target group for vaccination. Previous studies have shown that vaccine uptake for vaccine-preventable diseases such as influenza is low among HCWs (16, 17). Given the central role they play in treating COVID-19 and administering vaccinations, HCWs are uniquely positioned to influence vaccine uptake. Therefore, understanding the willingness of HCWs to be vaccinated against COVID-19, mainly if they are themselves hesitant, will be important in promoting vaccine uptake in the population.

We, therefore, assessed the willingness of healthcare workers in the Canton of Solothurn, Switzerland, to be vaccinated against influenza and COVID-19, and inquired about reasons for vaccine hesitancy among them.

METHODS

Study design

We conducted a cross-sectional web-based survey among HCWs in the Canton of Solothurn, Switzerland. We included adults aged 16 or older who work in the health care system in hospitals, medical practices, retirement and nursing homes, home care, pharmacies, and long-term care facilities.

Data collection

We developed and pilot tested a standardised questionnaire based on the study of Larson et al. (18) and the Vaccine Confidence Project at the London School of Hygiene and Tropical Medicine (www.vaccineconfidence.org). The electronic questionnaire collected information of three types: (i) demographic details such as sex, age, and profession; (ii) intention to be vaccinated against COVID-19 and reasons for being vaccinated/not being vaccinated or for being unsure, confidence in government, recommendation of the employer, and additional information needed to take a vaccination decision; (iii) history of influenza vaccination for influenza season 2020/21. We collected the data in mid-December 2020 before the first approval of a COVID-19 vaccine and associated campaigning using a web-based tool (www.findmind.ch).

All employees of the cantonal hospital in the Canton of Solothurn were invited to participate. The survey was also sent to the cantonal professional associations of physicians, nursing homes, long-term care facilities, and residential care, which invited their members to participate.

Definitions

The Swiss Federal Office of Public Health (FPOH) defined the following COVID-19 risk groups: persons over the age of 50; those with comorbidities including hypertension, chronic respiratory diseases, diabetes, being immunocompromised, cardiovascular disease, cancer, or obesity (BMI >30); persons who live in a nursing home or long-term care facility; and those who are pregnant.

The survey participants fell into three groups: persons willing to be vaccinated, those who were not willing to be vaccinated, and those who were hesitant or unsure about being vaccinated. Among those willing to be vaccinated against COVID-19, the reasons for deciding to do so could be characterised as self-protection, individual vaccination as a contribution to pandemic control, and membership in a risk group for severe COVID-19 disease. Reasons for unwillingness to be vaccinated included opposition to vaccines in general, the perception that COVID-19 is harmless or that PPE is sufficient, concerns about the effectiveness of the vaccine, its safety and side effects, bad experiences with previous vaccinations, fear of needles, and other reasons. People who were unsure about being vaccinated against COVID-19 were hesitant due to perceived inconsistent information, doubt about vaccine effectiveness, doubt about vaccine safety and fear of side effects, doubt about individual vaccination for pandemic control, and uncertainty among colleagues, and lack of information provided by the employer. Supplementary Table S1 provides further details.

Statistical analyses

We used descriptive statistics to characterise the study population by profession and intention to vaccinate against COVID-19. Differences between groups were assessed using chi-square, t-test, or Wilcoxon rank-sum tests as appropriate. We

calculated the proportions of intention to be vaccinated against COVID-19 with the corresponding 95% confidence intervals (95% CI).

We examined factors associated with the HCW's intention to be vaccinated against COVID-19 in univariate and multivariate logistic regression. Logistic models were adjusted for age group and profession, confidence in government reports, the employer's recommendation, and influenza vaccine uptake 2020/21. We grouped uncertainty about vaccination and having no intention of being vaccinated against COVID-19. We also performed a sensitivity analysis that combined those who were willing to be vaccinated and those who were not sure about vaccination. Finally, we compared the vaccine refusers and those who were unsure about COVID-19 vaccination. All analyses were performed in Stata (version 15.1, College Station, TX, USA).

Ethics statement

Data collection was anonymous. No ethical approval was needed, in line with the Swiss Human Research Act.

RESULTS

A total of 4,244 HCWs participated in the survey. Response rates were as follows: ~64% (2,679/~4,200) among hospital staff, ~38% (338/~900) among medical practice staff, ~11% (440/~4,000) among nursing home staff, and ~29% (470/~1,600) among long term care facility staff.

We excluded participants who did not fully complete the survey (n=421) and those for whom data on sex, age, or profession were missing (n=12). We further excluded participants younger than 16 years (n=18). The analyses thus included 3,793 HCWs ([Figure S1](#)).

Characteristics of the HCWs

The median age of HCW participants was 43 years (interquartile range [IQR] 31-53), 2,841 were female (74.9%). Among all HCWs, 2,445 (64.5%) worked in hospitals, 398 (10.5%) worked in long-term care facilities, 373 (9.8%) in retirement and nursing homes, 311 (8.2%) in medical practices, 232 (6.1%) in residential care, and 34 (0.9%) at pharmacies ([Table 1](#) and [Table S2](#)). Asked about COVID-19 vaccination, 1,511 HCWs (39.8%) reported they planned to take the COVID-19 vaccine, 1,114 (29.4%) said they were unsure, and 1,168 (30.8%) said they would not take the vaccination. Willingness to vaccinate was highest among medical doctors, among whom 467 of 617 (76.1%) were willing to be vaccinated, and it was lowest among nurses at 470 of 1690 (27.8%) ([Figure 1 A](#)). [Figure 1 B](#) shows the willingness to be vaccinated by setting.

The self-reported uptake for seasonal influenza 2020/2021 vaccination was 38.1% (1,446 of 3,793) while 118 (3.1%) were unsure whether they had been vaccinated, and 2,229 (58.8%) reported they were not vaccinated. The influenza

vaccination rate was highest among medical doctors (75.6%) and pharmacists (58.8%). Among the other professions, the vaccination rate ranged from 11.6% to 44.4% (Table 1).

Determinants of willingness to be vaccinated

The willingness among HCWs to be vaccinated against COVID-19 was lower among females than male participants (OR 0.33, 95%CI 0.28-0.38) and increased with age. Willingness was highest among the age group ≥ 60 years (OR 4.56, 95%CI 3.50-5.94) compared to the age group <30 years. HCWs reporting having been vaccinated against seasonal influenza 2020/21 were more willing to be vaccinated against COVID-19 (OR 6.30, 95%CI 5.45-7.23) compared to the unvaccinated group. Confidence in government reports and employers' vaccination recommendation were associated with willingness to be immunised against COVID-19: OR 4.12 (95%CI 3.37-5.03) and OR 14.19 (95%CI 11.53-17.47), respectively. In contrast, lack of such confidence was negatively associated with willingness to get vaccinated compared to the group with no opinion (Table 2). The willingness to be vaccinated against COVID-19 was higher among pharmacists (OR 6.22, 95%CI 2.96-13.13) and medical doctors (OR 8.24, 95%CI 6.66-12.21) compared to nurses.

In the multivariate analysis, willingness to be vaccinated was positively associated with confidence in government reports on the COVID-19 pandemic (aOR 1.59 95%CI 1.23-2.06) and negatively with lack of confidence in government (aOR 12.85 95%CI 10.11-16.33). Similarly, confidence in employer recommendations was positively associated, whereas lack of confidence was negatively associated. The analysis confirmed that HCWs who reported vaccination against seasonal influenza 2020/21 were more likely to be willing to be vaccinated against COVID-19 than those who were not vaccinated (aOR 2.70 95%CI 2.20-3.31) (Table 2).

In the sensitivity analysis grouping the willing and unsure together (rather than the refusers and unsure), results were similar to the primary analysis ([Table S3](#)).

Reasons for vaccine hesitancy

The main reasons for willingness to be vaccinated were personal protection, controlling the pandemic, and belonging to a risk group ([Figure 2A](#), [Table 1](#)). These reasons were different by profession ($p < 0.05$). The most frequent reasons for vaccine hesitancy among HCWs included concerns about vaccine safety and side effects (1073/1,168, 91.9%), the perception that personal protective equipment is sufficient (328/1,168, 28.1%), and that COVID-19 is harmless (231/1,168, 19.8% [Figure 2B](#) and [Table 1](#)). All reasons for vaccine hesitancy were similar in the different HCW professions, except for vaccine safety and side effects, which was mentioned more frequently as a reason for vaccine hesitancy among nurses ($p = 0.01$).

Among 1,114 HCWs who were unsure about a vaccination decision, 1,055 (94.7%) wanted more information on vaccine safety and side effects, 778 (69.6%) awaited reports from already vaccinated people, and 404 (36.3%) wanted an opinion from experts. The main reasons can be found in [Figure 2C](#), [Table 1](#). The reasons for being unsure were similar across the HCW professions, but pharmacists wanted more frequently information on the vaccine than medical doctors (100% vs. 88%, $p = 0.01$).

DISCUSSION

Less than half of participating HCWs reported willingness to be vaccinated against COVID-19. The most frequently given reason for vaccine hesitancy was concern about vaccine safety and side effects. Nurses were less likely to be willing to be vaccinated against COVID-19 than were medical doctors and pharmacists.

Vaccines are an effective control measure to reduce the burden of infectious disease. Poliomyelitis was eradicated, and we are close to eliminating measles, mumps, and rubella with vaccines (19). Vaccination will play an important role in the control of the COVID-19 pandemic. Even transient herd immunity in Switzerland will require an estimated 60% of the population to achieve immunity to SARS-CoV-2 either through infection and recovery or vaccination, not considering the potential impacts of SARS-CoV-2 variants of concern. This proportion varies depending on model assumptions (20).

Willingness to get vaccinated is central to achieving herd immunity. Several cross-sectional studies have assessed the willingness to get vaccinated. In seven countries, a European general population survey found that the willingness to be vaccinated against SARS-CoV-2 ranged from 62% to 80% (21). Similar results were found in the United Kingdom and Ireland, with 69% and 65% acceptance, respectively (22). In the USA, 37% of HCWs reported to be willing to get vaccinated against COVID-19 (23). These results are in line with our results (40%) in the Canton of Solothurn in Switzerland. These numbers cause concern because HCWs play an important role in vaccine uptake among the general population. A review on vaccine hesitancy has shown that vaccinated HCWs are more likely to recommend vaccination than unvaccinated HCWs (24). Furthermore, a vaccination

recommendation given by an HCW is frequently cited as the reason for vaccination acceptance (25). In contrast, the lack of such a recommendation was the most common reason for not being vaccinated (24, 26).

Some HCWs have expressed reservations about the safety and side effects of COVID-19 vaccines. One contributor to such reservations is the perception of undue haste in the development of COVID-19 vaccines compared to previous vaccines. In the twentieth century, it took a decade or longer to get vaccines to the market, for example, in the case of poliomyelitis (27). Many factors contributed to the comparatively short time it took for COVID-19 vaccines to be granted emergency use authorisations in many countries, with definitive licensing granted on a rolling basis. Researchers have been developing mRNA and viral vector-based vaccines for other diseases for more than a decade (28). As the pandemic spread, public willingness and even demand led to rapid enrolment in phase I through to phase III clinical trials with historically unprecedented speed. Simultaneously, the rapid spread of SARS-CoV-2 allowed endpoint-driven phase III trials to demonstrate vaccine efficacy sooner than was expected at their outset.

We found that confidence in the governmental authorities is associated with willingness to get vaccinated against COVID-19. During a health crisis or a pandemic, trust in the government and risk perception play a key role in vaccine acceptance (29, 30). A French survey among general practitioners showed they were more likely to recommend vaccines to patients when they trusted official sources of information (31). A global survey also reported an association between vaccine acceptance and participants' trust in government (32). High levels of trust in government was associated with willingness to follow governmental recommendations on preventive behaviors to contain swine flu (33) or compliance

with social distancing measures during the Ebola outbreak (34). In contrast, mistrust and misinformation reduced compliance with social and behavioural measures (35). A cross-sectional study during the COVID-19 pandemic among UK residents showed that residents who trusted the government to control the pandemic were more likely to follow government recommendations during the lockdown (36).

Strategies to increase vaccine coverage among HCWs should draw upon guidance from authorities or persons conceived as such by a target audience (in this case HCWs) and specific information about vaccine safety and efficacy. Thus, it is not surprising that many undecided HCWs want to have reports from fellow HCWs who have been vaccinated before they make up their minds. Peers provide important information that influences decision-making, but this source of information is often overlooked. HCWs may be the most effective promoters of vaccination for their fellow workers.

We observed that HCWs who were vaccinated against seasonal influenza are also more likely to willing to be vaccinated against COVID-19. Both diseases are contagious respiratory diseases caused by viruses, and they share some of the same symptoms (fever, cough) and approaches for prevention (hand hygiene, physical distancing, and masks). However, there are important differences. Superspreader events are more common for COVID-19 (37), and mortality rates are higher for COVID-19 than for influenza (38, 39). A recent study showed that over time, the uptake of the seasonal influenza vaccine in Switzerland had dropped overall and among older persons (≥ 65 years) and people with chronic disease (40). This reduced coverage might be explained by variable and lower seasonal influenza vaccine effectiveness, which can range from 30-60% (41, 42).

The limitations of this study include different response rates among the HCW professions and potential over-representation of hospital-based HCWs. We conducted a web-based cross-sectional online survey in which participation was voluntary but highly supported by the hospital-based personnel. Hospital-based HCWs may be over-represented. HCWs interested in the topic are more likely to respond to the questionnaire, and respondents with biases may select themselves into the sample. This might lead to an overestimation of vaccine hesitancy. Compared to other online surveys, this survey was distributed among specific groups of HCWs to cover this heterogeneous population better. Additionally, attitudes towards vaccination against COVID-19 and willingness to get vaccinated change over time. With more people being vaccinated, acceptance increases, and vaccine hesitancy might decrease.

This study's strength is the inclusion of diverse health care institutions and HCWs who play quite different roles within them. We included HCWs working with varying populations of risk ranging from nursing home residents, hospitalised patients to people living in long-term care facilities. Furthermore, the survey covered the three largest health care providers in the canton of Solothurn.

CONCLUSIONS

At this early stage of the COVID-19 vaccine campaign, the overall willingness to get vaccinated among HCWs is low. Balanced and transparent information on vaccine efficacy, safety and side effects should be provided to HCWs and dialogue on vaccine hesitancy initiated.

AUTHORS CONTRIBUTIONS

Conception and design LF, MH. Study coordination CM, MH, SM. Development of the questionnaire: CM, LF, MH, KZ. Data analysis: KZ, CM, LF, MH wrote the first draft of the paper and revised it based on comments from all authors. All authors reviewed and approved the final version of the manuscript.

ACKNOWLEDGEMENTS

We would like to thank all HCWs who were willing to participate. Further, we would like to thank all the institutions for distributing the survey to their employees. We thank Christopher Ritter for editorial assistance.

SOURCES OF SUPPORT

There was no specific funding for this project. KZ was supported by grant U01AI069924 from the U.S. National Institutes of Health's National Institute of Allergy and Infectious Diseases. ME was supported by special project funding (grant 189498) from the Swiss National Science Foundation.

COMPETING INTERESTS

All authors declare that they have no conflicts of interest.

REFERENCES

1. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *The New England journal of medicine*. 2020;382(8):727-33.
2. World Health Organization. Coronavirus Disease (COVID-19) Dashboard Geneva: World Health Organization; 2021 [cited 2021 13 Reburary]. Available from: <https://covid19.who.int/>.
3. Federal Office of Public Health (FOPH). Coronavirus: Situation Schweiz Bern Federal Office of Public Health; 2021 [Available from: <https://www.bag.admin.ch/bag/de/home/krankheiten/ausbrueche-epidemien-pandemien/aktuelle-ausbrueche-epidemien/novel-cov/situation-schweiz-und-international.html>].
4. Nicola M, Alsafi Z, Sohrabi C, Kerwan A, Al-Jabir A, Iosifidis C, et al. The socio-economic implications of the coronavirus pandemic (COVID-19): A review. *International journal of surgery (London, England)*. 2020;78:185-93.
5. Kim JH, Marks F, Clemens JD. Looking beyond COVID-19 vaccine phase 3 trials. *Nature Medicine*. 2021.
6. Parker EPK, Shrotri M, Kampmann B. Keeping track of the SARS-CoV-2 vaccine pipeline. *Nature reviews Immunology*. 2020;20(11):650.
7. London School of Hygiene & Tropical Medicine. COVID-19 vaccine tracker London: London School of Hygiene & Tropical Medicine; 2020 [Available from: https://vac-lshtm.shinyapps.io/ncov_vaccine_landscape/].
8. So AD, Woo J. Reserving coronavirus disease 2019 vaccines for global access: cross sectional analysis. *BMJ*. 2020;371:m4750.
9. Swissmedic. Swissmedic grants authorisation for the first COVID-19 vaccine in Switzerland Bern: Swissmedic; 2020 [Available from: https://www.swissmedic.ch/swissmedic/en/home/news/coronavirus-covid-19/covid-19-impfstoff_erstzulassung.html].
10. Swissmedic. Swissmedic grants authorisation for the COVID-19 vaccine from Moderna: Second COVID-19 vaccine authorised in Switzerland Bern: Swissmedic; 2021 [Available from: <https://www.swissmedic.ch/swissmedic/en/home/news/coronavirus-covid-19/zulassung-covid-19-impfstoff-moderna.html>].
11. Remuzzi A, Remuzzi G. COVID-19 and Italy: what next? *The Lancet*. 2020.
12. International Council of Nurses. International Council of Nurses High proportion of healthcare workers with COVID-19 in Italy is a stark warning to the world: protecting nurses and their colleagues must be the number one priority. 20 March, 2020. Geneva: International Council of Nurses, ; 2020 [Available from: <https://www.icn.ch/news/high-proportion-healthcare-workers-covid-19-italy-stark-warning-world-protecting-nurses-and>].
13. Garcia-Basteiro AL, Moncunill G, Tortajada M, Vidal M, Guinovart C, Jiménez A, et al. Seroprevalence of antibodies against SARS-CoV-2 among health care workers in a large Spanish reference hospital. *Nature Communications*. 2020;11(1):3500.
14. Folgueira MD, Muñoz-Ruipérez C, Alonso-López MÁ, Delgado R. SARS-CoV-2 infection in Health Care Workers in a large public hospital in Madrid, Spain, during March 2020. *medRxiv*. 2020:2020.04.07.20055723.
15. Thompson HA, Mousa A, Dighe A, Fu H, Arnedo-Pena A, Barrett P, et al. SARS-CoV-2 setting-specific transmission rates: a systematic review and meta-

- analysis. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2021.
16. Napolitano F, Bianco A, D'Alessandro A, Papadopoli R, Angelillo IF. Healthcare workers' knowledge, beliefs, and coverage regarding vaccinations in critical care units in Italy. *Vaccine*. 2019;37(46):6900-6.
17. Harrison N, Brand A, Forstner C, Tobudic S, Burgmann K, Burgmann H. Knowledge, risk perception and attitudes toward vaccination among Austrian health care workers: A cross-sectional study. *Human vaccines & immunotherapeutics*. 2016;12(9):2459-63.
18. Larson HJ, Schulz WS, Tucker JD, Smith DM. Measuring vaccine confidence: introducing a global vaccine confidence index. *PLoS currents*. 2015;7.
19. Andre FE, Booy R, Bock HL, Clemens J, Datta SK, John TJ, et al. Vaccination greatly reduces disease, disability, death and inequity worldwide. *Bulletin of the World Health Organization*. 2008;86(2):140-6.
20. Anderson RM, Vegvari C, Truscott J, Collyer BS. Challenges in creating herd immunity to SARS-CoV-2 infection by mass vaccination. *Lancet (London, England)*. 2020;396(10263):1614-6.
21. Neumann-Böhme S, Varghese NE, Sabat I, Barros PP, Brouwer W, van Exel J, et al. Once we have it, will we use it? A European survey on willingness to be vaccinated against COVID-19. *The European journal of health economics : HEPAC : health economics in prevention and care*. 2020;21(7):977-82.
22. Murphy J, Vallières F, Bentall RP, Shevlin M, McBride O, Hartman TK, et al. Psychological characteristics associated with COVID-19 vaccine hesitancy and resistance in Ireland and the United Kingdom. *Nature Communications*. 2021;12(1):29.
23. Shekhar R, Sheikh AB, Upadhyay S, Singh M, Kottewar S, Mir H, et al. COVID-19 Vaccine Acceptance Among Health Care Workers in the United States. *medRxiv*. 2021:2021.01.03.21249184.
24. Paterson P, Meurice F, Stanberry LR, Glismann S, Rosenthal SL, Larson HJ. Vaccine hesitancy and healthcare providers. *Vaccine*. 2016;34(52):6700-6.
25. Bianco A, Pileggi C, Iozzo F, Nobile CGA, Pavia M. Vaccination against Human Papilloma Virus infection in male adolescents: Knowledge, attitudes, and acceptability among parents in Italy. *Human vaccines & immunotherapeutics*. 2014;10(9):2536-42.
26. Smith LE, Amlôt R, Weinman J, Yiend J, Rubin GJ. A systematic review of factors affecting vaccine uptake in young children. *Vaccine*. 2017;35(45):6059-69.
27. Baicus A. History of polio vaccination. *World journal of virology*. 2012;1(4):108-14.
28. A gamble pays off in 'spectacular success': How the leading coronavirus vaccines made it to the finish line [press release]. Washington, United States The Washington Post2020.
29. Larson HJ, Clarke RM, Jarrett C, Eckersberger E, Levine Z, Schulz WS, et al. Measuring trust in vaccination: A systematic review. *Human vaccines & immunotherapeutics*. 2018;14(7):1599-609.
30. van der Weerd W, Timmermans DR, Beaujean DJ, Oudhoff J, van Steenbergen JE. Monitoring the level of government trust, risk perception and intention of the general public to adopt protective measures during the influenza A (H1N1) pandemic in The Netherlands. *BMC public health*. 2011;11:575.
31. Verger P, Fressard L, Collange F, Gautier A, Jestin C, Launay O, et al. Vaccine Hesitancy Among General Practitioners and Its Determinants During

- Controversies: A National Cross-sectional Survey in France. *EBioMedicine*. 2015;2(8):891-7.
32. Lazarus JV, Ratzan SC, Palayew A, Gostin LO, Larson HJ, Rabin K, et al. A global survey of potential acceptance of a COVID-19 vaccine. *Nat Med*. 2020:1-4.
 33. Rubin GJ, Amlôt R, Page L, Wessely S. Public perceptions, anxiety, and behaviour change in relation to the swine flu outbreak: cross sectional telephone survey. *BMJ*. 2009;339:b2651.
 34. Blair RA, Morse BS, Tsai LL. Public health and public trust: Survey evidence from the Ebola Virus Disease epidemic in Liberia. *Social Science & Medicine*. 2017;172:89-97.
 35. Vinck P, Pham PN, Bindu KK, Bedford J, Nilles EJ. Institutional trust and misinformation in the response to the 2018–19 Ebola outbreak in North Kivu, DR Congo: a population-based survey. *The Lancet Infectious Diseases*. 2019;19(5):529-36.
 36. Moxham-Hall V, Strang L. Public opinion and trust in government during a public health crisis London: King's College London; 2020 [Available from: <https://www.kcl.ac.uk/news/public-opinion-and-trust-in-government-during-a-public-health-crisis>.
 37. Adam DC, Wu P, Wong JY, Lau EHY, Tsang TK, Cauchemez S, et al. Clustering and superspreading potential of SARS-CoV-2 infections in Hong Kong. *Nature Medicine*. 2020;26(11):1714-9.
 38. Piroth L, Cottenet J, Mariet A-S, Bonniaud P, Blot M, Tubert-Bitter P, et al. Comparison of the characteristics, morbidity, and mortality of COVID-19 and seasonal influenza: a nationwide, population-based retrospective cohort study. *The Lancet Respiratory Medicine*. 2020.
 39. Petersen E. COVID-19 is not influenza. *The Lancet Respiratory Medicine*. 2020.
 40. Zürcher K, Zwahlen M, Berlin C, Egger M, Fenner L. Losing ground at the wrong time: trends in self-reported influenza vaccination uptake in Switzerland, Swiss Health Survey 2007–2017. *BMJ Open*. 2021;11(2):e041354.
 41. Sullivan SG, Chilver MB, Carville KS, Deng YM, Grant KA, Higgins G, et al. Low interim influenza vaccine effectiveness, Australia, 1 May to 24 September 2017. *Euro surveillance : bulletin European sur les maladies transmissibles = European communicable disease bulletin*. 2017;22(43).
 42. Sah P, Medlock J, Fitzpatrick MC, Singer BH, Galvani AP. Optimising the impact of low-efficacy influenza vaccines. *Proceedings of the National Academy of Sciences of the United States of America*. 2018.

TABLES AND FIGURE

Table 1: Characteristics of HCWs participating in the survey participants, overall and by profession.

	All HCWs n=3793	Nurses n=1690	Medical doctors n=614	Medical assistants n=63	Pharmacist and assistants n=34	Other medical staff ¹ n=380	Private caregivers n=224	Staff without patient contact n=788
Sex								
Male	952 (25.1)	202 (12.0)	358 (58.3)	1 (1.6)	10 (29.4)	57 (15.0)	77 (34.4)	247 (31.4)
Female	2,841 (74.9)	1,488 (88.0)	256 (41.7)	62 (98.4)	24 (70.6)	323 (85.0)	147 (65.6)	541 (68.7)
Age in years	43 (31-53)	39 (27-52)	46 (36-55)	39 (25-52)	49 (32-59)	43 (30.5-53)	41 (30-55)	48 (36-55)
Workplace								
Hospital	2,445 (64.5)	1,122 (66.4)	372 (60.6)	0	0	380 (100)	0	571 (72.5)
Medical practice	311 (8.2)	6 (0.4)	240 (39.1)	58 (92.1)	0	0	0	7 (0.9)
Retirement and nursing home	373 (9.8)	287 (17.0)	0	2 (3.2)	0	0	0	84 (10.7)
Residential care	323 (6.1)	193 (11.4)	0	2 (3.2)	0	0	0	37 (4.7)
Pharmacy	34 (0.9)	0	0	0	34 (100)	0	0	0
Long-term care facilities	398 (10.5)	82 (4.9)	2 (0.3)	1 (1.6)	0	0	224 (100)	89 (11.3)
Elderly	-	0	0	0	0	0	0	0
Willingness to get COVID-19 vaccine								
Yes	1,511 (39.8)	470 (27.8)	467 (76.1)	24 (38.1)	24 (70.6)	161 (42.4)	68 (30.4)	297 (37.7)
No	1,168 (30.8)	642 (38.0)	55 (9.0)	19 (30.2)	6 (17.6)	93 (24.5)	106 (47.3)	247 (31.3)
Unsure	1,114 (29.4)	578 (34.2)	92 (15.0)	20 (31.8)	4 (11.8)	126 (33.2)	50 (22.3)	244 (31.0)
Opinion on employer's recommendation								
No opinion	893 (23.5)	452 (26.8)	107 (17.4)	16 (25.4)	8 (23.5)	80 (21.1)	40 (17.9)	190 (24.1)
Completely and somewhat disagree	1,066 (28.1)	598 (35.4)	75 (12.2)	17 (27.0)	3 (8.8)	86 (22.6)	95 (42.4)	192 (24.4)
Completely and somewhat agree	1,834 (48.4)	640 (37.9)	432 (79.4)	30 (47.6)	23 (67.7)	214 (56.3)	89 (39.7)	406 (51.5)
Opinion on governments reports								
No opinion	645 (17.0)	337 (19.9)	47 (7.7)	12 (19.1)	5 (14.7)	55 (14.5)	45 (20.1)	144 (18.3)
Completely and somewhat disagree	868 (22.9)	489 (28.9)	48 (7.8)	18 (28.6)	6 (17.7)	57 (15.0)	71 (31.7)	179 (22.7)
Completely and somewhat agree	2,280 (60.1)	864 (51.1)	519 (84.5)	33 (52.4)	23 (67.7)	268 (70.5)	108 (48.2)	465 (59.0)
Had influenza vaccine 20/21								
No	2,229 (58.8)	1,101 (65.2)	113 (18.4)	34 (54.0)	11 (32.4)	217 (57.1)	193 (86.2)	560 (71.1)
Yes	1,446 (38.1)	550 (32.5)	464 (75.6)	28 (44.4)	20 (58.8)	151 (39.7)	26 (11.6)	207 (26.3)
Not sure	118 (3.1)	39 (2.3)	37 (6.0)	1 (1.6)	3 (8.8)	12 (3.2)	5 (2.2)	21 (2.7)
Reasons for vaccine willingness against COVID-19	1,511 (100)	470 (100)	467 (100)	24 (100)	24 (100)	161 (100)	68 (100)	297 (100)
Protection	1,461 (96.7)	449 (95.5)	460 (98.5)	23 (95.8)	21 (87.5)	151 (93.8)	65 (95.6)	292 (98.3)
Vaccine for pandemic control	1,328 (87.9)	395 (84.0)	434 (92.9)	24 (100)	21 (87.5)	143 (88.8)	55 (80.9)	256 (86.2)
Risk group	266 (17.6)	76 (16.2)	96 (20.6)	0	7 (29.2)	20 (12.4)	14 (20.6)	53 (17.8)
Reasons for vaccine hesitancy against COVID-19	1,168 (100)	642 (100)	55 (100)	19 (100)	6 (100)	93 (100)	106 (100)	247 (100)
Vaccine safety and side effects	1,073 (91.9)	603 (93.9)	51 (92.7)	17 (89.5)	4 (66.7)	87 (93.5)	90 (84.9)	221 (89.5)
PPE is sufficient	328 (28.1)	193 (30.1)	13 (23.6)	4 (21.1)	1 (16.7)	26 (28.0)	32 (30.2)	59 (23.9)
COVID-19 harmless	229 (19.6)	118 (18.4)	20 (36.4)	3 (15.8)	0	16 (17.2)	25 (23.6)	47 (19.0)
Bad experiences	201 (17.2)	111 (17.3)	12 (18.2)	2 (10.5)	0	11 (11.8)	20 (18.9)	45 (18.2)
Effectiveness of the vaccine	128 (11.0)	85 (13.2)	6 (10.9)	1 (5.3)	0	8 (8.6)	12 (11.3)	16 (6.5)

Opponents of vaccines in general	39 (3.3)	20 (3.1)	0	1 (5.3)	0	1 (1.1)	5 (4.7)	12 (4.9)
Fear of needles	40 (3.4)	20 (3.1)	1 (1.8)	0	1 (16.7)	4 (4.3)	4 (3.8)	10 (4.0)
Others	379 (32.4)	194 (30.2)	24 (43.6)	4 (21.1)	1 (16.7)	33 (35.5)	46 (43.4)	77 (31.2)
Reasons for being unsure of vaccination against COVID-19	1,114 (100)	578 (100)	92 (100)	20 (100)	4 (100)	126 (100)	50 (100)	244 (100)
Vaccine safety and side effects	1,058 (95.0)	538 (93.1)	88 (95.7)	18 (90.0)	4 (100)	124 (98.4)	50 (100)	236 (96.7)
Inconsistent information	329 (29.5)	192 (33.2)	29 (31.5)	1 (5.0)	1 (25.0)	40 (31.7)	17 (34.0)	51 (20.9)
Effectiveness of the vaccine	316 (28.4)	177 (30.6)	16 (17.4)	5 (25.0)	1 (25.0)	33 (26.2)	15 (30.0)	69 (28.3)
No education by my employer	210 (18.9)	126 (21.8)	14 (15.2)	5 (25.0)	0	29 (23.0)	4 (8.0)	32 (13.1)
Uncertainty by my colleagues	251 (22.5)	147 (25.4)	18 (19.6)	6 (30.0)	0	31 (24.6)	10 (20.0)	39 (16.0)
Vaccine for pandemic control	174 (15.6)	93 (16.1)	14 (15.2)	2 (10.0)	0	17 (13.5)	10 (20.0)	38 (15.6)
Information needs among unsure HCWs	1,114 (100)	578 (100)	92 (100)	20 (100)	4 (100)	126 (100)	50 (100)	244 (100)
Information about COVID-19	116 (10.4)	68 (11.8)	8 (8.7)	1 (5.0)	0	14 (11.1)	5 (10.0)	20 (8.2)
Information about the vaccine	1,055 (94.7)	550 (95.2)	81 (88.0)	19 (95.0)	4 (100)	121 (96.0)	47 (94.0)	233 (95.5)
Reports from vaccinated people	778 (69.8)	412 (71.3)	45 (48.9)	17 (85.0)	2 (50.0)	74 (58.7)	41 (82.0)	187 (76.6)
Opinion of experts	404 (36.3)	220 (38.1)	32 (34.8)	9 (45.0)	0	56 (44.4)	15 (30.0)	72 (29.5)

Abbreviations: PPE; personal protective equipment; HCW, health care worker

¹ Other medical staff includes physical therapists, occupational therapists, speech therapists, and midwives

Table 2: Characteristics of HCWs willingness to get vaccinated against COVID-19, vaccine hesitancy and unsure about vaccination and factors associated with willingness to get vaccinated.

Variables	Total Number of participants	Willingness to vaccinate against COVID-19			Comparing persons with willingness to be vaccinated against Covid-19 to persons with hesitancy/unsure			
		Yes	No	Unsure	unadjusted OR (95% CI)	p-value	adjusted OR (95% CI) *	p-value
Total	3,793	1,511	1,168	1,114				
Sex								
Male	952	561 (58.9)	196 (20.6)	195 (20.5)	1	<0.001		
Female	2841	950 (33.4)	972 (34.2)	919 (32.4)	0.33 (0.28-0.38)			
Age groups								
29 and younger	829	207 (24.9)	381 (46.0)	241 (29.1)	1		1	
30 to 39	807	285 (35.3)	257 (31.9)	265 (32.8)	1.64 (1.33-2.03)		0.93 (0.68-1.25)	
40 to 49	790	335 (42.4)	215 (27.2)	240 (30.4)	2.21 (1.79-2.73)		1.29 (0.96-1.76)	
50 to 59	1,012	470 (46.4)	254 (25.1)	288 (28.5)	2.61 (2.13-3.18)		1.78 (1.34-2.36)	
60 and older	355	214 (60.3)	61 (17.2)	80 (22.5)	4.56 (3.50-5.94)		2.45 (1.68-3.57)	
Profession								
Medical doctor	614	467 (76.0)	55 (9.0)	92 (15.0)	8.24 (6.66-12.21)	<0.001	4.88 (3.58-6.65)	<0.001
Nurse	1,690	470 (27.8)	642 (38.0)	578 (34.2)	1		1	
Medical assistant	63	24 (70.6)	6 (17.7)	4 (11.8)	1.60 (0.95-2.69)		1.45 (0.68-3.10)	
Pharmacist and pharmacist assistant	24	16 (66.7)	5 (20.8)	3 (12.5)	6.22 (2.96-13.13)		4.20 (1.53-11.52)	
Other medical staff ¹	380	161 (42.4)	93 (24.5)	126 (33.2)	1.91 (1.52-2.40)		1.22 (0.89-1.68)	
Not close patient contact	788	297 (37.7)	247 (31.3)	244 (31.9)	1.57 (1.31-1.88)		1.15 (0.89-1.48)	
Private caregiver	224	68 (30.4)	106 (47.3)	50 (22.3)	1.13 (0.83-1.53)		1.51 (0.98-2.33)	
Confidence in the government reports								
No opinion	645	150 (23.3)	238 (36.9)	257 (39.8)	1	<0.001	1	<0.001
Completely and somewhat disagree	868	95 (10.9)	533 (61.4)	240 (27.7)	0.41 (0.31-0.54)		0.58 (0.40-0.84)	
Completely and somewhat agree	2280	1,266 (55.5)	397 (17.4)	617 (27.1)	4.12 (3.37-5.03)		1.41 (1.07-1.85)	
Follow employer's recommendation								
No opinion	893	140 (15.7)	283 (31.7)	470 (52.6)	1	<0.001	1	<0.001
Completely and somewhat disagree	1,066	41 (3.9)	801 (75.1)	224 (21.0)	0.22 (0.15-0.31)		0.30 (0.20-0.44)	
Completely and somewhat agree	1,834	1,330 (72.5)	84 (4.6)	420 (22.9)	14.19 (11.53-17.47)		11.40 (8.87-14.65)	
Influenza vaccination 20/21								
No	2,347	555 (23.7)	1019 (43.4)	773 (32.9)	1	<0.001	1	<0.001
Yes	1,556	956 (66.1)	149 (10.3)	341 (23.6)	6.30 (5.45-7.23)		2.70 (2.20-3.31)	

* Model adjusted for age groups, profession, confidence in the government reports, following employer's recommendation, and influenza vaccination 20/21.

1 Other medical staff includes physical therapists, occupational therapists, speech therapists, and midwives

Figure 1: Number of people reporting the willingness to be vaccinated against COVID-19 in the Canton of Solothurn, Switzerland by healthcare profession (A) and by workplace (B). Percent with 95% confidence intervals are presented.

Figure 2: Reasons for vaccine willingness (A), vaccine hesitancy (B) and being uncertain (C) to Covid-19 among all HCWs. The exact numbers can be found in Table 1.

Supplementary Table and Figure:

Table S1: Grouping of the reasons for willingness, hesitancy and unsure to be vaccinated against COVID-19.

Reasons for willingness to be vaccinated	
To protect myself from COVID-19 infection.	Protection (myself, patients, and relatives)
Because I want to protect my colleagues at work and my patients.	
Because I want to protect my family members (parents, siblings, children, etc.).	
Because my vaccination is a contribution to the fight against the pandemic.	Vaccine for pandemic control
Because this is the only way to stop the pandemic.	
Because I belong to the risk groups.	Risk group
Reasons for hesitancy to be vaccinated	
I think vaccinations are generally unnecessary.	Opponents of vaccination
Vaccination against COVID-19 is not necessary.	COVID-19 is harmless
I don't think COVID-19 is dangerous for my health.	
COVID-19 is no worse than the flu.	Effectiveness of the vaccine
I don't think the vaccine is effective.	
I think the COVID-19 vaccine may not be safe.	Vaccine safety and side effects
I am concerned that the vaccine was approved too quickly and has not been tested enough	
I am afraid of possible side effects.	Bad experiences with vaccines
I have had a bad experience or reaction to a previous vaccination	
I had a bad experience with vaccinations at a previous physician or health clinic.	
I am afraid of needles.	Fear of needles
I am already protected because I comply with protective measures (masks, hand hygiene).	Personal protective equipment is sufficient
I protect my environment because I follow the protective measures, therefore vaccination is unnecessary.	
Because some colleagues do not want to be vaccinated either.	Other
Because the experts' opinions are inconsistent	
The best way is to let nature take its course.	
Religious reasons.	
Reasons for being unsure about being vaccinated	
The information is inconsistent	Inconsistent information
The vaccines were too fast/quickly approved Swissmedic	Vaccine safety and side effects
I fear possible side effects.	
I don't know if the vaccination will provide protection for me and/or others.	Effectiveness of the vaccine
I don't know if my vaccination will help contain the pandemic (herd immunity).	Vaccine for pandemic control
My colleagues are also unsure or against vaccination.	Uncertainty of my colleagues
I have not yet been educated by my employer.	Education by employer
Other	
Information needed to take a decision	
General information about the disease COVID-19.	Information about COVID-19
Information on the safety of the vaccination.	Information's about vaccine (safety, effectiveness, side effects)
Information about possible side effects of the vaccination.	
Information about the effectiveness of the vaccination (does it protect me?)	
Information about the vaccination procedure (e.g. how often vaccination is necessary).	
Experiences and reports from people who have already been vaccinated.	Reported experiences from vaccinated people
A statement from experts	Opinion by experts

Table S2: Participant characteristics by workplace.

	Total	Hospital	Medical practice	Nursing home	Home care organisation	Pharmacy	Long-term care facilities
Total	3975	2445	311	373	232	34	398
Sex							
Male	952 (25.1)	576 (23.6)	164 (52.7)	59 (15.8)	15 (6.5)	10 (29.4)	128 (6.5)
Female	2841 (74.9)	1,869 (76.4)	147 (47.3)	314 (84.2)	217 (93.5)	24 (70.6)	270 (67.8)
Age in years	43 (31-53)	40 (30-52)	52 (44-60)	46 (30-55)	49.5 (38.5-57)	49 (32-59)	44 (31-55)
Workplace							
Medical doctor	614 (16.2)	372 (15.2)	240 (77.2)	0	0	0	2 (0.5)
Nurse	1,690 (44.6)	1,122 (45.9)	6 (1.9)	287 (76.9)	193 (83.2)	0	82 (20.6)
Medical assistant	63 (1.7)	0	58 (18.7)	2 (0.5)	2 (0.9)	0	1 (0.3)
Other medical staff	380 (10.0)	380 (15.5)	0	0	0	0	0
Pharmacist	34 (0.9)	0	0	0	0	24 (70.6)	0
Not close patient contact	788 (20.8)	571 (23.4)	7 (2.3)	84 (22.5)	37 (16.0)	0	89 (22.4)
Willingness to get COVID-19 vaccine							
Yes	1511 (39.8)	936 (38.3)	226 (72.7)	126 (33.8)	79 (34.0)	24 (70.6)	120 (30.2)
No	1,168 (30.8)	726 (29.7)	32 (10.3)	137 (36.7)	87 (37.5)	6 (17.6)	180 (45.2)
Unsure	1,114 (29.4)	783 (32.0)	53 (17.0)	110 (29.5)	66 (28.5)	4 (11.8)	98 (24.6)
Follow my employer's recommendation							
Completely disagree	443 (11.7)	237 (9.7)	15 (4.8)	64 (17.2)	44 (19.0)	3 (8.8)	80 (20.1)
Somewhat disagree	623 (16.4)	403 (16.5)	32 (10.3)	61 (16.4)	47 (20.3)	0	80 (20.1)
No opinion	893 (23.5)	584 (23.9)	74 (23.8)	101 (27.1)	49 (21.1)	8 (23.5)	77 (19.4)
Somewhat agree	742 (19.6)	528 (21.6)	41 (13.2)	54 (14.5)	43 (18.5)	6 (17.7)	70 (17.6)
Completely agree	1,092 (28.8)	693 (28.3)	149 (47.9)	93 (24.9)	49 (21.1)	17 (50.0)	91 (22.9)
Confidence in the governments reports							
Completely disagree	230 (6.1)	129 (5.3)	7 (2.3)	37 (9.9)	14 (6.0)	0	43 (10.8)
Somewhat disagree	638 (16.8)	388 (15.9)	33 (10.6)	74 (19.8)	48 (20.7)	6 (17.7)	89 (22.4)
No opinion	645 (17.0)	409 (16.7)	31 (10.0)	76 (20.4)	47 (20.3)	5 (14.7)	77 (19.4)
Somewhat agree	955 (25.2)	645 (26.4)	71 (22.8)	74 (19.8)	70 (30.2)	11 (32.4)	84 (21.1)
Completely agree	1,325 (34.9)	874 (35.8)	169 (54.3)	112 (30.0)	53 (22.8)	12 (35.3)	105 (26.4)
Influenza vaccine 20/21							
No	2,229 (58.8)	1364 (55.8)	78 (25.1)	262 (70.2)	171 (73.7)	11 (32.4)	343 (86.2)
Yes	1,446 (38.1)	995 (40.7)	222 (71.4)	102 (27.4)	58 (25.0)	20 (58.8)	49 (12.3)
Unsure	118 (3.1)	86 (3.5)	11 (3.5)	9 (2.4)	1 (0.6)	3 (2.4)	6 (1.5)

1 Other medical staff includes physical therapists, occupational therapists, speech therapists, and midwives

Table S3: Sensitivity Analysis. Determinants for the willingness to be vaccinated against COVID-19 (yes/unsure) among HCWs compared with HCWs hesitancy.

Variables	Total Number of participants	Willingness to vaccinate against COVID-19		Comparing the willingness to vaccinate against COVID-19 (yes/unsure) to persons with hesitancy			
		Yes	No	unadjusted OR (95% CI)	p-value	adjusted OR (95% CI) *	p-value
Total	3,793	2,625	1,168				
Sex					<0.001		
Male	952	756 (79.4)	196 (20.6)	1			
Female	2,841	1,869 (65.8)	972 (34.2)	0.50 (0.42-0.59)			
Age groups					<0.001		<0.001
29 and younger	829	448 (54.0)	381 (46.0)	1		1	
30 to 39	807	550 (68.2)	257 (31.9)	1.64 (1.33-2.03)		1.48 (1.12-1.97)	
40 to 49	790	575 (72.8)	215 (27.2)	2.21 (1.79-2.73)		1.70 (1.23-2.29)	
50 to 59	1012	758 (74.9)	254 (25.1)	2.61 (2.13-3.18)		2.08 (1.58-2.73)	
60 and older	355	294 (82.8)	61 (17.2)	4.56 (3.50-5.94)		2.72 (1.79-4.13)	
Profession					<0.001		<0.001
Medical doctor	614	559 (91.0)	55 (9.0)	6.22 (4.64-8.35)		1.93 (1.31-2.84)	
Nurse	1,690	1,048 (62.0)	642 (40.0)	1		1	
Medical assistant	63	44 (69.4)	19 (30.2)	1.42 (0.82-2.45)		1.20 (0.56-2.57)	
Pharmacist and pharmacist assistant	34	28 (82.4)	6 (17.6)	2.86 (1.18-6.94)		0.85 (0.26-2.79)	
Other medical staff ¹	380	287 (75.5)	93 (24.5)	1.89 (1.47-2.44)		1.04 (0.73-1.47)	
Not close patient contact	788	541 (68.6)	246 (31.4)	1.34 (1.12-1.61)		0.62 (0.41-0.94)	
Private caregiver	224	118 (52.7)	106 (47.3)	0.68 (0.52-0.90)		0.80 (0.62-1.04)	
Confidence in the governments reports					<0.001		<0.001
No opinion	645	407 (63.1)	238 (36.9)	1		1	
Completely and somewhat disagree	868	335 (38.6)	533 (61.4)	0.37 (0.30-0.45)		0.82 (0.62-1.09)	
Completely and somewhat agree	2,280	1883 (82.6)	397 (17.4)	2.77 (2.29-3.36)		1.59 (1.23-2.06)	
Follow employer's recommendation					<0.001		<0.001
No opinion	893	610 (68.3)	283 (31.7)	1		1	
Completely and somewhat disagree	1,066	265 (24.9)	801 (75.1)	0.15 (0.13-0.19)		0.17 (0.13-0.21)	
Completely and somewhat agree	1,834	1,750 (95.4)	84 (4.6)	9.67 (7.45-12.54)		6.58 (4.94-8.76)	
Influenza Vaccination 20/21					<0.001		<0.001
No	2347	1,328 (56.7)	1019 (43.4)	1		1	
Yes	1,556	1,297 (89.7)	149 (10.3)	6.68 (5.53-8.06)		2.48 (1.94-3.17)	

* Model adjusted for age groups, profession, confidence in the government reports, following employer's recommendation, and influenza vaccination 20/21.

1 Other medical staff includes physical therapists, occupational therapists, speech therapists, and midwives

Figure S1: Flow chart.