Predictors of hospitalisation and death due to SARS-CoV-2 infection in Finland: a population-based register study with implications to vaccinations

Heini Salo^{1*}, Toni Lehtonen¹, Kari Auranen^{1,2}, Ulrike Baum¹, Tuija Leino¹

1. Infectious Disease Control and Vaccinations Unit, Department of Health Security, Finnish Institute for Health and Welfare (THL), Helsinki, Finland
2. The Center of Statistics, University of Turku, Turku, Finland

*Corresponding author

E-mail address: heini.salo@thl.fi

Article type: Research Article

Content: Original research

Keywords: SARS-CoV-2, COVID-19, risk factors, elderly, chronically ill

Abstract

Introduction. The aim of this study was to investigate how age and underlying medical conditions affect the risk of severe outcomes following SARS-CoV-2 infection and how they should be weighed while prioritising vaccinations against COVID-19.

Methods. This population-based register study includes all SARS-CoV-2 PCR-test-positive cases until 24 Feb 2021, based on the Finnish National Infectious Diseases Register. The cases were linked to other registers to identify presence of comorbidities and severe outcomes (hospitalisation, intensive care treatment, death). The odds of severe outcomes were compared in those with and without the pre-specified comorbidities using logistic regression. Furthermore, population-based rates were compared between those with a given comorbidity and those without any of the specified comorbidities using negative binomial regression.

Results. Age and various comorbidities were found to be predictors of severe COVID-19. Compared to 60–69-year-olds, the odds ratio (OR) of death was 7.1 for 70–79-year-olds, 26.7 for 80–89-year-olds, and 55.8 for ≥90-year-olds. Among the 20–69-year-olds, chronic renal disease (OR 9.4), malignant neoplasms (5.8), hematologic malignancy (5.6), chronic pulmonary disease (5.4), and cerebral palsy or other paralytic syndromes (4.6) were strongly associated with COVID-19 mortality; severe disorders of the immune system (8.0), organ or stem cell transplant (7.2), chronic renal disease (6.7), and diseases of myoneural junction and muscle (5.5) were strongly associated with COVID-19 hospitalisation. Type 2 diabetes and asthma, two very common comorbidities, were associated with all three outcomes, with ORs from 2.1 to 4.3. The population-based rate of SARS-CoV-2 infection decreased with age. Taking the 60–69-year-olds as reference,

the rate ratio was highest (3.0) for 20–29-year-olds but <1 for 70–79-year-olds and 80–89-year-olds.

Conclusion. Comorbidities predispose for severe COVID-19 among younger ages. In vaccine prioritisation both the risk of infection and the risk of severe outcomes, if infected, should be combined.

Introduction

Already during summer 2020, the first listings of predisposing conditions for severe COVID-19 were published to inform the public on the need to shield the most vulnerable.[1-6] Furthermore, while efficacy studies of COVID-19 vaccines were still being carried out, more detailed research was published on predisposing factors for severe infection outcomes leading to hospitalisation [7,8], intensive care [7,9,10], or death [7,8,11-14]. The implications from these studies have been valuable in planning how to allocate the limited number of vaccine doses optimally to prevent hospitalisations, need to intensive care and deaths.[15] For example, one of the questions was whether the elderly or individuals with certain background illnesses should be prioritised for COVID-19 vaccination.[15] Along with the other European Union countries, Finland joined a common procurement system and subsequently commenced COVID-19 vaccinations in late December 2020.

We carried out a country-level analysis of factors predisposing for severe COVID-19 outcomes, based on the possibility to link information across several national health care registers using the unique personal identifier. The specific aim of our study was to investigate how age and presence of underlying medical conditions affect the risk of severe COVID-19 outcomes and how age and medical conditions should be weighed against each other while prioritising vaccination. We thus

concentrated on assessing the risk of hospitalisation, intensive care treatment and death in those with SARS-CoV-2 infection, with specific attention to the elderly and persons with certain chronic illnesses and those taking potentially predisposing medications. We here report the results with a particular focus on how these data were used to support effective COVID-19 vaccination intervention in Finland.

Material and Methods

Data sources and case definitions

The total population of Finland in 2019 was 5.5 million, 51% of which were women. Age- and gender-specific population sizes were obtained from the Finnish Population Information System, a computerised national database which registers individual-based information for all Finnish residents, including name, gender, personal identity code, address, date of birth, and death. The personal identity code remains unchanged throughout the lifetime and is used for patient identification in all health care registers in Finland.

We established a dataset comprising virtually the entire Finnish population by linking data from multiple population-based and nationwide registers. This COVID-19 dataset included all individuals with a positive PCR test for SARS-CoV-2 identified between 1 January 2020 and 24 February 2021, based on the Finnish National Infectious Diseases Register. The SARS-CoV-2 cases were linked to other registers in order to identify severe outcomes. Three severe COVID-19 outcomes were defined as inpatient hospitalisation (Care Register for Health Care), admission to intensive care unit (ICU; the Finnish Intensive Care Consortium's Database, FICC), and death within 30 days after the positive PCR test. All cases hospitalised 14 days before or after the positive PCR test with a diagnosis indicating respiratory infection, were considered as COVID-19 outcomes.

83 Individuals with chronic diseases or medical conditions (from now on comorbidities) known as
 84 general risk factors of infectious diseases or with risk factors for severe COVID-19 from published
 85 studies were identified from the registers using a pre-defined list of diagnoses and codes (Table 1).
 86 The following registers were employed: the Care Register for Health Care, the Register of Primary
 87 Health Care Visits, as well as the Special Reimbursement Register for Medicine Expenses and
 88 Prescription Centre database, both maintained by the Social Insurance Institution of Finland (SII).
 89 The dataset covers for the most part all health care providers in Finland. The dataset was linked to
 90 the Finnish Population Information System in order to identify individuals without any of the
 91 comorbidities potentially predisposing to severe COVID-19 listed in Table 1.

92 Individuals entitled to reimbursement of medicine expenses from 1 January 2020 onwards were
 93 identified from the SII register for Special Reimbursements for Medicine Expenses. In Finland, all
 94 residents entitled to special reimbursement meet a set of specified medical criteria.

95 Individuals using medication predisposing to immunodeficiency or using medication for type 1 or 2
 96 diabetes were identified from the Prescription Centre database maintained by SII. Persons that
 97 had any of the prescribed medications of interest not more than two years prior to testing positive
 98 for SARS-CoV-2 were identified based on the Anatomical Therapeutic Chemical (ATC) codes.

99 We identified individuals who had used health care services due to any of the specified
 100 comorbidities not more than two years (cancer) or five years (all other comorbidities) prior to the
 101 positive PCR test. Individuals who had used secondary health care services due the specified
 102 comorbidities from the Care Register for Health Care with the International Classification of
 103 Diseases 10th revision (ICD-10) codes and the Nordic Classification of Surgical Procedures (NCSP)
 104 codes. Individuals who had used primary health care services were identified from the Register of

105 Primary Health Care Visits with ICD-10 codes and the International Classification of Primary Care
106 Second edition (ICPC-2) codes.

107 *Statistical analysis*

108 We estimate the incidence of SARS-CoV-2 infection and severe COVID-19 outcomes by age, sex
109 and comorbidity in individuals with a SARS-CoV-2 infection and in the Finnish population as a
110 whole. The three severe outcomes were hospitalisation, admission to ICU, and death. We first
111 estimated age-specific odds ratios (ORs) for each outcome (e.g. hospitalisation) in SARS-CoV-2
112 cases using logistic regression and adjusting for sex. We next estimated odds ratios, comparing
113 the odds of the outcome (e.g. hospitalisation) in SARS-CoV-2 cases with and without each of the
114 comorbidities, adjusting for age and sex. In addition, negative binomial regression was used to
115 estimate rate ratios (RR) for SARS-CoV-2 infection and the three severe outcomes, comparing
116 individuals with each of the comorbidities with individuals without any of the listed comorbidities
117 (Table 1). The negative binomial model was used instead of Poisson regression to account for
118 overdispersion in the rates. The analysis was adjusted for age and sex.

119 **Results**

120 *Characteristics of the study population*

121 In Finland, a total of 52 502 SARS-CoV-2 cases occurred from 1 March 2020 through 24 February
122 2021, 2743 (5.2%) of which were hospitalised, 538 (1.0%) were admitted to ICU, and 801 (1.5%)
123 died within 30 days after the positive PCR-test (Table 2). The corresponding incidence rates per
124 100 000 person-years were 822.4 for SARS-CoV-2 cases, 43 for hospitalised cases, 8.4 for ICU cases
125 and 12.5 for fatal cases. When restricting to 20–69-year-olds, there were 38 728 SARS-CoV-2

126 cases, 1 916 (4.9%) of which were hospitalised, 403 (1.0%) were admitted to ICU and 98 (0.3%)
 127 died within 30 days after the positive PCR-test.

128 The proportion of fatalities among SARS-CoV-2 cases increased with age (Figure 1). In those aged
 129 60–69 years, all fatal cases but one had at least one predisposing factor for severe COVID-19. By
 130 contrast, in those aged ≥ 70 fatalities occurred also in individuals without any predisposing factors.
 131 In 20–69-year-olds, 28% of SARS-CoV-2 cases, 64% of hospitalised cases, 72% of ICU cases, and
 132 87% of fatal cases had at least one predisposing factor.

133 The proportion of fatalities among SARS-CoV-2 cases increased with the number of comorbidities
 134 (Figure 1). In all age groups, the more comorbidities there were, the higher the mortality was
 135 among the infected. With three or more comorbidities the proportion of fatal cases varied from
 136 4% (65–69 years) to 32% (75–89 years). In the oldest the proportion of fatal cases was even
 137 higher, irrespective of the number of comorbidities.

138 During the study period, SARS-CoV-2 infections were most frequent among 20–29-year-olds,
 139 hospitalised cases among 50–59-year-olds, ICU cases among 60–69-year-olds, and fatal cases
 140 among 80–89-year-olds (Table 2). The age-specific incidence rates per 100 000 person-years
 141 peaked for SARS-CoV-2 infections in those aged 20–29 (1475), for hospitalised cases in those aged
 142 80–89 (79.6), for ICU cases in those aged 60–69 (17.6) and for fatal cases in those aged ≥ 90 (289.1)
 143 (Figure 2). Severe COVID-19 outcomes were thus clearly over-represented among the elderly. Only
 144 6% (3390/52502) of all SARS-CoV-2 cases were detected in individuals aged ≥ 70 (Table 2). Yet 28%
 145 (766/2 743) of hospitalised cases, 25% (132/538) of ICU cases and 88% (702/801) of the fatal cases
 146 occurred in this age group.

147 *Predictors of hospitalisation, ICU admission and death*

148 According to the logistic regression analysis, the risk of severe COVID-19 outcomes in an individual
 149 infected with SARS-CoV-2 increased with age. Compared to 60–69-year-olds, the odds ratio (OR) of
 150 death, adjusted for sex, was 7.1 (95% CI 5.3, 9.7) for the 70–79-year-olds, 26.7 (95% CI 20, 36) for
 151 the 80–89-year-olds, and 55.8 (95% CI 40.5, 77.9) for the ≥90-year-olds (Table 3a). Furthermore,
 152 the OR of hospitalisation was 2.0 (95% 1.8, 2.3) and of ICU admission 1.7 (95% CI 1.3, 2.2) for those
 153 aged 70–79 (Table 3a). In addition, male sex was identified as a risk factor for all severe outcomes:
 154 hospitalisation (OR 1.3; 95% CI 1.2, 1.4), ICU admission (OR 1.9; 95% CI 1.6, 2.3), and death (OR
 155 1.9; 95% CI 1.6, 2.2).

156 The more comorbidities a person had, the higher was their risk of severe COVID-19. The effect was
 157 even stronger when the analysis was restricted to 20–69-year-olds (Table 3b). Comparing SARS-
 158 CoV-2 cases with three or more comorbidities with cases without any of the comorbidities in
 159 question, the age- and sex-adjusted ORs of severe outcomes were 6.6 for hospitalisation (95% CI
 160 5.6, 7.7), 9.5 for ICU admission (95% CI 7.0, 12.9), and 16.1 for death (95% CI 8.1, 33.4).

161 Many of the individual comorbidities were clear risk factors for death. Among the 20–69-year-olds
 162 the strongest risk factors for death due to COVID-19 were severe chronic renal disease (OR 9.4;
 163 95% CI 4.2, 18.8), malignant neoplasms (OR 5.8; 95% CI 3.3, 9.7), hematologic malignancy (OR 5.6;
 164 95% CI 1.6, 14.1), chronic pulmonary disease (OR 5.4; 95% CI 2.0, 11.9), and cerebral palsy and
 165 other paralytic syndromes (OR 4.6; 95% CI 0.3, 22.6), when comparing the odds between
 166 individuals with and without the comorbidity in question (Table 3b). Likewise, cerebrovascular
 167 diseases, type 2 diabetes, autoimmune diseases treated with immunosuppressive drugs, use of
 168 glucocorticoids, secondary hypertension alone, ischaemic heart diseases and asthma increased the

169 risk of death, with ORs between 2 and 5. Some of the comorbidities were very rare, thus, it was
170 not possible to calculate reliable OR estimates.

171 Strong risk factors for COVID-19 hospitalisation among the 20–69-year-olds were severe disorders
172 of the immune system (OR 8.0; 95% CI 3.5, 17.2), organ or stem cell transplant (OR 7.2; 95% CI 4.0,
173 12.5), chronic renal disease (OR 6.7; 95% CI 4.6, 9.6), diseases of myoneural junction and muscle
174 (OR 5.5; 95% CI 2.9, 10.2) (Table 3b). Likewise, psychotic disorders, chronic pulmonary disease,
175 paralytic syndromes, asthma, hematologic malignancy, type 2 diabetes, cerebrovascular diseases,
176 cardiovascular diseases, autoimmune diseases treated with immunosuppressive drugs, sleep
177 apnea, and the use of clozapine, immunosuppressants or biologic drugs increased the risk of
178 hospitalisation, with ORs between 2 and 5. The severity of the comorbidity was associated with
179 severe disease outcomes. The risk of hospitalisation and admission to ICU was higher for asthma
180 and type 2 diabetes cases that had been treated in secondary health care (Figure 3).

181 According to the negative binomial regression model, the population-based rate of SARS-CoV-2
182 infection decreased with age. Compared to the 60–69-year-olds, the rate was 3.0-times higher in
183 20–29-year-olds (RR 3.0; 95% CI 2.8, 3.3) and 2.4-times higher in 30–39-year-olds (RR 2.4; 95% CI
184 2.2, 2.5), but somewhat lower in 70–79-year-olds (RR 0.7; 95% CI 0.6, 0.7) and 80–89-year-olds (RR
185 0.9; 95% CI 0.9, 1). The higher rate of SARS-CoV-2 infection in the ≥90-year-olds (RR 1.5; 95% CI
186 1.4, 1.7) can be explained by local epidemics which occurred in nursing homes. However, the rate
187 of SARS-CoV-2 infection did not differ between females and males. Compared to the 60–69-year-
188 olds, the population-based rate of hospitalisation was higher among the 50–59-year-olds (RR 1.3;
189 95% CI 1.2, 1.4) and the risk of ICU admission was higher among the 50–59-year-olds (RR 1.2; 95%
190 CI 1.0, 1.4).

191 In addition, for each comorbidity, we report the relative rates of hospitalisation, ICU admission
 192 and death in the 20–69-year-olds, comparing individuals with the comorbidity with those without
 193 any of the specified comorbidities (Table 4). The rate of SARS-CoV-2 infection was roughly the
 194 same for most of the comorbidities when comparing to those without any of the specified
 195 comorbidities (Table 4). However, some differences were obvious, e.g. those using clozapine (RR
 196 0.5; 95% CI 0.4, 0.7), with psychotic disorders (RR 0.6; 95% CI 0.5, 0.7) and heart failure, (RR 0.7;
 197 95% CI 0.6, 0.9) had lower rates of SARS-CoV-2 infection. By contrast, the relative rate of severe
 198 COVID-19 outcomes in the 20–69-year-olds, when comparing to those without any of the specified
 199 comorbidities varied, e.g. For hospitalisation it was 5.1–8.9-times higher for organ or stem cell
 200 transplant, severe chronic renal disease, severe disorders of the immune system, diseases of
 201 myoneural junction and muscle and type 2 diabetes.

202 At the upper part of Table 4, 60–69-year-olds is the reference category, and the rate ratios
 203 presented for each comorbidity and outcome at the bottom part of the table can be taken to
 204 present relative rates in that age group. The additional risk caused by a specific comorbidity at any
 205 other age category can be obtained by multiplying the rate ratio of interest at a lower part of the
 206 table by the rate ratio of the age group of interest. For almost all comorbidities, the relative risks
 207 of dying for 50–59-year-olds with comorbidities, were greater than for 60–69-year-olds without
 208 any comorbidities. For example, the risk for 55-year-old person with chronic pulmonary disease to
 209 die of COVID-19 is almost 10 times (0.5×19.5) of that of 65-year-old person without any of the
 210 listed conditions.

211 Discussion

212 In this population-based setting we estimated the risks for four entities: SARS-CoV-2 infection and
 213 COVID-19 related hospitalisation, ICU admission and death by age and various comorbidities or
 214 conditions. Age and certain comorbidities were found to be predictors of severe COVID-19 in cases
 215 of SARS-CoV2 infection. Old age was strongly associated with mortality, and having multiple
 216 comorbidities was associated with both mortality and hospitalisation. The estimation of
 217 population-based rates of hospitalisation and death related to SARS-CoV-2 infection allowed an
 218 even more detailed characterisation of the relative roles of age and comorbidities as risk factors
 219 for severe COVID-19. Based on this analysis, vaccination of the elderly aged ≥ 70 years was
 220 prioritised in Finland, followed by vaccination of those with highly predisposing medical
 221 conditions.[16]

222 As expected, immunosuppressive states such as organ or stem cell transplantation, severe
 223 disorders of the immune system or severe renal disease were associated with increased risks of
 224 hospitalisation and ICU admission due to COVID-19. Similar results were obtained in a population-
 225 based Danish study where organ transplantation and severe kidney disease were the two medical
 226 conditions associated with the highest risks of severe COVID-19.[7] In an English study, which
 227 recorded nearly 11 000 deaths in a cohort of more than 17 million SARS-CoV-2 positive patients,
 228 organ transplantation was associated with the highest risk of death due to COVID-19 (hazard ratio
 229 6.0; 95% CI 4.7, 7.6) among the studied medical conditions, after the analysis was adjusted for age
 230 and sex.[11] Furthermore, in our study patients with an autoimmune disease, those receiving
 231 immunosuppressive drug therapy were more prone to severe outcomes than patients without
 232 such therapy. Potential reasons for the increased risk are the immunosuppressive medication itself

233 and the increased severity of the underlying disease that necessitated an immunosuppressive drug
234 therapy.

235 Among SARS-CoV-2 cases aged 20-69, the relative risk of death was higher in patients with chronic
236 renal or liver disease, diseases of myoneural junction and muscle, cerebral palsy or other paralytic
237 syndromes as well as in those with psychotic disorders, especially with prescription for the
238 psychiatric medication clozapine. The association of psychiatric disorders and high COVID-19
239 mortality has been explained only partly by the presence of other comorbidities. [17] Other
240 reasons such as socioeconomic factors, delays in treatment seeking and certain deficits in cellular
241 immunity in schizophrenia have been implicated.[17] Although it is known that clozapine causes
242 neutropenia and predisposes for pneumonia[18] its role in COVID-19 infections has not been asessed
243 broadly.

244 In our analysis, asthma, similarly to other chronic pulmonary diseases, was associated with higher
245 risk of hospitalisation and intensive care treatment, but it was not associated with higher
246 mortality. The association with hospitalisation was weaker but present even when asthma had
247 been treated in primary health care only, indicating a presumably less severe manifestation.
248 However, in a review and meta-analysis where 76 studies and more than 17 million SARS-CoV-2
249 positive patients were included, asthma was not associated with severe COVID-19.[19] In our
250 study, 4.7% of 20–69-year-old SARS-CoV-2 cases were recognized as having asthma although the
251 overall prevalence in Finland is twice as high.[16] Our estimate is likely to represent the risk for
252 more severe asthma. In line with that, recent treatment of asthma with corticosteroids, indicating
253 a more severe manifestation, has been reported to be associated with an increased risk of severe
254 COVID-19 in some studies.[7,11]

255 Although they are no typical predictors for other infectious diseases, it has been widely noted that
 256 metabolic syndrome and obesity as well as type 2 diabetes are strongly associated with higher
 257 risks of severe COVID-19.[11,12,14,20] In our study, type 2 diabetes, which often occurs together
 258 with obesity and metabolic syndrome, was associated with higher risks of severe COVID-19 and
 259 death than the more serious disease, type 1 diabetes. In addition to obesity, regulation of blood
 260 glucose levels might play a role in the progression of SARS-CoV-2 infections among type 2 diabetes
 261 patients as high blood sugar levels have been shown to predict severe COVID-19.[11,20,21]
 262 Moreover, type 2 diabetes has been identified as an prognostic factor for survival among COVID-
 263 19 cases requiring critical care especially among the middle-aged and younger.[14]

264 In many studies, hypertension has been more prevalent among severe or fatal COVID-19 cases
 265 than among milder cases.[22] Also in our study, hypertension was more prevalent among severe
 266 cases; while only 7% of the SARS-CoV-2 positive 20-69-year-old individuals were hypertensive,
 267 25%, 32% and 36% of the hospitalised, ICU and fatal 20-69-year-old cases suffered from
 268 hypertension. However, patients who had only hypertension were not at increased risk of severe
 269 outcomes according to our study; similar results have been obtained elsewhere.[23] Since
 270 hypertension is common in patients who also have other comorbidities predisposing for severe
 271 COVID-19, such as obesity, hypertension alone is not an appropriate indicator for the prioritisation
 272 of COVID-19 vaccination.

273 The main strength of our study is the utilisation of multiple national registers, across which
 274 individual-level data were linked via a unique personal identifier. Information on possible
 275 predisposing comorbidities could therefore be verified from different sources. Furthermore, not
 276 only the conditional risk of hospitalisation given SARS-CoV-2 infection but also the rate of SARS-

277 CoV2 infection could be evaluated based on our nationwide dataset. It is a strength of our study
 278 that the effect of comorbidities on the risk on severe COVID-19 was analysed among the 20-69-
 279 year-olds, excluding the elderly, which should be prioritised by their age in any case.

280 Our study is solely based on register data. Therefore, we could not address the severity of the
 281 underlying comorbidities to any further extent than what could be deduced from the recorded
 282 diagnoses and medications. Another weakness is the rather limited number of cases, although we
 283 included all laboratory-confirmed SARS-CoV-2 cases that had occurred by 24 February 2021.

284 During the study period, non-pharmaceutical interventions reduced the spread of COVID-19 in
 285 Finland considerably. Additionally, not all national registers are adequate for real-time use. For
 286 example, the Finnish Cancer Register is updated only annually, so that, at the time of the study,
 287 data from 2019 and beyond were not yet available. Therefore, we could not utilise more detailed
 288 data from the Cancer register on the status or treatment of cancers and, therefore, could not
 289 identify cases receiving immunosuppressive treatments. Furthermore, the national registers used
 290 to identify individuals with the specified comorbidities did not contain private or occupational
 291 health care visits. However, if any disease entitling a special reimbursement is diagnosed at those
 292 visits, it is registered in the SII and thus included in our dataset.

293 When vaccinations are targeted to prevent severe COVID-19 among those with very common
 294 comorbidities, the number of vaccine doses needed is reasonably high. However, if we assume
 295 that vaccination prevents all severe cases (e. g. hospitalisations) and that the whole population,
 296 without vaccinations, would eventually acquire SARS-CoV-2 infection, the number needed to
 297 vaccinate to prevent one COVID-19 hospitalisation would remain small even for common diseases:

298 only 4.8 (1293 infections / 271 hospitalisations; Table 2) for type 2 diabetes, 6.1 for sleep apnea
299 and 7.3 for asthma.

300 According to our study, vaccination of the more critically ill, for example, patients with heart
301 failure or type 1 diabetes, is in fact slightly less effective in preventing severe COVID-19: the risks
302 of hospitalisation, intensive care treatment and death, conditional on SARS-CoV-2 infection, are
303 lower than among type 2 diabetes patients. Furthermore, in Finland patients with heart failure
304 and type 1 diabetes were less likely to have a SARS-CoV-2 infection than patients with type 2
305 diabetes or asthma. These findings may follow from effective shielding of the most vulnerable,
306 i.e., strict adherence to the nonpharmaceutical interventions recommended. If in the future we
307 are in a similar situation again (new pathogen, vaccine scarcity), this kind of study is the key.

308 **Authors' contributions**

309 HS, TL (Tuija), TL (Toni) conceptualized the study. TL (Toni) and UB conducted the statistical
310 analysis. TL (Tuija) reviewed the literature. HS, TL (Tuija) and KA drafted the manuscript. All
311 authors discussed the results and contributed to the final manuscript.

312 **Conflicts of interest**

313 All authors declare no conflict of interest

314

315

References

- [1] CDC COVID-19 Response Team. Preliminary Estimates of the Prevalence of Selected Underlying Health Conditions Among Patients with Coronavirus Disease 2019 - United States, February 12-March 28, 2020. MMWR Morb Mortal Wkly Rep 2020;69(13):382-6.
- [2] CDC COVID-19 Response Team. Severe Outcomes Among Patients with Coronavirus Disease 2019 (COVID-19) - United States, February 12-March 16, 2020. MMWR Morb Mortal Wkly Rep 2020;69(12):343-6.
- [3] Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX et al. Clinical Characteristics of Coronavirus Disease 2019 in China. N Engl J Med 2020;382(18):1708-20.
- [4] Guan WJ, Liang WH, Zhao Y, Liang HR, Chen ZS, Li YM et al. Comorbidity and its impact on 1590 patients with COVID-19 in China: a nationwide analysis. Eur Respir J 2020;55(5):10.1183/13993003.00547,2020. Print 2020 May.
- [5] WHO. COVID-19 and NCD risk factors. ;2021(July 1,).
- [6] European Centre for Disease Prevention and Control. Risk factors and risk groups. ;2021(July 1,).
- [7] Reilev M, Kristensen KB, Pottegård A, Lund LC, Hallas J, Ernst MT et al. Characteristics and predictors of hospitalization and death in the first 11 122 cases with a positive RT-PCR test for SARS-CoV-2 in Denmark: a nationwide cohort. Int J Epidemiol 2020;49(5):1468-81.
- [8] Petrilli CM, Jones SA, Yang J, Rajagopalan H, O'Donnell L, Chernyak Y et al. Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: prospective cohort study. BMJ 2020;369:m1966.
- [9] Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW et al. Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. JAMA 2020;323(20):2052-9.
- [10] Kim L, Garg S, O'Halloran A, Whitaker M, Pham H, Anderson EJ et al. Risk Factors for Intensive Care Unit Admission and In-hospital Mortality Among Hospitalized Adults Identified through the US Coronavirus Disease 2019 (COVID-19)-Associated Hospitalization Surveillance Network (COVID-NET). Clin Infect Dis 2021;72(9):e206-14.
- [11] Williamson EJ, Walker AJ, Bhaskaran K, Bacon S, Bates C, Morton CE et al. Factors associated with COVID-19-related death using OpenSAFELY. Nature 2020;584(7821):430-6.
- [12] Docherty AB, Harrison EM, Green CA, Hardwick HE, Pius R, Norman L et al. Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: prospective observational cohort study. BMJ 2020;369:m1985.

- 349 [13] Goodall JW, Reed TAN, Ardissino M, Bassett P, Whittington AM, Cohen DL et al. Risk factors
350 for severe disease in patients admitted with COVID-19 to a hospital in London, England: a
351 retrospective cohort study. Epidemiol Infect 2020;148:e251.
- 352 [14] Dennis JM, Mateen BA, Sonabend R, Thomas NJ, Patel KA, Hattersley AT et al. Type 2 Diabetes
353 and COVID-19-Related Mortality in the Critical Care Setting: A National Cohort Study in England,
354 March-July 2020. Diabetes Care 2021;44(1):50-7.
- 355 [15] Bubar KM, Reinholt K, Kissler SM, Lipsitch M, Cobey S, Grad YH et al. Model-informed COVID-
356 19 vaccine prioritization strategies by age and serostatus. Science 2021;371(6532):916-21.
- 357 [16] Finnish Institute for Health and Welfare. Vaccination order and at-risk groups for COVID-19. .
358 ;2021(July 1,).
- 359 [17] Nemani K, Li C, Olfson M, Blessing EM, Razavian N, Chen J et al. Association of Psychiatric
360 Disorders With Mortality Among Patients With COVID-19. JAMA Psychiatry 2021;78(4):380-6.
- 361 [18] de Leon J, Ruan CJ, Verdoux H, Wang C. Clozapine is strongly associated with the risk of
362 pneumonia and inflammation. Gen Psychiatr 2020;33(2):e100183,100183. eCollection 2020.
- 363 [19] Booth A, Reed AB, Ponzo S, Yassaee A, Aral M, Plans D et al. Population risk factors for severe
364 disease and mortality in COVID-19: A global systematic review and meta-analysis. PLoS One
365 2021;16(3):e0247461.
- 366 [20] Stefan N, Birkenfeld AL, Schulze MB. Global pandemics interconnected - obesity, impaired
367 metabolic health and COVID-19. Nat Rev Endocrinol 2021;17(3):135-49.
- 368 [21] Hamer M, Gale CR, Batty GD. Diabetes, glycaemic control, and risk of COVID-19
369 hospitalisation: Population-based, prospective cohort study. Metabolism 2020;112:154344.
- 370 [22] Gold MS, Sehayek D, Gabrielli S, Zhang X, McCusker C, Ben-Shoshan M. COVID-19 and
371 comorbidities: a systematic review and meta-analysis. Postgrad Med 2020;132(8):749-55.
- 372 [23] Clark CE, McDonagh STJ, McManus RJ, Martin U. COVID-19 and hypertension: risks and
373 management. A scientific statement on behalf of the British and Irish Hypertension Society. J Hum
374 Hypertens 2021;35(4):304-7.

375

376

377

378

379 Table 1. Comorbidities identified as potential risk factors predisposing to severe COVID-19

Comorbidities	Classification system	Codes	Register
Organ or stem cell transplant	ICD-10	T86, Z94	1,2
Malignant neoplasms	ICD-10	C00–C43, C45–C80, C97, D05.1, D39	1,2
Hematologic malignancy	ICD-10	C81–C85, C88, C90–C96	1,2
Severe disorders of the immune system	ICD-10	D70.8, D80–D84, E31.00	2
Severe chronic renal disease	ICD-10	I12, I13, N00–N05, N07, N08, N11, N14, N18, N19, E10.2, E11.2, E14.2	1,2
Severe chronic liver disease	ICD-10	K70.2, K70.3, K70.4, K71–K74	2
Asthma	ICD-10	J45, J46	2, 3
	ICPC-2	R96	3
Chronic pulmonary disease	ICD-10	J41–J44, J47	2
Essential (Primary) Hypertension	ICD-10	I10	1, 3
	ICPC-2	K86	3
Hypertensive heart and/or renal disease, secondary hypertension	ICD-10	I11.9, I12, I13.1, I13.9, I15	1,2
Ischaemic heart diseases	ICD-10	I20–I25	1,2
Heart failure	ICD-10	I11.0, I13.0, I13.2, I50	1,2
Type 1 diabetes	ICD-10	E10	2, 3
	ICPC-2	T89	3
	ATC	A10A	4
Type 2 diabetes	ICD-10	E11, E13, E14	2,3
	ICPC-2	T90	3
	ATC	A10B	4
Diseases of myoneural junction and muscle	ICD10	G70–G73	2
Cerebral palsy and other paralytic syndromes	ICD10	G80–G83	2
Cerebrovascular diseases	ICD-10	I60–I69	2
Sleep apnea	ICD-10	G47.3	2, 3
Continuous positive airway pressure therapy	NCSP	WX723, WX780	2
Clozapine (prescription for)	ATC	N05AH02	4
Psychotic disorders	ICD-10	F20–F29	2, 3
	ICPC-2	P72	3
Autoimmune disease	ICD-10	D86, K50, K51, L40, M02, M05–M07, M13.9, M45, M46.0, M46.1, M46.9, M94.1	1, 2
Glucocorticoids (prescription for)	ATC	H02AB02, H02AB04, H02AB06, H02AB07	4
Biologic drugs (entitlement to reimbursement of medical expenses)	ATC	L04AA24, L04AB01, L04AB02, L04AB04, L04AB05, L04AB06, L04AC03, L04AC05, L04AC07, L04AC10, L04AC12, L04AC13, L04AC14, L04AC16, L04AC18	1
Biologic drugs, Tumor necrosis factor alpha (TNF-α) inhibitors (Prescription for)	ATC	L04AB	4
Biologic drugs, other (prescription for)	ATC	L01XC02, L04AA24, L04AA26, L04AA33, L04AC	4
Other Immunosuppressant drugs	ATC	L01BA01, L04AA06, L04AA10, L04AA13, L04AA18, L04AA29, L04AA37, L04AD01, L04AD02, L04AX01, L04AX03	4

380 ATC, Anatomical Therapeutic Chemical Classification System; ICD-10, International Statistical Classification
 381 of Diseases and Related Health Problems, tenth revision; ICPC-2, International Classification of Primary
 382 Care, second edition; NCSP, Nordic Nomesco Classification of Surgical Procedures.

383 Registers: 1, Special Reimbursement Register for Medicine Expenses; 2, Care Register for Health Care; 3,
 384 Register of Primary Health Care Visits; 4, Prescription Centre database.

385

386 Table 2. Baseline characteristics of cases of SARS-CoV-2 infection

	SARS-CoV-2 PCR-positive n = 52502 (100%)	Hospitalised n = 2743 (100%)	Hospitalised, ICU n = 538 (100%)	Fatal cases n = 801 (100%)
Age, years				
0–9	3447 (6.6%)	33 (1.2%)	0 (0%)	<5
10–19	6937 (13.2%)	28 (1%)	<5	0 (0%)
20–29	11168 (21.3%)	120 (4.4%)	11 (2%)	<5
30–39	9376 (17.9%)	242 (8.8%)	39 (7.2%)	6 (0.7%)
40–49	7773 (14.8%)	389 (14.2%)	75 (13.9%)	7 (0.9%)
50–59	6837 (13%)	627 (22.9%)	134 (24.9%)	26 (3.2%)
60–69	3574 (6.8%)	538 (19.6%)	144 (26.8%)	58 (7.2%)
70–79	1782 (3.4%)	470 (17.1%)	115 (21.4%)	183 (22.8%)
80–89	1151 (2.2%)	245 (8.9%)	17 (3.2%)	324 (40.4%)
90+	457 (0.9%)	51 (1.9%)	0 (0%)	195 (24.3%)
20–29	38728 (73.8%)	1916 (69.9%)	403 (74.9%)	98 (12.2%)
Total	52502 (100%)	2743 (100%)	538 (100%)	801 (100%)
Sex				
male	26881 (51.2%)	1516 (55.3%)	353 (65.6%)	398 (49.7%)
female	25621 (48.8%)	1227 (44.7%)	185 (34.4%)	403 (50.3%)
Number of comorbidities, all ages				
0	37407 (71.2%)	808 (29.5%)	131 (24.3%)	45 (5.6%)
1	9200 (17.5%)	797 (29.1%)	150 (27.9%)	166 (20.7%)
2	3771 (7.2%)	594 (21.7%)	139 (25.8%)	272 (34%)
3+	2124 (4%)	544 (19.8%)	118 (21.9%)	318 (39.7%)
Number of comorbidities, 20–69-year-olds				
0	27940 (72.1%)	696 (36.3%)	112 (27.8%)	13 (13.3%)
1	7236 (18.7%)	589 (30.7%)	105 (26.1%)	29 (29.6%)
2	2429 (6.3%)	351 (18.3%)	97 (24.1%)	29 (29.6%)
3+	1123 (2.9%)	280 (14.6%)	89 (22.1%)	27 (27.6%)
Comorbidities, 20–69-year-olds				
Organ or stem cell transplant	59 (0.2%)	21 (1.1%)	9 (2.2%)	<5
Malignant neoplasms	596 (1.5%)	87 (4.5%)	18 (4.5%)	18 (18.4%)
Hematologic malignancy	112 (0.3%)	23 (1.2%)	7 (1.7%)	<5
Severe disorders of the immune system	36 (0.1%)	10 (0.5%)	<5	0 (0%)
Severe chronic renal disease	152 (0.4%)	53 (2.8%)	17 (4.2%)	9 (9.2%)
Severe chronic liver disease	65 (0.2%)	9 (0.5%)	<5	<5
Asthma	1823 (4.7%)	249 (13%)	77 (19.1%)	11 (11.2%)
Chronic pulmonary disease	122 (0.3%)	38 (2%)	12 (3%)	6 (6.1%)
Primary or secondary hypertension	2585 (6.7%)	455 (23.7%)	131 (32.5%)	35 (35.7%)
Essential (primary) hypertension alone	669 (1.7%)	54 (2.8%)	9 (2.2%)	<5
Secondary hypertension (hypertensive heart and/or renal disease) alone	878 (2.3%)	170 (8.9%)	63 (15.6%)	19 (19.4%)
Ischaemic heart diseases	498 (1.3%)	103 (5.4%)	33 (8.2%)	11 (11.2%)
Heart failure	66 (0.2%)	12 (0.6%)	<5	<5
Type 1 diabetes	312 (0.8%)	28 (1.5%)	5 (1.2%)	<5

Type 2 diabetes	1293 (3.3%)	271 (14.1%)	87 (21.6%)	25 (25.5%)
Diseases of myoneural junction and muscle	56 (0.1%)	16 (0.8%)	7 (1.7%)	<5
Cerebral palsy and other paralytic syndromes	70 (0.2%)	11 (0.6%)	<5	<5
Cerebrovascular diseases	247 (0.6%)	54 (2.8%)	15 (3.7%)	8 (8.2%)
Sleep apnea	1403 (3.6%)	230 (12%)	68 (16.9%)	11 (11.2%)
Psychotic disorders	60 (0.2%)	11 (0.6%)	<5	<5
Clozapine (prescription for)	362 (0.9%)	54 (2.8%)	5 (1.2%)	9 (9.2%)
Glucocorticoids (prescription for)	2989 (7.7%)	385 (20.1%)	84 (20.8%)	26 (26.5%)
Biologic drugs (entitlement to reimbursement of medical expenses)	163 (0.4%)	14 (0.7%)	<5	0 (0%)
Biologic drugs, Tumor necrosis factor alpha (TNF- α) inhibitors (prescription for)	135 (0.3%)	9 (0.5%)	<5	0 (0%)
Biologic drugs, other (prescription for)	50 (0.1%)	8 (0.4%)	<5	0 (0%)
Other Immunosuppressant drugs	486 (1.3%)	69 (3.6%)	16 (4%)	<5
Autoimmune disease with immunosuppressive drug therapy	709 (1.8%)	94 (4.9%)	18 (4.5%)	8 (8.2%)
Autoimmune disease without immunosuppressive drug therapy	651 (1.7%)	51 (2.7%)	13 (3.2%)	<5

387

388

389

390

391

392

393

394 Table 3a. Relative risk of severe COVID-19 (hospitalisation, ICU and death) in cases of SARS-Cov-2 infection
 395 by age group and sex. The reference category is 60–69-year-old females. Relative risks are presented in
 396 terms of odds ratios (OR) and their 95% confidence intervals (CI 95%).

OR (95% CI)						
	Hospitalised		Hospitalised, ICU		Fatal cases	
	Crude	Adjusted for sex	Crude	Adjusted for sex	Crude	Adjusted for sex
Age, years						
0–9	0.1 (0, 0.1)	0.1 (0, 0.1)	0 (0, 0)	0 (0, 0)	0 (0, 0.1)	0 (0, 0.1)
10–19	0 (0, 0)	0 (0, 0)	0 (0, 0)	0 (0, 0)	0 (0, 0)	0 (0, 0)
20–29	0.1 (0, 0.1)	0.1 (0, 0.1)	0 (0, 0)	0 (0, 0)	0 (0, 0)	0 (0, 0)
30–39	0.1 (0.1, 0.2)	0.1 (0.1, 0.2)	0.1 (0.1, 0.1)	0.1 (0.1, 0.1)	0 (0, 0.1)	0 (0, 0.1)
40–49	0.3 (0.3, 0.3)	0.3 (0.3, 0.3)	0.2 (0.2, 0.3)	0.2 (0.2, 0.3)	0.1 (0, 0.1)	0.1 (0, 0.1)
50–59	0.6 (0.5, 0.6)	0.6 (0.5, 0.6)	0.5 (0.4, 0.6)	0.5 (0.4, 0.6)	0.2 (0.1, 0.4)	0.2 (0.1, 0.4)
60–69	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
70–79	2 (1.8, 2.3)	2 (1.8, 2.3)	1.6 (1.3, 2.1)	1.7 (1.3, 2.2)	6.9 (5.2, 9.5)	7.1 (5.3, 9.7)
80–89	1.5 (1.3, 1.8)	1.6 (1.3, 1.9)	0.4 (0.2, 0.6)	0.4 (0.2, 0.6)	23.7 (17.9, 32)	26.7 (20, 36)
90+	0.7 (0.5, 1)	0.8 (0.6, 1)	0 (0, 0)	0 (0, 0)	45.1 (33, 62.5)	55.8 (40.5, 77.9)
Sex						
Male		1.3 (1.2, 1.4)		1.9 (1.6, 2.3)		1.9 (1.6, 2.2)
Female		1.0 (ref)		1.0 (ref)		1.0 (ref)

397

398 Table 3b. Relative risk of severe COVID-19 (hospitalisation, ICU and death) in cases of SARS-Cov-2 infection by number of comorbidities and by individual
399 comorbidities, adjusted for age and sex. Relative risks are presented in terms of odds ratios (OR) and their 95% confidence intervals (CI 95%).

	OR (95% CI)					
	Hospitalised		Hospitalised, ICU		Fatal cases	
	Crude	Adjusted for age and sex	Crude	Adjusted for age and sex	Crude	Adjusted for age and sex
Number of comorbidities, all ages¹						
1	4.3 (3.9, 4.8)	2.4 (2.2, 2.7)	4.7 (3.7, 6)	2.5 (2, 3.2)	15.3 (11.1, 21.5)	2.6 (1.9, 3.8)
2	8.5 (7.6, 9.5)	3.5 (3.1, 4)	10.9 (8.6, 13.9)	4.5 (3.4, 5.8)	64.5 (47.5, 89.8)	4.4 (3.1, 6.3)
3+	15.6 (13.8, 17.6)	5.3 (4.6, 6.1)	16.7 (13, 21.5)	6.2 (4.6, 8.2)	146.2 (107.8, 203)	6.2 (4.4, 9)
Number of comorbidities, 20–69-year-olds¹						
1	3.5 (3.1, 3.9)	2.5 (2.2, 2.8)	3.7 (2.8, 4.8)	2.4 (1.8, 3.2)	8.6 (4.6, 17.2)	4.7 (2.5, 9.6)
2	6.6 (5.8, 7.6)	3.8 (3.3, 4.4)	10.3 (7.8, 13.6)	5.4 (4, 7.2)	26 (13.8, 51.7)	9.9 (5.1, 20.3)
3+	13 (11.1, 15.2)	6.6 (5.6, 7.7)	21.4 (16.1, 28.4)	9.5 (7, 12.9)	52.9 (27.8, 106.2)	16.1 (8.1, 33.4)
Comorbidities, 20–69-year-olds²						
Organ or stem cell transplant	10.7 (6.2, 18.1)	7.2 (4, 12.5)	17.5 (8, 34.1)	9.8 (4.3, 19.8)	6.9 (0.4, 31.6)	2.8 (0.2, 13.5)
Malignant neoplasms	3.4 (2.7, 4.3)	1.7 (1.4, 2.2)	3.1 (1.8, 4.8)	1.4 (0.9, 2.3)	14.8 (8.6, 24.3)	5.8 (3.3, 9.7)
Hematologic malignancy	5 (3.1, 7.8)	2.9 (1.7, 4.6)	6.4 (2.7, 12.9)	3.1 (1.3, 6.4)	15.2 (4.6, 37.1)	5.6 (1.6, 14.1)

Severe disorders of the immune system	7.4 (3.4, 14.9)	8 (3.5, 17.2)	2.7 (0.2, 12.6)	2.5 (0.1, 12.4)	0 (0, 5211.9)	0 (0, 4720.9)
Severe chronic renal disease	10.5 (7.5, 14.7)	6.7 (4.6, 9.6)	12.5 (7.2, 20.2)	6.1 (3.5, 10.2)	27.2 (12.5, 52.2)	9.4 (4.2, 18.8)
Severe chronic liver disease	3.1 (1.4, 6)	1.5 (0.7, 3)	4.6 (1.1, 12.5)	2.1 (0.5, 5.7)	26.9 (8.1, 67)	9.8 (2.9, 25.7)
Asthma	3.3 (2.9, 3.8)	3 (2.5, 3.4)	4.9 (3.8, 6.3)	4.3 (3.3, 5.6)	2.6 (1.3, 4.6)	2 (1, 3.7)
Chronic pulmonary disease	8.8 (6, 12.9)	3.7 (2.5, 5.5)	10.7 (5.5, 18.7)	3.8 (1.9, 6.7)	21.7 (8.3, 46.5)	5.4 (2, 11.9)
Primary or secondary hypertension	5.1 (4.5, 5.7)	2.3 (2.1, 2.7)	7 (5.7, 8.7)	2.9 (2.3, 3.7)	7.9 (5.1, 11.8)	2.2 (1.4, 3.4)
Essential (primary) hypertension alone	1.7 (1.3, 2.2)	0.8 (0.6, 1.1)	1.3 (0.6, 2.4)	0.6 (0.3, 1.1)	1.2 (0.2, 3.8)	0.5 (0.1, 1.5)
Secondary hypertension (hypertensive heart and/or renal disease) alone	5 (4.2, 5.9)	2.1 (1.7, 2.5)	8.5 (6.4, 11.2)	3.2 (2.4, 4.3)	10.6 (6.2, 17.1)	2.7 (1.6, 4.5)
Ischaemic heart diseases	5.2 (4.2, 6.5)	2 (1.6, 2.6)	7.3 (4.9, 10.3)	2.4 (1.6, 3.4)	9.9 (5, 17.9)	2.1 (1, 3.9)
Heart failure	4.3 (2.2, 7.7)	2.3 (1.1, 4.3)	3 (0.5, 9.6)	1.3 (0.2, 4.1)	6.1 (0.3, 28.1)	1.8 (0.1, 8.6)
Type 1 diabetes	1.9 (1.3, 2.8)	1.9 (1.2, 2.8)	1.6 (0.6, 3.4)	1.4 (0.5, 3.1)	1.3 (0.1, 5.7)	1 (0.1, 4.8)
Type 2 diabetes	5.8 (5, 6.6)	2.8 (2.4, 3.3)	8.5 (6.6, 10.8)	3.7 (2.8, 4.7)	10.1 (6.3, 15.7)	3.2 (2, 5.1)
Diseases of myoneural junction and muscle	7.7 (4.2, 13.6)	5.5 (2.9, 10.2)	13.8 (5.7, 28.7)	8.1 (3.2, 17.9)	14.9 (2.4, 48.7)	5.8 (0.9, 20.4)
Cerebral palsy and other paralytic syndromes	3.6 (1.8, 6.6)	3.3 (1.6, 6.2)	5.8 (1.8, 14.1)	4.9 (1.4, 12.4)	5.8 (0.3, 26.4)	4.6 (0.3, 22.6)
Cerebrovascular diseases	5.5 (4, 7.4)	2.5 (1.8, 3.4)	6.3 (3.6, 10.4)	2.4 (1.3, 4)	14.3 (6.3, 28)	3.8 (1.7, 7.6)
Sleep apnea	4.1 (3.6, 4.8)	2.2 (1.9, 2.6)	5.6 (4.3, 7.3)	2.7 (2, 3.5)	3.4 (1.7, 6.1)	1.2 (0.6, 2.2)
Psychotic disorders	3.4 (2.5, 4.6)	3.4 (2.5, 4.7)	1.3 (0.5, 2.9)	1.2 (0.4, 2.7)	11 (5.1, 20.8)	10.2 (4.6, 20)
Clozapine (prescription for)	4.3 (2.1, 8)	4.2 (2, 8)	3.3 (0.5, 10.6)	2.9 (0.5, 9.7)	6.7 (0.4, 31)	6.1 (0.3, 30.6)
Glucocorticoids (prescription for)	3.3 (2.9, 3.7)	2.6 (2.3, 2.9)	3.2 (2.5, 4.1)	2.4 (1.8, 3)	4.3 (2.7, 6.7)	3 (1.9, 4.7)
Biologic drugs (entitlement to reimbursement of medical expenses)	1.8 (1, 3)	1.7 (0.9, 2.9)	0.6 (0, 2.6)	0.5 (0, 2.4)	0 (0, 0.8)	0 (0, 76.3)

Biologic drugs, Tumor necrosis factor alpha (TNF- α) inhibitors (prescription for)	1.4 (0.6, 2.6)	1.4 (0.6, 2.6)	0.7 (0, 3.2)	0.7 (0, 3.3)	0 (0, 2.4)	0 (0, 462.3)
Biologic drugs, other (prescription for)	3.7 (1.6, 7.4)	2.9 (1.2, 5.9)	1.9 (0.1, 8.9)	1.4 (0.1, 6.5)	0 (0, 46.4)	0 (0, 0)
Other Immunosuppressant drugs	3.3 (2.5, 4.2)	2.5 (1.9, 3.3)	3.3 (1.9, 5.4)	2.4 (1.4, 3.9)	3.4 (1, 8.1)	2.2 (0.7, 5.3)
Autoimmune disease with immunosuppressive drug therapy	3 (2.4, 3.8)	2.3 (1.8, 2.9)	2.5 (1.5, 4)	1.8 (1.1, 2.8)	4.8 (2.1, 9.3)	3.1 (1.4, 6.1)
Autoimmune disease without immunosuppressive drug therapy	1.7 (1.2, 2.2)	1.2 (0.9, 1.6)	2 (1.1, 3.3)	1.4 (0.8, 2.4)	0.6 (0, 2.7)	0.4 (0, 1.7)

400
401
402

¹ Compared to the individuals without any of the comorbidities in question.
² Comparing the odds of each outcome with and without the comorbidities in question.

403

404

405 Table 4. Relative rates of SARS-Cov-2 infection and severe COVID-19 outcomes (hospitalisation, admission to ICU and death) in the Finnish population aged
406 20–29, based on a negative binomial regression model. The table presents rate ratios (RR) and 95% confidence intervals (CI 95%).

	RR (95% CI)			
	SARS-CoV-2 PCR-positive	Hospitalised	Hospitalised, ICU	Fatal cases
Age, years				
20–29	3.1 (2.9, 3.3)	0.4 (0.3, 0.5)	0.2 (0.1, 0.3)	0.1 (0, 0.3)
30–39	2.4 (2.3, 2.5)	0.8 (0.7, 0.9)	0.5 (0.4, 0.7)	0.2 (0.1, 0.3)
40–49	2.3 (2.2, 2.4)	1.2 (1, 1.3)	0.9 (0.8, 1.2)	0.3 (0.2, 0.4)
50–59	1.8 (1.8, 1.9)	1.3 (1.2, 1.5)	1.2 (1, 1.4)	0.5 (0.4, 0.7)
60–69	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Sex				
Male	1 (1, 1)	1.3 (1.2, 1.4)	1.2 (1.1, 1.3)	2.1 (1.8, 2.3)
Female	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Comorbidities, 20–69-year-olds¹				
Organ or stem cell transplant	1 (0.8, 1.3)	8.9 (5.5, 13.8)	20.7 (9.7, 38.6)	15.2 (0.8, 76.6)
Malignant neoplasms	1 (0.9, 1.1)	3.1 (2.3, 4)	3.6 (2.1, 5.7)	20.7 (10, 43.9)
Hematologic malignancy	1 (0.8, 1.2)	4.7 (2.9, 7.2)	7.6 (3.2, 15.3)	25.9 (7.2, 73.9)
Severe disorders of the immune system	0.9 (0.6, 1.2)	7.5 (3.7, 13.5)	4.7 (0.3, 20.8)	NA
Severe chronic renal disease	1 (0.8, 1.1)	8.3 (6, 11.3)	14.3 (8.2, 23.2)	46.4 (19, 108.7)
Severe chronic liver disease	0.9 (0.7, 1.1)	2.7 (1.3, 4.9)	4.8 (1.2, 12.7)	38.6 (10.8, 110.3)
Asthma	1.1 (1.1, 1.2)	4.8 (4, 5.9)	9.5 (7.1, 12.7)	9.8 (4.3, 22.1)
Chronic pulmonary disease	0.8 (0.7, 1)	4.6 (3.1, 6.6)	7.4 (3.9, 13)	19.5 (6.8, 50.1)
Primary or secondary hypertension	0.9 (0.9, 1)	3.5 (2.8, 4.2)	4.9 (3.8, 6.4)	7.5 (4, 15)
Essential (primary) hypertension alone	1 (0.9, 1.1)	1.8 (1.3, 2.4)	1.6 (0.8, 3)	2.2 (0.4, 8.2)
Secondary hypertension (hypertensive heart and/or renal disease) alone	0.8 (0.7, 0.9)	2.9 (2.3, 3.7)	5.6 (4, 7.7)	9.1 (4.4, 19.1)
Ischaemic heart diseases	1 (0.9, 1.1)	3.4 (2.6, 4.5)	5.2 (3.4, 7.7)	8.9 (3.8, 20.2)
Heart failure	0.7 (0.6, 0.9)	2.8 (1.5, 4.9)	2.4 (0.4, 7.4)	6.6 (0.4, 33.6)

Type 1 diabetes	0.8 (0.7, 0.9)	2.5 (1.6, 3.7)	2.6 (0.9, 5.8)	4.1 (0.2, 20.7)
Type 2 diabetes	1.1 (1.1, 1.2)	5.1 (4.1, 6.3)	7.9 (5.9, 10.5)	12.8 (6.5, 26.1)
Diseases of myoneural junction and muscle	0.9 (0.7, 1.1)	7 (4, 11.3)	17 (7.2, 34)	31.5 (4.9, 114.8)
Cerebral palsy and other paralytic syndromes	1 (0.7, 1.2)	5 (2.5, 8.7)	10.7 (3.3, 25.5)	19.9 (1.1, 100.2)
Cerebrovascular diseases	0.9 (0.8, 1.1)	4.2 (3, 5.8)	6 (3.3, 10)	17.9 (7, 43.1)
Sleep apnea	1 (0.9, 1)	3.7 (3, 4.6)	5.6 (4.1, 7.6)	6.1 (2.6, 13.7)
Psychotic disorders	0.6 (0.5, 0.7)	3.1 (2.2, 4.1)	1.7 (0.6, 3.7)	24.6 (10.2, 57.2)
Clozapine (prescription for)	0.5 (0.4, 0.7)	2.9 (1.5, 5)	3 (0.5, 9.5)	12.7 (0.7, 63.7)
Glucocorticoids (prescription for)	1.1 (1, 1.1)	3.9 (3.2, 4.7)	5.1 (3.8, 6.8)	11.5 (6, 23.2)
Biologic drugs (entitlement to reimbursement of medical expenses)	0.9 (0.8, 1.1)	2.4 (1.3, 4)	1 (0.1, 4.6)	NA
Biologic drugs, Tumor necrosis factor alpha (TNF- α) inhibitors (prescription for)	0.9 (0.8, 1.1)	1.9 (0.9, 3.5)	1.3 (0.1, 5.7)	NA
Biologic drugs, other (prescription for)	1 (0.8, 1.3)	4.7 (2.1, 8.9)	3.4 (0.2, 15)	NA
Other Immunosuppressant drugs	0.8 (0.8, 0.9)	3.3 (2.4, 4.3)	4.4 (2.5, 7.3)	7.9 (2.2, 22.3)
Autoimmune disease with immunosuppressive drug therapy	0.9 (0.8, 0.9)	3.1 (2.4, 4.1)	3.6 (2.1, 5.7)	11.4 (4.5, 27.2)
Autoimmune disease without immunosuppressive drug therapy	0.9 (0.8, 0.9)	1.9 (1.4, 2.6)	2.8 (1.5, 4.8)	1.6 (0.1, 8)

¹Comparing the rate of each outcome in individuals with the comorbidity in question to those without any of the specified comorbidities

407
408

409 Figure 1. Proportion (%) of fatal outcome in cases of SARS-CoV-2 infection by age group and number of
410 comorbidities.

411

412

413

414

Figure 2. Incidence (per 100 000 person-years) of the SARS-Cov-2 infection and COVID-19 outcomes by age group in the Finnish population. SARS-Cov-2 cases on the left side Y axis and the remainder on right side Y axis.

422 Figure 3. Predictors of severe COVID-19 (hospitalisation and ICU) in cases of SARS-Cov-2 infection. Odds
423 ratios (OR) and 95% confidence intervals (CI 95%).

424