

**Supplemental Material for** *Nonresponse adjustment using clinical and perioperative patient characteristics is critical for understanding post-discharge opioid consumption*

Chris J. Kennedy, PhD<sup>1,2</sup>; Jayson S. Marwaha, MD<sup>1,2</sup>; P. Nina Scalise, MD<sup>2</sup>; Kortney A.

Robinson, MD, MPH<sup>2</sup>; Brandon Booth, BS<sup>2</sup>; Aaron Fleishman, MPH<sup>2</sup>; Larry A. Nathanson, MD<sup>3</sup>; Gabriel A. Brat, MD, MPH<sup>1,2</sup>

<sup>1</sup>Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA

<sup>2</sup>Department of Surgery, Beth Israel Deaconess Medical Center, Boston, MA, USA

<sup>3</sup>Department of Emergency Medicine, Beth Israel Deaconess Medical Center, Boston, MA, USA

## **Supplemental Study Details**

### **Missing data imputation**

Missing values were imputed using generalized low-rank models (GLRM), a generalization of principal component analysis that linearly decomposes and reconstructs the original dataset while reducing random variation (noise) in the variable measurements.<sup>1</sup> GLRM hyperparameters were optimized via a grid search based on reconstruction error on a 25% holdout set.<sup>2</sup> The hyperparameter grid consisted of 75 combinations: number of components  $\in \{36, 38, 40\}$ , X regularization  $\in \{\text{none, quadratic, L1}\}$ , Y regularization  $\in \{\text{none, quadratic, L1}\}$ , X gamma  $\in \{0, 4, 8\}$ , Y  $\in \{0, 4, 8\}$ . The optimal hyperparameters were: components = 40, X regularization = quadratic, Y regularization = quadratic, X gamma = 8, Y gamma = 4. A performance comparison of GLRM-based imputation versus median/mode imputation is provided in Supplemental Table 5.

### **Targeted learning estimation**

The SuperLearner library for the outcome regression (prediction of MMEs consumed, among those prescribed opioids) was: the outcome mean, stratification on prescribed MMEs and discharge day MMEs, ordinary least squares (OLS), lasso, and random forest. The latter three algorithms were tested with all predictors and with predictors restricted to those linearly correlated with the outcome at a p-value  $< 0.1$ . The SuperLearner library for the propensity score estimation (measurement of opioid consumption) was: the outcome mean, ordinary least squares (OLS), logistic regression, lasso, random forest, and Bayesian additive regression trees (BART). The latter five algorithms were tested with all predictors and with predictors restricted to those linearly correlated with the outcome at a p-value  $< 0.1$ .

## **References**

1. Udell M. Generalized Low Rank Models. Boyd S, Mackey L, Van Roy B, eds. Published

online 2015.

<http://search.proquest.com/dissertations-theses/generalized-low-rank-models/docview/2507949861/se-2>

2. Kennedy CJ, Mark DG, Huang J, van der Laan MJ, Hubbard AE, Reed ME. Development of an ensemble machine learning prognostic model to predict 60-day risk of major adverse cardiac events in adults with chest pain. *medRxiv*. Published online 2021.

### **Supplemental Table 1.** SMS-to-web survey details.

#### *Web Survey Instrument*

1. On what part of your body was the surgery? (More than 1 may apply. Choose 1.)
  - a. Head or Neck (brain, skull, ears, face, mouth, thyroid, parathyroid, throat, carotid)
  - b. Back (spine, vertebrae)
  - c. Chest (lungs, breast, esophagus, heart)
  - d. Abdomen (stomach, intestines, appendix, gallbladder, colon, hernia, liver, pancreas, transplant, kidney, prostate, bladder)
  - e. Leg or Arm (toes, foot, ankle, knee, hip, fingers, hand, wrist, elbow, axilla/underarm, shoulder)
  - f. Genitals and Buttock (penis, vagina, abscess hemorrhoid, fistula, fissure, pilonidal cyst)
  - g. Blood Vessels (angiogram, aorta, vascular repair)

Opioids are pain medications such as oxycodone, Vicodin, Percocet, hydrocodone, Dilaudid, and tramadol. Questions about opioids refer to these types of medications.

2. If you were prescribed an opioid, have you stopped taking your prescription?
  - a. I was not prescribed an opioid pain medication
  - b. No, I am still taking the opioid medication
  - c. I am no longer taking the opioid pain medication (I may still be taking other pain medications)
  - d. I never filled the opioid prescription given to me

[If Q2 = c, skip to Q4. If Q2 = a or d, end immediately.]

3. May we text you back in a week to see if your pain has improved?
  - a. Yes
  - b. No

[Survey completed]

=====

[Questions 4 - 9 are repeated for each opioid-class drug prescribed, between 0 and 3 times.]

4. You were prescribed {number} {medication-name} {medication-strength} dose). Is this correct?
  - a. Yes
  - b. No

[If Q4 = Yes, skip to Q6]

5. Please tell us what drug, dose (mg), and number of pills you were prescribed.
  - a. [Free text input]
  
6. Did you receive any refills for this prescription?
  - a. Yes
  - b. No
  
7. Was this prescription a pill or liquid?
  - a. Pill
  - b. Liquid

[If Q7 = Liquid, skip to Q9]

8. How many of these pills do you have left?
  - a. [Free text input]

[Skip to Q10]

9. Please estimate the percent of liquid you have left.
  - a. [Free text input] (%)

=====

10. How would you rate your overall pain control after discharge home from the hospital?
  - a. Excellent
  - b. Manageable
  - c. Intolerable

11. How would you describe your pain?
  - a. More than expected
  - b. As expected
  - c. Less than expected
  
12. How well did your providers explain how much pain you experienced?
  - a. Very well
  - b. Reasonably well
  - c. Not well at all
  
13. Have you used other medications for pain including but not limited to: (Please check all that apply)
  - a. Tylenol (acetaminophen)
  - b. Aspirin
  - c. Motrin (ibuprofen)
  - d. Aleve (naproxen)
  - e. Neurontin (gabapentin)
  - f. Lyrica (pregabalin)
  - g. Other
  - h. None of the above

[If Q13 != Other, Skip to Q15]

14. Please List Other Medication
  - a. [Free text input]
  
15. Prior to surgery, I was taking medications for: (Please check all that apply)
  - a. Anxiety
  - b. Depression
  - c. Sleep
  - d. ADHD or ADD
  - e. Drug dependence
  - f. None of the above
  
16. For pain prior to surgery, were you taking
  - a. Opioids
  - b. Non-opioid medications
  
17. How did you manage your leftover opioid prescription?
  - a. Kept them in an unlocked space

- b. Kept them in a locked space
- c. Took them to dropboxes (police stations and pharmacies) and/or to collection days
- d. Flushed them
- e. Threw them in the garbage
- f. Combined them with coffee grounds, kitty litter or dirt and put in garbage
- g. Didn't have excess medication
- h. Took them to my doctor
- i. Other

*SMS Message Content*

Please help BIDMC fight the opioid crisis. Click on the link to fill out our survey on patient opioid use.

LINK.

Reply to this text with STOP to opt out of future communications.

*SMS Message Timing*

Text messages were sent at 1pm Eastern Time.

**Supplemental Table 2.** Summary listing of predictor variables.

Variable	Type	Unique values	Missing rate	Mode	Mean	Median	Min	Max
<b>Opioids</b>								
1 discharge_day_mme	Continuous	105	0.0%	0	3	0.0	0.00	65.0
2 opioid_drugs_prescribed	Integer	4	0.0%	1	1	1.0	0.00	3.0
3 prescribed_mme	Continuous	166	0.2%	0	140	75.0	0.00	7,680.0
<b>Demographics</b>								
4 age	Integer	78	0.0%	64	56	58.0	18.00	95.0
5 age_group	Ordinal int.	4	0.0%	(45,60]				
6 employment_status	Categorical	3	13.5%	Occupation				
7 gender	Categorical	2	0.0%	F				
8 hispanic_ind	Categorical	5	0.3%	N				
9 race_group	Categorical	5	0.0%	White				
<b>Comorbidities</b>								
10 alcohol	Ordinal int.	3	13.5%	Occasionally				
11 bmi	Continuous	1,784	0.3%	26.63	29	28.2	8.52	67.9
12 current_smoker	Binary	2	0.0%	0	0	0.0		
13 daily_narcotic_use	Binary	2	13.5%	0	0	0.0		
14 hx_opioid_abuse	Binary	2	0.0%	0	0	0.0		
15 recreational_drugs	Binary	2	13.5%	0	0	0.0		
16 tobacco	Ordinal int.	4	13.5%	Never used				
<b>Surgical</b>								
17 prim_surg_svc	Categorical	12	10.0%	ORTHO				
18 wound_class	Categorical	5	0.5%	CL				
<b>Anesthesia</b>								
19 anesth_type	Categorical	7	0.1%	GEN				
20 asa_class	Integer	5	0.1%	2	2	2.0	1.00	5.0
21 asa_class_e	Binary	2	0.0%	0	0	0.0		
22 pain_quality	Categorical	8	13.5%	None				
23 pain_score	Integer	11	13.5%	0	2	0.0	0.00	10.0
<b>Hospital</b>								
24 case_type	Categorical	3	0.1%	S				
25 ibax_adm_type	Categorical	3	0.3%	EL				
26 ibax_disch_disp	Categorical	5	0.3%	01				
27 inout_cd	Categorical	2	0.3%	O				
28 length_of_stay	Integer	16	0.0%	0	2	1.0	0.00	15.0
29 los_group	Ordinal int.	3	0.0%	Outpatient				
30 pat_type_bidmc	Categorical	3	0.1%	INP				
31 pat_type_sched	Categorical	4	0.1%	OP				
32 payor_grp	Categorical	16	0.3%	MR				
<b>Missing</b>								
33 miss_anesth_type	Binary	2	0.0%	0	0	0.0		
34 miss_asa_class	Binary	2	0.0%	0	0	0.0		
35 miss_bmi	Binary	2	0.0%	0	0	0.0		
36 miss_inout_cd	Binary	2	0.0%	0	0	0.0		
37 miss_prescribed_mme	Binary	2	0.0%	0	0	0.0		
38 miss_prim_surg_svc	Binary	2	0.0%	0	0	0.0		
39 miss_tobacco	Binary	2	0.0%	0	0	0.0		
40 miss_wound_class	Binary	2	0.0%	0	0	0.0		

**Supplemental Table 3.** Demographic comparison between phone and SMS survey responders.

<b>Characteristic</b>	<b>Value</b>	<b>Phone</b>	<b>SMS</b>	<b>P-value</b>
Age, years	18 - 45	22.6%	17.9%	0.0001
	46 - 60	29.9%	33.4%	
	61 - 70	28.3%	31.7%	
	71 - 98	19.2%	16.9%	
Sex*	Female	47.4%	54.3%	<0.0001
	Male	50.7%	45.7%	
Race	Asian	2.2%	2.5%	0.0617
	Black	9.1%	6.8%	
	Other	10.8%	10.8%	
	White	77.9%	79.9%	
Body mass index	< 18.5	0.9%	0.8%	0.5643
	18.5 - 24.9	23.0%	26.7%	
	25 - 29.9	27.9%	34.8%	
	30+	29.9%	37.7%	
	Missing	18.3%	0.1%	

\*The sex response option “other” was included in the phone survey instrument but not in the SMS-recruited version of the survey instrument, so that row is not included in this table.

**Supplemental Table 4.** Opioid consumption in MMEs for the top 10 surgical procedure groups by volume, examining the difference between nonresponse bias adjusted and unadjusted survey results, and comparing to the earlier phone survey.

Surgical approach	n	Median				75th percentile			
		Prescribed	Survey Response	Targeted	Phone	Prescribed	Survey Response	Targeted	Phone
Thoracic/Lumbar/Sacral Discectomy/Laminectomy/Fusion	393	300	157	173	217	450	432	449	435
Anterior Cervical Discectomy and Fusion	150	225	80	134	208	300	225	261	318
Thoracic/Lumbar/Sacral Microdiscectomy	128	225	30	33	142	228	150	149	225
Sternotomy	464	160	30	2	22	300	157	157	200
Mammoplasty	173	90	16	0	82	150	65	56	155
Inguinal Hernia	289	45	7	9	11	75	37	42	48
Cholecystectomy	382	37	15	16	15	75	37	37	38
Appendectomy	151	37	7	11	22	56	35	33	46
Carpal Tunnel Release	145	0	0	0	10	37	37	33	38
Thyroidectomy	122	0	0	0	0	37	20	10	22



**Supplemental Table 5.** Imputation performance comparison of generalized low-rank models with median/mode imputation. The average percent reduction in error across all variables was 46%.

<b>Variable</b>	<b>Missingness (%)</b>	<b>Error GLRM</b>	<b>Error Std.</b>	<b>Percent reduction</b>
employment_status	13.5	0.04	0.34	89.5
alcohol	13.5	0.37	0.37	0.0
daily_narcotic_use	13.5	0.77	0.25	-204.0
recreational_drugs	13.5	0.36	0.36	0.0
tobacco2	13.5	0.91	0.45	-101.1
pain_quality	13.5	0.09	0.34	72.2
pain_score	13.5	1.03	3.97	74.2
prim_surg_svc	10.0	0.15	0.70	78.0
wound_class	0.5	0.07	0.28	75.8
hispanic_ind	0.3	0.10	0.16	38.2
ibax_adm_type	0.3	0.01	0.15	91.5
ibax_disch_disp	0.3	0.04	0.31	86.5
inout_cd	0.3	0.73	0.69	-5.5
payor_grp	0.3	0.24	0.75	68.1
bmi	0.3	1.09	6.83	84.0
prescribed_mme	0.2	92.9	241.3	61.5
asa_class	0.1	0.08	0.86	91.1
anesth_type	0.1	0.04	0.26	83.5
case_type	0.1	0.02	0.09	82.8
pat_type_bidmc	0.1	0.00	0.52	100.0
pat_type_sched	0.1	0.02	0.61	97.1