1 **Brief Report**

2 **Title:** Prospective cohort study of workers diagnosed with COVID-19 and subsequent
3 unemployment

4 **Running title:** COVID-19 and unemployment

5

6 **Authors:**

7 Tomohisa Nagata¹, Masako Nagata¹, Ayako Hino², Seiichiro Tateishi³, Akira Ogami⁴,
8 Mayumi Tsuji⁵, Shinya Matsuda⁶, Yoshihisa Fujino⁷, Koji Mori¹, MD, PhD; for the
9 CORoNaWork project

10 **Author contributions:**

11 T.N. created the questionnaire, conducted the analysis, and wrote the manuscript. M.N., A.H.,
12 S.T., A.O., M.T., S.M., and K.M. reviewed drafts of the manuscript, provided advice on the
13 interpretation of the data, and procured funding for the research. Y.F. carried out the overall
14 survey planning, created the questionnaire, and reviewed drafts of the manuscript.

15 **Affiliations:**

16 ¹Department of Occupational Health Practice and Management, Institute of Industrial
17 Ecological Sciences, University of Occupational and Environmental Health, Japan

18 ²Department of Mental Health, Institute of Industrial Ecological Sciences, University of
19 Occupational and Environmental Health, Japan

20 ³Department of Occupational Medicine, School of Medicine, University of Occupational and

21 Environmental Health, Japan

22 ⁴Department of Work Systems and Health, Institute of Industrial Ecological Sciences,

23 University of Occupational and Environmental Health, Japan

24 ⁵Department of Environmental Health, School of Medicine, University of Occupational and

25 Environmental Health, Japan

26 ⁶Department of Public Health, School of Medicine, University of Occupational and

27 Environmental Health, Japan

28 ⁷Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences,

29 University of Occupational and Environmental Health, Japan

30

31 **Correspondence:**

32 Tomohisa Nagata, MD, PhD

33 Department of Occupational Health Practice and Management, Institute of Industrial

34 Ecological Sciences, University of Occupational and Environmental Health, Japan, 1-1

35 Iseigaoka, Yahatanishi-ku, Kitakyushu 807-8555, Japan.

36 email: tomohisa@med.uoeh-u.ac.jp; Tel: +81-93-603-1611; Fax: +81-93-603-2155

37

38

39 Abstract

40 **Objectives:** The purpose of this study was to investigate the relationships of workers being
41 diagnosed with coronavirus disease 2019 (COVID-19) and being identified as close contacts
42 of infected persons with unemployment in Japan.

43 **Methods:** This was a prospective cohort study using questionnaires about COVID-19
44 administered to Japanese workers. A baseline survey conducted on December 22–25, 2020,
45 was used to determine history of being diagnosed with COVID-19 or being identified as a
46 close contact of an infected person. Unemployment since the baseline survey was ascertained
47 with a follow-up survey on February 18 and 19, 2021. The odds ratios of unemployment were
48 estimated using a multilevel logistic model with adjusted covariates nested in prefecture of
49 residence.

50 **Results:** Women ($n = 8,771$) accounted for 44% of the total sample ($n = 19,941$), and the
51 mean age was 48.0 years. In terms of unemployment because of negative reasons, the
52 multivariate analysis showed that the odds ratio of unemployment associated with being
53 diagnosed with COVID-19 was 2.18 (95% CI: 1.05–4.55) and that the odds ratio associated
54 with being identified as a close contact was 0.93 (95% CI: 0.29–2.95).

55 **Conclusions:** There is an association between workers being diagnosed with COVID-19 and
56 unemployment. Occupational health professionals should follow up closely with workers
57 diagnosed with COVID-19 after they return to work to prevent them from becoming

58 unemployed against their will.

59 **Keywords:** unemployment, COVID-19, SARS-CoV-2, workplace, Japan

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78 **Introduction**

79 The spread of coronavirus disease 2019 (COVID-19) has become a global issue. The
80 COVID-19 pandemic has had a profound impact not only on people's daily lives, but also on
81 the economy. The introduction of strong restrictive measures such as lockdowns in response
82 to the pandemic has influenced the global economy, including by leading to an increase in
83 unemployment rates worldwide.¹ In Japan, the total unemployment rate gradually increased
84 as COVID-19 spread, reaching 2.9% for men and women together in February 2021.²

85 Patients diagnosed with COVID-19 and individuals identified as close contacts of
86 infected persons are isolated from those who are not sick. When these isolated people are
87 workers, they are not able to go to work for a certain period of time. In Japan, workers with
88 COVID-19 cannot go to work until at least 10 days have passed since symptom onset, at least
89 72 hours have passed since resolution of fever, and there has been an improvement in other
90 symptoms.³ However, it is unclear how the subsequent continued employment of workers
91 who are diagnosed with COVID-19 or identified as close contacts of infected persons is.

92 The purpose of this study was to investigate the relationship between workers being
93 diagnosed with COVID-19 or identified as close contacts of infected persons and
94 unemployment in Japan.

95 **Materials and methods**

96 This prospective cohort study about COVID-19 among Japanese workers was conducted

under the CORoNaWork (Collaborative Online Research on the Novel-coronavirus and Work) Project. The details of the study protocol have been described elsewhere.⁴ Briefly, we administered a baseline questionnaire on December 22–25, 2020, and a follow-up questionnaire on February 18 and 19, 2021, when Japan was in its third wave of the pandemic.

For the baseline survey, on December 22–25, 2020, a total of 33,087 workers were recruited throughout Japan from 605,381 randomly selected panelists who were registered with an Internet survey company. The inclusion criteria for participants were currently working and being aged 20–65 years. This study did not invite healthcare professionals or caregivers to participate. We used cluster sampling with stratification by sex, job type, and region. We excluded 6,051 invalid responses because of response time < 6 minutes, body weight < 30 kg or height < 140 cm, inconsistent answers to similar questions, and incorrect answers to questions intended to identify fraudulent responses. We distributed the follow-up questionnaire to the 27,036 people with valid responses to the baseline questionnaire on February 18 and 19, 2021. In total, 19,941 participants completed both questionnaires (follow-up rate = 73.8%).

The present study was approved by the Ethics Committee of the University of Occupational and Environmental Health, Japan (reference No. R2-079 and R3-006). Informed consent was obtained from all participants.

116 Explanatory variables

117 The baseline survey asked two questions about the participants' COVID-19 diagnostic history.

118 First, the participants were asked, "Have you had COVID-19?" with the response options of

119 *yes* or *no*. If the answer to this question was *yes*, the respondent was considered a worker

120 being diagnosed with COVID-19. If the answer to the question was *no*, to determine whether

121 they had been identified as a close contact of an infected person, the respondent was asked,

122 "Have you been identified as a close contact of a person diagnosed with COVID-19?" with

123 the response options of *yes* or *no*. Workers answering *yes* to this question were identified as

124 close contacts. Workers reporting that they had been diagnosed with COVID-19 or identified

125 as close contacts should have been classified in this way according to public health center

126 instructions, so we believe that these questions allowed us to identify infected and exposed

127 people with a high degree of accuracy.

128 Outcomes

129 Unemployment was ascertained in the following way. First, the baseline survey included only

130 people who were employed at the time of response. In the follow-up survey, in response to

131 the question "Have you changed your place of work since December 2020?" respondents

132 were asked to select one of the following six options: *no change, I was transferred to another*

133 *company, I resigned and got a new job right away, I stopped working and was unemployed*

134 *for a period of time but am now working, I stopped working and started a business (e.g.,*

managing a company, running a sole proprietorship, or engaging in self-employment), and *I stopped working and am currently not working (including job seeking)*. We wanted to identify unemployment because of negative reasons as the primary outcome to achieve the research objectives, so we defined unemployment using the responses of *I stopped working and was unemployed for a period of time but am now working* and *I stopped working and am currently not working (including job seeking)*. As a secondary outcome, we also considered unemployment regardless of the reason, comparing respondents who chose *no change* or *I was transferred to another company* with those who chose any other option.

Control variables

We retrieved the following data from the baseline survey for inclusion as control variables: age, sex, annual household income, educational attainment, marital status, job type, self-rated health, and region.

Age was treated as a continuous variable. Education was classified into five as follow: junior high school, high school, junior college or technical school, university, and graduate school.

Yearly household income was classified into four as follow: less than 2.50 million Japanese yen (JPY), 2.50–3.75 million JPY, 3.75–5.25 million JPY, and greater than 5.25 million JPY.

Marital status was classified into three as follow: married, divorced or widowed, and never married. Job type was classified into three categories: mainly desk work, work mainly involving interpersonal communication, and mainly physical work. The following question

was asked to assess self-rated health: “Please choose the option that best describes your overall health condition, including both physical and mental health” with the following five response options: *extremely good, very good, good, fair, and bad*.

In addition, prefecture of residence was included as a community-level variable.

Statistical analysis

The odds ratios (ORs) of unemployment (primary and secondary outcome) for infected workers and persons identified as close contacts of infected persons were estimated using a multilevel logistic model nested in prefectures of residence. The multivariate model was adjusted for age and sex, educational attainment, annual household income, marital status, job type, and self-rated health. The incidence rate of COVID-19 by prefecture was included as a prefecture-level variable. *P*-values less than 0.05 were considered statistically significant. All analyses were conducted using Stata (Stata Statistical Software release SE16.1; StataCorp LLC, College Station, TX, USA).

Results

The basic characteristics of the respondents are shown in Table 1. There were 8,771 women in the sample, accounting for 44% of the total ($n = 19,941$). The mean age was 48.0 years, with 454 (2.3%) reporting experiencing unemployment because of negative reasons and 725 (3.6%) reporting experiencing unemployment for any reason.

Table 2 shows the associations of being diagnosed with COVID-19 and being identified as a

close contact of someone with COVID-19 with unemployment. For the primary outcome of unemployment because of negative reasons, the multivariate analysis showed that the OR of unemployment associated with being diagnosed with COVID-19 was 2.18 (95% CI: 1.05–4.55) and that the OR associated with being identified as a close contact was 0.93 (95% CI: 0.29–2.95).

For the secondary outcome (unemployment regardless of the reason), the multivariate analysis showed that the OR of unemployment associated with being diagnosed with COVID-19 was 3.47 (95% CI: 2.09–5.74) and that the OR associated with being identified as a close contact was 1.19 (95% CI: 0.52–2.73).

Discussion

We evaluated the relationships of workers having COVID-19 and being identified as close contacts with unemployment. We observed an association between workers being diagnosed with COVID-19 and unemployment but not between being identified as close contacts and unemployment.

There are several possible reasons why workers diagnosed with COVID-19 subsequently became unemployed. First, workers with COVID-19 may have recovered once but then been unable to work as a result of suffering from the long-term effects of COVID-19. A recent longitudinal prospective cohort study of post-COVID-19 patients found that 30% had at least one symptom persisting for nine months or longer.⁵ The sequelae of COVID-19

are referred to as “post-COVID syndrome” or “COVID long haul syndrome.”⁶ However, in our analysis, the relationship between being diagnosed with COVID-19 and unemployment remained significant even after adjusting for self-rated health; thus, poor health may not be the reason for unemployment.

Second, workers who had COVID-19 may have resigned as a result of feeling uncomfortable in the workplace. Workers with COVID-19 have to be away from work for about two weeks. If coworkers are in close contact with the infected people, the coworkers must also take time off work. Workers with COVID-19 may have felt bad about having a major impact on their workplace and therefore have resigned. This situation may also have led to bullying and harassment in the workplace. Such harassment has been reported to be a common occurrence among healthcare workers⁷, but it can also occur in the general workforce. Workplace bullying, in addition to increasing mental distress and even work-related suicide, also has socioeconomic consequences related to sick leave and unemployment.⁸ Although workers who were identified as close contacts of infected people also needed to take a certain amount of time off work, they did not spread the infection to others in the workplace; this may have prevented workers identified as close contacts from being forced into resigning.

Some limitations of this study warrant mention. First, exposure was assessed by asking about the existence of a history of COVID-19 in the baseline survey (at the end of

211 December 2020), but it was not clear when respondents had been diagnosed with COVID-19
 212 from January 16, 2020 (when the first case of COVID-19 was reported in Japan) and
 213 December 2020 (when the baseline survey was conducted). However, the number of people
 214 in Japan diagnosed with COVID-19 was overwhelmingly higher after October 2020, so it is
 215 very likely that most respondents in this study who had COVID-19 were diagnosed from
 216 October to December 2020. Second, information on past diagnosis was self-reported.
 217 However, almost all COVID-19 patients and people identified as close contacts are diagnosed
 218 after undergoing polymerase chain reaction testing. This process is rigorously followed to
 219 prevent the spread of infection, so the information is likely to be fairly accurate. Third, we
 220 did not investigate the reasons for unemployment except for starting their own business. As a
 221 result, we were unable to fully distinguish between cases in which workers resigned
 222 voluntarily and those in which their employment was terminated against their will. Future
 223 research should conduct analyses using survey data that allow for a thorough assessment of
 224 the reasons for unemployment.

225 In conclusion, there is an association between workers being diagnosed with
 226 COVID-19 and unemployment. Occupational health professionals need to follow up closely
 227 with workers diagnosed with COVID-19 after their return to work to prevent them from
 228 becoming unemployed against their will.

229

Acknowledgments

This study was supported and partly funded by the University of Occupational and Environmental Health, Japan; General Incorporated Foundation (Anshin Zaidan); The Development of Educational Materials on Mental Health Measures for Managers at Small-sized Enterprises; Health, Labour and Welfare Sciences Research Grants; Comprehensive Research for Women's Healthcare (H30-josei-ippan-002); Research for the Establishment of an Occupational Health System in Times of Disaster (H30-roudou-ippan-007), scholarship donations from Chugai Pharmaceutical Co., Ltd., the Collabo-Health Study Group, and Hitachi Systems, Ltd.

The current members of the CORoNaWork Project, in alphabetical order, are as follows: Dr. Yoshihisa Fujino (present chairperson of the study group), Dr. Akira Ogami, Dr. Arisa Harada, Dr. Ayako Hino, Dr. Hajime Ando, Dr. Hisashi Eguchi, Dr. Kazunori Ikegami, Dr. Kei Tokutsu, Dr. Keiji Muramatsu, Dr. Koji Mori, Dr. Kosuke Mafune, Dr. Kyoko Kitagawa, Dr. Masako Nagata, Dr. Mayumi Tsuji, Ms. Ning Liu, Dr. Rie Tanaka, Dr. Ryutaro Matsugaki, Dr. Seiichiro Tateishi, Dr. Shinya Matsuda, Dr. Tomohiro Ishimaru, and Dr. Tomohisa Nagata. All members are affiliated with the University of Occupational and Environmental Health, Japan.

Disclosure

Ethical approval: This study was approved by the ethics committee of the University of

Occupational and Environmental Health, Japan (reference No. R2-079 and R3-006) .

Informed Consent: Informed consent was obtained in the form of the website.

Registry and the Registration No. of the study/Trial: N/A

Animal Studies: N/A

Conflict of Interest: The authors declare no conflicts of interest associated with this manuscript.

Data Availability Statement: The data that support the findings of this study are available from the corresponding author, Tomohisa Nagata, upon reasonable request.

References

1. International Labor Organization. News.

[https://www.ilo.org/global/about-the-ilo/newsroom/news/WCMS_738742/lang--en/index.](https://www.ilo.org/global/about-the-ilo/newsroom/news/WCMS_738742/lang--en/index.htm)

[htm](https://www.ilo.org/global/about-the-ilo/newsroom/news/WCMS_738742/lang--en/index.htm). Accessed July 6, 2021.

2. The Japan Institute for Labour Policy and Training.

<https://www.jil.go.jp/kokunai/statistics/covid-19/c06.html#c06-2>. Accessed July 6, 2021.

3. Nangaku M, Kadowaki T, Yotsuyanagi H, et al. The Japanese Medical Science Federation

COVID-19 Expert Opinion English Version. JMA J. 2021; 4(2): 148-162. doi:

268 10.31662/jmaj.2021-0002.

269 4. Fujino Y, Ishimaru T, Eguchi H, et al. Protocol for a Nationwide Internet-based Health

270 Survey of Workers During the COVID-19 Pandemic in 2020. Journal of UOEH. 2020;

271 43(2), 217–225.

272 5. Logue JK, Franko NM, McCulloch DJ, et al. Sequelae in Adults at 6 Months After

273 SARS-CoV-2 infection. JAMA Netw Open. 2021;4:e210830.

274 6. Vanichkachorn G, Newcomb R, Cowl CT, et al. Post-COVID-19 Syndrome (Long Haul

275 Syndrome): Description of a Multidisciplinary Clinic at Mayo Clinic and Characteristics

276 of the Initial Patient Cohort. Mayo Clin Proc. 2021; 96(7): 1782-1791. doi:

277 10.1016/j.mayocp.2021.04.024.

278 7. Dye TD, Alcantara L, Siddiqi S, et al. Risk of COVID-19-related bullying, harassment

279 and stigma among healthcare workers: an analytical cross-sectional global study. BMJ

280 Open. 2020; 10(12): e046620. doi: 10.1136/bmjopen-2020-046620.

281 8. Sansone RA, Sansone LA. Workplace bullying: a tale of adverse consequences. Innov

282 Clin Neurosci. 2015; 12(1-2): 32-7.

283

284

285

286

Table 1. Basic characteristics of study respondents

	Workers who were not infected	Workers identified as close contacts ^a	Workers diagnosed with COVID-19
Number of respondents	19,646	141	154
Age, mean (SD)	48.0 (10.1)	45.7 (10.6)	43.8 (11.5)
Men	10,998 (56.0%)	87 (61.7%)	85 (55.2%)
Education			
Junior high or high school	5300 (27.0%)	23 (16.3%)	37 (24.0%)
Vocational school, junior college, or technical school	4513 (23.0%)	37 (26.2%)	35 (22.7%)
University	8742 (44.5%)	70 (49.6%)	68 (44.2%)
Graduate school	1091 (5.6%)	11 (7.8%)	14 (9.1%)
Annual income (million JPY)			
< 2.50	4183 (21.3%)	17 (12.1%)	33 (21.4%)
≥ 2.50 and < 3.75	5388 (27.4%)	34 (24.1%)	36 (23.4%)
≥ 3.75 and < 4.99	4750 (24.2%)	35 (24.8%)	36 (23.4%)
≥ 4.99	5325 (27.1%)	55 (39.0%)	49 (31.8%)
Marital status			
Married	11,004 (56.0%)	85 (60.3%)	96 (62.3%)
Widowed/divorced	2012 (10.2%)	10 (7.1%)	14 (9.1%)
Unmarried	6630 (33.7%)	46 (32.6%)	44 (28.6%)
Job type			
Mainly desk work	10,135 (51.6%)	61 (43.3%)	72 (46.8%)
Work mainly involving interpersonal communication	4847 (24.7%)	44 (31.2%)	46 (29.9%)
Mainly physical work	4664 (23.7%)	36 (25.5%)	36 (23.4%)
Self-rated health			
Extremely good	542 (2.8%)	6 (4.3%)	22 (14.3%)
Very good	1938 (9.9%)	19 (13.5%)	35 (22.7%)
Good	7374 (37.5%)	50 (35.5%)	42 (27.3%)
Fair	7270 (37.0%)	49 (34.8%)	45 (29.2%)
Bad	2522 (12.8%)	17 (12.1%)	10 (6.5%)
Unemployment			
Unemployment because of negative reasons	443 (2.3%)	3 (2.1%)	8 (5.2%)
Unemployment regardless of the reason	700 (3.6%)	6 (4.3%)	19 (12.3%)

^a Workers identified as close contacts of people who were diagnosed with COVID-19

Abbreviations: JPY, Japanese Yen

296

297

298

299

300

301 Table 2. Associations of being diagnosed with COVID-19 and being identified as a close contact of an infected person with unemployment

		Unemployment because of negative reasons							Unemployment regardless of the reason							
		Crude				Mutivariable			Crude				Mutivariable			
		OR	95% CI	<i>p</i> -value		OR	95% CI	<i>p</i> -value	OR	95% CI	<i>p</i> -value	OR	95% CI	<i>p</i> -value		
COVID-19																
Workers who were not infected	reference				reference				reference			reference				
Workers identified as close contacts ^a	0.93	0.29	2.93	0.898	0.93	0.29	2.95	0.899	1.17	0.52	2.67	0.703	1.19	0.52	2.73	0.684
Workers diagnosed with COVID-19	2.37	1.16	4.87	0.018	2.18	1.05	4.55	0.038	3.77	2.32	6.14	< 0.001	3.47	2.09	5.74	< 0.001

304 Note: Multivariable model included: sex, age, educational background, annual income, marital status, job type, and self-rated health.

305 Abbreviations: OR, odds ratio; CI, confidential interval

306 ^a Workers identified as close contacts of people who were diagnosed with COVID-19

307