

## Supplementary Materials

### 1. Substantive deviations from pre-registered protocol

The AR(1) covariance structure was used for all generalised estimating equation (GEE) models, as the most representative correlation structure between timepoints. The final GEE model was originally to include only the explanatory variables that were significant in three separate GEE models containing sociodemographic, COVID-19-related or behavioural explanatory variables, but was changed to contain all explanatory variables as the aim was to identify significant variables whilst adjusting for all other variables to guide health policy, rather than to build the best predictive model for a change in self-reported weight/BMI. GEE models containing sociodemographic, COVID-19-related and behavioural explanatory variables were therefore redundant and not computed. In the sensitivity analysis for a change in BMI, a change of 5% was used as the cut-off for an 'increase' or 'decrease' in BMI instead of a change in BMI categories because fully adjusted GEE models using the categorical change outcome did not compute due to zero 'all other' ethnicity participants self-reporting a categorical increase or decrease in BMI. The 5% cut-off reflected the 5% cut-off used in the sensitivity analysis for weight.

The pre-registered protocol in RQ1 stated that the proportions of the sample increasing or decreasing (i) weight and (ii) BMI between timepoints and mean change (with 95% confidence intervals) in (i) weight and (ii) BMI (as a continuous variable) in those increasing or decreasing between timepoints would be obtained via an unadjusted, unweighted GEE model. This was instead obtained using simple descriptive statistics.

The alcohol consumption explanatory variable was changed from the pre-registered Alcohol Use Disorder Identification Test for Consumption (AUDIT-C) score to a binary measure of weekly unit frequency due to an error in the survey that did not allow for an AUDIT-C score to be computed at the 3-months follow-up survey. The alcohol variable was changed to high-risk or low-risk alcohol consumption, based on UK government recommendations. The categorical baseline BMI explanatory variable was changed to a continuous baseline BMI predictor variable to increase power and simplify interpretation.

Bayes Factors were calculated for non-significant findings for both weight and BMI outcomes.

#### Bayes Factors Robustness checks

Robustness checks were conducted for Bayes Factors using alternative priors. For gender, a prior mean difference between -0.23kg and 6.1kg produced the same outcome of inconclusive evidence for an effect. For reduced physical activity, a mean difference within -0.390kg and 0.490kg produced the same outcome of inconclusive evidence for an effect for the same standard error.

## Testing linearity of logit in the binary sensitivity analysis models

Linearity of logit in the continuous explanatory variables was tested in the binary sensitivity analysis models for an increase/decrease in weight/BMI vs all other. Continuous explanatory variables were Box-Tidwell transformed. Continuous explanatory variables and transformed variables were then all entered into binary logistic GEE models for an increase/decrease in weight/BMI vs all other. The linearity of logit assumptions were violated in the 'Decrease weight vs all other' model for the baseline BMI, HFSS snacks intake, fruit and vegetable intake, and  $\Delta$ HFSS snacks change score continuous variables. In the 'Increase BMI vs all other' GEE model, linearity of logit assumptions were violated for the baseline BMI continuous variable. The above variables were replaced where assumptions were violated in univariate and fully adjusted models (Supplementary tables S8, S9, S10 and S11), with the following categorical alternatives:

- Baseline BMI:
  - 'Decrease weight vs all other': two categories based on WHO classifications; <25,  $\geq$ 25.
  - 'Increase BMI vs all other' (with <25 BMI): two categories based on WHO classifications; underweight, normal weight.
- HFSS snacks intake: the median binary cut-off as used in Figure 2.
- Fruit and vegetable intake: consuming a few portions per day vs all other.
- $\Delta$ HFSS snacks change score: increased intake vs all other.

## Detailed description of study explanatory variables

The full wording of questions can be found at: <https://osf.io/bja7g/>

Time-invariant explanatory variables are obtained from participants' baseline responses. Time-variant explanatory variables are collected at baseline, 3-months and 6-months surveys.

### Time-invariant

#### Socio-demographic

- **Gender** (2 levels) '*Which of the following best describes how you think of yourself?*' **female** (female), **all others** (male, non-binary, prefer not to say)
- **BMI at baseline**. Continuous. As described in 2.4.1.
- **Age**. Continuous. '*What is your age? (years)*'.
- **Ethnicity** (2 levels). '*What is your ethnic group?*' **white** (1 white), **all others** (Mixed/Multiple ethnic groups, Asian/Asian British, Black/African/Caribbean/Black British, Chinese, Arab, Other ethnic group, prefer not to say)

- **Occupation and working from home (3 levels) Unemployed, Employed from home, Employed not from home.**
  - All participants were asked, *‘What is your current MAIN occupation (during COVID-19)?’* Laid off during COVID-19, unemployed since before COVID-19, retired, homemaker, full-time parent or carer, unable to work due to disability, other, employed (full or part-time), self-employed (full or part-time), student, furloughed during COVID-19.
  - Participants who answered as employed (full or part-time), self-employed (full or part-time), student, furloughed during COVID-19 to the question *‘What is your current MAIN occupation (during COVID-19)?’* were asked, *‘Can you do your work or study from home?’* Yes, I can do all the work or study from home, I can only do some work or study from home, or No, my work or study cannot be done at home.
  - **Unemployed** (Laid off during COVID-19, Unemployed since before COVID-19, Retired, Homemaker, Full-time parent or carer, Unable to work due to disability, Other), **Employed from home** (Employed AND Yes, I can do all the work or study from home, OR I can only do some work or study from home) **Employed not from home** (Employed AND No, my work or study cannot be done at home).
- **Composite socioeconomic score:** Sum score of education level and combined housing tenure and household income (Sum score 0-3):
  - **Education level.** *‘What is the highest level of education that you have completed?’* **0**, below A-level (No formal qualification, GCSE/School certificate/O-level/CSE, Vocational qualifications (e.g. NVQ1+2)), or **1**, A-level or above (A-level/Higher school certificate or equivalent (e.g. NVQ3), Bachelor degree or equivalent (e.g. NVQ4), Masters/PhD/PGCE or equivalent, Other).
  - **Housing and Income.** *‘What is your housing tenure?’* Owned outright, Mortgage Rented from local authority, Rented from private landlord, Belongs to housing association, Shared ownership (part owned, part rented), Other. *‘What was your household annual income BEFORE COVID- 19?’* up to 13 499 GBP, 13 500-24 999 GBP, 25 000-49 999 GBP, ≥50 000 GBP, prefer not to say. **2** (Housing = Owned outright OR mortgage AND Income = 50 000+ GBP), **1** (Housing = All other AND Income = 50 000+ GBP), **1** (Housing = Owned outright OR mortgage AND Income = All other), **0** (Housing = All other AND Income = All other).

## COVID-19 related (impact on well-being and lifestyle)

- **Living arrangements (3 levels): alone, with children (with or without adults), with adults only.**
  - **Alone.** Participants responding ‘I live on my own’ when asked *‘How many persons other than yourself (including children) LIVE WITH YOU NOW in the same flat or house?’* Participants answering, ‘I live with 1 or more people’ were

then asked: *'Do you live with any of these persons below?'* Being able to select multiple responses: With my partner, husband/wife, boyfriend/girlfriend, with children 0-5 years old, with children 6-15 years old, with persons aged 16-69 (family or friends), with persons aged 70+ (family or friends), persons who you believe may be vulnerable to COVID-19 for any reason, persons who are in poor health

- **With any children (with or without adults)** (if participants respond with 'children 0-5 years old' or 'with children 6-15 years old')
- **Adults only** (all other participants who did not say 'I live on my own' or did not select 'with children 0-5 years old' or 'with children 6-15 years old')

## Time-variant

### COVID-19 related (impact on well-being and lifestyle)

- **Isolation** *'Which type of COVID-19 induced isolation are you experiencing?'* (2 levels) **total and some** (Total isolation/quarantine, Some isolation), **general and no isolation** (General isolation, No isolation)
- **Composite wellbeing, social and living measure:** At baseline: *'How would you rate the below aspects of your life in the since COVID-19?'* At follow-up: *'How would you rate the below aspects of your life in the past month?'* 1-5: 1 poor, 2, 3 average, 4, 5 excellent. Continuous mean score of living + wellbeing + social + family (1-5)

## Behavioural: time-variant

### Diet:

- **HFSS snacks intake** continuous
- **HFSS meals intake** continuous
- **Fruit and vegetable intake** continuous
  - Participants were asked at each survey, *'In the past month, how often did you usually eat or drink...'* for nine food items: eight HFSS food items (ready meals, fast food and takeaways, sugary or sweetened drinks, sweets or chocolate, cakes and biscuits, desserts and savoury snacks), and one item for fruit and vegetable intake. Participants were also asked at baseline: *'BEFORE COVID-19, how often did you usually eat or drink...'* for the nine food items. Respondents could answer on a 7-point scale: A few times per day, Once a day, A few times per week, Once a week, A few times per month, Once a month, Less often/never. Apart from for the retrospective 'before COVID-19' question, participants could also answer 'not sure'. To estimate monthly consumption, responses for the food item questions were converted into monthly portion frequencies. Assuming a minimum of four weeks per calendar month, an answer of 'a few times per day' was recorded as 56 portions per month (i.e., 2

portions x 7 days x 4 weeks), 'Once a day' as 28 portions per month, 'Few times per week' as 12 portions per month, 'Once a week' as four portions per month, 'Few times per month' as two portions per month, 'Once a month' as one portion per month, 'Less often/never' as 0.5 portions per month. Participants responding as 'not sure' were excluded.

- Sugary or sweetened drinks, sweets or chocolate, cakes and biscuits, desserts and savoury snacks were summed to produce a 'HFSS snacks intake' monthly frequency outcome variable. Ready meals, fast food and takeaways monthly frequencies were summed to produce a 'HFSS meals intake' monthly frequency outcome variable.
- HFSS snacks intake, HFSS meals intake and fruit and vegetable intake at baseline, 3- and 6-months follow-up surveys were combined to create time-varying **HFSS snacks intake** and **HFSS meals intake** and **fruit and vegetable intake** continuous variables.
- **Physical activity (2 levels) 1** (decrease in both or, decrease in either but not an increase in the other), **0** (all other)
  - Physical activity was a binary score comparing strength and aerobic physical activity levels at baseline, 3- and 6-months follow-ups to the retrospectively reported pre-COVID-19 strengthening and aerobic physical activity levels. A reduction in physical activity was defined as a decrease by one or more strength training sessions per week, and/or a decrease by 20 or more minutes per week of aerobic physical activity at each timepoint compared to pre-COVID-19 physical activity levels. A decrease in one, but an increase in the other was treated as 'all other'.
  - **Weekly strength training.** *'BEFORE COVID-19, on average, on HOW MANY DAYS PER WEEK did you do STRENGTH TRAINING? 'SINCE COVID-19, on average, on HOW MANY DAYS PER WEEK have you done STRENGTH TRAINING?'* 0 days per week, 1 day per week, 2 days per week, 3 days per week, 4 days per week or more. Compared to the before COVID-19 response, a decrease by 1 or more days per week or all other at each timepoint.
  - **Weekly aerobic training.** *'(In the month) BEFORE COVID-19, HOW LONG (in MINUTES) was your average session of moderate or vigorous AEROBIC PHYSICAL ACTIVITY?'*, *'SINCE COVID-19, HOW LONG (in MINUTES) was your average session of moderate or vigorous AEROBIC PHYSICAL ACTIVITY?'* (15-480 minutes) and *'(In the month) BEFORE COVID-19 on average, HOW MANY TIMES PER WEEK did you do at least 15 MINUTES or more of moderate or vigorous AEROBIC PHYSICAL ACTIVITY?'*, *'SINCE COVID-19 on average, HOW MANY TIMES PER WEEK have you done 15 MINUTES or more of moderate or vigorous AEROBIC PHYSICAL ACTIVITY?'* (0,1,2 to 13 times per week, or 14+ times per week). Compared to the before COVID-19 response, a decrease by 20 or more minutes per week or all other at each timepoint.
- **Alcohol consumption. Binary.** Participants were asked *'How often did you have a drink containing alcohol in the past month?'*, *'In the past month, how many units of*

*alcohol did you drink on a typical day when you were drinking?'. Scores were multiplied to compute total weekly units of alcohol consumed, and converted into a binary measure of weekly consumption, categorised into  $\leq 14$  and  $>14$  alcohol units.*

- **Smoking status. Binary.** At baseline, *'Which statement about tobacco use and cigarette smoking best describes you?'* At follow-up, *'2 MONTHS ago/previously, we asked about your tobacco use. Your situation may have changed since then, or not. Which of the following best applies to you now?'* (2 levels) **Yes** (At baseline: I smoke cigarettes (including hand-rolled) every day, I smoke cigarettes (including hand-rolled), but not every day, I do not smoke cigarettes at all, but I do smoke tobacco of some kind (eg. pipe, cigar or shisha). At follow-up: I smoke cigarettes (including hand-rolled) every day, I smoke cigarettes (including hand-rolled) but not every day, I do not smoke cigarettes at all, but I do smoke tobacco of some kind (e.g. pipe, cigar or shisha)), **No** (At baseline: I have stopped smoking completely in the last year, I stopped smoking completely more than a year ago, I have never smoked any cigarettes. At follow-up: I have stopped smoking or using tobacco completely in the past 2/3 months, I stopped smoking completely more than 2/3 months ago, I have never been a regular smoker (i.e. smoked for a year or more).

### **Behavioural, time-invariant**

Time invariant  $\Delta$ HFSS snacks intake change score,  $\Delta$ HFSS meals intake change score and  $\Delta$ fruit and vegetable intake change score were computed as described below.

- **$\Delta$ HFSS snacks intake change score.** A continuous change score was calculated from the difference between the before-COVID-19 and the since-COVID-19 monthly HFSS snacks frequencies reported at baseline.
- **$\Delta$ HFSS meals intake change score.** A continuous change score was calculated from the difference between the before-COVID-19 and the since-COVID-19 monthly HFSS meals frequencies reported at baseline.
- **$\Delta$ Fruit and vegetable intake change score.** A continuous change score was calculated from the difference between the before-COVID-19 and the since-COVID-19 monthly fruit and vegetable intake frequencies reported at baseline.

## Supplementary tables

**Table S1.** Weighted baseline participant characteristics for the total, included and excluded samples.

**Table S2.** Means with 95% confidence intervals for weight and BMI at each timepoint using complete cases.

**Table S3.** Mean difference in weight/BMI with 95% confidence intervals between timepoints using complete cases.

**Table S4.** Univariate unadjusted and adjusted models for each predictor variable and change in self-reported weight and BMI.

**Table S5.** Unweighted and weighted proportions of underweight, normal weight, overweight and obesity at baseline (May-June 2020), 3 months (August-September 2020) and 6 months follow-up (November-December 2020) using complete cases.

**Table S6.** Univariate unadjusted and adjusted models for each predictor variable and change in self-reported weight and BMI using complete cases.

**Table S7.** Full GEE model containing all predictor variables and the full GEE model containing all predictor variables and significant predictor\*time interactions from univariate models adjusted for time using complete cases.

**Table S8.** Univariate unadjusted and adjusted models for each predictor variable and change in self-reported weight as a binary outcome using complete cases.

**Table S9.** Full GEE model containing all predictor variables and the full GEE model containing all predictor variables and significant predictor\*time interactions from univariate models adjusted for time, for a change in self-reported weight as a binary outcome using complete cases.

**Table S10.** Univariate unadjusted and adjusted models for each predictor variable and change in self-reported BMI as a binary outcome using complete cases.

**Table S11.** Full GEE model containing all predictor variables and the full GEE model containing all predictor variables and significant predictor\*time interactions from univariate models adjusted for time, for a change in self-reported BMI as a binary outcome using complete cases.

**Table S1.** Weighted baseline participant characteristics for the total, included and excluded samples.

	Total Sample	Included sample	Excluded sample	
	Weighted %	Weighted %	Weighted %	p
<b>N</b>	2805	1631	1174	
<b>Gender</b>				0.120
All other	51.7	53.0	50.0	
Female	48.3	47.0	50.0	
<b>Ethnicity</b>				<0.001
All other	11.5	7.5	17.1	
White	88.5	92.5	82.9	
<b>Mean BMI [SD] (N=2614)</b>	26.4 [5.2]	26.7 [5.1]	26.0 [5.3]	<b>0.001</b>
<b>Mean Age [SD]</b>	48.3 [16.7]	53.1 [14.8]	41.6 [16.8]	<b>&lt;0.001</b>
<b>Occupation and work from home</b>				<b>&lt;0.001</b>
Unemployed (including retired persons and students)	30.9	36.2	23.5	
Employed working from home	40.4	35.8	46.8	
Employed not working from home	28.7	28.0	29.6	
<b>Socioeconomic score</b>				<b>&lt;0.001</b>
Income <£50k, unowned Housing and no higher education	13.0	11.5	15.2	
1 of: ≥£50K income, housing ownership/mortgage or higher education	40.5	38.6	43.2	
2 of: ≥£50K income, housing ownership/mortgage or higher education	34.7	37.2	31.2	
All of: ≥£50K income, housing ownership/mortgage and higher education	11.8	12.8	10.5	
<b>Living conditions</b>				<b>&lt;0.001</b>

Alone	18.8%	19.3%	18.2%	
With children (with or without adults)	17.3%	14.8%	20.7%	
With adults only	63.9%	65.9%	61.1%	
<b>Isolation status (N=2763)</b>				<b>0.027</b>
Total or some isolation	75.8%	77.3%	73.6%	
General or no isolation	24.2%	22.7%	26.4%	
<b>Mean Quality of Life [SD] (1-5) (N=2704)</b>	3.2 [0.9]	3.3 [0.9]	3.1 [0.9]	<b>&lt;0.001</b>
<b>Mean HFSS snacks (portions per month) [SD] (N=2395)</b>	61.1 [49.9]	60.9 [50.7]	61.5 [48.5]	0.790
<b>Mean HFSS meals (portions per month) [SD] (N=2409)</b>	7.4 [9.2]	6.6 [7.4]	8.8 [11.5]	<b>&lt;0.001</b>
<b>Mean Fruit and vegetables (portions per month) [SD] (N=2438)</b>	39.4 [19.7]	40.6 [19.2]	37.5 [20.2]	<b>&lt;0.001</b>
<b>Mean ΔHFSS snacks change score (portions per month) [SD] (N=2395)</b>	8.3 [38.8]	9.8 [39.5]	5.9 [37.3]	<b>0.018</b>
<b>Mean ΔHFSS meals change score (portions per month) [SD] (N=2409)</b>	-1.4 [8.4]	-1.5 [7.8]	-1.3 [9.4]	0.649
<b>Mean ΔFruit and vegetables change score (portions per month) [SD] (N=2438)</b>	0.2 [12.6]	0.1 [11.7]	0.3 [14.0]	0.751
<b>Physical activity (N=2618)</b>				0.679
All other	75.8%	75.5%	76.2%	
Reduced	24.2%	24.5%	23.8%	
<b>Alcohol consumption (N=2541)</b>				0.436
≤14 weekly units	80.8%	81.3%	80.0%	
>14 weekly units	19.2%	18.7%	20.0%	
<b>Smoking status</b>				<b>&lt;0.001</b>
Yes	26.5%	19.7%	35.9%	
No	68.9%	80.3%	64.1%	

*SD: Standard Deviation.*

**Table S2.** Means with 95% confidence intervals for weight and BMI at each timepoint using complete cases.

Complete Cases	Self-reported weight			Self-reported BMI	
	N	Mean [95% CI]	SE	Mean [95% CI]	SE
May-June 2020	1543	74.8 [74.0,75.6]	0.41408	26.2 [25.9,26.4]	0.12693
August-September 2020	1543	75.1 [74.3,76.0]	0.41781	26.3 [26.0,26.6]	0.12905
November-December 2020	1543	74.9 [74.0,75.7]	0.41442	26.2 [26.0,26.5]	0.12820

*SE: Standard Error. CI: Confidence Interval.*

**Table S3.** Mean difference in weight/BMI with 95% confidence intervals between timepoints using complete cases.

Complete Cases	Self-reported weight (kg)				Self-reported BMI (kg/m <sup>2</sup> )		
	N	Mean difference [95% CI]	SE	p	Mean difference [95% CI]	SE	p
Baseline - 3 mo.	1543	<b>0.37 [0.14,0.60]</b>	<b>0.09514</b>	<b>&lt;0.001</b>	<b>0.13 [0.06,0.21]</b>	<b>0.03319</b>	<b>&lt;0.001</b>
3 mo. - 6 mo.	1543	<b>-0.28 [-0.48,-0.09]</b>	<b>0.08603</b>	<b>0.002</b>	<b>-0.10 [-0.16,-0.03]</b>	<b>0.03000</b>	<b>0.002</b>
Baseline - 6 mo.	1543	0.08 [-0.14,0.30]	0.11253	0.469	0.04 [-0.04,0.11]	0.03970	0.354

*SE: Standard Error, CI: Confidence Interval, mo: months.*

**Table S4.** Univariate unadjusted and adjusted models for each predictor variable and change in self-reported weight and BMI.

Predictor	Change in self-reported weight				Change in self-reported BMI			
	Unadjusted models		Adjusted models w/interaction		Unadjusted models		Adjusted models w/interaction	
	W $\chi^2$	p	W $\chi^2$	p	W $\chi^2$	p	W $\chi^2$	p
Gender	2.030	0.154	1.883	0.170	2.591	0.107	2.469	0.116
Time			<b>10.243</b>	<b>0.001</b>			<b>10.632</b>	<b>0.001</b>
Time*Gender			1.669	0.196			1.176	0.278
Ethnicity	0.069	0.792	0.034	0.853	0.074	0.785	0.038	0.845
Time			<b>5.811</b>	<b>0.016</b>			<b>5.890</b>	<b>0.015</b>
Time*Ethnicity			0.779	0.377			0.910	0.340
Baseline BMI	<b>11.759</b>	<b>0.001</b>	<b>10.529</b>	<b>0.001</b>	<b>11.353</b>	<b>0.001</b>	<b>10.187</b>	<b>0.001</b>
Time			<b>7.856</b>	<b>0.005</b>			<b>7.563</b>	<b>0.006</b>
Time*Baseline BMI			<b>11.001</b>	<b>0.001</b>			<b>10.596</b>	<b>0.001</b>
Age	0.231	0.631	0.146	0.702	0.165	0.684	0.094	0.760
Time			2.145	0.143			2.237	0.135
Time*Age			0.459	0.498			0.491	0.483
Occupation and work from home	0.313	0.855	0.280	0.869	0.198	0.906	0.179	0.914
Time			<b>8.504</b>	<b>0.004</b>			<b>8.365</b>	<b>0.004</b>
Time*Occupation and work from home			0.488	0.783			0.279	0.870

Socioeconomic score	2.514	0.473	2.338	0.505	2.891	0.409	2.696	0.441
Time			3.703	0.054			3.515	0.061
Time*Socioeconomic score			2.073	0.557			2.394	0.495
Living arrangements	0.344	0.842	0.389	0.823	0.430	0.806	0.470	0.791
Time			<b>4.445</b>	<b>0.035</b>			<b>4.985</b>	<b>0.026</b>
Time*Living arrangements			0.870	0.647			0.557	0.757
Isolation status	0.680	0.410	0.448	0.503	0.634	0.426	0.533	0.465
Time			<b>12.203</b>	<b>&lt;0.001</b>			<b>12.975</b>	<b>&lt;0.001</b>
Time*Isolation status			0.978	0.323			1.379	0.240
Quality of life	0.174	0.676	0.038	0.846	0.219	0.640	0.023	0.878
Time			0.259	0.611			0.411	0.522
Time*Quality of life			0.008	0.928			0.003	0.955
HFSS snacks intake	<b>15.593</b>	<b>&lt;0.001</b>	<b>13.775</b>	<b>&lt;0.001</b>	<b>15.772</b>	<b>&lt;0.001</b>	<b>14.328</b>	<b>&lt;0.001</b>
Time			<b>29.291</b>	<b>&lt;0.001</b>			<b>28.201</b>	<b>&lt;0.001</b>
Time*HFSS snacks intake			<b>13.221</b>	<b>&lt;0.001</b>			<b>12.206</b>	<b>&lt;0.001</b>
HFSS meals intake	2.002	0.157	1.851	0.174	1.600	0.206	1.321	0.250
Time			<b>19.85</b>	<b>&lt;0.001</b>			<b>21.041</b>	<b>&lt;0.001</b>
Time*HFSS meals intake			<b>4.257</b>	<b>0.039</b>			<b>4.839</b>	<b>0.028</b>

Fruit and vegetables intake	<b>5.397</b>	<b>0.020</b>	<b>5.456</b>	<b>0.020</b>	<b>5.315</b>	<b>0.021</b>	<b>5.425</b>	<b>0.020</b>
Time			0.792	0.374			0.855	0.355
Time*Fruit and vegetables intake			<b>4.475</b>	<b>0.034</b>			<b>4.776</b>	<b>0.029</b>
$\Delta$ HFSS snacks change score	<b>4.296</b>	<b>0.038</b>	<b>4.532</b>	<b>0.033</b>	<b>4.599</b>	<b>0.032</b>	<b>4.881</b>	<b>0.027</b>
Time			<b>7.368</b>	<b>0.007</b>			<b>7.003</b>	<b>0.008</b>
Time* $\Delta$ HFSS snacks change score			0.185	0.667			0.199	0.655
$\Delta$ HFSS meals change score	0.008	0.929	0.004	0.950	0.003	0.954	0.001	0.979
Time			<b>10.258</b>	<b>0.001</b>			<b>10.276</b>	<b>0.001</b>
Time* $\Delta$ HFSS meals change score			0.018	0.894			0.024	0.878
$\Delta$ Fruit and vegetables change score	1.037	0.309	1.080	0.299	1.303	0.254	1.356	0.244
Time			<b>9.285</b>	<b>0.002</b>			<b>9.084</b>	<b>0.003</b>
Time* $\Delta$ Fruit and vegetables change score			0.025	0.875			0.038	0.845
Physical activity	0.002	0.963	<0.001	0.999	0.032	0.859	0.051	0.821
Time			<b>7.562</b>	<b>0.006</b>			<b>7.110</b>	<b>0.008</b>
Time*Physical activity			0.236	0.627			0.341	0.559
Alcohol consumption	2.946	0.086	2.373	0.123	2.289	0.130	1.778	0.182
Time			1.514	0.219			1.751	0.186
Time*Alcohol consumption			<b>9.351</b>	<b>0.002</b>			<b>8.974</b>	<b>0.003</b>
Smoking status	0.092	0.762	0.072	0.789	0.092	0.762	0.071	0.790

Time	2.854	0.091	2.906	0.088
Time*Smoking status	0.077	0.782	0.079	0.778

*Missingness with each model depends on if participants responded to survey questions relating to the predictor variable. W  $\chi^2$ : Wald Chi-square.*

**Table S5.** Unweighted and weighted proportions of underweight, normal weight, overweight and obesity at baseline (May-June 2020), 3 months (August-September 2020) and 6 months follow-up (November-December 2020) using complete cases.

	Unweighted			Weighted		
	Apr-Jun	Jul-Sep	Nov-Dec	Apr-Jun	Jul-Sep	Nov-Dec
<b>N</b>	1543	1543	1543	1333	1392	1358
<b>BMI Category (%)</b>						
Underweight	2.0	2.0	2.1	2.6	2.9	2.6
Normal weight	45.0	44.1	44.9	38.3	38.8	39.9
Overweight	32.8	32.9	32.8	37.0	35.7	35.5
Obesity	20.2	21.0	20.2	22.1	22.7	22.0

**Table S6.** Univariate unadjusted and adjusted models for each predictor variable and change in self-reported weight and BMI using complete cases.

Predictor	Change in self-reported weight				Change in self-reported BMI			
	Unadjusted models		Adjusted models w/interaction		Unadjusted models		Adjusted models w/interaction	
	W $\chi^2$	p	W $\chi^2$	p	W $\chi^2$	p	W $\chi^2$	p
Gender	2.761	0.097	2.761	0.097	3.287	0.070	3.287	0.070
Time			<b>11.000</b>	<b>0.001</b>			<b>11.232</b>	<b>0.001</b>
Time*Gender			2.017	0.156			1.429	0.232
Ethnicity	0.001	0.978	0.001	0.978	0.012	0.912	0.012	0.912
Time			<b>5.915</b>	<b>0.015</b>			<b>6.159</b>	<b>0.013</b>
Time*Ethnicity			0.396	0.529			0.669	0.413
Baseline BMI	<b>9.913</b>	<b>0.002</b>	<b>9.913</b>	<b>0.002</b>	<b>9.680</b>	<b>0.002</b>	<b>9.680</b>	<b>0.002</b>
Time			<b>8.070</b>	<b>0.005</b>			<b>7.835</b>	<b>0.005</b>
Time*Baseline BMI			<b>11.286</b>	<b>0.001</b>			<b>10.918</b>	<b>0.001</b>
Age	0.354	0.552	0.354	0.552	0.344	0.558	0.344	0.558
Time			2.390	0.122			2.537	0.111
Time*Age			0.574	0.449			0.642	0.423
Occupation and work from home	0.888	0.642	0.888	0.642	0.738	0.691	0.738	0.691
Time			<b>8.411</b>	<b>0.004</b>			<b>8.155</b>	<b>0.004</b>

Time*Occupation and work from home			0.804	0.669			0.552	0.759
Socioeconomic score	1.087	0.780	1.087	0.780	1.395	0.707	1.395	0.707
Time			3.878	0.049			3.403	0.065
Time*Socioeconomic score			1.553	0.670			1.905	0.592
Living arrangements	0.510	0.775	0.510	0.775	0.730	0.694	0.730	0.694
Time			<b>5.219</b>	<b>0.022</b>			<b>5.691</b>	<b>0.017</b>
Time*Living arrangements			0.803	0.669			0.526	0.769
Isolation status	1.058	0.304	0.250	0.617	0.939	0.332	0.327	0.567
Time			<b>12.113</b>	<b>0.001</b>			<b>12.802</b>	<b>&lt;0.001</b>
Time*Isolation status			1.233	0.267			1.633	0.201
Quality of life	0.036	0.850	0.216	0.642	0.053	0.819	0.171	0.679
Time			0.138	0.710			0.248	0.618
Time*Quality of life			0.058	0.809			0.009	0.925
HFSS snacks intake	<b>13.217</b>	<b>&lt;0.001</b>	<b>13.025</b>	<b>&lt;0.001</b>	<b>12.956</b>	<b>&lt;0.001</b>	<b>13.064</b>	<b>&lt;0.001</b>
Time			<b>29.276</b>	<b>&lt;0.001</b>			<b>28.001</b>	<b>&lt;0.001</b>
Time*HFSS snacks intake			<b>13.198</b>	<b>&lt;0.001</b>			<b>12.030</b>	<b>0.001</b>
HFSS meals intake	2.522	0.112	2.878	0.090	1.732	0.188	1.927	0.165
Time			<b>21.490</b>	<b>&lt;0.001</b>			<b>22.116</b>	<b>&lt;0.001</b>
Time*HFSS meals intake			<b>4.842</b>	<b>0.028</b>			<b>5.124</b>	<b>0.024</b>

Fruit and vegetables intake	<b>6.377</b>	<b>0.012</b>	<b>6.922</b>	<b>0.009</b>	<b>6.206</b>	<b>0.013</b>	<b>6.772</b>	<b>0.009</b>
Time			1.136	0.287			1.207	0.272
Time*Fruit and vegetables intake			<b>5.352</b>	<b>0.021</b>			<b>5.671</b>	<b>0.017</b>
$\Delta$ HFSS snacks change score	2.058	0.151	2.058	0.151	2.177	0.140	2.177	0.140
Time			<b>7.896</b>	<b>0.005</b>			<b>7.424</b>	<b>0.006</b>
Time* $\Delta$ HFSS snacks change score			0.436	0.509			0.452	0.502
$\Delta$ HFSS meals change score	0.544	0.461	0.544	0.461	0.389	0.533	0.389	0.533
Time			<b>12.149</b>	<b>&lt;0.001</b>			<b>11.977</b>	<b>0.001</b>
Time* $\Delta$ HFSS meals change score			0.159	0.691			0.139	0.709
$\Delta$ Fruit and vegetables change score	0.122	0.727	0.122	0.727	0.199	0.655	0.199	0.655
Time			<b>10.692</b>	<b>0.001</b>			<b>10.345</b>	<b>0.001</b>
Time* $\Delta$ Fruit and vegetables change score			0.031	0.861			0.019	0.890
Physical activity	0.150	0.903	<b>0.010</b>	0.920	0.015	0.903	0.020	0.887
Time			<b>7.685</b>	<b>0.006</b>			<b>7.215</b>	<b>0.007</b>
Time*Physical activity			0.30	0.584			0.398	0.528
Alcohol consumption	2.122	0.145	1.934	0.164	1.674	0.196	1.503	0.220
Time			2.136	0.144			2.357	0.125
Time*Alcohol consumption			<b>8.811</b>	<b>0.003</b>			<b>8.527</b>	<b>0.003</b>

Smoking status	0.066	0.797	0.052	0.820	0.048	0.826	0.036	0.850
Time			2.910	0.088			2.872	0.090
Time*Smoking status			0.079	0.778			0.091	0.762

*Missingness with each model depends on if participants responded to survey questions relating to the predictor variable.  $W\chi^2$ : Wald Chi-square.*

**Table S7.** Full GEE model containing all predictor variables and the full GEE model containing all predictor variables and significant predictor\*time interactions from univariate models adjusted for time using complete cases.

Complete cases All predictors (N=1416)	Change in self-reported weight QIC = 43324.151				Change in self-reported BMI QIC = 5392.741			
	W $\chi^2$	p	B [95% CI]	SE	W $\chi^2$	p	B [95% CI]	SE
Gender	3.073	0.080			3.564	0.059		
All other			Reference				Reference	
Female			0.418 [-0.049,0.886]	0.2387			0.147 [-0.006,0.300]	0.0780
Ethnicity	0.072	0.788			0.056	0.812		
All other			Reference				Reference	
White			-0.088 [-0.728,0.553]	0.3268			-0.023 [-0.268,0.215]	0.1252
Baseline BMI	<b>9.865</b>	<b>0.002</b>	<b>-0.091 [-0.148,-0.034]</b>	<b>0.0291</b>	<b>9.823</b>	<b>0.002</b>	<b>-0.032 [-0.052,-0.012]</b>	<b>0.0103</b>
Age	1.814	0.178	0.011 [-0.005,0.026]	0.0080	2.091	0.148	0.004 [-0.001,0.009]	0.0028
Occupation and work from home	0.058	0.971			0.011	0.995		
Unemployed			Reference				Reference	
Employed working from home			-0.068 [-0.619,0.484]	0.2814			-0.010 [-0.206,0.185]	0.0998
Employed not working from home			-0.049 [-0.722,0.624]	0.3434			-0.008 [-0.242,0.227]	0.1197
Socioeconomic score	1.448	0.694			1.650	0.648		
Income <£50k, unowned Housing and no higher education			Reference				Reference	
1 of: ≥£50K income, housing ownership/mortgage or higher education			0.523 [-0.815,1.860]	0.6826			0.111 [-0.349,0.570]	0.2346
2 of: ≥£50K income, housing ownership/mortgage or higher education			0.282 [-0.990,1.554]	0.6490			0.015 [-0.420,0.449]	0.2218
All of: ≥£50K income, housing ownership/mortgage and higher education			0.152 [-1.148,1.453]	0.6635			-0.033 [-0.477,0.411]	0.2267
Living conditions	0.100	0.951			0.135	0.935		
Alone			Reference				Reference	
With children (with or without adults)			0.097 [-0.532,0.725]	0.3206			0.036 [-0.185,0.256]	0.1124
With adults only			0.072 [-0.440,0.584]	0.2611			0.033 [-0.148,0.214]	0.0923
Isolation status	0.339	0.561			0.373	0.541		
Total or some isolation			Reference				Reference	

General or no isolation			-0.092 [-0.403,0.218]	0.1584			-0.034 [-0.143,0.075]	0.0558
Quality of Life	<0.001	0.996	<0.001 [-0.220,0.219]	0.1121	0.015	0.904	0.005 [-0.075,0.085]	0.0406
HFSS snacks	<b>11.656</b>	<b>0.001</b>	<b>0.009 [0.004,0.014]</b>	<b>0.0027</b>	<b>11.929</b>	<b>0.001</b>	<b>0.003 [0.001,0.005]</b>	<b>0.0010</b>
HFSS meals	3.677	0.055	0.021 [<0.001,0.043]	0.0110	2.777	0.096	0.006 [-0.001,0.014]	0.0039
Fruit and vegetables	<b>5.215</b>	<b>0.022</b>	<b>-0.011 [-0.021,-0.002]</b>	<b>0.0050</b>	<b>5.043</b>	<b>0.025</b>	<b>-0.004 [-0.007,&lt;0.001]</b>	<b>0.0018</b>
ΔHFSS snacks change score	2.054	0.152	0.008 [-0.003,0.019]	0.0057	2.228	0.136	0.003 [-0.001,0.006]	0.0018
ΔHFSS meals change score	0.366	0.545	-0.010 [-0.044,0.023]	0.0172	0.285	0.593	-0.003 [-0.015,0.008]	0.0058
ΔFruit and vegetables change score	0.062	0.804	-0.002 [-0.020,0.016]	0.0091	0.090	0.764	-0.001 [-0.007,0.005]	0.0031
Physical activity	0.031	0.861			0.007	0.932		
All other			Reference				Reference	
Reduced			0.028 [-0.283,0.339]	0.1586			-0.005 [-0.115,0.105]	0.0560
Alcohol consumption	3.707	0.054			3.330	0.068		
≤14 weekly units			Reference				Reference	
>14 weekly units			0.390 [-0.007,0.787]	0.2024			0.121 [-0.009,0.252]	0.0665
Smoking status	0.045	0.833			0.018	0.893		
Yes			Reference				Reference	
No			0.064 [-0.528,0.656]	0.3020			0.014 [-0.196,.225]	0.1072

Change in self-reported weight QIC = 43090.196

Change in self-reported BMI QIC = 5366.550

All predictors + significant time interactions (N=1416)	W $\chi^2$	p	W $\chi^2$	p
Time*Baseline BMI	<b>13.202</b>	<b>&lt;0.001</b>	<b>12.46</b>	<b>&lt;0.001</b>
Time*HFSS snacks	<b>16.997</b>	<b>&lt;0.001</b>	<b>15.63</b>	<b>&lt;0.001</b>
Time*Alcohol consumption	<b>12.908</b>	<b>&lt;0.001</b>	<b>12.66</b>	<b>0.001</b>

*Models also included Time as a covariate. For 'All predictors + significant time interactions', Type III tests for the predictor\*time interactions are shown only. There were no material changes in significance of main effects with the addition of the time interactions, except for alcohol consumption (p=0.046) and HFSS meals intake (p=0.049) which became significant for a change in weight. QIC is a relative, 'lower is better' measure of goodness of fit. W  $\chi^2$ : Wald Chi-square, B: Beta parameter, SE: Standard Error, CI: Confidence Interval.*

**Table S8.** Univariate unadjusted and adjusted models for each predictor variable and change in self-reported weight as a binary outcome using complete cases.

Binary Weight	Increase weight vs all other				Decrease weight vs all other			
	Unadjusted models		Adjusted models w/interaction		Unadjusted models		Adjusted models w/interaction	
	W $\chi^2$	p	W $\chi^2$	p	W $\chi^2$	p	W $\chi^2$	p
Gender	1.064	0.302	1.011	0.315	3.034	0.082	2.925	0.087
Time			1.512	0.219			<b>30.785</b>	<b>&lt;0.001</b>
Time*Gender			0.925	0.336			0.019	0.889
Ethnicity	0.002	0.969	0.002	0.968	0.686	0.407	**	**
Time			<0.001	0.992			**	**
Time*Ethnicity			0.420	0.517			**	**
Baseline BMI	2.290	0.130	2.322	0.128	<b>13.351</b>	<b>&lt;0.001</b>	<b>12.163</b>	<b>&lt;0.001</b>
Time			0.666	0.414			28.750	<b>&lt;0.001</b>
Time*Baseline BMI			0.222	0.638			0.063	0.802
Age	<b>5.106</b>	<b>0.024</b>	<b>5.146</b>	<b>0.023</b>	1.282	0.257	1.303	0.254
Time			0.113	0.736			1.880	0.170
Time*Age			0.032	0.859			0.076	0.783
Occupation and work from home	0.912	0.634	0.834	0.659	0.731	0.694	1.402	0.496
Time			<b>4.035</b>	<b>0.045</b>			<b>25.886</b>	<b>&lt;0.001</b>
Time*Occupation and work from home			1.506	0.471			4.449	0.108

Socioeconomic score	<b>14.341</b>	<b>0.002</b>	<b>13.948</b>	<b>0.003</b>	6.622	0.085	**	**
Time			2.528	0.112			**	**
Time*Socioeconomic score			1.847	0.605			**	**
Living arrangements	1.140	0.565	1.105	0.576	1.878	0.391	1.462	0.481
Time			<b>2.948</b>	<b>0.086</b>			<b>25.406</b>	<b>&lt;0.001</b>
Time*Living arrangements			0.883	0.643			0.367	0.832
Isolation status	0.064	0.800	0.076	0.783	<b>9.201</b>	<b>0.002</b>	0.427	0.513
Time			1.846	0.174			<b>20.967</b>	<b>&lt;0.001</b>
Time*Isolation status			1.288	0.256			0.308	0.579
Quality of life	2.156	0.142	1.360	0.244	<b>9.909</b>	<b>0.002</b>	<b>6.578</b>	<b>0.010</b>
Time			0.138	0.711			0.610	0.435
Time*Quality of life			0.522	0.470			<b>4.123</b>	<b>0.042</b>
HFSS snacks	3.350	0.067	1.000	0.317	<b>4.307</b>	<b>0.038</b>	3.636	0.057
Time			0.470	0.493			<b>34.552</b>	<b>&lt;0.001</b>
Time*HFSS snacks			<b>3.996</b>	<b>0.046</b>			0.911	0.340
HFSS meals	1.814	0.178	0.488	0.485	0.001	0.980	0.009	0.924
Time			0.003	0.959			<b>29.907</b>	<b>&lt;0.001</b>
Time*HFSS meals			2.790	0.095			3.059	0.080

Fruit and vegetables	<b>13.939</b>	<b>&lt;0.001</b>	<b>12.561</b>	<b>&lt;0.001</b>	0.989	0.320	0.813	0.367
Time			<b>1.458</b>	<b>0.227</b>			17.885	<0.001
Time*Fruit and vegetables			0.566	0.452			2.322	0.128
$\Delta$ HFSS snacks change score	2.848	0.091	2.832	0.092	0.415	0.519	0.704	0.401
Time			2.119	0.145			31.601	<0.001
Time* $\Delta$ HFSS snacks change score			0.006	0.937			1.280	0.258
$\Delta$ HFSS meals change score	0.154	0.695	0.138	0.710	2.428	0.119	1.866	0.172
Time			1.768	0.184			<b>35.365</b>	<b>&lt;0.001</b>
Time* $\Delta$ HFSS meals change score			0.147	0.701			1.633	0.201
$\Delta$ Fruit and vegetables change score	0.184	0.668	0.177	0.674	0.011	0.916	0.027	0.869
Time			2.480	0.115			<b>33.273</b>	<b>&lt;0.001</b>
Time* $\Delta$ Fruit and vegetables change score			0.006	0.938			0.341	0.559
Physical activity	0.850	0.357	0.763	0.382	0.321	0.571	0.019	0.890
Time			1.429	0.232			<b>24.086</b>	<b>&lt;0.001</b>
Time*Physical activity			0.978	0.323			2.438	0.118
Alcohol consumption	0.027	0.870	0.005	0.941	<b>5.755</b>	<b>0.016</b>	<b>4.314</b>	<b>0.038</b>
Time			2.934	0.087			<b>12.509</b>	<b>&lt;0.001</b>
Time*Alcohol consumption			0.838	0.360			<b>4.854</b>	<b>0.028</b>
Smoking status	2.974	0.085	2.774	0.096	<b>8.076</b>	<b>0.004</b>	<b>7.192</b>	<b>0.007</b>

Time	2.901	0.089	<b>13.501</b>	<b>&lt;0.001</b>
Time*Smoking status	0.227	0.634	1.705	0.192

*\*\*Model did not compute due to numerical errors. In the 'Decrease weight vs all other' model, baseline BMI, HFSS snacks intake, fruit and vegetable intake, and HFSS snacks change score continuous variables were converted to categorical variables due to violating the linearity of logit assumption. Missingness with each model depends on if participants responded to survey questions relating to the predictor variable. W  $\chi^2$ : Wald Chi-square.*

**Table S9.** Full GEE model containing all predictor variables and the full GEE model containing all predictor variables and significant predictor\*time interactions from univariate models adjusted for time, for a change in self-reported weight as a binary outcome using complete cases.

Binary weight outcome variable		Increase weight vs all other			Decrease weight vs all other		
Model	Predictor	QIC	Type III Test		QIC	Type III Test	
			W $\chi^2$	p		W $\chi^2$	p
All predictors (N=1416)		1806.499			1416.847		
	Gender		1.037	0.309		<b>7.545</b>	<b>0.006</b>
	Ethnicity		0.008	0.928		0.766	0.381
	Baseline BMI		1.922	0.166		<b>13.646</b>	<b>&lt;0.001</b>
	Age		<b>6.570</b>	<b>0.010</b>		2.856	0.091
	Occupation and work from home		3.829	0.147		0.848	0.654
	Socioeconomic score		4.761	0.190		3.461	0.326
	Living conditions		0.876	0.645		1.402	0.496
	Isolation status		0.002	0.963		0.837	0.360
	Quality of Life		0.069	0.792		1.361	0.243
	HFSS snacks intake		1.519	0.218		<b>5.345</b>	<b>0.021</b>
	HFSS meals intake		0.019	0.890		1.040	0.308
	Fruit and vegetables intake		<b>9.368</b>	<b>0.002</b>		3.000	0.083
	$\Delta$ HFSS snacks change score		3.585	0.058		0.105	0.746
	$\Delta$ HFSS meals change score		0.333	0.564		2.047	0.152
	$\Delta$ Fruit and vegetables change score		0.243	0.622		0.012	0.912
	Physical activity		0.674	0.412		0.016	0.899
	Alcohol consumption		0.022	0.883		<b>11.158</b>	<b>0.001</b>
	Smoking status		0.026	0.873		2.306	0.129

All predictors and significant time interactions  
(N=1416)

1803.355

1416.522

Time\*HFSS snacks intake

3.775 0.052

Time\*Alcohol consumption

**5.078 0.024**

---

*Models also included Time as a covariate. Type III tests are shown only. There were no material changes in significance of main effects. In the 'Decrease weight vs all other' model, baseline BMI, HFSS snacks intake, fruit and vegetable intake, and HFSS snacks change score continuous variables were converted to categorical variables due to violating the linearity of logit assumption. QIC is a relative, 'lower is better' measure of goodness of fit.  $W \chi^2$ : Wald Chi-square.*

**Table S10.** Univariate unadjusted and adjusted models for each predictor variable and change in self-reported BMI as a binary outcome using complete cases.

Binary BMI outcome variable	Increase BMI vs all other (<25 BMI)				Decrease BMI vs all other (≥25 BMI)				
	Unadjusted models		Adjusted models w/interaction		Unadjusted models		Adjusted models w/interaction		
Predictor	W	χ <sup>2</sup>	p	W	χ <sup>2</sup>	p	W	χ <sup>2</sup>	p
Gender	0.005		0.945	0.013		0.909	0.387		0.534
Time				<b>5.261</b>		<b>0.022</b>			<b>23.65</b>
Time*Gender				0.675		0.411			0.150
Ethnicity	0.110		0.741	0.115		0.735	2.098		0.147
Time				0.585		0.444			**
Time*Ethnicity				0.014		0.906			**
Baseline BMI	<b>12.444</b>		<b>&lt;0.001</b>	<b>12.274</b>		<b>&lt;0.001</b>	1.160		0.281
Time				2.712		0.100			1.067
Time*Baseline BMI				0.061		0.804			3.070
Age	0.507		0.476	0.294		0.588	2.094		0.148
Time				0.639		0.424			0.032
Time*Age				1.615		0.204			1.233
Occupation and work from home	1.250		0.535	1.119		0.572	2.551		0.279
Time				3.553		0.059			<b>21.051</b>
									<b>&lt;0.001</b>

Time*Occupation and work from home			1.103	0.576			4.483	0.106
Socioeconomic score	<b>8.415</b>	<b>0.038</b>	7.772	0.051	3.773	0.287	4.420	0.220
Time			<b>3.905</b>	<b>0.048</b>			<b>13.021</b>	<b>&lt;0.001</b>
Time*Socioeconomic score			2.811	0.422			2.022	0.568
Living arrangements	1.504	0.471	1.376	0.502	1.221	0.543	0.546	0.761
Time			<b>4.719</b>	<b>0.030</b>			<b>23.423</b>	<b>&lt;0.001</b>
Time*Living arrangements			0.868	0.648			3.006	0.222
Isolation status	2.858	0.091	0.662	0.416	2.764	0.096	0.553	0.457
Time			1.988	0.159			<b>18.484</b>	<b>&lt;0.001</b>
Time*Isolation status			0.017	0.898			1.885	0.170
Quality of life	0.162	0.687	0.001	0.980	<b>5.064</b>	<b>0.024</b>	2.430	0.119
Time			2.208	0.137			0.290	0.590
Time*Quality of life			3.74	0.053			0.264	0.607
HFSS snacks intake	1.213	0.271	0.533	0.466	<b>5.843</b>	<b>0.016</b>	<b>6.411</b>	<b>0.011</b>
Time			0.493	0.483			<b>15.768</b>	<b>&lt;0.001</b>
Time*HFSS snacks intake			0.416	0.519			0.784	0.376
HFSS meals intake	1.095	0.295	1.281	0.258	1.512	0.219	1.204	0.273
Time			3.188	0.074			<b>20.456</b>	<b>&lt;0.001</b>
Time*HFSS meals intake			0.136	0.713			1.537	0.215

Fruit and vegetables intake	<b>8.678</b>	<b>0.003</b>	<b>7.481</b>	<b>0.006</b>	<b>5.980</b>	<b>0.014</b>	<b>6.847</b>	<b>0.009</b>
Time			0.628	0.428			2.681	0.102
Time*Fruit and vegetables intake			0.070	0.791			0.060	0.807
$\Delta$ HFSS snacks change score	0.751	0.386	0.824	0.364	0.515	0.473	0.047	0.829
Time			<b>5.343</b>	<b>0.021</b>			<b>17.772</b>	<b>&lt;0.001</b>
Time* $\Delta$ HFSS snacks change score			1.898	0.168			2.031	0.154
$\Delta$ HFSS meals change score	2.814	0.093	2.514	0.113	1.079	0.299	0.726	0.394
Time			3.232	0.072			<b>26.727</b>	<b>&lt;0.001</b>
Time* $\Delta$ HFSS meals change score			0.344	0.558			2.756	0.097
$\Delta$ Fruit and vegetables change score	0.052	0.820	0.079	0.778	0.054	0.816	0.065	0.799
Time			<b>4.413</b>	<b>0.036</b>			<b>25.119</b>	<b>&lt;0.001</b>
Time* $\Delta$ Fruit and vegetables change score			0.095	0.758			0.041	0.840
Physical activity	0.623	0.430	0.477	0.490	0.204	0.652	0.021	0.884
Time			1.716	0.190			<b>15.761</b>	<b>&lt;0.001</b>
Time*Physical activity			1.324	0.250			2.465	0.116
Alcohol consumption	0.222	0.638	0.155	0.694	<b>6.014</b>	<b>0.014</b>	<b>4.769</b>	<b>0.029</b>
Time			3.336	0.068			<b>8.708</b>	<b>0.003</b>
Time*Alcohol consumption			0.475	0.491			2.549	0.110

Smoking status	0.958	0.328	0.682	0.409	<b>5.260</b>	<b>0.022</b>	<b>4.397</b>	<b>0.036</b>
Time			<b>4.700</b>	<b>0.030</b>			<b>9.806</b>	<b>0.002</b>
Time*Smoking status			0.766	0.381			1.886	0.170

*\*\*Model did not compute due to numerical errors. In the 'Increase BMI vs all other' GEE model, the baseline BMI continuous variable was converted to a categorical variable, due to violating the linearity of logit assumption. Missingness with each model depends on if participants responded to survey questions relating to the predictor variable. W  $\chi^2$ : Wald Chi-square.*

**Table S11.** Full GEE model containing all predictor variables and the full GEE model containing all predictor variables and significant predictor\*time interactions from univariate models adjusted for time, for a change in self-reported BMI as a binary outcome using complete cases.

Binary BMI outcome variable		Increase BMI vs all other (<25 BMI)			Decrease BMI vs all other (>=25 BMI)		
Model	Predictor	QIC	Type III Test		QIC	Type III Test	
			W $\chi^2$	p		W $\chi^2$	p
All predictors (N=673/743)		838.478			871.439		
	Gender		0.010	0.921		<b>4.132</b>	<b>0.042</b>
	Ethnicity		0.131	0.718		2.434	0.119
	Baseline BMI		<b>8.589</b>	<b>0.003</b>		0.842	0.359
	Age		2.782	0.095		<b>4.286</b>	<b>0.038</b>
	Occupation and work from home		4.436	0.109		2.555	0.279
	Socioeconomic score		4.434	0.218		1.976	0.577
	Living conditions		1.567	0.457		2.749	0.253
	Isolation status		0.279	0.597		0.241	0.624
	Quality of Life		0.407	0.523		1.167	0.280
	HFSS snacks intake		1.073	0.300		<b>7.113</b>	<b>0.008</b>
	HFSS meals intake		1.729	0.189		0.132	0.716
	Fruit and vegetables intake		<b>5.628</b>	<b>0.018</b>		<b>5.052</b>	<b>0.025</b>
	$\Delta$ HFSS snacks change score		1.881	0.170		1.272	0.259
	$\Delta$ HFSS meals change score		2.430	0.119		0.581	0.446
	$\Delta$ Fruit and vegetables change score		0.100	0.752		0.007	0.932
	Physical activity		0.047	0.828		0.015	0.904
	Alcohol consumption		0.397	0.528		<b>12.353</b>	<b>&lt;0.001</b>
	Smoking status		<0.001	0.999		1.709	0.191

*Models also included Time as a covariate. Type III tests are shown only. In the 'Increase BMI vs all other' GEE model, the baseline BMI continuous variable was converted to a categorical variable, due to violating the linearity of logit assumption. There were no material changes in significance of main effects. QIC is a relative, 'lower is better' measure of goodness of fit.  $W \chi^2$ : Wald Chi-square.*