STUDY OF BLOOD GROUP ANALYSIS AND ITS CORRELATION WITH LYMPHOPENIA IN COVID-19 INFECTED CASES – OUR EXPERIENCE IN TERTIARY CARE HOSPITAL

Authors:

Swarupa Ravuri ¹
Assistant Professor,
Department of Pathology,
Government Medical College,
Suryapet, Telangana, India.
Email Id: swarupa.ravuri@gmail.com

Corresponding Authors: Saritha Cigiri ², Harika Kalangi³, Anunayi Jeshtadi⁴, K.Nithesh Kumar⁵, Nampally Vamshi Krishna⁶, Srilatha.Bacchu⁷, Sudhakar.K MBBS⁸

²Associate Professor, Department of Pathology, Government Medical College, Suryapet, Telangana, India; Email: csaritha7319@gmail.com

³ Junior Resident, Suryapet, Telangana, India; Email Id: harikakalangi1693@gmail.com

⁴Professor and HOD, Department of Pathology, Government Medical College, Suryapet, Telangana, India; Email: anunayi@1990yahoo.com

⁵Professor in Community Medicine, Mamata Medical Collage, Khammam, Telangana, India; Email Id: ramyanithesh@gmail.com

⁶Junior resident, Suryapet, Telangana, India; Email: Vamshi260500@gmail.com

⁷CAS, Area hospital, Suryapet, Telangana, India; Email: Srilatha.happy2help@gmail.com

⁸Dr.K.Sudhakar, MBBS Principal Investigator, Clinical Research, Hyderabad, India; Email Id: sudhakar,omc@gmail.com.

ABSTRACT:

AIMS AND OBJECTIVES

To study the distribution and frequencies of ABO and Rhesus (Rh) blood groups among confirmed cases of Covid-19 infection. We also studied the relation between ABO blood group system and lymphopenia and studied the gender association in COVID-19 patients.

METHODES

A hospital based retrospective study was conducted at Government Medical College Suryapet from 01 Aug 2020 to 30 Sep 2020. A total of 200 Covid cases were included in the study who came to the hospital with the complaints of Fever, Sore throat, Body Pains, Cough, Breathlessness, Diarrhoea etc. Patients confirmed Covid infection was tested for blood grouping and RH typing by using forward blood grouping with the help of commercially available standard monoclonal antisera. CBP was processed in sysmax 5 part Haematology

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

STUDY OF BLOOD GROUP ANALYSIS AND ITS CORRELATION WITH LYMPHOPENIA IN COVID-19 INFECTED CASES – OUR EXPERIENCE IN TERTIARY CARE HOSPITAL

analyser. Blood group frequency was tested also assessed the gender association, Covid patients presents with lymphopenia the relation between the ABO blood group and lymphocyte count was determined.

RESULTS

Males were more compared to the females .Middle aged group male patients were more commonly involved. Most predominant blood group was group B 79(39.5%), group O 78(39%),group A 37(18.5%), group AB 6(3%),most of them were 190 (95%)Rh positive, only 10 Rh negative (5%).To assess the Lymphopenia in our study we divided the absolute lymphocyte count into 5 groups. Group 1 cases are more 58 (29%), Group 2 91(45.5%), Group 3 30 (15%), Group 4 16(8%), Group 5 5(2.5%).

CONCLUSION

Male patients with blood group B were more compared to other blood groups however more number of studies are necessary to confirm these findings in a larger sample and among individuals of different ethnicities.

Keywords: ABO, BLOOD GROUPS, Rhesus(Rh) typing.

INTRODUCTION:

Corona virus disease also named Novel Corona Virus. First case was reported in December 2019 in Wuhan city and it was spread rapidly, more than 1,00,000 people were infected by the end of March 2020. This began in Wuhan wet-market, deals with animal trade business of different kinds of horse, bats, poultry, snakes, marmots. Recent reports have highlighted the role of zoonotic links, cross-species jumping and spill over conjuncture between animals and human transmission, before acquiring direct human-to-human contact. WHO announced a public health emergency of international concern. There are currently 7 known coronavirus that can infect human, Severe Acute Respiratory Syndrome (SARS), Middle East Respiratory Syndrome (MERS). Exact mode of transmission was unknown. Major routes of transmission person to person are through respiratory droplets and close contact with patients. Incubation period is 2-14 days, clinically patients with symptoms like fever, fatigue, dry cough and in severe cases presents with acute respiratory distress, septic shock, even death. Corona virus

STUDY OF BLOOD GROUP ANALYSIS AND ITS CORRELATION WITH LYMPHOPENIA IN COVID-19 INFECTED CASES – OUR EXPERIENCE IN TERTIARY CARE HOSPITAL

can affect the various systems in our body like Respiratory, Enteric, Neurological, Hepatic and renal.

Pathogenesis:

The mode of entry of this virulent pathogens is through respiratory droplets, tears, body fluids, mucous membranes of the eyes, mouth or nose & another mode of transmission through feco-oral route of transmission. Virus shows Tropism once it enters the body, type 2 pneumocystis and ciliated bronchial epithelial cells through angiotensin-converting enzyme 2 (ACE2) receptors and the immune cells like the dendritic cells and the macrophages. The incubation period is about 02 to 14 days post infection. By eliciting clinical data and details characteristics and determining, what were the risk factors may be associated with severity progression of the disease from the virus. Angiotensin converting enzyme 2 (ACE-2) host cell receptor plays a critical role in the entry of the virus into the cell to cause the infection. Variable expression of ACE-2 expression in the airway epithelial cells is one of the hypothesis for variable susceptibility to disease. In one study Gullion et al investigated the relation between natural antibodies of ABO blood system and ACE2 receptors, how these antibodies inhibit the adhesion of virus protein to the ACE receptors, he observed that S protein / ACE 2 dependent adhesion of Chinese hamster ovary cells to an ACE 2 expressing cell line could be inhibited by either monoclonal or anti-A antibodies may block the interaction between Corona virus and ACE 2 receptors providing the protection against the infection into the host. Blood tests have plays an important role in diagnosing the disease.

ABO blood grouping, discovered in 1900 by Austin immunologist Karl Landsteiner to explain the phenomena of red blood cell (RBC) agglutination, is well-documented hypothesis. Landsteiner carbohydrate moieties are genetically inherited. These moieties suggest correlation between ABO blood type, cardiovascular disease, cancer and certain infections including SARS virus. In humans 4 types of ABO blood groups namely A, AB, B, O, located on chromosome 9 (9q34.2). Blood group antigens are one of the main antigens named human histo-blood group antigens (HBG'S) located on surface of RBC'S. Difference in the expression of antigen can increase or decrease host susceptibility to many infections. Blood groups O have significantly high risk of hepatitis B. HBV infection more in Rh-Positive Donors. Rotavirus gastro enteritis was significantly more prevalent among blood group A patients, malaria patients with blood group A had high risk of anaemia compared to

STUDY OF BLOOD GROUP ANALYSIS AND ITS CORRELATION WITH LYMPHOPENIA IN COVID-19 INFECTED CASES – OUR EXPERIENCE IN TERTIARY CARE HOSPITAL

another blood groups. AB blood group had high risk of developing Dengue haemorrhagic fever O blood group patients more susceptible to norovirus infection. SARS Corona virus is a new virus we have to find out which blood group individuals susceptible to Covid 19 infection. A retrospective study was conducted to explore the relationship and also lymphopenia is a common feature of Covid 19 patients associated with severity disease. The association between ABO blood groups and lymphocytes count was also investigated.

Materials & Methods:

Study design:	Descriptive, Retrospective & prospective study
Study population:	Patients who referred to department of pathology
Location of study:	Government Medical College/Hospital, Suryapet, Telangana, India
Study duration:	2 Months
Inclusion criteria:	Patients with above 18 years were included
Exclusion criteria:	Patients with below 18 years were excluded
Sampling method:	Consecutive sampling method
Sample size:	A total of 200 were included

METHODS:

A retrospective study was performed at Government Medical College/ Hospital Suryapet, Telangana. In this study 200 patients who have admitted in the hospital with the symptoms of Covid 19 at least one symptom or sign of cough, respiratory distress, fever or features that are unexplained by any other disease, history of travel to other country, contact with the relatives or close contact with the patients that who are confirmed positive for Covid 19 by Real Time PCR (RT-PCR) in the last 14 days. Patients with any of the above said symptoms of Covid 19 comes to us during the study period from 01 Aug 2020 to 30 Sep 2020 were enrolled in Government Medical College Suryapet, all were subjected to demographic, clinical feature, laboratory findings. Demographics include age, gender, hypertension, diabetics and heart diseases. Laboratory investigations include Throat swab test and RT-PCR confirmed cases were further investigated CBP, NLR, PLR, blood grouping and RH typing, LFT, Creatinine, LDH. Only the results of RT-PCR, CBP, blood grouping were used for this study. CBP was

STUDY OF BLOOD GROUP ANALYSIS AND ITS CORRELATION WITH LYMPHOPENIA IN COVID-19 INFECTED CASES – OUR EXPERIENCE IN TERTIARY CARE HOSPITAL

processed in Sysmax 5 part analyser, and blood grouping was performed with standard protocols. Blood grouping was determined by forward grouping (cell grouping) by agglutination methods with the help of commercially available standard monoclonal antisera A, antisera B, antisera AB and antisera D manufactured by Tulip Diagnostics (p) Ltd Verna Goa. All the blood group O were tested by anti-H lectin Tulip Diagnostics (p) Ltd Verna Goa to rule out Bombay blood group. These antisera mixed with patient blood and check for agglutination accordingly blood group is determined. Rh typing was done to see the Rh factor on the surface of the red blood cells. Rh positive if cells have this surface protein, Rh Negative if you do not have this cell surface protein. All information was obtained and analysed with standard excel programme.

ASSOCIATION ANALYSIS:

The association between different blood types and Covid 19. According to gender, subgroups were stratified to assess whether there was a significant difference between blood type and the incidence of COVID 19. Correlation analysis between blood group and lymphocytes count was also performed.

STATISTICAL ANALYSIS:

Statistical analysis was carried out using the Statistical Package for Social Sciences (SPSS) VERSION 21.0. Independent sample t-tests were used for age, white blood cell count, lymphocyte count, neutrophil lymphocyte ratio. Chi-square test with 95% confidence interval is used to test ABO blood group frequency in all populations and different gender subgroup was tested. Analysis association between the ABO blood grouping and the lymphocyte count was performed with analysis of variance (ANOVA) AND A LINEAR REGRESSION MODEL. A $p < 0.05$ was considered significant.

RESULTS:

A total of 200 patients were included in this study where Males were 135 (67.5%) & females were 65 (32.5%). Age distribution was shown in the table No.01. Most of the patients belong to 50 years (25.5%) of age group & 60 years (21%). Maximum age was 85 years, minimum age 19 years and mean age was 50 years.

STUDY OF BLOOD GROUP ANALYSIS AND ITS CORRELATION WITH LYMPHOPENIA IN COVID-19 INFECTED CASES – OUR EXPERIENCE IN TERTIARY CARE HOSPITAL

Descriptive Statistics				
	<i>Mean</i>	<i>SD</i>	<i>Maximum</i>	<i>Minimum</i>
Age	50.91	14.67	85.00	19.00
Neutrophil lymphocytes ratio	06.72	05.81	34.60	00.60
Platelet lymphocytes Ratio	246.90	183.04	1903.00	34.50
MPV	10.02	01.08	17.90	07.80
Platelet Count	292846.17	128954.314	751000.00	233.00
Lymphocytes	1.62	0.94	5.70	0.04
WBC Count	10089.62	4968.40	3600.00	11.80
HB	12.90	2.09	18.09	6.50

Table No.01 shows the age distribution of patients.

Blood group distribution shown in the PIE diagram No. 01

STUDY OF BLOOD GROUP ANALYSIS AND ITS CORRELATION WITH LYMPHOPENIA IN COVID-19 INFECTED CASES – OUR EXPERIENCE IN TERTIARY CARE HOSPITAL

Most predominant blood group was group B 79(39.5%), group O 78(39%), group A 37(18.5%), group AB 6 (3%), most of them were Rh positive only 10 Rh negative (5%).

In addition lymphocyte count depletion is related to severity of the disease we also studied the relation between lymphopenia and blood group association shown in Table 2

	Blood group	N	Mean	SD	P value
Lymphocytes	A	37	1.91	1.04	0.1
	AB+	6	1.10	0.40	
	B+	79	1.61	0.87	
	O+	78	1.53	0.96	
	Total	200	1.62	0.94	

WBC count	A	37	9638.16	4264.97	0.8
	AB+	6	9200.00	2920.27	
	B+	79	10374.84	5075.20	
	O+	78	10083.33	5329.44	
	Total	200	10089.62	4968.40	

Table No.2 showing the lymphocyte count and WBC count in association between the different blood groups.

Lymphopenia in our study we divided the absolute lymphocyte count into 5 groups. Group 1 0- 1.0 Group 2 1.1 – 2.0, Group 3 2.1-3.0, Group 4 3.1-4.0, Group 5 >4

Group 1 cases are more 58 (29%), Group 2 91(45.5%), Group 3 30 (15%), Group 4 16(8%), Group 5 5(2.5%). Mean lymphocyte count was 1.62 minimum was 0.04 and maximum was 5.70. P-value counted in different blood groups it was 0.1 found to be significant. Association between lymphocyte count WBC count and blood groups had shown in box diagrams 01& 02.

STUDY OF BLOOD GROUP ANALYSIS AND ITS CORRELATION WITH LYMPHOPENIA IN COVID-19 INFECTED CASES – OUR EXPERIENCE IN TERTIARY CARE HOSPITAL

Correlations									
	Age	Neutrophil lymphocytes ratio	Platelet lymphocytes Ratio	Ratio	MPV	Platelet Count	Lymphocytes	WBC count	HB
Age	Pearson Correlation	0.106	0.131	0.106	-0.002	0.051	-0.11	0.037	-0.221*
	Sig. (2-tailed)	0.064	0.064	0.136	0.762	0.475	0.123	0.603	0.002
Neutrophil lymphocytes ratio	Pearson Correlation	0.131	0.106	0.611**	-0.068	0.003	-.414**	.331*	-0.041
	Sig. (2-tailed)	0.064	0.106	0	0.338	0.966	0	0	0.562
Platelet lymphocytes Ratio	Pearson Correlation	0.106	.611**	1	-.182*	.348**	-.392**	-0.007	-0.066
	Sig. (2-tailed)	0.136	0		0.01	0	0	0.922	0.352
MPV	Pearson Correlation	-0.022	-0.068	-.182*	1	-.378**	-0.034	-0.089	0.124
	Sig. (2-tailed)	0.762	0.338	0.01		0	0.635	0.21	0.079

STUDY OF BLOOD GROUP ANALYSIS AND ITS CORRELATION WITH LYMPHOPENIA IN COVID-19 INFECTED CASES – OUR EXPERIENCE IN TERTIARY CARE HOSPITAL

Platelet count	Pearson Correlation	0.051	0.003	.348*	-.378*	1	.227**	.324*	-0.104
	Sig. (2-tailed)	0.475	0.966	0	0		0.001	0	0.144
Lymphocytes	Pearson Correlation	-0.11	-.414**	-.392*	-0.034	.227**	1	.247*	0.088
	Sig. (2-tailed)	0.123	0	0	0.635	0.001		0	0.216
WBC Count	Pearson Correlation	0.037	.331**	-0.007	-0.089	.324**	.247**	1	-0.051
	Sig. (2-tailed)	0.603	0	0.922	0.21	0	0		0.47
HB	Pearson Correlation	-.221**	-0.041	-0.066	0.124	-0.104	0.088	-0.051	1
	Sig. (2-tailed)	0.002	0.562	0.352	0.079	0.144	0.216	0.47	
** Correlation is significant at the 0.01 level (2-tailed).									
* Correlation is significant at the 0.05 level (2-tailed).									

Table 3 shows Pearson correlation between different parameters and values less than 0.01 found to be significant.

STUDY OF BLOOD GROUP ANALYSIS AND ITS CORRELATION WITH LYMPHOPENIA IN COVID-19 INFECTED CASES – OUR EXPERIENCE IN TERTIARY CARE HOSPITAL

Descriptives					
	Blood group	N	Mean	SD	P value
Neutrophil lymphocytes ratio	A	37	6.41	7.01	0.8
	AB+	6	8.17	6.30	
	B+	79	6.57	5.09	
	O+	78	6.92	5.95	
	Total	200	6.72	5.81	
Platelet lymphocytes Ratio	A	37	260.79	302.00	0.8
	AB+	6	291.00	178.85	
	B+	79	238.85	129.74	
	O+	78	245.07	156.54	
	Total	200	246.90	183.04	
MPV	A	37	9.88	0.95	0.5
	AB+	6	9.70	0.28	
	B+	79	10.01	0.93	
	O+	78	10.13	1.30	
	Total	200	10.02	1.08	
Platelet count	A	37	314783.78	153810.51	0.7
	AB+	6	290500.00	94256.57	
	B+	79	289835.44	124779.34	
	O+	78	285669.65	123587.72	
	Total	200	292846.17	128954.31	
Lymphocytes	A	37	1.91	1.04	0.1
	AB+	6	1.10	0.40	
	B+	79	1.61	0.87	
	O+	78	1.53	0.96	
	Total	200	1.62	0.94	
WBC count	A	37	9638.16	4264.97	0.8
	AB+	6	9200.00	2920.27	
	B+	79	10374.84	5075.20	
	O+	78	10083.33	5329.44	
	Total	200	10089.62	4968.40	
HB	A	37	12.75	2.51	0.7
	AB+	6	12.60	1.80	
	B+	79	13.10	1.96	
	O+	78	12.78	2.04	
	Total	200	12.90	2.09	

STUDY OF BLOOD GROUP ANALYSIS AND ITS CORRELATION WITH LYMPHOPENIA IN COVID-19 INFECTED CASES – OUR EXPERIENCE IN TERTIARY CARE HOSPITAL

STATE/ UNION/ TERRITORIES	POPULATION (CENSUS) 2011	NUMBER OF COVID- 19 INFECTED CASES	INFECTED CASES PER MILLION OF POPULATION	NUMBER OF DEATHS DUE TO COVID- 19	DEATH RATE PER MILLION OF POPULATION	BLOOD GROUP - A - B - O - AB NUMBER (%)	BLOOD GROUP - A - B - O - AB NUMBER (%)	BLOOD GROUP - A - B - O - AB NUMBER (%)	BLOOD GROUP - A - B - O - AB NUMBER (%)
ANDHRA PRADESH	49386799	5269	106.688	78	1.579	30052 (38.926)	18784 (24.330)	23971 (31.049)	4395 (5.692)
GUJARAT	60439692	21521	356.073	1347	22.286	18428 (32.131)	13602 (23.717)	20270 (35.343)	5051 (8.807)
HARYANA	25351462	5579	220.066	52	2.051	1769 (29.851)	1335 (22.527)	2269 (38.288)	553 (9.331)
HIMACHAL PRADESH	6864602	451	65.699	6	0.874	1230 (25.802)	1175 (24.648)	1707 (35.808)	655 (13.740)
PUNJAB	27743338	2805	101.105	55	1.982	5411 (34.085)	2904 (18.292)	6038 (38.034)	1522 (9.587)
RAJASTHAN	68548437	11600	169.223	259	3.778	5260 (33.889)	3479 (22.414)	5575 (35.919)	1207 (7.776)
TAMILNADU	72147030	36841	510.637	326	4.518	5925 (33.234)	3528 (19.789)	6489 (36.397)	1886 (10.578)
TELENGANA	35193978	4111	116.809	156	4.432	9525 (41.582)	4368 (19.069)	7704 (33.633)	1309 (5.714)
TRIPURA	3673917	895	243.609	1	0.272	331 (30.876)	288 (26.865)	329 (30.690)	124 (11.567)
UTTAR PRADESH	199812341	11610	58.104	321	1.606	41489 (29.328)	30246 (21.380)	56430 (39.889)	13300 (9.401)
UTTARAKHAND	10086292	1562	154.863	15	1.487	6239 (27.625)	6648 (29.436)	7204 (31.898)	2493 (11.038)

STUDY OF BLOOD GROUP ANALYSIS AND ITS CORRELATION WITH LYMPHOPENIA IN COVID-19 INFECTED CASES – OUR EXPERIENCE IN TERTIARY CARE HOSPITAL

Studies from other countries Li et al, Zhao et al studies they found that the blood group A patients infected with SARS CoV 2 higher than the healthy controls and blood group O in SARS CoV 2 was low, our study not correlating with these two studies, another study conducted by Michel zietz study blood group B association with covid19 infection is more than the other blood groups.

Zhao et al 2020 conducted a study with 2173 patients with covid confirmed patients it was meta-analysis study they concluded that patient's ABO blood group and the patients with blood group A were associated with high risk when compared to other non A blood groups, O blood group was associated with lower risk for the infection compared to others.

Yamamoto et al 2020 study concluded that s protein produced in A, AB, B blood group individuals with their antigens present on the surface of the red blood cells would produce corresponding respective antibodies. These antibodies would block the interaction with S-protein and ACE receptors.

Tanigawa et al in 2020 reviewed the current literature on COVID19 and associated phenotypes. It was a study conducted in UK bio bank with 337579 cases, in this study they aggregated human leucocyte antigen and ABO blood type frequencies. They found that blood group O show consistent risk reduction in accordance with Zhao et al study. The mechanism of association and susceptibility of patients with Blood group B among patients it is unknown. Normally interaction between the spike proteins and ACE-2 receptors is hampered by the antibody type A, which gives the explanation of protection of blood group O against the COVID 19 infection and mortality. The role of ABO antibodies altering the interaction between the spike proteins of SARS Cov 2 and ACE 2 RECEPTORS still unknown. For the multiplication of virus and spread of infection to another hosts, the spike protein of virion carries A, B, AB glycan antigen depends on the blood group of carrier. Blood group O patients contains both A & B antibodies, infection rate with A, B & AB antigens would be reduced. Protection also depends on the ABO antibody titers. These variation of observations on association of blood groups and COVID 19 infection could be due to different strains of SARS Cov 2 with varying pathogenicity and discrepancies in the investigated population.

STUDY OF BLOOD GROUP ANALYSIS AND ITS CORRELATION WITH LYMPHOPENIA IN COVID-19 INFECTED CASES – OUR EXPERIENCE IN TERTIARY CARE HOSPITAL

In our research we also studied the relationship between ABO blood grouping & lymphocytes Count. We also further investigated the importance of Lymphocyte count in the evaluation of severity of Covid 19 infection. The lymphocyte count in B type blood group patients mean was 1.91. AB blood group patients were 6 in number this blood group patients present with low lymphocyte count 1.10 with high NLR values and P-value calculated 0.1 found to be significant.

Preventive measures such as social distancing, hand hygiene, home or hospital quarantine as well as restrictions on movement (lockdown) help in preventing the spread of the disease. Isolation and supportive therapy, including ventilator support, are the mainstays of treatment of infected patients.

Limitations

Meanwhile several drawbacks existed in our study, limited sample size of Covid – 19 patients, second regional selection bias. This is an observational study attempts were made to control for confounding.

Conclusion:

Male patients with blood group B were more effective when compared to other blood groups, however more number of studies are necessary to confirm these findings in a larger sample and among individuals of different ethnicities.

BIBLIOGRAPHY

1. Wang, D., Hu, B., Hu, C., Zhu, F., Liu, X., Zhang, J., et al. (2020). Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. JAMA. 323, 1061–1069. doi: 10.1001/jama.2020.1585
2. Wang, D., Hu, B., Hu, C., Zhu, F., Liu, X., Zhang, J., et al. (2020). Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. JAMA. 323, 1061–1069. doi: 10.1001/jama.2020.1585
3. Wu, Z., and McGoogan, J. M. (2020). Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. JAMA. 323, 1239–1242. doi: 10.1001/jama.2020.2648
4. Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., et al. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. Lancet 395, 497–506. doi: 10.1016/S0140-6736(20)30183-5
5. Bennett J, Dolin R, Blaser MJ. Principles and Practice of Infectious Disease . 8th Edition Elsevier / Saunders , PA, USA (2014).

STUDY OF BLOOD GROUP ANALYSIS AND ITS CORRELATION WITH LYMPHOPENIA IN COVID-19 INFECTED CASES – OUR EXPERIENCE IN TERTIARY CARE HOSPITAL

5. Meng J, Xiao G, Zhang J, He X, Ou M, Bi J, Yang R, diW,WangZ, Li Z, Gao H, Liu L, Zhang G (2020) Renin-angiotensin systeminhibitors improve the clinical outcomes of COVID-19 patients with hypertension. *Emerg Microbes Infect* 9(1):757–760.

6. Hussain A, Bhowmik B, do Vale Moreira NC (2020) COVID-19and diabetes: knowledge in progress. *Diabetes Res Clin Pract* 162:108142.

7. GuoW, Li M, Dong Y, et al (2020) Diabetes is a risk factor for theprogression and prognosis of COVID-19 [published online aheadof print, 2020 Mar 31]. *DiabetesMetab Res Rev*. <https://doi.org/10.1002/dmrr.3319>.

8. Zhao Q,Meng M, Kumar R, et al (2020) The impact of COPD andsmoking history on the severity of COVID-19: A systemic reviewand meta-analysis. *J Med Virol*. <https://doi.org/10.1002/jmv.25889>.

9 Wrapp, D., Wang, N., Corbett, K. S., Goldsmith, J. A., Hsieh, C. L., Abiona, O., et al. (2020). Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* 367, 1260–1263. doi: 10.1126/science.abb2507

10 Leung JM, Yang CX, Tam A, Shaipanich T, Hackett TL, SingheraGK, Dorscheid DR, Sin DD (2020) ACE-2 expression in the smallairway epithelia of smokers and COPD patients: implications forCOVID-19. *Eur Respir J* 55:2000688.

11. Guillon, P., Clement, M., Seville, V., Rivain, J. G., Chou, C. F., Ruvoen-Clouet, N.,et al. (2008). Inhibition of the interaction between the SARS-CoV spike proteinand its cellular receptor by anti-histo-blood group antibodies. *Glycobiology* 18,1085–1093. doi: 10.1093/glycob/cwn093

12 Melzer, D., Perry, J. R., Hernandez, D., Corsi, A. M., Stevens, K., Rafferty, I., et al. (2008). A genome-wide association study identifies protein quantitative trait loci (pQTLs). *PLoS Genet*. 4:e1000072. doi: 10.1371/journal.pgen.1000072.

13.Wiggins, K. L., Smith, N. L., Glazer,N. L., Rosendaal, F. R., Heckbert, S. R., Psaty, B.M., et al. (2009). ABO genotype and risk of thrombotic events and hemorrhagicstroke. *J. Thromb. Haemost.* 7, 263–269. doi: 10.1111/j.1538-7836.2008.03243.

14.Mohammadali, F., and Pourfathollah, A. (2014). Association of ABO and Rh bloodgroups to blood-borne infections among blood donors in Tehran-Iran. *Iran. J.Public Health* 43, 981–989.

15. Elnady, H. G., Abdel, S. O., Saleh,M. T., Sherif, L. S., Abdalmoneam, N., Kholoussi,N. M., et al. (2017). ABO blood grouping in Egyptian children with rotavirus gastroenteritis. *Prz. Gastroenterol.* 12, 175–180. doi: 10.5114/pg.2017.70469.

16 .Murugananthan, K., Subramaniyam, S., Kumanan, T., Owens, L., Ketheesan,N., and Noordeen, F. (2018). Blood group AB is associated with severe forms of dengue virus infection. *Virusdisease* 29, 103–105.doi: 10.1007/s13337-018-0426-8.

17.ABObloodgroupsystemisassociatedwithCOVID19mortality:AnepidemiologicalinvestigationintheIndianpopulationSunali Padhi1,SubhamSuvankar1,DebabrataDash,VenketeshK.Panda,AbhijitPati,JogeswarPanigrahi,AdityaK.Panda*Departmentof BioscienceandBioinformatics,KhallikoteUniversity,TransitCampus:GMaxBuilding,Konisi,Berhampur,761008Odisha,India. <https://doi.org/10.1016/j.traccli.2020.08.00912467820>/©2020Soci'et'efranc aisedetransfusionsanguine(SFTS).PublishedbyElsevierMassonSAS.Allrightsreserved. *transfusion Clinique et biologie*27(2020)253- 258.

18. Li, J., Wang, X., Chen, J., Cai, Y., Deng, A., and Yang, M. (2020). Associationbetween ABO blood groups and risk of SARS-CoV-2 pneumonia. *Br. J.Haematol.* 190, 24–27. doi: 10.1111/bjh.16797.

19 Zhao, J., Yang, Y., Huang, H., Li, D., and Gu, D., Lu, X., et al. (2020). Relationshipbetween the ABO blood group and the COVID-19 susceptibility. *MedRxiv*.doi: 10.1101/2020.03.11.20031096. [Epub ahead of print].

20. Yamamoto F. ABO Blood Groups and SARS-CoV-2 Infection; 2020.

21. Saxena S. Coronavirus disease-2019: A brief compilation of facts. *J Oral Maxillofac Pathol* 2020;24:5-7.

22. BreimanA,Ruvën-ClouetN,LePendJ.Harnessingthenaturalanti-glycanimmuneresponse to limit the transmission of enveloped viruses suchasSARS-CoV-2.*PLoSPathog*2020;16:e1008556.

MAMATA MEDICAL COLLEGE, KHAMMAM, TELANGANA STATE
Communication of Decision of the Institutional Ethics Committee (IEC)/
Institutional Review Board

(IRB)IEC/IRB No: 3

Protocol Title:		
STUDY OF BLOOD GROUP ANALYSIS AND ITS CORRELATION WITH LYMPHOPENIA IN COVID-19 INFECTED CASES - OUR EXPERIENCE IN TERTIARY CARE HOSPITAL		
Principal Investigator: <u>Dr. RAVURI SWARUPA</u> CO INVESTIGATOR: <u>Dr. K. NITHESH KUMAR</u>		
Name & Address of Institution:		
MAMATA MEDICAL COLLEGE, KHAMMAM		
☒ New review	☐ Revised review	☐ Expedited review
Date of review (DD/MM/YY)		
Date of previous review if revised application: <u>26/6/2021</u>		
Decision of the IEWC / IRB		
☒ Recommended	☐ Recommended with suggestions	
☐ Revision		
Suggestions / Reasons / Remarks:		
Recommended for a period of: <u>one year</u>		

Please note ☆

- Inform IEC/IRB immediately in case of any adverse events and serious adverse events.
- Inform IEC/ IRB in case of any change of study procedure, site and investigator
- This permission is only for period mentioned above. Annual report to be submitted to IEC/IRB.
- Members of IEC/IRB have right to monitor the trial with prior intimation.

Anu N
 Signature of Member Secretary
 IEC/IRB

PROFESSOR & HEAD
DEPARTMENT OF MICROBIOLOGY
MAMATA MEDICAL COLLEGE
KHAMMAM - 507 002 (A.P.)