Exploring the relationship between resting state intra-network connectivity and accelerometer-measured physical activity in pediatric concussion: A cohort study

Bhanu Sharma, Joyce Obeid, Carol DeMatteo, Michael D. Noseworthy, Brian W. Timmons

Bhanu Sharma¹

Joyce Obeid¹

Carol DeMatteo^{2,3}

Michael D. Noseworthy^{4,5,6,7}

Brian W. Timmons^{1*}

¹Child Health and Exercise Medicine Program, Department of Pediatrics, McMaster University

² School of Rehabilitation Science, McMaster University

³ CanChild Centre for Childhood Disability Research, McMaster University

⁴ Imaging Research Centre, St. Joseph's Healthcare, Hamilton, ON, Canada

⁵ Department of Electrical & Computer Engineering, McMaster University

⁶ McMaster School of Biomedical Engineering, McMaster University

⁷ Department of Radiology, McMaster University

* 1280 Main Street West, Hamilton, Ontario, L8S4L8; timmonbw@mcmaster.ca; (905) 521-2100 x 77218

ABSTRACT

Objectives: To explore the association between resting state functional connectivity and accelerometer-measured physical activity in pediatric concussion.

Methods: Fourteen children with concussion (aged 14.54 ± 2.39 years, 8 female) were included in this secondary data-analysis. Participants had neuroimaging at 15.3 ± 6.7 days post-injury and subsequently a mean of 11.1 ± 5.0 days of accelerometer data. Intra-network connectivity of the default mode network (DMN), sensorimotor network (SMN), salience network (SN), and fronto-parietal network (FPN) was computed.

Results: Per general linear models, only intra-network connectivity of the DMN was associated with habitual physical activity levels. More specifically, increased intra-network connectivity of the DMN was significantly associated with higher levels of subsequent accelerometer-measured light physical activity ($F_{(2, 11)} = 7.053$, $p = 0.011$, $R_a^2 = 0.562$; $\beta = 0.469$), moderate physical activity ($F_{(2, 11)} = 7.053$, $p = 0.011$, $R_a^2 = 0.562$; $\beta = 0.725$), and vigorous physical activity ($F_{(2, 11)} = 10.855$, $p = 0.002$, $R_a^2 = 0.664$; $\beta = 0.79$). Intra-network connectivity of the DMN did not significantly predict sedentary time. Likewise, the SMN, SA, and FPN were not significantly associated with either sedentary time or physical activity.

Conclusion: These findings suggest that there is a positive association between the intra-network connectivity of the DMN and device-measured physical activity in children with concussion. Given that DMN impairment can be commonplace following concussion, this may be associated with lower levels of habitual physical activity, which can preclude children from experiencing the symptom-improving benefits of sub-maximal physical activity.

KEY FINDINGS

What are the new findings?

- Intra-network connectivity of the default mode network is associated with subsequent accelerometer-measured light, moderate, and vigorous physical activity within the first-month of pediatric concussion
- Similar associations with physical activity are not observed when examining the intra-network connectivity of the sensorimotor network, salience network, or fronto-parietal network
- Improved connectivity within the default mode network may lead to increased participation in light to vigorous physical activity in pediatric concussion

How might it impact on clinical practice in the future?

- Default mode network impairment is commonplace in concussion, and this may limit children from experiencing the symptom-improving benefits of physical activity
- Adjunctive interventions (e.g., mindfulness) that improve the health of the default mode network should be further studied in pediatric concussion

INTRODUCTION

Concussions are mild traumatic brain injuries that result in functional neurological disturbance in the absence of gross structural damage¹. While on the mild end of the brain injury spectrum, concussions can nonetheless result in enduring symptoms that negatively impact multiple aspects of daily living². In children, a population in which rates of concussion are increasing rapidly³⁻⁵, nearly 30% experience symptoms that persist beyond the expected recovery window of four weeks^{1,6,7}. These symptoms can lead to quality-of-life deficits that can last for months⁸⁻¹¹.

A growing body of research has used resting-state functional magnetic resonance imaging (rs-fMRI). In this study, rs-fMRI was used to provide insight into areas of the brain that are synchronously active at rest in the absence of any particular cognitive stimuli) to explore the functional neuropathology of concussion. While the brain regions and/or networks studied, time of imaging, and rs-fMRI analysis methods themselves have varied across studies, studies consistently show that there are rs-fMRI disturbances in children with concussion in comparison to their healthy peers¹²⁻²³. Collectively, these studies have shown patterns of both hyper- and hypo-connectivity within the initial weeks of injury²⁴, abnormal rs-fMRI activity in children with protracted symptoms¹⁵, and a persistence of functional impairment in asymptomatic children who have been medically cleared to return-to-sport²¹.

With rs-fMRI disturbances in pediatric concussion now better characterized, studies have aimed at understanding the relationship between functional connectivity and other relevant and widely studied clinical outcomes including symptoms, sleep, cognition, mood¹⁵⁻

^{17,25}. However, the associations between resting state brain activity and other salient features of pediatric concussion are still understudied. In particular, the impact of rs-fMRI activity on habitual physical activity in pediatric concussion remains unexplored. Within the current landscape of concussion research and clinical management, wherein the importance of exercise has been increasingly recognized and established in recent years^{26,27}, this represents a considerable knowledge gap. There is now a shift away from prolonged rest^{28,29}, which was the former *status quo* in concussion management³⁰. Conversely, sub-maximal aerobic exercise studies in concussion—and meta-analyses of them—suggest that engaging in physical activity within two-weeks of injury improves symptoms^{26,27,31}. Therefore, there is now considerable evidence that exercise, and conversely prolonged rest, have an impact on concussion recovery. What is not known is how the widely established rs-fMRI disturbances observed in pediatric concussion relate to sedentary time or physical activity levels post-injury.

With the recent wave of research on exercise in pediatric concussion, understanding the relationship between functional brain impairment and physical activity has become as germane as understanding the relationship between rs-fMRI and symptomatology or other clinical features. Therefore, in this study, we examined the relationship between intra-network rs-fMRI activity (within four widely researched and validated resting state networks) and accelerometer-measured habitual physical activity and sedentary time up to one-month post-injury in children with concussion. It was hypothesized that reduced intra-network rs-fMRI activity (a measure of functional neuropathology) is associated with increased sedentary time and reduced physical activity levels.

METHODS

The Hamilton Integrated Research Ethics Board approved this study.

Design

The data reported here were initially collected as part of the *Back-to-Play* study, a larger cohort study (led by the senior authors) with the goal of informing return-to-activity guidelines for children with concussion. This report is a secondary data analysis of accelerometer and rs-fMRI data collected as part of the larger cohort. The present sample is comprised only of participants from the parent study with both neuroimaging data and subsequently collected accelerometer data (up to 1-month post-injury).

Participants

Participants in the parent study were recruited at McMaster Children's Hospital and/or its affiliated rehabilitation and sports medicine clinics. Patients who were diagnosed with concussion by a member of our clinical team were informed about the *Back to Play* study. Those interested in participating were referred to our research team for more information about its objectives, risks, and potential benefits. After this initial discussion, those intent on participating were consented (or assented, along with parental consent, if aged under 16 years) recruited. An intake assessment was then scheduled by the research team as soon after the initial clinical consultation as possible. Exclusion criteria for the larger study included more severe injuries or those requiring more complex care. For the present study, patients were required to have neuroimaging and then subsequently at least 5 days of valid accelerometer wear.

Data collection procedures

Neuroimaging

All neuroimaging data were collected using a 3-Tesla GE Discovery 750 MRI scanner (with a 32-channel phased array receive coil) at the Imaging Research Centre (IRC) at St. Joseph's Healthcare, Hamilton. A screening questionnaire was performed by the IRC imaging technologist to ensure that the scan could be performed safely, and to inform patients and their families about the MRI procedure. The technologist then positioned the patient in the MRI, immobilizing their head with foam pads to minimize motion and to improve patient comfort.

The neuroimaging battery began with a 3-plane localizer with calibration sequences. Anatomical images were then collected, per a 3D IR-prepped fast SPGR T1-weighted sequence (TR/TE=11/36/4.25ms, flip angle=12°, interpolated 512x512 matrix, 22 cm FOV). Immediately prior to the resting state scan, a fieldmap was acquired (to correct for magnetic field inhomogeneities) using the same geometry as the functional scan (as follows). Resting state functional data were collected using axial 2D acquisition, gradient echo EPI, TR/TE=2000/35ms, flip angle=90°, 64x64 matrix, 300 time points (10min), 22 cm FOV). During the resting state scan, patients were asked to remain awake with their eyes open, and not think of "anything in particular".

Accelerometry

Patients were given a compact and light-weight waist-worn tri-axial accelerometer at their intake assessment to wear until self-reported symptom resolution, at which point the

accelerometer was mailed back to the study team and acceleration data were downloaded. The accelerometer used in the *Back-to-Play* study was the ActiGraph GT3x (Pensacola, FL), which has demonstrated high reproducibility in measuring physical activity in acquired brain injury³². Per the parent study, movement was recorded continuously at 30-Hz and downloaded into 3-second epochs. Patients were also given a logbook, in which they noted when the device was put on and off in the morning and evening, respectively, as well as any other times the device was removed (when participating in water-based activities, for example). Patients were instructed by the research team at the intake assessment to wear the accelerometer on the right hip during all waking hours, except for water activities, and how to use the logbook.

Data processing and analyses

Neuroimaging

All MRI pre-processing and analyses were performed in CONN 19c³³, which was run using SPM12 and MATLAB 2020a (Mathworks, Natick, MA). The only exception in the pre-processing pipeline was that functional data were unwarped using the B_0 maps acquired immediately prior to the rs-fMRI scan outside of CONN 19c using *epiunwarp*³⁴, which draws on functionality in FSL^{35,36}. The B_0 -corrected maps were then uploaded into CONN 19c along with respective anatomical data for pre-processing.

The following steps were involved with pre-processing, and were guided by recommendations within CONN 19c and associated publications^{37,38}. First, functional data were realigned and co-registered to a reference image (and adjusted to the movement-defined deformation field associated with the reference image). Second, slice-timing correction was

performed to time-shift and resample the rs-fMRI data to coincide with the mid-point of each TR. Third, using SPM's Artifact Detection Tool (ART)³⁹, outlier scans in the functional data were flagged based on subject-motion exceeding a 0.9mm framewise displacement or BOLD signal fluctuations > 5 standard deviations from the mean signal. Fourth, functional data are normalized/registered to MNI space, based on posterior tissue probability maps. Fifth, spatial smoothing was performed, with a Gaussian kernel of full-width at half-maximum of 6mm. Subsequently, de-noising was performed in CONN 19c which involved two steps: 1) an anatomical component-based noise correction procedure (**aCompCorr**) to “regress out” noise components associated with cerebral white matter/cerebrospinal regions, outlier scans, subject motion (based on a 12-parameter affine transformation of the anatomical to functional data)⁴⁰⁻⁴², and 2) temporal filtering, using a filter ranging from 0.008Hz and above 0.01Hz. After both pre-processing and de-noising, data were inspected visually and per the quality assurance metrics in CONN 19c.

Once data were ready for analysis, intra-network connectivity of four widely studied and validated brain networks was computed^{43,44}, namely the default mode network (DMN), sensorimotor network (SMN), salience network (SN) and fronto-parietal network (FPN) and habitual physical activity in children with concussion. Intra-network connectivity was selected as the measure of choice for this preliminary investigation as it is representative of a resting state network holistically, as has been shown in publications related to other neurological populations⁴⁵⁻⁴⁸.

Average within-network connectivity were computed using the Matlab command line script *conn_withinbetweenROIttest*, which extracted and exported single-subject intra-

network correlation values which were Fisher Z-transformed to improve normality. Seed regions are listed in **Table 1**.

[insert Table 1 here]

Accelerometry

Data were downloaded from the ActiGraph GT3x devices using the accompanying software package, ActiLife Version 6 (Pensicola, FL). Triaxial accelerations collected at 30-Hz were downloaded into 3-sec sampling intervals or epochs, which were selected to reflect the median bout duration for high-intensity activities in children⁴⁹. Once downloaded and converted into a 3-second epoch, wear-time validation was performed in ActiLife. First, a semi-automated procedure was performed to define a non-wear period of minimum length of 5-minutes. These periods were then inspected against the on- and off-times recorded in the patient logbooks, with non-wear periods excluded if they matched records from the patient log books. Days of accelerometer data without an associated log book entry were excluded. Data were then scored to determine activity intensity according to widely used Evenson cut-points⁵⁰ which have the following activity count ranges: sedentary time (0-25 counts/15s), light physical activity (LPA; 26-573 counts/15s), moderate physical activity (MPA; 574-1002 counts/15s), and vigorous physical activity (VPA; 1003+ counts/15s).

Cleaned and scored data were then exported to SPSS Version 27 (IBM Corp. Released 2020. IBM SPSS Statistics for Windows, Version 27.0. Armonk, NY). Only children with more than 4-wear days of data, with each day comprised of more than 600 minutes of wear time,

were included in the analysis⁵¹. Any accelerometer data beyond 30-days post-injury were excluded from the current analysis, given that the scope of the study.

Statistical analyses

Average daily sedentary time, as well as average daily LPA, MPA, and VPA were calculated for cleaned and validated accelerometer cases in SPSS. These summary data were then imported into CONN 19c, wherein the *Calculator* tool was used to build a general linear model (GLM) with intra-network connectivity and age as predictors of average daily activity (of the various intensities) or sedentary time. Therefore, for each network of interest, four models were built to examine the association between intra-network rs-fMRI connectivity and average time per day in: LPA, MPA, VPA, and sedentary. Given the sample size of our study, additional variables were not included in the model; age was selectively chosen given that connectivity of resting state networks has been shown to vary with age⁵².

RESULTS

Overview

Fourteen children with concussion (aged 14.54 ± 2.39 years, 8 female) were imaged at 15.3 ± 6.7 days post-injury and had an average of 11.1 ± 5.0 days of accelerometer data (with ≥ 600 minutes of wear-time per day) thereafter. Average daily monitoring time, sedentary time, and time spent in activity is summarized in **Table 2**.

[insert Table 2 here]

The groupwise average intra-network connectivity values for each of the networks studied are presented in **Table 3** with associated standard deviations and ranges.

[insert Table 3 here]

Relating brain activity and physical activity

From the series of GLMs, the only significant models (and the only network-related beta-coefficients that were significant) pertained to models including the DMN. The intra-network connectivity of the SMN, SA, and FPN did not predict accelerometer-measured sedentary time, LPA, MPA, or VPA. **Figure 1** shows the model-predicted activity levels (along with 95% confidence intervals) by activity type against intra-network connectivity. The figures show that with respect to the DMN, increased intra-network connectivity is closely associated with increased levels of activity, in particular MPA and VPA. Intra-network connectivity of the DMN was not, however, associated with average daily sedentary time.

[insert Figure 1 here]

The DMN-specific models and their relevant parameters for all intensities of activity are summarized in **Table 4**. Both unstandardized beta coefficients (B) and standardized beta coefficients (β) are provided for interpretation. Overall, there was a trend for decreased DMN intra-network connectivity predicting more sedentary time, and conversely, increased intra-network connectivity of the DMN was significantly associated with increased LPA, MPA, and VPA. For significant models, the residuals were normally distributed.

DISCUSSION

This is the first study to examine the relationship between rs-fMRI impairment and subsequent activity levels in pediatric—or adult—concussion. We demonstrate that intra-network connectivity within the DMN—but not the SMN, SA, or FPN—is associated with accelerometer-measured LPA, MPA, and VPA within the first month of injury in pediatric concussion. Adding to the literature demonstrating associations between DMN impairment and concussion symptoms, depression, anxiety, and sleep-impairment^{15,16}, we show that intra-network connectivity within DMN may be implicated with habitual physical activity within the first month of injury in children.

The DMN and physical activity

Originally, the DMN was considered to be a “day dreaming network”, active in the absence of external stimuli as a type of neural baseline⁵³. However, more recent research on the DMN has shown that it is also associated with higher-order cognitive processes and can be active during goal-oriented thoughts and tasks. More specifically, studies now show that the DMN can be engaged when individuals imagine or plan for the future⁵⁴. Other studies suggest that the DMN is important in cognitive processes that result in immediate action or interfere with present behavioural goals⁵⁵. In adults, the DMN has also been linked to “internal mentation”, or introspective thoughts about constructs such as the future or personal intentions^{56,57}. Together, these findings would suggest that DMN impairments may have the capacity to influence planned behaviours, such as participation in physical activity, and that DMN impairments can alter how information about physical activity behaviours are perceived,

processed, and planned. This would help explain our findings, wherein only the DMN was found to be associated with subsequent levels of physical activity. Further research is needed to build on this possibility. Prior research shows that in young adults with concussion, in comparison to healthy controls, a sub-maximal aerobic fitness test resulted in functional disturbance of the DMN^{58,59}. More specifically, said test reduced inter-network connectivity between seed regions of the DMN, namely the PCC and the lateral parietal ROI, PCC and the right lateral parietal ROI, and the PCC and MPFC⁵⁹. Adding to this, our study suggests a bi-directional relationship, wherein intra-network impairment of the DMN may have an impact subsequent physical activity in children with concussion. Given that many studies have demonstrated DMN impairment in pediatric concussion^{14-19,21,22}, and our recent work demonstrates that children with concussion are less active than their healthy peers, the current findings suggest that widespread DMN impairment is associated with reduced levels of activity in children with concussion. Larger pediatric concussion cohorts are required to more definitively characterize the relationship between resting state brain activity and physical activity.

Exercise and concussion management

The status quo in concussion management is changing. The traditional “rest-is-best” approach is being supplanted by an “exercise is medicine” mindset³⁰. However, the present study suggests that in pediatric concussion, DMN impairment (which is a common in concussion) has a moderate association with physical activity. This underscores the need for physicians to actively advise pediatric concussion patients to engage in safe, sub-maximal aerobic exercise after a short (24-48 hour) period of rest, as suggested by current guidelines¹. Otherwise, patients with DMN impairment may not be physically active, and this precludes

them from experiencing the benefits of light physical activity on concussion symptoms.

Further, in the ultimate interest of promoting physical activity, interventions such mindfulness can be prescribed acutely given their positive impact on the functional connectivity of the DMN⁶⁰⁻⁶⁴. Mindfulness is not a contraindication to any other medications that may be prescribed after concussion, and it has other established benefits in brain injury, including improved self-efficacy^{65,66}. Future research is needed in this area.

Generalizability

Our findings need to be studied in adults to understand if they generalize outside of a pediatric context, as several factors may make our findings pediatric-specific. For example, DMN functional connectivity is age-dependent, peaking in adulthood while being less coherent during childhood and senescence⁵²; physical activity also decreases with age⁶⁷. This would suggest that the association between intra-network functional connectivity of the DMN and physical activity may be variable in children when compared to adults.. Further, cardiorespiratory fitness is also related to the functional integrity of multiple brain networks, including the DMN⁶⁸. While the present study did not control for cardiorespiratory fitness (though all participants had sport-related injuries and were physically active prior to injury), similar studies in adults should control for this metric given that it declines non-linearly and in an age-dependent rate in adults⁶⁹, which may be confounding in adult samples with large age variance.

LIMITATIONS

This secondary data analysis is limited by the sample size. Larger studies examining the relationship between the DMN and post-concussion physical activity are warranted. We also cannot rule out the possibility that greater DMN impairment is associated with more severe symptoms that may, in turn, reduce participation in physical activity. This study design does not permit inference on causality. Further, this study did not exhaustively study all resting state networks and/or anatomical regions; future research should expand on the regions of interest. There are also other rs-fMRI and accelerometer analysis methods that can be used to characterize the relationship between brain activity and physical activity. The impact of physical activity on resting state networks in pediatric concussion also warrants study.

CONCLUSIONS

This exploratory study is the first to examine the association between intra-network connectivity and accelerometer-measured physical activity in pediatric concussion. We found that intra-network connectivity of the DMN—but not the SMN, SA, or FPN—was significantly associated with levels of LPA, MPA, and VPA performed within the first month after pediatric concussion. Given the increasing role of aerobic exercise in concussion management, and that the DMN is commonly perturbed following injury, children with concussion may be less likely to be physically active. Clinical management should continue to encourage participation in sub-maximal aerobic activity in the acute stages of injury.

STATEMENTS

Competing interests: The authors have no competing interests to declare.

Contributorship: BW Timmons, MD Noseworthy, and C DeMatteo conceptualized and designed the parent study. C DeMatteo had general oversight over the parent study, with J Obeid and MD Noseworthy maintaining oversight over accelerometer and neuroimaging data collection, respectively. B Sharma created the research question, cleaned and analyzed all accelerometer and neuroimaging data, and prepared the manuscript draft and subsequent revisions.

Acknowledgements: The study team would like to thank all research participants for their time and interest in our research.

Funding, grant, and award info: This research was supported by the Canadian Institutes of Health Research (#31257), as well as Doctoral support to B Sharma from the Canadian Institutes of Health Research (CIHR-CGS-D, #157864). BW Timmons is the Canada Research Chair in Child Health & Exercise Medicine.

Ethical approval information: The study was approved by the Hamilton Integrated Research Ethics Board (www.hireb.ca) # 14-376.

Data sharing agreement: Data are available upon request.

Patient involvement: Given that this study is a secondary data analysis, patient and public involvement was not part of this analysis.

REFERENCES

1. McCrory P, Meeuwisse W, Dvorak J, et al. Consensus statement on concussion in sport—the 5th international conference on concussion in sport held in Berlin, October 2016. *British journal of sports medicine*. 2017;51(11):838-847.
2. Voormolen DC, Polinder S, Von Steinbuechel N, et al. The association between post-concussion symptoms and health-related quality of life in patients with mild traumatic brain injury. *Injury*. 2019;50(5):1068-1074.
3. Fridman L, Scolnik M, Macpherson A, et al. Annual trends in follow-up visits for pediatric concussion in emergency departments and physicians' offices. *The Journal of pediatrics*. 2018;192:184-188.
4. Meehan III WP, Mannix R. Pediatric concussions in United States emergency departments in the years 2002 to 2006. *The Journal of pediatrics*. 2010;157(6):889-893.
5. Veliz P, McCabe SE, Eckner JT, et al. Trends in the Prevalence of Concussion Reported by US Adolescents, 2016-2020. *JAMA*. 2021;325(17):1789-1791.
6. Barlow KM. Postconcussion syndrome: a review. *Journal of child neurology*. 2016;31(1):57-67.
7. Barlow KM, Crawford S, Stevenson A, et al. Epidemiology of postconcussion syndrome in pediatric mild traumatic brain injury. *Pediatrics*. 2010;126(2):e374-e381.
8. Novak Z, Aglipay M, Barrowman N, et al. Association of persistent postconcussion symptoms with pediatric quality of life. *JAMA pediatrics*. 2016;170(12):e162900-e162900.
9. Plourde V, Yeates KO, Brooks BL. Predictors of long-term psychosocial functioning and health-related quality of life in children and adolescents with prior concussions. *Journal of the International Neuropsychological Society*. 2018;24(6):540-548.
10. Fineblit S, Selci E, Loewen H, et al. Health-related quality of life after pediatric mild traumatic brain injury/concussion: a systematic review. *Journal of neurotrauma*. 2016;33(17):1561-1568.
11. Russell K, Selci E, Chu S, et al. Longitudinal assessment of health-related quality of life following adolescent sports-related concussion. *Journal of neurotrauma*. 2017;34(13):2147-2153.
12. Abbas K, Shenk TE, Poole VN, et al. Effects of repetitive sub-concussive brain injury on the functional connectivity of default mode network in high school football athletes. *Developmental neuropsychology*. 2015;40(1):51-56.
13. Borich M, Babul A-N, Yuan PH, et al. Alterations in resting-state brain networks in concussed adolescent athletes. *Journal of neurotrauma*. 2015;32(4):265-271.
14. Dona O, Noseworthy MD, DeMatteo C, et al. Fractal analysis of brain blood oxygenation level dependent (BOLD) signals from children with mild traumatic brain injury (mTBI). *PLoS One*. 2017;12(1):e0169647.
15. Iyer KK, Barlow KM, Brooks B, et al. Relating brain connectivity with persistent symptoms in pediatric concussion. *Annals of clinical and translational neurology*. 2019;6(5):954-961.

16. Iyer KK, Zalesky A, Barlow KM, et al. Default mode network anatomy and function is linked to pediatric concussion recovery. *Annals of clinical and translational neurology*. 2019;6(12):2544-2554.
17. Kaushal M, España LY, Nencka AS, et al. Resting-state functional connectivity after concussion is associated with clinical recovery. *Human brain mapping*. 2019;40(4):1211-1220.
18. Manning KY, Schranz A, Bartha R, et al. Multiparametric MRI changes persist beyond recovery in concussed adolescent hockey players. *Neurology*. 2017;89(21):2157-2166.
19. Meier TB, Giraldo-Chica M, Espana LY, et al. Resting-state fMRI metrics in acute sport-related concussion and their association with clinical recovery: a study from the NCAA-DOD CARE Consortium. *Journal of neurotrauma*. 2020;37(1):152-162.
20. Murdaugh DL, King TZ, Sun B, et al. Longitudinal changes in resting state connectivity and white matter integrity in adolescents with sports-related concussion. *Journal of the International Neuropsychological Society: JINS*. 2018;24(8):781-792.
21. Newsome MR, Li X, Lin X, et al. Functional connectivity is altered in concussed adolescent athletes despite medical clearance to return to play: a preliminary report. *Frontiers in neurology*. 2016;7:116.
22. Plourde V, Rohr CS, Virani S, et al. Default mode network functional connectivity after multiple concussions in children and adolescents. In: Oxford University Press; 2020.
23. Stephenson DD, Meier TB, Pabbathi Reddy S, et al. Resting-State Power and Regional Connectivity After Pediatric Mild Traumatic Brain Injury. *Journal of magnetic resonance imaging*. 2020;52(6):1701-1713.
24. Rausa VC, Shapiro J, Seal ML, et al. Neuroimaging in paediatric mild traumatic brain injury: a systematic review. *Neuroscience & Biobehavioral Reviews*. 2020.
25. Gornall A, Takagi M, Morawakage T, et al. Mental health after paediatric concussion: a systematic review and meta-analysis. *British journal of sports medicine*. 2021.
26. Carter KM, Pauhl AN, Christie AD. The Role of Active Rehabilitation in Concussion Management: A Systematic Review and Meta-analysis. *Medicine and science in sports and exercise*. 2021.
27. Langevin P, Frémont P, Fait P, et al. Aerobic Exercise for Sport-related Concussion: A Systematic Review and Meta-analysis. *Medicine and science in sports and exercise*. 2020.
28. Grool AM, Aglipay M, Momoli F, et al. Association between early participation in physical activity following acute concussion and persistent postconcussive symptoms in children and adolescents. *Jama*. 2016;316(23):2504-2514.
29. Thomas DG, Apps JN, Hoffmann RG, et al. Benefits of strict rest after acute concussion: a randomized controlled trial. *Pediatrics*. 2015;135(2):213-223.
30. Leddy JJ, Haider MN, Ellis M, et al. Exercise is medicine for concussion. *Current sports medicine reports*. 2018;17(8):262.
31. Lal A, Kolakowsky-Hayner SA, Ghajar J, et al. The Effect of Physical Exercise after a Concussion: A Systematic Review and Meta-Analysis. *The American journal of sports medicine*. 2017;0363546517706137.
32. Baque E, Barber L, Sakzewski L, et al. Reproducibility in measuring physical activity in children and adolescents with an acquired brain injury. *Brain Inj*. 2016;30(13-14):1692-1698.

33. Whitfield-Gabrieli S, Nieto-Castanon A. Conn: a functional connectivity toolbox for correlated and anticorrelated brain networks. *Brain connectivity*. 2012;2(3):125-141.
34. Davis AD, Noseworthy MD. Motion and distortion correction of skeletal muscle echo planar images. *Magnetic resonance imaging*. 2016;34(6):832-838.
35. Smith SM, Jenkinson M, Woolrich MW, et al. Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage*. 2004;23:S208-S219.
36. Jenkinson M, Beckmann CF, Behrens T, et al. Smith SM. *FSL neuroimage*. 2012;62:782-790.
37. Andersson JL, Hutton C, Ashburner J, et al. Modeling geometric deformations in EPI time series. *Neuroimage*. 2001;13(5):903-919.
38. Henson R, Buechel C, Josephs O, et al. The slice-timing problem in event-related fMRI. *NeuroImage*. 1999;9:125-.
39. Ashburner J, Barnes G, Chen C-C, et al. SPM12 manual. *Wellcome Trust Centre for Neuroimaging, London, UK*. 2014;2464.
40. Behzadi Y, Restom K, Liao J, et al. A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage*. 2007;37(1):90-101.
41. Friston KJ, Williams S, Howard R, et al. Movement-related effects in fMRI time-series. *Magnetic resonance in medicine*. 1996;35(3):346-355.
42. Power JD, Mitra A, Laumann TO, et al. Methods to detect, characterize, and remove motion artifact in resting state fMRI. *Neuroimage*. 2014;84:320-341.
43. Fornito A, Zalesky A, Bullmore E. *Fundamentals of brain network analysis*. Academic Press; 2016.
44. Power JD. Resting-State fMRI: Preclinical Foundations. In: *fMRI*. Springer; 2020:47-63.
45. Houck JM, Çetin MS, Mayer AR, et al. Magnetoencephalographic and functional MRI connectomics in schizophrenia via intra-and inter-network connectivity. *Neuroimage*. 2017;145:96-106.
46. Ke J, Zhang L, Qi R, et al. Typhoon-related post-traumatic stress disorder and trauma might lead to functional integration abnormalities in intra-and inter-resting state networks: a resting-state fMRI independent component analysis. *Cellular Physiology and Biochemistry*. 2018;48(1):99-110.
47. Lang S, Yoon EJ, Kibreab M, et al. Mild behavioral impairment in Parkinson's disease is associated with altered corticostriatal connectivity. *NeuroImage: Clinical*. 2020;26:102252.
48. Zhu H, Zhou P, Alcauter S, et al. Changes of intranetwork and internetwork functional connectivity in Alzheimer's disease and mild cognitive impairment. *Journal of neural engineering*. 2016;13(4):046008.
49. Baquet G, Stratton G, Van Praagh E, et al. Improving physical activity assessment in prepubertal children with high-frequency accelerometry monitoring: a methodological issue. *Preventive medicine*. 2007;44(2):143-147.
50. Evenson KR, Catellier DJ, Gill K, et al. Calibration of two objective measures of physical activity for children. *Journal of sports sciences*. 2008;26(14):1557-1565.
51. Rich C, Geraci M, Griffiths L, et al. Quality control methods in accelerometer data processing: defining minimum wear time. *PloS one*. 2013;8(6):e67206.

52. Mak LE, Minuzzi L, MacQueen G, et al. The default mode network in healthy individuals: a systematic review and meta-analysis. *Brain connectivity*. 2017;7(1):25-33.
53. Buckner RL, Andrews-Hanna JR, Schacter DL. The brain's default network: anatomy, function, and relevance to disease. 2008.
54. Konishi M, McLaren DG, Engen H, et al. Shaped by the past: the default mode network supports cognition that is independent of immediate perceptual input. *PloS one*. 2015;10(6):e0132209.
55. Smallwood J, Tipper C, Brown K, et al. Escaping the here and now: evidence for a role of the default mode network in perceptually decoupled thought. *Neuroimage*. 2013;69:120-125.
56. Andrews-Hanna JR. The brain's default network and its adaptive role in internal mentation. *The Neuroscientist*. 2012;18(3):251-270.
57. Stawarczyk D, D'Argembeau A. The dynamics of memory retrieval for internal mentation. *Scientific reports*. 2019;9(1):1-13.
58. Slobounov S, Gay M, Zhang K, et al. Alteration of brain functional network at rest and in response to YMCA physical stress test in concussed athletes: RsfMRI study. *Neuroimage*. 2011;55(4):1716-1727.
59. Zhang K, Johnson B, Gay M, et al. Default mode network in concussed individuals in response to the YMCA physical stress test. *Journal of neurotrauma*. 2012;29(5):756-765.
60. Bauer CC, Rozenkrantz L, Caballero C, et al. Mindfulness training preserves sustained attention and resting state anticorrelation between default-mode network and dorsolateral prefrontal cortex: A randomized controlled trial. *Human brain mapping*. 2020;41(18):5356-5369.
61. Berkovich-Ohana A, Glicksohn J, Goldstein A. Mindfulness-induced changes in gamma band activity—implications for the default mode network, self-reference and attention. *Clinical neurophysiology*. 2012;123(4):700-710.
62. Brewer JA, Worhunsky PD, Gray JR, et al. Meditation experience is associated with differences in default mode network activity and connectivity. *Proceedings of the National Academy of Sciences*. 2011;108(50):20254-20259.
63. Doll A, Hölzel BK, Boucard CC, et al. Mindfulness is associated with intrinsic functional connectivity between default mode and salience networks. *Frontiers in human neuroscience*. 2015;9:461.
64. Wang X, Xu M, Song Y, et al. The network property of the thalamus in the default mode network is correlated with trait mindfulness. *Neuroscience*. 2014;278:291-301.
65. Azulay J, Smart CM, Mott T, et al. A pilot study examining the effect of mindfulness-based stress reduction on symptoms of chronic mild traumatic brain injury/postconcussive syndrome. *The Journal of head trauma rehabilitation*. 2013;28(4):323-331.
66. Panicia M, Knafo R, Thomas S, et al. Mindfulness-based yoga for youth with persistent concussion: a pilot study. *American Journal of Occupational Therapy*. 2019;73(1):7301205040p7301205041-7301205040p7301205011.
67. Guthold R, Stevens GA, Riley LM, et al. Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1·9 million participants. *The lancet global health*. 2018;6(10):e1077-e1086.

68. Voss MW, Weng TB, Burzynska AZ, et al. Fitness, but not physical activity, is related to functional integrity of brain networks associated with aging. *Neuroimage*. 2016;131:113-125.
69. Jackson AS, Sui X, Hébert JR, et al. Role of lifestyle and aging on the longitudinal change in cardiorespiratory fitness. *Archives of internal medicine*. 2009;169(19):1781-1787.

TABLES

Table 1: Seed regions used to calculate the intra-network connectivity of the four resting state networks of interest.

Network	Seeds
	MNI co-ordinates (x, y, z)
DMN	Medial pre-frontal cortex (1, 55, -3)
	Lateral parietal, left (-39, -77, 33)
	Lateral parietal, right (47, -67, 29)
	Posterior cingulate cortex (1, -61, 38)
SMN	Lateral seed, left (-55, -12, 29)
	Lateral seed, right (56, -10, 29)
	Superior seed (0, -31, 67)
SN	Anterior cingulate cortex (0, 22, 35)
	Anterior insula, left (-44, 13, 1)
	Anterior insula, right (47, 14, 0)
	Rostral pre-frontal cortex, left (-32, 45, 27)
	Rostral pre-frontal cortex, right (32, 46, 27)
	Supramarginal gyrus, left (-60, -39, 31)
	Supramarginal gyrus, right (62, -35, 32)
FPN	Lateral pre-frontal cortex, left (-43, 33, 28)
	Lateral pre-frontal cortex, right

	(41, 38, 30)
	Posterior cingulate cortex, left
	(-46, -58, 49)
	Posterior cingulate cortex, right
	(52, -52, 45)

Table 2: Average daily (in minutes/day) monitoring time and activity by intensity in our cohort.
(LPA = light physical activity, MPA = moderate physical activity, VPA = vigorous physical activity).

	Mean	SD	Range
Monitoring time	798.3	68.0	686.8-914.5
Sedentary	631.0	75.9	504.2-733.2
LPA	116.1	33.9	68.9-172.2
MPA	28.6	11.3	13.4-48.6
VPA	22.7	16.9	3.3-57.8

Table 3: Average intra-network connectivity across the four networks of study (as Fisher Z-transformed correlation coefficients). (DMN = default mode network, SMN = sensorimotor network, SN = salience network, and FPN = fronto-parietal network).

	Mean	SD	Range
DMN	0.50	0.15	0.21-0.89
SMN	0.60	0.21	0.18-1.06
SA	0.48	0.11	0.23-0.70
FPC	0.58	0.16	0.34-0.81

Table 4: General linear model parameters for DMN-specific analyses. B=unstandardized beta coefficient; β =standardized beta coefficient. (LPA = light physical activity, MPA = moderate physical activity, VPA = vigorous physical activity).

Sedentary time					
Predictor	B	95% CI for B	β	t	p
DMN	-71.633	-311.907, 168.640	-0.146	-0.656	0.525
Age	20.998	5.601, 36.395	0.667	3.002	0.012
Model	$F_{(2, 11)} = 3.653$, $p = 0.061$, $R_a^2 = 0.458$				
LPA					
Predictor	B	95% CI for B	β	t	p
DMN	102.900	86.319, 289.261	0.469	2.346	0.039
Age	-8.486	-14.671, -2.301	-0.603	-3.020	0.012
Model	$F_{(2, 11)} = 7.053$, $p = 0.011$, $R_a^2 = 0.562$				
MPA					
Predictor	B	95% CI for B	β	t	p
DMN	52.908	19.627, 86.189	0.725	3.499	0.005
Age	-0.394	-2.527, 1.739	-0.084	-0.407	0.692
Model	$F_{(2, 11)} = 6.159$, $p = 0.016$, $R_a^2 = 0.528$				
VPA					
Predictor	B	95% CI for B	β	t	p
DMN	88.755	44.565, 128.945	0.792	4.526	0.001
Age	1.148	-1.555, 3.852	0.164	0.935	0.370
Model	$F_{(2, 11)} = 10.855$, $p = 0.002$, $R_a^2 = 0.664$				

FIGURE LEGENDS

Figure 1: Model-predicted activity levels (in minutes/day) plotted against intra-network connectivity (Fisher Z-transformed correlation coefficients) and a linear fitted line. Only the models highlighted with a red box were significant. (LPA = light physical activity, MPA = moderate physical activity, VPA = vigorous physical activity).