Genetic, structural and clinical analysis of spastic paraplegia 4

Parizad Varghaei^{1,2}, MD, Mehrdad A Estiar^{2,3}, MSc, Setareh Ashtiani⁴, MSc, Simon Veyron⁵, PhD, Kheireddin Mufti^{2,3}, MSc, Etienne Leveille⁶, MD, Eric Yu^{2,3}, BSc, Dan Spiegelman, MSc², Marie-France Rioux⁷, MD, Grace Yoon⁸, MD, Mark Tarnopolsky⁹, MD, PhD, Kym M. Boycott¹⁰, MD, Nicolas Dupre^{11,12}, MD, MSc, Oksana Suchowersky^{4,13}, MD, Jean-François Trempe⁵, PhD, Guy A. Rouleau^{2,3,14}, MD, PhD, Ziv Gan-Or^{2,3,14}, MD, PhD

1. Division of Experimental Medicine, Department of Medicine, McGill University, Montreal, Quebec, Canada
2. The Neuro (Montreal Neurological Institute-Hospital), McGill University, Montreal, Quebec, Canada
3. Department of Human Genetics, McGill University, Montréal, Québec, Canada
4. Alberta Children's Hospital, Medical Genetics, Calgary, Alberta, Canada
5. Department of Pharmacology & Therapeutics and Centre de Recherche en Biologie Structurale - FRQS, McGill University, Montréal, Canada
6. Faculty of Medicine, McGill University, Montreal, QC, Canada
7. Department of Neurology, Université de Sherbrooke, Sherbrooke, Québec, Canada.
8. Divisions of Neurology and Clinical and Metabolic Genetics, Department of Paediatrics, University of Toronto, The Hospital for Sick Children, Toronto, Ontario, Canada
9. Department of Pediatrics, McMaster University, Hamilton, Ontario, Canada
10. Children's Hospital of Eastern Ontario Research Institute, University of Ottawa, Ottawa, Ontario, Canada
11. Department of Medicine, Faculty of Medicine, Université Laval, Québec City, Quebec, Canada
12. Neuroscience Axis, CHU de Québec-Université Laval, Québec City, Québec, Canada
13. Departments of Medicine (Neurology) and Medical Genetics, University of Alberta, Edmonton, Alberta, Canada
14. Department of Neurology and Neurosurgery, McGill University, Montréal, Québec, Canada

Correspondence to:

Guy A. Rouleau,
 Director, Montréal Neurological Institute and hospital,
 Address: 3801, University Street, Office 636
 Montréal, Québec H3A 2B4
 Phone: +1-514-398-2690,
 Fax: +1-514 398-8248
 Email: guy.rouleau@mcgill.ca

Ziv Gan-Or
 Department of Neurology and Neurosurgery
 McGill University
 1033 Pine Avenue, West,
 Ludmer Pavilion, room 312
 Montreal, QC, H3A 1A1,

Phone: +1-514-398-5845
Fax. +1-514 398-8248
Email: ziv.gan-or@mcgill.ca

Word count: 3689

Running title: Genetic, structural and clinical aspects of SPG4

Keywords: Spastic paraplegia, HSP, SPG4, SPAST

Relevant conflicts of interest/financial disclosures: The authors report no competing interests.

Funding sources for the study: This study was funded by CIHR Emerging Team Grant, in collaboration with the Canadian Organization for Rare Disorders (CORD), grant number RN127580 – 260005, and by a CIHR Foundation grant granted to GAR. MAE is funded by the Fonds de Recherche du Québec–Santé (FRQS). GAR holds the Wilder Penfield Chair in Neurosciences. ZGO is supported by the Fonds de recherche du Québec–Santé Chercheur-Boursier award and is a William Dawson Scholar.

Abstract

Background: Spastic paraplegia type 4 (SPG4), resulting from heterozygous mutations in the *SPAST* gene, is the most common form among the heterogeneous group of hereditary spastic paraplegias (HSPs).

Objective: To study genetic and clinical characteristics of SPG4 across Canada.

Methods: The *SPAST* gene was analyzed in a total of 696 HSP patients from 431 families by either HSP-gene panel sequencing or whole exome sequencing (WES). We used Multiplex ligation-dependent probe amplification to analyze copy number variations (CNVs), and performed *in silico* structural analysis of selected mutations. Clinical characteristics of patients were assessed, and long-term follow-up was done to study genotype-phenotype correlations.

Results: We identified 157 SPG4 patients from 65 families who carried 41 different *SPAST* mutations, six of which are novel and six are CNVs. We report novel aspects of mutations occurring in Arg499, a case with homozygous mutation, a family with probable compound heterozygous mutations, three patients with *de novo* mutations, three cases with pathogenic synonymous mutation, co-occurrence of SPG4 and multiple sclerosis, and novel or rarely reported signs and symptoms seen in SPG4 patients.

Conclusion: Our study demonstrates that SPG4 is a heterogeneous type of HSP, with diverse genetic features and clinical manifestations. In rare cases, biallelic inheritance, *de novo* mutation, pathogenic synonymous mutations and CNVs should be considered.

Introduction

Spastic paraplegia type 4 (SPG4, OMIM #182601) is the most frequent form of either sporadic or familial hereditary spastic paraplegias (HSPs), caused by heterozygous mutations in the *SPAST* gene.¹⁻³ With more than 80 potentially causative loci or genes reported to date,⁴ HSPs are known to affect 1-10/100,000 of the population,⁵ and autosomal dominant (AD) HSPs comprise 43%-80% of them.⁵⁻¹¹ Among all AD-HSPs, 70-80% are categorized as “pure”¹² with a phenotype limited to pyramidal signs in the lower limbs, with or without deep sensory loss and sphincter disturbances.¹²⁻¹⁵ Of all pure AD-HSPs, about 40% are caused by *SPAST* mutations.¹²

SPAST encodes spastin, which is a protein from the AAA (ATPase associated with various cellular activities) family of ATPases.^{16, 17} Spastin controls different aspects of microtubule dynamics (e.g. microtubule number, motility, length, disassembly and remodeling), and hydrolyses ATP to cleave microtubules,^{18, 19} a necessary step in axonal transport.²⁰ Its mechanism involves binding to the C-terminus of tubulin and severing tubulin subunits from the microtubule in an ATP hydrolysis-dependent manner.¹⁹ The structure of human spastin residues 323-610 was solved by cryoelectron microscopy.²¹ The structure reveals an AAA+ ATPase homohexamer (Figure 1A), with ATPase active sites located at the interface between every adjacent subunit (Figure 1B).^{22, 23} A tubulin peptide runs through the channel, suggesting a hand-over-hand mechanism of substrate translocation, with five subunits interacting with the peptide forming a spiral staircase and one displaced from the peptide/substrate (Figure 1A).²¹

To date, more than 200 *SPAST* mutations have been found.^{3, 16, 24-26} Deletion/insertions, nonsense, and splice site mutations are distributed throughout the gene, while missense mutations are mostly clustered in the AAA domain.³ Exon deletions may account for 20% of cases in whom point mutations are not detected.^{27, 28}

The penetrance of SPG4 is up to 80-90% and is age-dependent,³ with age at onset (AAO) that may range from infancy to the eighth decade of life.^{1, 29} Most cases present as juvenile or adult-onset pure spastic paraplegia with urinary sphincter disturbances, pes cavus, and dysarthria. However, *SPAST* mutations are known to cause clinically heterogeneous manifestations, and a high variety of signs and symptoms is reported among carriers of different *SPAST* mutations. This clinical heterogeneity occurs even among patients harboring the same mutations,³⁰ suggesting that other factors affect the clinical presentation of SPG4.

Although previous studies have deciphered some aspects of this variability, the genotype-phenotype correlations and some rare features of SPG4 are not fully understood. In this study, we analyzed a large cohort of SPG4 patients from Canada to better clarify the genetic and clinical spectrum of the disease.

Methods

Population

A total of 696 HSP patients from 431 families were recruited in eight medical centers across Canada (Montreal, Quebec, Ottawa, Toronto, Hamilton, Calgary, Edmonton, and Vancouver) as part of CanHSP, a Canadian consortium for the study of HSP. Details about the diagnosis and recruitment process has been previously reported.³¹ Clinical assessments were done including family history, demographic data, developmental history, AAO, and HSP core symptoms (lower extremity weakness and spasticity, extensor plantar responses, hyperreflexia, and bladder dysfunction). Other neurological, as well as non-neurological clinical presentations of HSP, were also assessed. For a group of the patients, the Spastic Paraplegia Rating Scale (SPRS), which is

an indicator of severity in HSP, was measured.^{32, 33} For another subset of patients, brain and/or spine MRI were performed. The disease was deemed pure unless the patient had at least one sign not attributable to the lateral corticospinal tract or pyramidal tract, including ataxia, intellectual disability, cognitive decline, language development delay, extrapyramidal signs, visual dysfunction, epilepsy, deafness, dysarthria, optic atrophy, peripheral neuropathy, and dystonia; in which case the disease was classified as complex.

All the data is stored in a central database at McGill University. All patients signed informed consent forms and the institutional review board approved the study protocol.

Genetic and data analysis

DNA was extracted from peripheral blood using a standard procedure.³⁴ Initially, HSP-gene panel sequencing was performed on 379 patients. Then, 194 genetically undiagnosed patients, and additional 206 patients who were not analyzed with panel sequencing (400 patients in total from 291 families), went through whole exome sequencing (WES).

For WES, the Agilent SureSelect Human All Exon v4 kit for capture and targeted enrichment of the exome was used. To analyze the WES data, we used a list of 785 HSP-related genes or genes associated with similar neurological disorders which cause spasticity (Supplementary Table 1). Illumina HiSeq 2000/2500/4000 system was used for sequencing captured samples. Using Burrows-Wheeler Aligner (BWA), the sequence reads were then aligned against the human genome (GRCh37 assembly).³⁵ We used the Genome Analysis ToolKit (GATK)³⁶ and Annotate Variation (ANNOVAR)³⁷ for variant calling and annotation, respectively. We excluded variant calls with a genotype quality less than 97 and less than 30x depth of coverage. Integrated Genomics Viewer was used to visually inspect the detected

variants, and suspicious variants were validated by Sanger Sequencing. Sanger Sequencing was also used for assessing sporadic patients' parents, to determine if the proband had a *de novo* mutation.

SPAST variants (NM_014946) were initially selected based on identifying missense and LoF alleles, including frame-shift, splice-site, nonsense, and copy number variations (CNVs) with a minor allele frequency less than 0.01 in gnomAD.³⁸ The variants' pathogenicity has been determined using VarSome,³⁹ according to the American College of Medical Genetics and Genomics (ACMG) guideline. Variants classified as “Benign” and “Likely Benign”, as well as intronic splice site variants higher than ± 3 were excluded from the analysis. To detect CNVs, ExomeDepth⁴⁰ was used on WES data, followed by 48 selected samples that went through Multiplex ligation-dependent probe amplification (MLPA) testing (MRC Holland, Amsterdam, The Netherlands) to confirm or exclude suspected *SPAST* CNVs. InterPro⁴¹ was applied to identify domains and corresponding sites in the protein.

Statistical analysis

To determine the association between two categorical variables, one categorical variable with one continuous variable, and two continuous variables, Pearson chi-squared test, Mann-Whitney U test, and Spearman's rank correlation coefficient were used, respectively. *P*-value was set at <0.05 and Bonferroni correction for multiple comparisons was applied when necessary. SPSS was used to perform all statistical analyses.

***In silico* structural analysis**

The atomic coordinates of human spastin bound to a glutamate-rich peptide, ADP, BeF₃ and Mg²⁺ and *D. melanogaster* spastin bound to a glutamate-rich peptide, ADP, ATP and Mg²⁺ were

downloaded from the Protein Data Bank (ID 6PEN and 6P07). The effect induced by each mutation was evaluated using the “mutagenesis” toolbox in The PyMOL Molecular Graphics System, Version 2.4.0 Schrödinger, LLC. and the DynaMut server <http://biosig.unimelb.edu.au/dynamut/>.⁴²

Results

Cohort characteristics

We identified 65 families (15.1% of the families in HSP cohort), and a total of 157 patients (22.5% of HSP patients) with SPG4. Mean AAO was 22 years (0-67, SD: 19.89), and it followed a bimodal distribution; the first peak in the first 5 years of life, and the second peak from 35 to 44 years of age (Supplementary Figure 1). Mean age at examination was 43.4, and mean disease duration was 21 years. Patients with longer disease duration presented with higher SPRS scores (Spearman’s correlation coefficient; $p = 0.048$). In contrast, age at onset was not associated with the severity of the disease (Spearman’s correlation coefficient; $p = 0.934$). Complex HSP was seen in 24/65 (36.9%) of the probands, and 37/157 (23.6%) of all the patients. No significant differences were seen between male and female patients in AAO and other clinical manifestations. Table 1 details the clinical presentation of the patients.

We detected 41 different *SPAST* mutations in our cohort (Table 2). Most mutations (22/34, 64.7%) occurred between amino acids 374 and 567, in the AAA cassette. Among the missense mutations, 15/20 (75%) were clustered in AAA cassette, while LoF mutations were more evenly distributed across gene (Figure 2). Presentation of the disease did not differ significantly between the two types of mutation (Table 1).

Novel *SPAST* mutations

Of the 41 different *SPAST* mutations in our cohort, six were novel, including p.(Trp77Ter), p.(Glu366AspfsTer28), p.(Gly382_Pro383insArg), p.(Phe403Leu), p.(Arg498AlafsTer30), and p.(Ser597ThrfsTer3). We performed *in silico* analysis to investigate the impact of these mutations on the structure and activity of spastin, with exception of p.(Trp77Ter), as Trp77 is not visible in any structure of spastin and could not be investigated.

Glu366 is located in the loop₃₆₅₋₃₇₇ and the frameshift mutation p.(Glu366AspfsTer28) changes all the residue subsequent to residue 365 and inserts a stop codon at position 394. This mutation truncates residues 394 to 616 which includes residues involved in oligomerization and ATP-binding, which will abrogate its enzymatic activity. Gly382 and Pro383 are located in the nucleotide-binding loop₃₈₂₋₃₈₉.⁴³ Residues Gly385 and Asn386 interact directly with the ADP, in a conformation that is shaped by the Pro383 and Pro384 (Figure 1C). The mutation p.(Gly382_Pro383insArg) lengthens this loop and likely destabilizes the interaction between the protein and the nucleotide. Phe403 forms a pi-stacking interaction with Phe439 and hydrophobic interactions with neighbouring residues (Figure 1D). Thus, mutation p.(Phe403Leu) would destabilize the domain and impair the ATPase activity.

In this structure, Arg498 and Arg499 are both involved in coordinating ADP. Arg498 is making direct interaction with the β phosphate of the nucleotide located in the active site, while Arg499 stabilizes helix₄₅₅₋₄₇₀ and coordinates BeF₃, a phosphate structural analog (Figure 1E). By homology with *D. melanogaster* spastin structure (PDB ID: 6P07), those residues seem to be also involved in ATP β and γ phosphate coordination (Figure 1F). As a result, we can

hypothesize that the BeF₃ in the active site is mimicking the γ phosphate on an ATP. Mutation p.(Arg498AlafsTer30) causes residues 498 and 499 to be serine and valine residues, which would prevent ATP stabilization. In addition, this frame shift terminates the protein after residue 528, further disrupting the ATP-binding domain. This mutation will therefore abolish the ATPase activity. Finally, mutation p.(Ser597ThrfsTer3) will induce a frame shift at residue 597 and truncate the protein at residue 600. The missing C-terminal residues form a helix involved in intersubunit interactions. This mutation would therefore destabilize the hexameric assembly, which would disrupt the ATPase activity.

Founder French-Canadian mutations and known CNVs

The most frequent *SPAST* mutation in our cohort, p.(Gly559Asp), which has previously been suggested to be a founder mutation in French-Canadian population⁴⁴, was carried by 8 families (12.3%) and 15 patients in total (9.5%). Seven out of 24 French-Canadian probands (29%) harbored this mutation, while among all the patients with other or unknown ancestral backgrounds, this mutation was present in only one out of 29 probands (3.4%, $p=0.011$). Moreover, the mutation p.(Phe403Leu) was only detected in patients with French-Canadian ancestral background, in 5 patients from 2 families, suggesting it may be a founder mutation. Seven patients (4.5%) from six families (9.2%) carried a CNV (Supplementary Table 2), all of which have been previously reported.²⁷

Earlier age at onset and specific clinical features in patients with mutations in *SPAST* p.Arg499

The range of AAO in all 4 patients from 4 different families, two of which have previously been reported⁴⁵ with the p.(Arg499His) mutation was 1-3 years (Mann-Whitney U test; $p = 0.003$). The age range at which these four patients started to show symptoms was 1-5. Patients with this mutation in our cohort were more likely to present with motor delay, speech delay, dysarthria, learning disability, progressive cognitive deficits, and upper extremity weakness (Supplementary Table3).

Although statistically insignificant, cases who carried another mutation affecting the same amino acid residue, p.(Arg499Cys), also showed symptoms at a younger age (the range of AAO was 1-5 in three patients and 11-15 in one patient, $p = 0.111$). When combined together, the two mutations occurred in Arg499 locus were associated with a younger AAO (Mann-Whitney U test; $p = 0.004$). *In silico* analysis of Arg499 is discussed above, along with Arg498. Furthermore, as Arg499 stabilizes the γ phosphate, mutations p.(Arg499Cys) or p.(Arg499His) would also impede ATP binding and catalysis.

Possible biallelic inheritance in SPG4

The *SPAST* p.(Ser44Leu) variant has been previously suggested to play modifier role in SPG4.^{1, 46, 47} This variant was detected in one out of the 36 families that had undergone WES. From the three affected siblings, two had undergone WES, and both were heterozygous for the novel pathogenic mutation p.(Ser597ThrfsTer3), as well as for the p.(Ser44Leu) variant. The three siblings had complex HSP with disease onset in early childhood (clinical presentation of these patients is detailed in Supplementary Table 4). One unaffected parent only carried the p.(Ser44Leu) polymorphism, and the other unaffected parent was heterozygous for the

pathogenic p.(Ser597ThrfsTer3) mutation and wild-type for p.(Ser44Leu) (Supplementary Figure 2). The last physical examination of the parent with the pathogenic mutation at the age between 56 to 60 revealed completely normal findings, including normal reflexes, gait, and motor exam, suggesting that the p.(Ser44Leu) variant may be a modifier of the disease, or that extreme anticipation with undetected clinical effects in the mother exist in this family.

Another patient with consanguineous parents, carried a homozygous mutation, *SPAST* p.(Tyr51Ter). The patient had started to show symptoms at an age range of 1-5 years and suffered from core HSP symptoms, motor and speech delay, swallowing difficulty, dysarthria, upper extremity weakness, pes cavus, and skeletal abnormalities. Her brain and total spine MRI were normal and her SPRS score was 40 when examined at an age between 16 and 20 years. The patient's parents were both asymptomatic heterozygous carriers.

Patients with novel or rarely reported clinical manifestations

In our cohort, four patients (Supplementary Table 5) presented with deafness, one of whom had a *de novo SPAST* p.(Arg499His) mutation, and the remaining three had familial SPG4. We also report a patient with ocular movement abnormality, in whom extraocular movements showed slightly hypometric saccades but pursuit was smooth, and a mild horizontal gaze evoked rotatory nystagmus. The disease-causing mutation in this patient was p.(Gln434Ter).

One of the signs less reported to date in SPG4 is upper extremity intention tremor, seen in two of the patients in the current study. The causative mutations in these two patients were p.(Arg503Trp) and p.(Leu371Pro). Their respective range of AAO was 46-50 and 26-30, and both presented with a severe form of the disease (respective SPRS scores: 34 and 39). The first

patient also had swallowing difficulty and peripheral neuropathy, and both had upper extremity ataxia.

Another clinical feature seen in two of the patients in our cohort was seizures, which is quite rare in SPG4. One of the patients who carried the mutation p.(Pro489Leu), had a range of AAO of 21-25 years, and suffered from generalized seizures starting at 41-45 years of age. The second patient harbored the splice-site mutation c.1321+1G>A, with AAO of between 1 and 5 years, had episodes of atypical seizures at 56-60.

De novo cases

The parents of the three sporadic cases in our cohort did not carry the causative *SPAST* mutations. The first patient, with *SPAST* p.(Ser407Asn), had an AAO of 1-5 years. This case presented with lower extremity weakness and spasticity, upper and lower limb hyperreflexia, extensor plantar response, ankle clonus, and sensory abnormalities. She also showed signs of upper extremity weakness, motor and speech delay, dysarthria, learning disability, and swallowing difficulty. Their brain MRI was normal. The second patient had *de novo SPAST* p.(Arg499His). This patient started to show symptoms at age range of 1-5, and her symptoms included lower extremity weakness, spasticity and hyperreflexia, extensor plantar responses, and ankle clonus. Furthermore, they had motor and speech delay, dysarthria, and learning disability. They had a normal brain MRI, and their SPRS score was 37. The last *de novo* patient, also with a p.(Arg499His) mutation, also with an AAO range of 1-5 years, had an SPRS score of 28, and apart from core symptoms of HSP, had motor and speech delay, deafness, and dysarthria. The two latter patients have been previously reported.⁴⁵

Cases with synonymous mutation and co-occurrence of multiple sclerosis and SPG4

We report three patients from two unrelated families who carried the synonymous mutation *SPAST* p.(Lys414Lys), previously reported to be pathogenic.⁴⁸ The first patient had an AAO of between 6-10 years, and presented with lower extremity spasticity, hyperreflexia, and extensor plantar responses. The second family had two affected individuals, parent and offspring. The parent started to have difficulty walking at the age of 61-65 which led to using a cane, had marked spasticity and brisk reflexes in the lower limbs along with ankle clonus, upgoing plantar responses, urge incontinency, and mild decrease of vibration sensation in the ankles and toes. The offspring noticed difficulty walking at 21-25 years of age, which progressed slowly. This case had upper and lower limb weakness, bilateral Babinski sign and ankle clonus, and positive Hoffman sign. At an age between 31 and 35, lumbar puncture carried out due to diplopia was positive for oligoclonal bands and suggested a diagnosis of clinically isolated syndrome (CIS), which is considered as the first clinical episode in multiple sclerosis (MS).⁴⁹ They were treated with methylprednisone, and after one month, the diplopia resolved.

Discussion

In this large-scale analysis of SPG4 from CanHSP, we report novel *SPAST* mutations, the possibility of a founder mutations in the French-Canadian population, novel characteristics of mutations occurring in Arg499, and potential biallelic inheritance. We also report patients with rare or novel clinical manifestations, and co-occurrence of SPG4 and MS.

Using *in silico* analysis, we predict that the novel mutations reported in this study cause a loss-of-function of spastin, either by substantially shortening the protein, affecting its ATPase activity, or disturbing the formation of a ternary and quaternary structure necessary for catalysis. Spastin, a microtubule-cleaving enzyme involved in the cytoskeletal rearrangement,⁵⁰⁻⁵² also has a role in intracellular trafficking, cytokinesis regulation and resealing of nuclear membrane.⁵³⁻⁵⁵ In neurons, it is involved in the regeneration of axons and axonal transport.^{51, 56-58} Mutations in *SPAST* could result in a disruption of normal organelle trafficking and distribution, and thus micro-organelle accumulation in axons and swelling of axons⁵⁹⁻⁶³ which could result in HSP phenotype. We looked further into the three patients identified as sporadic and found out that all carried *de novo* mutations. Consistent with previous reports of more severe manifestations in patients with *de novo* mutations,⁶⁴ all patients showed a severe form of the disease. However, it is important to note that the mutation in two of the *de novo* patients (Arg499His), which is also carried by 33% of *de novo* patients,⁶⁴ is associated with a younger AAO, and infantile-onset ascending spastic paraplegia (IAHSP); regardless of the mode of inheritance.^{65, 66} Patients harboring this mutation in our cohort did not show all criteria to be diagnosed as IAHSP (spastic paraplegia progressing to tetraplegia, ocular gaze paralysis, pseudobulbar palsy⁶⁷), however, they showed a higher probability of suffering from upper extremity weakness, dysarthria, speech and motor delay, learning disabilities and cognitive delay. It has been suggested that amino acid 499 is located in one of the highly conserved regions of the *SPAST* gene,¹⁸ and mutations that occur in this region could have a significant effect on the hydrophobicity of the protein and its ability to sever microtubules.⁴⁵

The mutation in our homozygous patient has been reported previously in a heterozygous patient;⁶⁸ however, the presence of an additional mutation or CNVs could not have been

determined in that study, as only *SPAST* had been sequenced. Our patient with the homozygous mutation *SPAST* p.(Tyr51Ter) showed a severe form of the disease. Previously, there have been two reports of homozygous variants in *SPAST*. One, with the nonsense mutation p.(Ser545Ter), which also reports a complex and severe form of the disease,⁶⁹ and another, with the mutation p.(Leu534Pro), who showed a pure spastic paraparesis at the age of 39.⁷⁰ The patient from our cohort and the previously reported patient with p.(Ser545Ter) presented with a more severe form of the disease, probably because they carried nonsense mutations, while the latter patient had a missense mutation. Furthermore, in the latter study, the parents of the proband were asymptomatic carriers of the mutation.

The mode of inheritance observed in the patient with homozygous mutation and the family carrying the potential modifier mutation *SPAST* p.(Ser44Leu), may be explained by incomplete penetrance and/or anticipation⁷¹⁻⁷³ rather than biallelic inheritance, which could be considered in prenatal diagnosis and genetic counselling. The synonymous mutation *SPAST* p.(Lys414Lys) we report in three patients has been previously reported to underlie the disease, and minigene assay suggested that this variant leads to an aberrant splicing effect.⁴⁸ Although we did not find other synonymous *SPAST* mutations in additional HSP patients, it is important to not exclude synonymous variants when analyzing HSP specifically and other diseases in general.

The AAO distribution in our cohort followed the same trend as previously reported,^{1, 68} however, our results did not replicate previous findings which suggested that the underlying mutation and the patient sex can modify the disease, and that patients with later onset of the disease can have faster progression.^{1, 24} In our data, we did not find an association between mutation type, sex, or late onset, with the disease course. Table 3 compares the frequency of symptoms not attributable to pure form of the disease in our cohort with previous studies. As

shown, we report deafness in SPG4 for the first time. The co-occurrence of SPG4 and MS has been reported before;^{74, 75} however, the results of studies assessing an association between HSP-related mutations and MS are controversial^{76, 77} and further studies are required to verify such a relation.

Our study has several limitations. For instance, in spite of being one of the largest HSP cohorts, the number of SPG4 patients was limited, especially for a genotype-phenotype correlation study. Furthermore, the families interested in participating in this study were probably those who had remained undiagnosed after the primary *SPAST* testing; therefore, the proportion of SPG4 patients in this study may be underestimated. In addition, clinical signs and symptoms for some of the patients were missing.

In conclusion, we report one of the largest SPG4 cohorts, with 41 different mutations including 6 novel mutations. We suggest that modes of inheritance other than autosomal dominant could be involved in SPG4. Our study sheds light on some rarely reported or unreported aspects of the disease and helps improve genetic counseling and clinical trials conducted on SPG4. Towards this goal, larger efforts and international collaborations are required.

Acknowledgments

We thank the patients and their families for participating in this study. This study was funded by CIHR Emerging Team Grant, in collaboration with the Canadian Organization for Rare Disorders (CORD), grant number RN127580 – 260005, and by a CIHR Foundation grant granted to GAR. MAE is funded by the Fonds de Recherche du Québec–Santé (FRQS). GAR holds the Wilder Penfield Chair in Neurosciences. ZGO is supported by the Fonds de recherche du Québec–Santé Chercheur-Boursier award and is a William Dawson Scholar. We thank Patrick Dion, Daniel Rochefort, Helene Catoire, and Vessela Zaharieva for their assistance.

Author's Roles

P.V.: Research project conception, organization, and execution; statistical analysis design and execution, writing of first draft.

M.A.E.: Research project conception, organization and execution; manuscript review and critique.

S.A., M.F.R., G.Y., M.T., K.M.B., N.D., O.S., and E.L.: Patient recruitment and data collection.

E.L., E.Y., DS, S.V., and J.F.T.: Data analysis.

K.F.: Manuscript preparation.

G.A.R.: Research project conception and organization; manuscript review and critique.

Z.G.O.: Research project conception and organization; statistical analysis design, review and critique; manuscript review and critique.

Financial Disclosures of All Authors (for the Preceding 12 Months)

M.A.E. received a doctoral award from Fonds de la recherche en santé du Quebec (FRQS). O.S. has received non-related grants – Abbvie, Roche, WaveLifeSciences, and Vaccinex and honoraria – UpToDate and Springer. J.F.T. is a paid consultant and member of the scientific advisory board for Mitokinin Inc. He also received a grant from the Agora Open Science Trust (Grant Number: AOST-M4ND-001). He contributed to three inventions; Report of invention, McGill University Online Disclosure, Track code D2019-0117, Title: The small molecule PRT062607 targets PINK1; Report of invention, McGill University Online Disclosure, Track code D2020-0190, Title: Indole-2-carboxylic acid derivatives bind to Parkin; Report of invention, McGill University Online Disclosure, Track code D2021-0004, Title: Method for the expression and purification of Taq DNA polymerase. G.A.R. received grants from the following agencies and organizations: CIHR Foundation Scheme, Genome Quebec, ERANET PerMed/FRSQ, ALS Canada-Brain Canada, ALS Canada, Radala Foundation, CQDM, ALS Canada-Brain Canada, Brain Canada. Z.G.O. is supported by the Fonds de recherche du Quebec-Sante (FRQS) Chercheurs-boursiers award, and is a Parkinson's Disease Canada New Investigator awardee. He received consultancy fees from Ono Therapeutics, Handl Therapeutics, Neuron23, Lysosomal Therapeutics Inc., Bial Biotech Inc., Deerfield, Lighthouse, and Idorsia, all unrelated to the current study. All other authors report nothing to disclose.

References

1. Parodi L, Fenu S, Barbier M, et al. Spastic paraplegia due to SPAST mutations is modified by the underlying mutation and sex. *Brain* 2018;141(12):3331-3342.
2. Giudice TL, Lombardi F, Santorelli FM, Kawarai T, Orlacchio A. Hereditary spastic paraplegia: clinical-genetic characteristics and evolving molecular mechanisms. *Experimental neurology* 2014;261:518-539.
3. Solowska JM, Baas PW. Hereditary spastic paraplegia SPG4: what is known and not known about the disease. *Brain* 2015;138(9):2471-2484.
4. Estiar MA, Yu E, Salem IH, et al. Evidence for non-Mendelian inheritance in spastic paraplegia 7. *medRxiv* 2020.
5. Ruano L, Melo C, Silva MC, Coutinho P. The global epidemiology of hereditary ataxia and spastic paraplegia: a systematic review of prevalence studies. *Neuroepidemiology* 2014;42(3):174-183.
6. Skre H. Hereditary spastic paraplegia in Western Norway. *Clinical genetics* 1974;6(3):165-183.
7. McMonagle P, Webb S, Hutchinson M. The prevalence of "pure" autosomal dominant hereditary spastic paraparesis in the island of Ireland. *Journal of Neurology, Neurosurgery & Psychiatry* 2002;72(1):43-46.
8. Polo J, Calleja J, Combarros O, Berciano J. Hereditary ataxias and paraplegias in Cantabria, Spain: an epidemiological and clinical study. *Brain* 1991;114(2):855-866.
9. Harding A. Hereditary "pure" spastic paraplegia: a clinical and genetic study of 22 families. *Journal of Neurology, Neurosurgery & Psychiatry* 1981;44(10):871-883.
10. Schüle R, Wiethoff S, Martus P, et al. Hereditary spastic paraplegia: clinicogenetic lessons from 608 patients. *Annals of neurology* 2016;79(4):646-658.
11. Koh K, Ishiura H, Tsuji S, Takiyama Y. JASPAC: Japan spastic paraplegia research consortium. *Brain sciences* 2018;8(8):153.
12. Blackstone C. Hereditary spastic paraplegia. *Handbook of clinical neurology* 2018;148:633-652.
13. Parodi L, Fenu S, Stevanin G, Durr A. Hereditary spastic paraplegia: more than an upper motor neuron disease. *Revue neurologique* 2017;173(5):352-360.
14. Martinuzzi A, Montanaro D, Vavla M, et al. Clinical and paraclinical indicators of motor system impairment in hereditary spastic paraplegia: a pilot study. *PLoS One* 2016;11(4):e0153283.
15. Klebe S, Stevanin G, Depienne C. Clinical and genetic heterogeneity in hereditary spastic paraplegias: from SPG1 to SPG72 and still counting. *Revue neurologique* 2015;171(6-7):505-530.
16. Hazan J, Fonknechten N, Mavel D, et al. Spastin, a new AAA protein, is altered in the most frequent form of autosomal dominant spastic paraplegia. *Nature genetics* 1999;23(3):296-303.
17. Depienne C, Stevanin G, Brice A, Durr A. Hereditary spastic paraplegias: an update. *Current opinion in neurology* 2007;20(6):674-680.
18. Errico A, Ballabio A, Rugarli EI. Spastin, the protein mutated in autosomal dominant hereditary spastic paraplegia, is involved in microtubule dynamics. *Human molecular genetics* 2002;11(2):153-163.

19. Vemu A, Szczesna E, Zehr EA, et al. Severing enzymes amplify microtubule arrays through lattice GTP-tubulin incorporation. *Science* 2018;361(6404).
20. Salinas S, Carazo-Salas RE, Proukakis C, Schiavo G, Warner TT. Spastin and microtubules: Functions in health and disease. *Journal of neuroscience research* 2007;85(12):2778-2782.
21. Han H, Schubert HL, McCullough J, et al. Structure of spastin bound to a glutamate-rich peptide implies a hand-over-hand mechanism of substrate translocation. *Journal of Biological Chemistry* 2020;295(2):435-443.
22. Zehr E, Szyk A, Piszczek G, Szczesna E, Zuo X, Roll-Mecak A. Katanin spiral and ring structures shed light on power stroke for microtubule severing. *Nature structural & molecular biology* 2017;24(9):717.
23. Sandate CR, Szyk A, Zehr EA, Lander GC, Roll-Mecak A. An allosteric network in spastin couples multiple activities required for microtubule severing. *Nature structural & molecular biology* 2019;26(8):671-678.
24. Fonknechten N, Mavel D, Byrne P, et al. Spectrum of SPG4 mutations in autosomal dominant spastic paraplegia. *Human molecular genetics* 2000;9(4):637-644.
25. Lindsey J, Lusher M, McDermott C, et al. Mutation analysis of the spastin gene (SPG4) in patients with hereditary spastic paraparesis. *Journal of medical genetics* 2000;37(10):759-765.
26. Shoukier M, Neesen J, Sauter SM, et al. Expansion of mutation spectrum, determination of mutation cluster regions and predictive structural classification of SPAST mutations in hereditary spastic paraplegia. *European journal of human genetics* 2009;17(2):187-194.
27. Depienne C, Fedirko E, Forlani S, et al. Exon deletions of SPG4 are a frequent cause of hereditary spastic paraplegia. *Journal of medical genetics* 2007;44(4):281-284.
28. Beetz C, Nygren A, Schickel J, et al. High frequency of partial SPAST deletions in autosomal dominant hereditary spastic paraplegia. *Neurology* 2006;67(11):1926-1930.
29. Fink JK. Hereditary spastic paraplegia: clinico-pathologic features and emerging molecular mechanisms. *Acta neuropathologica* 2013;126(3):307-328.
30. de Souza PVS, de Rezende Pinto WBV, de Rezende Batistella GN, Bortholin T, Oliveira ASB. Hereditary spastic paraplegia: clinical and genetic hallmarks. *The Cerebellum* 2017;16(2):525-551.
31. Chrestian N, Dupré N, Gan-Or Z, et al. Clinical and genetic study of hereditary spastic paraplegia in Canada. *Neurology Genetics* 2017;3(1).
32. Schüle R, Holland-Letz T, Klimpe S, et al. The Spastic Paraplegia Rating Scale (SPRS): a reliable and valid measure of disease severity. *Neurology* 2006;67(3):430-434.
33. Adry RARdC, Lins CC, Kruschewsky RdA, Castro Filho BG. Comparison between the spastic paraplegia rating scale, Kurtzke scale, and Osame scale in the tropical spastic paraparesis/myelopathy associated with HTLV. *Revista da Sociedade Brasileira de Medicina Tropical* 2012;45(3):309-312.
34. Miller S, Dykes D, Polesky H. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic acids research* 1988;16(3):1215.
35. Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler transform. *bioinformatics* 2009;25(14):1754-1760.
36. McKenna A, Hanna M, Banks E, et al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome research* 2010;20(9):1297-1303.

37. Wang K, Li M, Hakonarson H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic acids research* 2010;38(16):e164-e164.
38. Karczewski KJ, Francioli LC, Tiao G, et al. The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature* 2020;581(7809):434-443.
39. Kopanos C, Tsiolkas V, Kouris A, et al. VarSome: the human genomic variant search engine. *Bioinformatics* 2019;35(11):1978.
40. Plagnol V, Curtis J, Epstein M, et al. A robust model for read count data in exome sequencing experiments and implications for copy number variant calling. *Bioinformatics* 2012;28(21):2747-2754.
41. Mitchell AL, Attwood TK, Babbitt PC, et al. InterPro in 2019: improving coverage, classification and access to protein sequence annotations. *Nucleic acids research* 2019;47(D1):D351-D360.
42. Rodrigues CH, Pires DE, Ascher DB. DynaMut: predicting the impact of mutations on protein conformation, flexibility and stability. *Nucleic acids research* 2018;46(W1):W350-W355.
43. Evans KJ, Gomes ER, Reisenweber SM, Gundersen GG, Luring BP. Linking axonal degeneration to microtubule remodeling by Spastin-mediated microtubule severing. *The Journal of cell biology* 2005;168(4):599-606.
44. Meijer IA, Dupré N, Brais B, et al. SPG4 founder effect in French Canadians with hereditary spastic paraplegia. *Canadian journal of neurological sciences* 2007;34(2):211-214.
45. Gillespie MK, Humphreys P, McMillan HJ, Boycott KM. Association of early-onset spasticity and risk for cognitive impairment with mutations at amino acid 499 in SPAST. *Journal of child neurology* 2018;33(5):329-332.
46. McDermott C, Burness C, Kirby J, et al. Clinical features of hereditary spastic paraplegia due to spastin mutation. *Neurology* 2006;67(1):45-51.
47. Svenson IK, Kloos MT, Gaskell PC, et al. Intragenic modifiers of hereditary spastic paraplegia due to spastin gene mutations. *Neurogenetics* 2004;5(3):157-164.
48. Svenson IK, Ashley-Koch AE, Gaskell PC, et al. Identification and expression analysis of spastin gene mutations in hereditary spastic paraplegia. *The American Journal of Human Genetics* 2001;68(5):1077-1085.
49. Sorensen PS, Sellebjerg F, Hartung H-P, Montalban X, Comi G, Tintoré M. The apparently milder course of multiple sclerosis: changes in the diagnostic criteria, therapy and natural history. *Brain* 2020;143(9):2637-2652.
50. Lumb JH, Connell JW, Allison R, Reid E. The AAA ATPase spastin links microtubule severing to membrane modelling. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research* 2012;1823(1):192-197.
51. Sardina F, Pisciotanni A, Ferrara M, et al. Spastin recovery in hereditary spastic paraplegia by preventing neddylation-dependent degradation. *Life science alliance* 2020;3(12).
52. Connell JW, Allison RJ, Rodger CE, Pearson G, Zlamalova E, Reid E. ESCRT-III-associated proteins and spastin inhibit protrudin-dependent polarised membrane traffic. *Cellular and Molecular Life Sciences* 2020;77(13):2641-2658.
53. Connell JW, Lindon C, Luzio JP, Reid E. Spastin couples microtubule severing to membrane traffic in completion of cytokinesis and secretion. *Traffic* 2009;10(1):42-56.
54. Allison R, Lumb JH, Fassier C, et al. An ESCRT–spastin interaction promotes fission of recycling tubules from the endosome. *Journal of Cell Biology* 2013;202(3):527-543.
55. Vietri M, Schink KO, Campsteijn C, et al. Spastin and ESCRT-III coordinate mitotic spindle disassembly and nuclear envelope sealing. *Nature* 2015;522(7555):231-235.

56. Trotta N, Orso G, Rossetto MG, Daga A, Broadie K. The hereditary spastic paraplegia gene, spastin, regulates microtubule stability to modulate synaptic structure and function. *Current biology* 2004;14(13):1135-1147.
57. Riano E, Martignoni M, Mancuso G, et al. Pleiotropic effects of spastin on neurite growth depending on expression levels. *Journal of neurochemistry* 2009;108(5):1277-1288.
58. Stone MC, Rao K, Gheres KW, et al. Normal spastin gene dosage is specifically required for axon regeneration. *Cell reports* 2012;2(5):1340-1350.
59. Tarrade A, Fassier C, Courageot S, et al. A mutation of spastin is responsible for swellings and impairment of transport in a region of axon characterized by changes in microtubule composition. *Human molecular genetics* 2006;15(24):3544-3558.
60. Kasher PR, De Vos KJ, Wharton SB, et al. Direct evidence for axonal transport defects in a novel mouse model of mutant spastin-induced hereditary spastic paraplegia (HSP) and human HSP patients. *Journal of neurochemistry* 2009;110(1):34-44.
61. Denton KR, Lei L, Grenier J, Rodionov V, Blackstone C, Li XJ. Loss of spastin function results in disease-specific axonal defects in human pluripotent stem cell-based models of hereditary spastic paraplegia. *Stem cells* 2014;32(2):414-423.
62. Havlicek S, Kohl Z, Mishra HK, et al. Gene dosage-dependent rescue of HSP neurite defects in SPG4 patients' neurons. *Human molecular genetics* 2014;23(10):2527-2541.
63. TARRADE A, Peris L, Courageot S, et al. Microtubule-targeting drugs rescue axonal swellings in cortical neurons from spastin knockout mice. *Disease Models & Mechanisms* 1 (6), 72-83(2013) 2013.
64. Schieving JH, de Bot ST, van de Pol LA, et al. De novo SPAST mutations may cause a complex SPG4 phenotype. *Brain* 2019;142(7):e31-e31.
65. Souza PVSd, Bortholin T, Naylor FGM, Pinto WBVdR, Oliveira ASB. Infantile-onset ascending spastic paraplegia phenotype associated with SPAST mutation. *Journal of the neurological sciences* 2016.
66. Ogasawara M, Saito T, Koshimizu E, Akasaka N, Sasaki M. A p. Arg499His Mutation in SPAST Is Associated with Infantile Onset Ascending Spastic Paralysis Complicated with Dysarthria and Anarthria. *Neuropediatrics* 2019;50(06):391-394.
67. Lesca G, Eymard-Pierre E, Santorelli F, et al. Infantile ascending hereditary spastic paralysis (IAHSP): clinical features in 11 families. *Neurology* 2003;60(4):674-682.
68. de Bot ST, van den Elzen RT, Mensenkamp A, et al. Hereditary spastic paraplegia due to SPAST mutations in 151 Dutch patients: new clinical aspects and 27 novel mutations. *Journal of Neurology, Neurosurgery & Psychiatry* 2010;81(10):1073-1078.
69. Cruz-Camino H, Vázquez-Cantú M, Vázquez-Cantú DL, et al. Clinical Characterization of 2 Siblings with a Homozygous SPAST Variant. *The American Journal of Case Reports* 2020;21:e919463-919461.
70. de Bot ST, Van Den Elzen R, Mensenkamp A, et al. Hereditary spastic paraplegia due to SPAST mutations in 151 Dutch patients: new clinical aspects and 27 novel mutations. *Journal of Neurology, Neurosurgery & Psychiatry* 2010;81(10):1073-1078.
71. Bürger J, Metzke H, Paternotte C, Schilling F, Hazan J, Reis A. Autosomal dominant spastic paraplegia with anticipation maps to a 4-cM interval on chromosome 2p21-p24 in a large German family. *Human genetics* 1996;98(3):371-375.
72. Reddy PL, Seltzer WK, Grewal RP. Possible anticipation in hereditary spastic paraplegia type 4 (SPG4). *Canadian Journal of Neurological Sciences* 2007;34(2):208-210.

73. Kawarai T, Montecchiani C, Miyamoto R, et al. Spastic paraplegia type 4: A novel SPAST splice site donor mutation and expansion of the phenotype variability. *Journal of the neurological sciences* 2017;380:92-97.
74. Boucher JJ, Counihan TJ. Co-incident primary progressive multiple sclerosis and hereditary spastic paraplegia (SPG4)—a case report. *Multiple Sclerosis and Related Disorders* 2020;44:102375.
75. Mead S, Proukakis C, Wood N, Crosby A, Plant G, Warner T. A large family with hereditary spastic paraparesis due to a frame shift mutation of the spastin (SPG4) gene: association with multiple sclerosis in two affected siblings and epilepsy in other affected family members. *Journal of Neurology, Neurosurgery & Psychiatry* 2001;71(6):788-791.
76. Jia X, Madireddy L, Caillier S, et al. Genome sequencing uncovers phenocopies in primary progressive multiple sclerosis. *Annals of neurology* 2018;84(1):51-63.
77. DeLuca G, Ramagopalan SV, Cader M, et al. The role of hereditary spastic paraplegia related genes in multiple sclerosis. *Journal of neurology* 2007;254(9):1221-1226.
78. Murphy S, Gorman G, Beetz C, et al. Dementia in SPG4 hereditary spastic paraplegia: clinical, genetic, and neuropathologic evidence. *Neurology* 2009;73(5):378-384.
79. Ribai P, Depienne C, Fedirko E, et al. Mental deficiency in three families with SPG4 spastic paraplegia. *European journal of human genetics* 2008;16(1):97-104.
80. Nielsen J, Johnsen B, Koefoed P, et al. Hereditary spastic paraplegia with cerebellar ataxia: a complex phenotype associated with a new SPG4 gene mutation. *European journal of neurology* 2004;11(12):817-824.
81. Tallaksen CM, Guichart-Gomez E, Verpillat P, et al. Subtle cognitive impairment but no dementia in patients with spastin mutations. *Archives of neurology* 2003;60(8):1113-1118.
82. Chamard L, Ferreira S, Pijoff A, Silvestre M, Berger E, Magnin E. Cognitive impairment involving social cognition in SPG4 hereditary spastic paraplegia. *Behavioural neurology* 2016;2016.
83. Orlacchio A, Kawarai T, Totaro A, et al. Hereditary spastic paraplegia: clinical genetic study of 15 families. *Archives of Neurology* 2004;61(6):849-855.
84. Orlacchio A, Gaudiello F, Totaro A, et al. A new SPG4 mutation in a variant form of spastic paraplegia with congenital arachnoid cysts. *Neurology* 2004;62(10):1875-1878.
85. Guthrie G, Pfeffer G, Bailie M, et al. The neurological and ophthalmological manifestations of SPG4-related hereditary spastic paraplegia. *Journal of neurology* 2013;260(3):906-909.
86. Kumar KR, Sue CM, Burke D, Ng K. Peripheral neuropathy in hereditary spastic paraplegia due to spastin (SPG4) mutation—a neurophysiological study using excitability techniques. *Clinical neurophysiology* 2012;123(7):1454-1459.

Tables

Table 1. Frequency of signs and symptoms among different categories of gender, mutation type and protein domain.

Clinical presentation	Frequency	Gender			Type of mutation			Protein domain (missense mutations)					
		Male (n=72, 45.9%)	Female (n=85, 54.1%)	P value	LoF (n=21/41)	Missense (n=20/41)	P value	MIT ^a	AAA_ATPase ^b	AAA_lid_3 ^c	Vps4_C ^d	Other domains	P value
Mean Age at onset ±SD	-	25.9±19.6	19±20	.040	24±20	27.4±20.2	.465	38	22.9±19.2	32.5±21.7	29.3±23.3	30.5±19.1	.626
Mean SPRS score ±SD	-	23.1±12.5	22±14.7	.853	22±8.7	22.8±15.6	.696	28	20.1±14.3	6.0±2.8	3	34.6±6.6	.066
Lower extremity weakness	53/102 (52%)	25/44 (56.8%)	28/58 (48.3%)	.392	10/24 (41.6%)	35/50 (70%)	.019	-	24/32 (75%)	3/10 (30%)	1/1 (100%)	7/7 (100%)	.010
Lower extremity spasticity	92/109 (84.4%)	44/48 (91.7%)	48/61 (78.7%)	.064	25/28 (89.3%)	45/49 (91.8%)	.708	-	32/32 (100%)	7/10 (70%)	-	6/7 (85.7%)	.008
Lower extremity hyperreflexia	102/111 (91.9%)	46/49 (93.9%)	56/62 (90.3%)	.496	25/25 (100%)	51/53 (96.2%)	.325	-	32/33 (97%)	13/13 (100%)	-	6/7 (85.7%)	.261
Extensor plantar response	82/105 (78.1%)	38/44 (86.4%)	44/61 (72.1%)	.082	19/24 (79.2%)	46/50 (92%)	.114	-	30/31 (96.8%)	10/12 (83.3%)	-	6/7 (85.7%)	.278
Abnormal bladder function	52/100 (52%)	25/43 (58.1%)	27/57 (47.4%)	.286	15/22 (68.2%)	28/51 (54.9%)	.290	-	15/32 (46.9%)	6/11 (54.5%)	1/1 (100%)	6/7 (85.7%)	.227
Ankle clonus	59/97 (60.8%)	25/40 (62.5%)	34/57 (59.6%)	.777	14/22 (63.6%)	34/47 (72.3%)	.464	-	19/28 (67.9%)	11/12 (91.7%)	-	4/7 (57.1%)	.189
Motor delay	9/50 (18%)	3/24 (12.5%)	6/26 (23.1%)	.331	0/10 (0%)	7/34 (20.6%)	.118	-	6/22 (27.3%)	0/6 (0%)	0/1 (0%)	1/5 (20%)	.490
Speech delay or abnormality	8/47 (17%)	3/24 (12.5%)	5/23 (21.7%)	.400	0/11 (0%)	8/31 (25.8%)	.061	-	6/20 (30%)	1/6 (16.7%)	0/1 (0%)	1/4 (25%)	.851
Learning disability	10/51 (19.6%)	3/25 (12%)	7/26 (26.9%)	.180	0/11 (0%)	8/34 (23.5%)	.076	-	7/22 (31.8%)	0/6 (0%)	0/1 (0%)	1/5 (20%)	.387
Progressive cognitive deficiency	6/73 (8.2%)	5/36 (13.9%)	1/37 (2.7%)	.082	1/19 (5.3%)	3/42 (7.1%)	.784	-	3/28 (10.7%)	0/6 (0%)	0/1 (0%)	0/7 (0%)	.656
Retinopathy or optic	1/71 (1.4%)	0/34 (0%)	1/38 (2.6%)	.348	-	-	-	-	-	-	-	-	-

atrophy													
Ocular movement abnormalities	1/72 (1.4%)	0/33 (0%)	1/38 (2.6%)	.341	1 /19 (5.3%)	0/41 (0%)	.139	-	-	-	-	-	-
Deafness	4/71 (5.6%)	3/34 (8.8%)	1/37 (2.7%)	.264	0 /19 (0%)	4/41 (9.8%)	.159	-	2/27 (7.4%)	0/6 (0%)	1/1 (100%)	1/7 (14.3%)	.017
Swallowing difficulty	4/72 (5.6%)	1/34 (2.9 %)	3/38 (7.9%)	.360	0 /20 (0%)	4/40 (10%)	.143	-	3/27 (11.1%)	0/6 (0%)	0/1 (0%)	1/6 (16.7%)	.774
Dysarthria	12/73 (16.4%)	5/35 (14.3%)	7/38 (18.4%)	.634	3 /20 (15%)	8/41 (19.5%)	.667	-	6/27 (22.2%)	1/6 (16.7%)	0/1 (0%)	1/7 (14.3%)	.914
Upper extremity weakness	5/72 (6.9%)	2/34 (5.9 %)	3/38 (7.7%)	.737	0 /19 (0%)	4/41 (9.8%)	.159	-	3/27 (11.1%)	0/6 (0%)	0/1 (0%)	1/7 (14.3%)	.807
Upper extremity hyperreflexia	31/77 (40.3%)	15/36 (41.7%)	16/41 (39%)	.814	11 /21 (52.4%)	15/44 (34.1%)	.159	-	8/27 (29.6%)	3/9 (33.3%)	1/1 (100%)	3/7 (42.9%)	.491
Sensory abnormalities	33/73 (45.2%)	15/35 (42.9%)	18/38 (47.4%)	.699	10 /21 (47.6%)	19/40 (47.5%)	.993	-	12/26 (46.2%)	3/6 (50%)	1/1 (100%)	3/7 (42.9%)	.753
Peripheral neuropathy	10/70 (14.3%)	4/35 (11.4%)	6/35 (17.1%)	.495	2 /20 (10%)	7/39 (17.9%)	.421	-	5/24 (20.8%)	1/8 (12.5%)	0/1 (0%)	1/6 (16.7%)	.914
Pes cavus	16/73 (21.9%)	9/35 (25.7%)	7/38 (18.4%)	.452	6 /19 (31.6%)	5/41 (12.2%)	.071	-	4/27 (14.8%)	0/6 (0%)	0/1 (0%)	1/7 (14.3%)	.759
Upper extremity ataxia	4/72 (5.6%)	1/35 (2.8%)	3/37 (8.1%)	.331	0 /19 (0%)	4/42 (9.5%)	.162	-	2/27 (7.4%)	1/7 (14.3%)	0/1 (0%)	1/7 (14.3%)	.893
Upper extremity Intent tremor	2/72 (2.8%)	0/35 (0%)	2/37 (5.4%)	.227	0 /19 (0%)	2/42 (4.8%)	.490	-	1/27 (3.7%)	0/7 (0%)	0/1 (0%)	1/7 (14.3%)	.331
Seizures	2/71 (2.8%)	0/34 (0%)	2/37 (5.4%)	.334	0 /19 (0%)	1/41 (2.4%)	.492	-	1/27 (3.7%)	0/6 (0%)	0/1 (0%)	0/7 (0%)	.912
Skeletal abnormality	4/69 (5.8%)	2/33 (6.1%)	2/36 (5.5%)	.929	0 /19 (0%)	3/40 (7.5%)	.220	-	2/27 (7.4%)	0/6 (0%)	-	1/7 (14.3%)	.621
Amyotrophy or lower motor neuron features	6/71 (8.5%)	5/35 (14.3%)	1/36 (2.8%)	.081	1/12 (8.3%)	3/19 (15.8%)	.542	-	3/33 (9.1%)	0/6 (0%)	-	1/8 (12.5%)	.823
Abnormal brain MRI	6/48 (12.5%)	4/28 (14.3%)	2/20 (10%)	.658	3 /13 (23.1%)	3/32 (9.4%)	.220	-	2/20 (10%)	0/5 (0%)	-	1/7 (14.3%)	.696
Abnormal spine MRI	8/53 (15.1%)	4/30 (13.3%)	4/23 (17.4%)	.683	2 /15 (13.3%)	6/32 (18.8%)	.645	-	4/17 (23.5%)	1/9 (11.1%)	-	1/6 (16.7%)	.735

Bonferroni corrected p value: 0.0017.

SD, standard deviation; SPRS, Spastic Paraplegia Rating Scale; MRI, magnetic resonance imaging; n, number; LoF, loss of function, MIT, microtubule interacting and trafficking; AAA, ATPase associated with various cellular activities; ATP, adenosine triphosphate.

^a InterPro (IPR) number: IPR036181; ^b IPR003593; ^c IPR041569; ^d IPR015415

Table 2. Mutations identified in the current study and their characteristics.

Nucleotide Change	Amino acid Change	Number of Families	Number of Patients	AF in gnomAD	Pathogenicity (VarSome)
c.127G>T	p.(Glu43Ter)	1	1	NR	Pathogenic
c.153C>G	p.(Tyr51Ter)	1	1	NR	Pathogenic
c.231G>A	p.(Trp77Ter)	1	1	NR	Pathogenic
c.562delG	p.(Ala188Profs Ter8)	1	1	NR	Pathogenic
c.687delT	p.(Ser229Argfs Ter11)	1	1	NR	Pathogenic
c.869A>G	p.(Lys290Arg)	1	1	NR	Likely Pathogenic
c.1098delG	p.(Glu366Aspfs Ter28)	1	1	NR	Pathogenic
c.1111C>T	p.(Leu371Phe)	2	2	NR	Pathogenic
c.1112T>C	p.(Leu371Pro)	1	1	NR	Pathogenic
c.1139T>C	p.(Leu380Pro)	1	1	NR	Likely Pathogenic
c.1144_1145 insGTC	p.(Gly382_Pro383 insArg)	1	2	NR	-
c.1158T>G	p.(Asn386Lys)	1	1	NR	Pathogenic
c.1196C>T	p.(Ser399Leu)	3	8	NR	Pathogenic
c.1209C>A	p.(Phe403Leu)	2	5	NR	Likely Pathogenic
c.1212_1216del	p.(Asn405Lysfs Ter36)	1	1	NR	-
c.1220G>A	p.(Ser407Asn)	1	1	NR	Pathogenic
c.1242A>G	p.(Lys414Lys)	2	3	NR	Uncertain significance
c.1276C>G	p.(Leu426Val)	1	1	NR	Pathogenic
c.1300C>T	p.(Gln434Ter)	1	1	NR	Pathogenic
c.1321+1G>A		1	2	NR	Pathogenic
c.1356_1357 insGGG	p.(Gly452dup)	1	1	NR	-
c.1378C>T	p.(Arg460Cys)	1	1	NR	Pathogenic
c.1385A>G	p.(Lys462Arg)	1	1	NR	Pathogenic
c.1466C>T	p.(Pro489Leu)	1	4	NR	Likely Pathogenic
c.1492_1493 del	p.(Arg498Alafs Ter30)	1	1	NR	Pathogenic
c.1493+2T>A		1	1	NR	Pathogenic

c.1495C>T	p.(Arg499Cys)	2	4	NR	Pathogenic
c.1496G>A	p.(Arg499His)	4	4	NR	Pathogenic
c.1507C>T	p.(Arg503Trp)	3	4	4.82E-06	Pathogenic
c.1537-2A>G		1	7	NR	Pathogenic
c.1610T>G	p.(Leu537Arg)	1	1	NR	Pathogenic
c.1676G>A	p.(Gly559Asp)	8	15	NR	Pathogenic
c.1684C>T	p.(Arg562Ter)	1	2	NR	Pathogenic
c.1685G>A	p.(Arg562Gln)	1	1	NR	Pathogenic
c.1729-1G>C		2	2	NR	Pathogenic
c.1741C>T	p.(Arg581Ter)	2	2	4.81E-06	Pathogenic
c.1790delG	p.(Ser597Thrfs Ter3)	1	3	NR	Pathogenic
c.1844C>T	p.(Thr615Ile)	1	1	NR	Likely Pathogenic
del exon 1		3	3	NR	Pathogenic
del exons 5-7		1	2	NR	Pathogenic
del exons 16-17		1	1	NR	Pathogenic

AAO, age at onset; UNK, unknown; NR, not reported; AF, allele frequency.

Table 3. Frequency of symptoms not attributable to the pure form of hereditary spastic paraplegia in the present study compared to previous studies.

Sign/ Symptom	Frequency in the current study	Previous studies
Learning disabilities	19.6%	4.2% intellectual impairment ¹ 1.5% Intellectual disability ⁶⁸
Speech delay or abnormality	17%	4/62 (6.5%) ⁶⁸
Dysarthria	16.4%	3% ⁶⁸
Cognitive deficiency	8.2%	Most frequent additional manifestation ⁷⁸⁻⁸²
Upper extremity intention tremor	2.8%	Mild arm tremor in 10% of patients ⁶⁸
Seizure	2.8%	Case reports ^{69, 75, 83, 84}
Ocular movement abnormality	1.4%	Case reports ⁸⁵
Swallowing difficulty	5.6%	3% ⁶⁸
Upper limb weakness	6.9%	Case reports ⁸⁰
Upper limb ataxia	5.6%	Case reports ⁸⁰
Peripheral neuropathy	14.3%	Case reports ⁸⁶
Deafness	5.6%	-

Figures

Figure 1. Structural analysis of human SPG4. A: Structure of SPG4 (PDB: 6PEN) forming a homohexameric assembly. The substrate peptide is shown as violet spheres. B: The interface between subunit A (green) and B (cyan) forms an active site. C: Loop₃₈₂₋₃₈₉ is part of the nucleotide binding pocket. D: β-strand₄₀₂₋₄₀₆ stabilizes helix₃₈₇₋₃₉₉ and the C-terminal residues of this structure. E: Arg498 and Arg499 interact with ADP, BeF₃, Asp470 and Ala495's main chain, stabilizing the nucleotide and the active site. Polar interactions are shown as yellow dotted lines. F: *D. melanogaster* Spastin structure (PDB: 6P07) was solved with an ATP molecule in the active site. Arg640 and Arg641, homologous to Arg498 and Arg499 in human Spastin, are coordinating ATP in the active site.

Supplementary Tables and Figures

Supplementary Table 1. Genes that are known or suspected to be involved in HSP or other similar disorders.

Supplementary Table 2. Copy number variations (CNVs) in the current study.

Supplementary Table 3. Signs and symptoms seen more frequently in patients harboring the p.(Arg499His) mutation.

Supplementary Table 4. Clinical manifestations of the family with probable biallelic inheritance. Patients 1 and 2 had undergone whole exome sequencing and were compound heterozygous carriers of the novel pathogenic mutation p.(Ser597ThrfsTer3), and the p.(Ser44Leu) variant. Sample from patient 3 was not available.

Supplementary Table 5. Characteristics of SPG4 patients with deafness.

Supplementary Figure 1. Distribution of age at onset in the cohort. The onset of SPG4 mostly occurs in the first five years of life, and between 35-44 years of age.

Supplementary Figure 2. Sanger sequencing results of family with modifier mutation. Mother and father carry the p.(Ser597ThrfsTer3) and p.(Ser44Leu) mutations respectively. Siblings are heterozygote for both mutations.