Title: RNA viromics of Southern California wastewater and detection of SARS-CoV-2 single nucleotide variants.

Running title: RNA viromics and SARS-CoV-2 detection in wastewater.

Jason A. Rothman^{a#}, Theresa B. Loveless^b, Joseph Kapcia III^a, Eric D. Adams^a, Joshua A. Steele^c, Amity G. Zimmer-Faust^c, Kylie Langlois^c, David Wanless^c, Madison Griffith^c, Lucy Mao^c, Jeffrey Chokry^c, John F. Griffith^c, Katrine L. Whiteson^{a#}

^a Department of Molecular Biology and Biochemistry, University of California, Irvine, Irvine, CA, USA.

^b Department of Biomedical Engineering, University of California, Irvine, Irvine, CA, USA.

^c Southern California Coastal Water Research Project, Costa Mesa, CA, USA

Co-corresponding authors: Jason A. Rothman, University of California, Irvine, Irvine, CA, 92697, (949) 824-3509, rothmanj@uci.edu. Katrine L. Whiteson, University of California, Irvine, Irvine, CA, 92697, (949) 824-9032, katrine@uci.edu.

Abstract:

Municipal wastewater provides an integrated sample of a diversity of human-associated microbes across a sewershed, including viruses. Wastewater-based epidemiology (WBE) is a promising strategy to detect pathogens and may serve as an early-warning system for disease outbreaks. Notably, WBE has garnered substantial interest during the COVID-19 pandemic to track disease burden through analyses of SARS-CoV-2 RNA. Throughout the COVID-19 outbreak, tracking SARS-CoV-2 in wastewater has been an important tool for understanding the spread of the virus. Unlike traditional sequencing of SARS-CoV-2 isolated from clinical samples, which adds testing burden to the healthcare system, in this study, metatranscriptomics was used to sequence virus directly from wastewater.

Here, we present a study in which we explored RNA viral diversity through sequencing 94 wastewater influent samples across seven treatment plants (WTPs), collected August 2020 – January 2021, representing approximately 16 million people in Southern California. Enriched viral libraries identified a wide diversity of RNA viruses that differed between WTPs and over time, with detected viruses including coronaviruses, influenza A, and noroviruses. Furthermore, single nucleotide variants (SNVs) of SARS-CoV-2 were identified in wastewater and we measured proportions of overall virus and SNVs across several months. We detected several SNVs that are markers for clinically-important SARS-CoV-2 variants, along with SNVs of unknown function, prevalence, or epidemiological consequence.

Our study shows the potential of WBE to detect viruses in wastewater and to track the diversity and spread of viral variants in urban and suburban locations, which may aid public health efforts to monitor disease outbreaks.

41 Importance:

42 Wastewater based epidemiology (WBE) can detect pathogens across sewersheds, which
 43 represents the collective waste of human populations. As there is a wide diversity of RNA
 44 viruses in wastewater, monitoring the presence of these viruses is useful for public health,
 45 industry, and ecological studies. Specific to public health, WBE has proven valuable during the
 46 COVID-19 pandemic to track the spread of SARS-CoV-2 without adding burden to healthcare
 47 systems. In this study, we used metatranscriptomics and RT-ddPCR to assay RNA viruses across
 48 Southern California wastewater from August 2020 – January 2021, representing approximately
 49 16 million people from Los Angeles, Orange, and San Diego counties. We found that SARS-
 50 CoV-2 quantification in wastewater correlates well with county-wide COVID-19 case data, and
 51 that we can detect SARS-CoV-2 single nucleotide variants through sequencing. Likewise, WTPs
 52 harbored different viromes, and we detected other human pathogens such as noroviruses and
 53 adenoviruses, furthering our understanding of wastewater viral ecology.

Introduction

Municipal wastewater represents a matrix containing a wide diversity of microbes and is representative of the collective waste of a human population across a catchment area. (1). The microbial content of wastewater can be useful in determining the levels of biological contamination of an area, including the presence of human and animal feces, antimicrobial resistance genes, pathogenic bacteria, and viruses (1–6). In regard to viruses specifically, wastewater often contains high titers of bacteriophages and plant-infecting viruses, along with generally smaller proportions of viruses that infect animals – including humans (7–10). Many studies have used metagenomics to characterize the viral content of wastewater, but these studies typically rely on extracted DNA, which is unable to capture the wide diversity of RNA-based viruses (11–13). As RNA viruses can be important pathogens of humans and agricultural organisms, using metatranscriptomic sequencing to study these diverse viruses in wastewater is relevant to public health and industry and may allow for a greater understanding of the ecological processes that occur in wastewater (2, 8, 14–16).

Wastewater based epidemiology (WBE) is a useful method to detect the presence of human pathogens and may serve as an early-warning system for disease outbreaks (17). For example, WBE has been used to track the prevalence of enteric viruses such as norovirus, rotavirus, adenovirus, poliovirus, influenza, and SARS coronaviruses (18–23), with the added benefit that WBE does not rely on public health and clinical resources (17). Aside from basic detection, it has been shown that the viral load of wastewater often precedes clinical outbreaks, and may offer a forecast of the severity of localized disease outbreaks (18, 24–26). Infectious diseases are often underreported in clinical settings - likely due to asymptomatic cases, avoidance of healthcare, and incorrect diagnoses - which prevents accurate disease surveillance

and incidence reporting, potentially hampering public health responses (27, 28). While not all pathogens are excreted into wastewater at detectable levels, the presence of detectable viruses in wastewater and human fecal samples indicates that WBE may still provide useful information in monitoring disease (21, 29, 30).

The coronavirus disease 2019 (COVID-19) pandemic has placed an intense strain on healthcare systems worldwide and has resulted in over 4 million human deaths (31). Caused by the 2019 emergence of the enveloped +ssRNA Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) (32, 33), this virus has been reported in direct nasal swab patient samples, human feces, and wastewater samples (15, 23–25, 29, 34–36). While detection and quantification of SARS-CoV-2 are clearly important, the predominant method of detection, RT-qPCR, does not allow for the characterization of viral variants, which is critical to monitor during the COVID-19 pandemic to track the evolution of SARS-CoV-2 (37, 38). In light of the limitations of RT-qPCR, thousands of patient-derived SARS-CoV-2 genomes have been sequenced, and important viral variants have been discovered (37, 39–41). However, the sequencing of patient samples mainly relies on clinical samples (42), which were difficult to obtain during the COVID-19 pandemic (43, 44). As wastewater represents a composite of human waste from the total catchment area, metagenomic sequencing of these samples may allow for the detection of viral variants and pathogen diversity across larger populations and regions without further burdening healthcare workers (11, 13, 45, 46).

Given the importance of monitoring the COVID-19 pandemic and exploring RNA viral diversity, in this study metatranscriptomic sequencing and droplet digital PCR (ddPCR) were conducted in parallel to characterize the RNA viromes and viral load of SARS-CoV-2 of 94 influent samples from seven wastewater treatment plants (WTPs) representing a total population

100 of 16 million individuals across Southern California. Through our study, we investigated several
101 lines of inquiry: First, what is the diversity of RNA viruses across Southern California
102 wastewater, and does viral abundance change longitudinally? Second, can we detect human-
103 infecting viruses - including coronaviruses - in wastewater, and how does SARS-CoV-2
104 quantification from wastewater samples compare to county-level COVID-19 case counts? Third,
105 can we use metatranscriptomics to detect and track the emergence of SARS-CoV-2 strain
106 variants in wastewater over time?

Results

ddPCR quantification of SARS-CoV-2 and correlation with daily countywide COVID-19 cases

We quantified SARS-CoV-2 viral load in 85 influent wastewater samples from August 2020 – early January 2021 across seven WTPs in the California counties of Los Angeles, Orange, and San Diego counties. Because of COVID-19 reporting at the county level, we correlated SARS-CoV-2 N1 gene copies per liter of influent wastewater with cases within counties only (i.e. the Hyperion WTP was correlated with Los Angeles county COVID-19 cases; supplemental file SF1). Generally, the viral load of each individual WTP correlated well with the seven-day rolling average of reported countywide COVID-19 cases: Hyperion ($\rho = 0.84$, $P < 0.001$), Joint Water Pollution Control Plant ($\rho = 0.83$, $P < 0.001$), North City ($\rho = 0.94$, $P = 0.03$), South Bay ($\rho = 0.37$, $P = 0.41$), Orange County ($\rho = 0.86$, $P < 0.001$), Point Loma ($\rho = 0.82$, $P < 0.001$), and San Jose Creek (Los Angeles county, $\rho = -0.15$, $P = 0.62$) (Fig. 1).

Sequencing library characteristics

We obtained a total of 1,119,674,084 quality-filtered deduplicated paired reads across 180 libraries (90 unique samples), and as we used two different library preparation strategies, we report the statistics separately as follows: For unenriched libraries, we obtained (794,773,512) reads (average: 84,55,037 range: 2,096,056 – 17,005,378) of which 40,512,123 (average: 430,980 [2.1%], range: 3,560 [0.05%] – 1,675,525 [13.5%]) were viruses and an average of 2.4% (range 0.06% - 25.1%) were human reads. For enriched libraries, we obtained 324,900,572 reads (average: 3,777,914; range: 1,182,078 - 14,392,582) of which 11,660,490 (average: 135,587 [4.7%], range: 2,580 [0.09%] – 484,944 [17.8%]) were viruses and an average of 2.0%

(range 0.06% - 19.1%) were human reads. We were able to pair 172 libraries (86 unique samples) that had both unenriched and enriched samples successfully sequenced.

Viral composition of wastewater and detection of selected pathogens

As enrichment changed the viral composition of wastewater samples, we analyzed the unenriched and enriched sample data separately. Unenriched samples contained sequences from 2,495 viruses, and the top ten most proportionally abundant viruses accounted for an average of 97.6% of the total abundance. The average relative abundances $\pm$ standard deviation of these “top ten” viruses were as follows: Tomato brown rugose fruit virus ($66.0 \pm 7.4\%$), Pepper mild mottle virus ($10.6 \pm 3.7\%$), Cucumber green mottle mosaic virus ($10.4 \pm 3.9\%$), Tomato mosaic virus ($4.8 \pm 2.9\%$), Tobacco mild green mosaic virus ($2.1 \pm 1.2\%$), Tropical soda apple mosaic virus ($1.5 \pm 0.8\%$), Tomato mottle mosaic virus ($1.4 \pm 1.8\%$), crAssphage ($0.4 \pm 0.8\%$), Opuntia virus 2 ($0.2 \pm 0.4\%$), and Melon necrotic spot virus ($0.2 \pm 0.2\%$) (Fig. 2).

We analyzed the data from IRV enriched samples as above, and found sequences from 2,215 viruses, with the top ten most proportionally abundant viruses accounting for an average of 97.2% of total abundance. The average relative abundances $\pm$ standard deviation of these “top ten” viruses were as follows: Tomato brown rugose fruit virus ($60.2 \pm 8.6\%$), Pepper mild mottle virus ($13.0 \pm 4.4\%$), Cucumber green mottle mosaic virus ($11.5 \pm 4.3\%$), Tomato mosaic virus ($5.0 \pm 3.0\%$), Tobacco mild green mosaic virus ($2.6 \pm 1.6\%$), Tropical soda apple mosaic virus ($1.8 \pm 1.4\%$), Tomato mottle mosaic virus ($1.5 \pm 1.4\%$), SARS-CoV-2 virus ($0.9 \pm 2.3\%$), crAssphage ($0.4 \pm 0.7\%$), and Opuntia virus 2 ($0.3 \pm 0.6\%$) (Fig. 2). We were able to detect many more reads of SARS-CoV-2 with respiratory virus enrichment, as there were only 337

SARS-CoV-2 reads (an average proportional abundance of 0.0004%) in unenriched samples, while across enriched samples, we detected 124,135 SARS-CoV-2 reads.

The Illumina Respiratory Virus Oligo Panel enriches for 40 viruses (including SARS-CoV-2), so we were able to compare viral detection in 86 enriched and unenriched samples, along with two non-respiratory viruses often detected in wastewater: norovirus and PMMoV. In enriched libraries, we detected the presence of SARS-CoV-2 in 68 samples, Human coronavirus (HCoV)-OC43 in 22 samples, HCoV-229E in one sample, Influenza A in 10 samples, human adenoviruses in 15 samples, human bocaviruses in 13 samples, and noroviruses in 52 samples. In unenriched samples, we detected SARS-CoV-2 in 24 samples, HCoV-OC43 in two samples, and noroviruses in 59 samples. Likewise, we detected PMMoV in all samples regardless of enrichment (Fig. 3), indicating that sequencing was successful.

Wastewater viral ecology

We analyzed the Shannon indexes for unenriched samples and found that overall alpha diversity was significantly different between wastewater sampling sites ($F_{(6,87)} = 7.5$, $P < 0.001$), then used Tukey's HSD post-hoc pairwise comparison testing to show that only NC-HTP, PL-HTP, PL-JWPCP, PL-OC, and PL-SJ alpha diversities were different from each other ($P_{\text{adj}} < 0.05$, Fig. 4). We also compared the Bray-Curtis dissimilarities of the samples with Adonis and found that overall treatment plants' beta diversity was significantly different ($R^2 = 0.41$, $P < 0.001$, Fig. 4). Additionally, we used ANCOM to test for differential abundance of viruses at > 0.0001 average relative abundance between treatment plants. This resulted in 16 viruses being differentially abundant between treatment plants ($W > 46$, $P_{\text{adj}} < 0.05$ each, Fig. 4). In enriched samples, we compared the proportional abundances of human respiratory viruses at > 0.0001

average relative abundance and show that SARS-CoV-2 and HCoV-OC43 were significantly different across treatment plants ($W > 46$, $P_{adj} < 0.05$ each, Fig. 4).

As we sampled the treatment plants multiple times over our study (up to 145 days from first sample), we were able to study how the viromes changed longitudinally. We compared diversity measures over time with LMEs using treatment plant as a random effect and show that both alpha and beta diversity remained stable across the sampling periods in unenriched samples (Shannon: $t = 0.21$, $P = 0.84$, Bray-Curtis: $t = -0.31$, $P = 0.76$). We also used LME on viruses present at > 0.0001 average relative abundance with treatment plant as a random effect and showed that 19 viruses' relative abundances changed over time ($P_{adj} = < 0.05$, Fig. S1). We specifically were interested in SARS-CoV-2 in enriched samples and used LMEs with treatment plant as a random effect, and found that the relative abundance of this virus increased over August 2020 – January 2021 ($t = 4.0$, $P < 0.001$, Fig. S1).

Sequencing SARS-CoV-2 SNVs in wastewater samples

We were interested in reads mapping specifically to the SARS-CoV-2 genome as a way to detect single nucleotide variants (SNVs) in wastewater. Across the 68 enriched samples that had detectable SARS-CoV-2 reads, we obtained an average breadth of genomic coverage of 24.0% (range 0.2% - 99.8%) per sample at an average sequencing depth of 8.1 reads per base (range 0.002 – 177.6) (Fig. 5 and supplemental file SF2). After masking the likely problematic nucleotide sites (as suggested by <https://virological.org/t/masking-strategies-for-sars-cov-2-alignments/480/14>, March 2021 update), we obtained 2558 SNVs (2002 unique) across all samples, however due to the low breadth of coverage in many samples, many of these sites may be spurious or unresolved. After applying a more stringent cutoff of 50% breadth of coverage

across the SARS-CoV-2 genome, we obtained 2060 SNVs (1656 unique) across 14 samples (Fig. 5 and supplemental file SF3).

As we took samples at multiple time points per treatment plant, we were able to track the proportion of SNVs per nucleotide position over time, most notably through samples taken from the Hyperion (HTP) facility, likely due to higher viral load leading to higher sequencing depth. Within HTP, we plotted the proportion of SNVs (as compared to the reference strain) over time at nucleotide positions obtained from at least three samples with greater than 50% breadth of genomic coverage. Three of the detected SNVs are apparently fixed in the viral population (sites 241, 14408, and 23403 are all 100% SNV), while the other 17 SNVs appear to vary widely over the sampling dates (overall average %CV = 103%, range 0 – 161%) with no apparent directionality (Fig. 6).

Discussion

Our composite wastewater samples contained a diversity of RNA viruses - mainly plant-infecting viruses along with lower relative abundances of animal-infecting viruses - that differed between wastewater treatment plants (WTPs), supporting previous studies indicating that location affects the presence of viruses (13). Furthermore, several individual viruses varied over time while overall diversity remained unchanged, likely due to localized infections within WTP catchment areas (3, 9, 47). We detected several human pathogenic viruses across all WTPs, supporting the hypothesis that wastewater-based epidemiology (WBE) has the potential to inform public health and researchers about viral presence and distribution without relying on standard healthcare practices (17). Through respiratory-virus-enriched library preparation and sequencing, we detected the presence of SARS-CoV-2 at every WTP, along with several SNVs across the SARS-CoV-2 genome. This suggests that WBE can reveal the pool of potential viral variants across large geographic areas, again without adding stress to healthcare systems, with the benefit that composite sampling can collect wastewater 24 hours per day (45, 46, 48). Lastly, we show that SARS-CoV-2 viral load at WTPs generally correlates with county-level COVID-19 case counts, indicating that WBE can be useful in monitoring the severity and dynamics of disease spread (18, 19, 23–25, 49).

The vast majority of viruses present in our samples were plant viruses, mainly those infecting tomatoes, peppers, and cucumbers in the genus *Tobamovirus*. This result is consistent with other studies, which suggests that these viruses are diverse and widespread in wastewater, likely originating from agricultural runoff or human feces (3, 7, 50). Even though these viruses were ubiquitous throughout our samples, WTPs had significantly different overall viromes, indicating there may be signatures of location and wastewater catchment throughout Southern

California, which has been suggested from other sampling locations (13). Aside from overall diversity, several plant- or arthropod-infecting viruses were differentially abundant between WTPs, possibly due to differences in peoples' diet (and thus viral excretion), infected plant growth, and localized infections of arthropods (i.e. Hubei picorna-like viruses and Beihai permutotetra-like viruses) (3, 9, 47). Likewise, we found that several viruses' relative abundances varied over time, regardless of treatment plant, suggesting that there are infection/subsidence dynamics or seasonal trends as previously suggested (7, 10, 49, 51). As the day-to-day relative abundance of viruses varied, we suggest that future wastewater surveys incorporate longitudinal and composite sampling to accurately capture viral diversity.

We detected several viruses pathogenic to humans across WTPs, including noroviruses adenoviruses, bocaviruses, coronaviruses, and influenza A - which agrees with other studies - indicating that WBE is robust and applicable to multiple viruses (18–23). By applying metagenomic or metatranscriptomic sequencing to wastewater, we can simultaneously detect a diversity of viruses, potentially alerting public health to unknown or underreported infections or new viral strains (27, 28, 46, 52). As we prepared sequencing libraries using two methods (unenriched and Illumina Respiratory Virus Oligo Panel enriched shotgun metatranscriptomics), we could compare the effects of enrichment on virus detection, and report that viral enrichment greatly improved our ability to detect influenza A and coronaviruses (especially SARS-CoV-2). We suggest that if researchers and the public health community are specifically interested in respiratory viruses, enrichment and the proper wastewater concentration /extraction methods should be employed for appropriate sensitivity and specificity to viruses of interest (23, 25, 34, 46, 53). Alongside sequencing, we also used ddPCR to quantify SARS-CoV-2 in wastewater and show that our results generally correlated well with county-reported 7-day rolling average

COVID-19 cases which agrees with previous studies (23, 24, 54). We note that SARS-CoV-2 viral load and case counts were not significantly correlated at every WTP indicating that there is likely unknown variability in the survival of RNA within wastewater, or that there may be variable influent flow or water quality affecting the viral load at specific WTPs (55, 56). We also note that our comparison is limited to county-level data, rather than being a comparison of the areas contributing to the influent stream and that this could be obscuring the relationship to case data at some treatment plants.

Our respiratory-virus-enriched metatranscriptomic sequencing detected the presence of SARS-CoV-2 at every WTP - and in most of our samples - although we note that could not accurately quantify viruses through sequencing. Instead, the power of metatranscriptomic sequencing lies in our ability to detect SNVs across the SARS-CoV-2 genome (46), and depending on the sample, we often sequenced >50% of the genome at appreciable read depth. For example, we detected the GISAID clade GH (lineage B.1*) markers 241C>T, 3037C>T, 14408C>T, 23403A>G, and 25563G>T at 100% prevalence where those regions of the genome were sequenced (39, 57, 58). Likewise, as we detected over 1000 SARS-CoV-2 SNVs across our samples, we found many SNVs of putatively unknown function that have been detected in patient samples, such as 6285C>T and 9891C>T (found in, but does not solely define, variants B.1.525 and B.1.1.318 respectively), 28854C>T and 28887C>T (40, 59, 60). We also detected many SNVs also found in wastewater samples from Northern California (46) and SNVs that to the best of our knowledge, have yet to be sequenced (41, 46, 48). As our sequencing data are not quantitative, our study suggests that sequencing wastewater is useful for SNV detection across wide catchment areas, but not the true prevalence of SNVs (46). Furthermore, we recognize that wastewater likely represents a collection of different viruses' RNA rather than one intact virion,

as SARS-CoV-2 may be relatively fragile in wastewater (55, 61). Lastly, we note that most SNVs came from samples with the highest number of SARS-CoV-2 reads, and suggest sequencing samples deeply to obtain a useful breadth and depth of viral genomes for variant detection (46, 62).

Conclusion

Wastewater-based epidemiology has the potential to aid public health in monitoring the spread and severity of disease outbreaks. Our research contributes to the growing field of WBE by showing that Southern California wastewater harbors a diversity of RNA viruses, and that these viral populations vary over time and between WTPs. Likewise, we are able to detect human viral pathogens without increasing the burden on local healthcare systems, further supporting the benefit of WBE. Through ddPCR and metatranscriptomic sequencing, we were able to measure the viral load of SARS-CoV-2 in wastewater and identify potentially novel SNVs which may assist in monitoring viral evolution and the emergence of new variants. We suggest that future researchers employ longitudinal metatranscriptomic sequencing on wastewater samples to further understand the spread of RNA viruses and how these viruses change over time.

Materials and Methods

Sample collection

We collected 94 1-liter 24-hour composite influent wastewater samples at seven WTPs across Southern California between August 2020 – January 2021 (Supplemental file SF4). The samples were aliquoted into 50mL tubes, then stored at 4°C until sample processing. Note that extractions were performed independently for ddPCR quantification and viromic sequencing.

Wastewater sample processing for SARS-CoV-2 ddPCR quantification

We prepared influent samples for ddPCR following the method described in Steele et al., 2021 (63). Briefly, we first added bovine coronavirus vaccine (BoCoV, Bovilis Merck & Co, Kenilworth, NJ) to 20 mL of wastewater as a sample processing control to assess viral RNA recovery. We then added MgCl₂ to a final concentration of 25mM and adjusted the pH to < 3.5 with 20% HCl on a mixed cellulose ester membrane (Type HA: Millipore, Bedford, MA) in replicates of six. We then transferred the HA filters to pre-loaded 2 mL ZR BashingBead Lysis tubes (Zymo, Irvine, CA), and bead beat the samples with a BioSpec beadbeater (BioSpec Products, Bartlesville, OK) for one minute. We then extracted total nucleic acids with a BioMerieux NucliSENS extraction kit with magnetic bead capture (BioMerieux, Durham, NC) following the manufacturer's protocol.

SARS-CoV-2 reverse transcriptase droplet digital PCR quantification and correlation with countywide COVID-19 cases

We used one-step reverse transcription droplet digital PCR (RT-ddPCR) to quantify the N1 region of the SARS-CoV-2 N gene with primer and probe sequences designed by the CDC

(63, 64) and we quantified the bovine coronavirus using previously designed primers (63, 65). We set up RT-ddPCR reactions following the manufacturer's instructions on a Bio-Rad Qx200 (Bio-Rad, Hercules, CA). For all assays, a minimum of two reactions and a total of $\geq 20,000$ droplets were generated per sample and at least five no template control (NTC) reactions and two positive control reactions were run per 96-well plate as well as extraction-specific NTCs. Each sample was required to have a minimum of three positive droplets (66) to be included in further analyses. We also assessed RNA recovery using the BoCoV exogenous control, and samples with $< 3\%$ recovery were excluded from further analyses.

We used the SARS-CoV-2 ddPCR quantification data and compared viral loads with county-level COVID-19 reported case data from the State of California Health and Human Services Agency (<https://data.chhs.ca.gov/dataset/covid-19-time-series-metrics-by-county-and-state>). We calculated rolling 7-day averages for COVID-19 cases in Los Angeles, Orange, and San Diego counties with the R package "zoo" (67), and ran Pearson correlations between viral load and reported COVID-19 cases with the R package "Hmisc" (68) at each time point where we had both data points.

Wastewater sample processing for metatranscriptomic sequencing

We followed a similar protocol as Crits-Cristoph et al 2021 and Wu et al 2020 (23, 46) to concentrate viruses and extract RNA. We pasteurized 50 mL of wastewater in a 65 °C water bath for 90 minutes, then filtered the samples through a sterile 0.22 um vacuum filter (VWR, Radnor, PA) to remove solids. We then concentrated the filtrate through ultracentrifugation at 3,000xg with 10kDa Amicon filters (MilliporeSigma, Burlington, MA) by successively centrifuging then discarding the flowthrough until the entire 50 mL sample was processed. This resulted in final

volumes of < 500uL for each sample, which we then stored at -80 °C until RNA extraction. We thawed the wastewater concentrate on ice, then used an Invitrogen PureLink RNA Mini Kit with DNase (Invitrogen, Waltham, MA) to extract RNA by following the manufacturer’s protocol, quantified the resulting RNA with an AccuBlue Broad Range RNA Quantification kit (Biotium, Fremont, CA) kit on a DeNovix QFX Fluorometer (DeNovix, Wilmington, DE) spectrophotometer, then stored the RNA at -80 °C.

Next generation sequencing library preparation

Sample library preparation and next-generation sequencing was performed by the UC Irvine Genomics High Throughput Facility (GHTF). The GHTF used two separate library preparation strategies per sample: One library was prepared with the Illumina RNA Prep with Enrichment kit (Illumina, San Diego, CA), and the other was prepared using the same preparation kit with the addition of the Illumina Respiratory Virus Oligo Panel to enrich for human respiratory viruses. The GHTF then sequenced the paired-end libraries either 2x100bp or 2x150bp (see supplemental file SF4) on an Illumina NovaSeq 6000 with an S4 300 cycle kit and sent the data as demultiplexed FASTQ files.

Bioinformatics and sequence data processing

We used the UC Irvine High Performance Community Computing Cluster (HPC3) for all data processing on the provided FASTQ files. We removed primers, adapter sequences, and low-quality bases with “bbduk” in the BBTools software package (69), then removed PCR duplicates with BBTools “dedupe.” We downloaded the NCBI Virus RefSeq Genome database and used Bowtie2 (70) to build an index of the viral genomes, then mapped the cleaned FASTQ reads

against this index. We used “Samtools” (71) to convert the resulting SAM files to sorted BAM files and finally used “inStrain” (72) to calculate all viral abundances and profile viral variants on read pairs with >90% average nucleotide identity to their respective reference genomes. Lastly, we used “mosdepth” (73) and “bedtools” (74) to calculate genome coverage and breadth across all SARS-CoV-2 sequences.

For community diversity analyses, we tabulated viral abundances, then normalized the reads into within-sample relative abundances in R (75). We used this table to generate Shannon Diversity indices and Bray-Curtis dissimilarity matrices with the R package “vegan” (76). We also ran Adonis PERMANOVA tests on the distance matrices, and performed non-metric multidimensional scaling on the data, compared the proportional abundances of viruses with ANCOM (77), and compared the relative abundances of viruses over time with “lmerTest” using WTP as a random effect (78). Lastly, we plotted all figures with “ggplot2” (79), “gggenes” (80), or “pheatmap” (81).

Data Availability

Raw sequencing data has been deposited on the NCBI Sequence Read Archive under accession number PRJNA729801, and representative code can be found at github.com/jasonarothman/wastewater_viroomics_sarscov2.

Acknowledgements

We would like to thank the staff of City of Los Angeles Sanitation and Environment, Los Angeles County Sanitation District, Orange County Sanitation District, and City of San Diego Public Utilities for collecting influent. We would also like to thank E. Macias for assistance with samples. This research was supported by Emergency COVID-19 Research Seed Funding through the University of California Office of the President Research Grants Program Office (awards #R01RG3732 and #R00RG2814) awarded to JAR, TBL, and KWL, and a Hewitt Foundation for Biomedical Research postdoctoral fellowship to JAR. This work was made possible, in part, through access to the Genomics High Throughput Facility Shared Resource of the Cancer Center Support Grant (P30CA-062203) at the University of California, Irvine and NIH shared instrumentation grants 1S10RR025496-01, 1S10OD010794-01, and 1S10OD021718-01, and access to computing resources from the UCI High Performance Cloud Computing Center.

394

395

396 Figure 1: Pearson correlations of SARS-CoV-2 wastewater viral loads as measured by RT-
397 ddPCR and the seven-day rolling average of COVID-19 case counts within county of WTP
398 faceted by WTP. COVID-19 rolling averages and wastewater viral load were generally
399 significantly correlated: HTP ($\rho = 0.84$, $P < 0.001$), JWPCP ($\rho = 0.83$, $P < 0.001$), NC ($\rho = 0.94$,
400 $P = 0.03$), SB ($\rho = 0.37$, $P = 0.41$), OC ($\rho = 0.86$, $P < 0.001$), PL ($\rho = 0.82$, $P < 0.001$), and SJ (ρ
401 $= -0.15$, $P = 0.62$).

Figure 2: Stacked barplot indicating the ten most proportionally abundant viruses across all samples, with the relative abundance of SARS-CoV-2 being highest in HTP samples from 12/29/20. Abbreviations denote WTP, “enriched” are samples enriched for respiratory viruses, and dates across the “x” axis signify sampling date.

409 Figure 3: Heatmap indicating the presence or absence of reads mapping to each listed virus. Heat
 410 color denotes if a virus was detected in unenriched samples, respiratory virus enriched samples,
 411 both, or neither type of sample, and abbreviations signify WTP.

Figure 4: A) Nonmetric multidimensional scaling (NMDS) ordination of the Bray-Curtis dissimilarities of all unenriched samples colored by WTP. Treatment plants were significantly different from each other ($R^2 = 0.41$, $P < 0.001$). B) Boxplots of the Shannon diversity indexes colored by WTP. Overall alpha diversity was significantly different ($F_{(6,87)} = 7.5$, $P < 0.001$). C) Boxplots of virus relative abundances that significantly differed between treatment plants as

419 tested with ANCOM, with the right bottom two panels (SARS-CoV-2 and HCoV OC43)
420 representing enriched samples *only*.

Figure 5: Multipanel figure showing sequence reads per base across all enriched samples, the position and proportion of SNVs across samples with >50% genomic breadth only, and an ORF map of the SARS-CoV-2 genome.

426

427

428

429 Figure 6: Line and dot plot of SNVs sequenced at least three times at the Hyperion WTP faceted
 430 by SARS-CoV-2 genomic position over time in samples with > 50% sequencing breadth.
 431 “Proportion of SNVs” refers to the proportion of SNVs versus the SARS-CoV-2 reference strain
 432 (i.e. 1 SNV and 3 reference reads = 0.25 proportion).

References:

1. Newton RJ, McClary JS. 2019. The flux and impact of wastewater infrastructure microorganisms on human and ecosystem health. *Curr Opin Biotechnol* 57:145–150.
2. Wu L, Ning D, Zhang B, Li Y, Zhang P, Shan X, Zhang Q, Brown MR, Li Z, Van Nostrand JD, Ling F, Xiao N, Zhang Y, Vierheilig J, Wells GF, Yang Y, Deng Y, Tu Q, Wang A, Global Water Microbiome Consortium, Zhang T, He Z, Keller J, Nielsen PH, Alvarez PJJ, Criddle CS, Wagner M, Tiedje JM, He Q, Curtis TP, Stahl DA, Alvarez-Cohen L, Rittmann BE, Wen X, Zhou J. 2019. Global diversity and biogeography of bacterial communities in wastewater treatment plants. *Nat Microbiol* 4:1183–1195.
3. Symonds EM, Griffin DW, Breitbart M. 2009. Eukaryotic viruses in wastewater samples from the United States. *Appl Environ Microbiol* 75:1402–1409.
4. McLellan SL, Huse SM, Mueller-Spitz SR, Andreishcheva EN, Sogin ML. 2010. Diversity and population structure of sewage-derived microorganisms in wastewater treatment plant influent. *Environ Microbiol* 12:378–392.
5. Karkman A, Do TT, Walsh F, Virta MPJ. 2018. Antibiotic-resistance genes in waste water. *Trends Microbiol* 26:220–228.
6. Edwards RA, Vega AA, Norman HM, Ohaeri M, Levi K, Dinsdale EA, Cinek O, Aziz RK, McNair K, Barr JJ, Bibby K, Brouns SJJ, Cazares A, de Jonge PA, Desnues C, Díaz Muñoz SL, Fineran PC, Kurilshikov A, Lavigne R, Mazankova K, McCarthy DT, Nobrega FL, Reyes Muñoz A, Tapia G, Trefault N, Tyakht AV, Vinuesa P, Wagemans J, Zhernakova A, Aarestrup FM, Ahmadov G, Alassaf A, Anton J, Asangba A, Billings EK, Cantu VA, Carlton JM, Cazares D, Cho G-S, Condeff T, Cortés P, Cranfield M, Cuevas DA, De la Iglesia R, Decewicz P, Doane MP, Dominy NJ, Dziewit L, Elwasila BM, Eren AM, Franz C, Fu J, Garcia-Aljaro C, Ghedin E, Gulino KM, Haggerty JM, Head SR, Hendriksen RS, Hill C, Hyöty H, Ilina EN, Irwin MT, Jeffries TC, Jofre J, Junge RE, Kelley ST, Khan Mirzaei M, Kowalewski M, Kumaresan D, Leigh SR, Lipson D, Lisitsyna ES, Llagostera M, Maritz JM, Marr LC, McCann A, Molshanski-Mor S, Monteiro S, Moreira-Grez B, Morris M, Mugisha L, Muniesa M, Neve H, Nguyen N-P, Nigro OD, Nilsson AS, O’Connell T, Odeh R, Oliver A, Piuri M, Prussin AJ Ii, Qimron U, Quan Z-X, Rainetova P, Ramírez-Rojas A, Raya R, Reasor K, Rice GAO, Rossi A, Santos R, Shimashita J, Stachler EN, Stene LC, Strain R, Stumpf R, Torres PJ, Twaddle A, Ugochi Ibekwe M, Villagra N, Wandro S, White B, Whiteley A, Whiteson KL, Wijmenga C, Zambrano MM, Zschach H, Dutilh BE. 2019. Global phylogeography and ancient evolution of the widespread human gut virus crAssphage. *Nat Microbiol* 4:1727–1736.
7. Cantalupo PG, Calgua B, Zhao G, Hundesa A, Wier AD, Katz JP, Grabe M, Hendrix RW, Girones R, Wang D, Pipas JM. 2011. Raw sewage harbors diverse viral populations. *MBio* 2.
8. Rosario K, Nilsson C, Lim YW, Ruan Y, Breitbart M. 2009. Metagenomic analysis of viruses in reclaimed water. *Environ Microbiol* 11:2806–2820.

- 473 9. Kitajima M, Sassi HP, Torrey JR. 2018. Pepper mild mottle virus as a water quality
474 indicator. *npj Clean Water* 1:19.
- 475 10. Eftim SE, Hong T, Soller J, Boehm A, Warren I, Ichida A, Nappier SP. 2017. Occurrence of
476 norovirus in raw sewage - A systematic literature review and meta-analysis. *Water Res*
477 111:366–374.
- 478 11. Martínez-Puchol S, Rusiñol M, Fernández-Cassi X, Timoneda N, Itarte M, Andrés C, Antón
479 A, Abril JF, Girones R, Bofill-Mas S. 2020. Characterisation of the sewage virome:
480 comparison of NGS tools and occurrence of significant pathogens. *Sci Total Environ*
481 713:136604.
- 482 12. Ng TFF, Marine R, Wang C, Simmonds P, Kapusinszky B, Bodhidatta L, Oderinde BS,
483 Wommack KE, Delwart E. 2012. High variety of known and new RNA and DNA viruses of
484 diverse origins in untreated sewage. *J Virol* 86:12161–12175.
- 485 13. Bibby K, Peccia J. 2013. Identification of viral pathogen diversity in sewage sludge by
486 metagenome analysis. *Environ Sci Technol* 47:1945–1951.
- 487 14. Ferrera I, Sánchez O. 2016. Insights into microbial diversity in wastewater treatment
488 systems: How far have we come? *Biotechnol Adv* 34:790–802.
- 489 15. Bivins A, North D, Ahmad A, Ahmed W, Alm E, Been F, Bhattacharya P, Bijlsma L,
490 Boehm AB, Brown J, Buttiglieri G, Calabro V, Carducci A, Castiglioni S, Cetecioglu Gurol
491 Z, Chakraborty S, Costa F, Curcio S, De Los Reyes FL, Delgado Vela J, Farkas K,
492 Fernandez-Casi X, Gerba C, Gerrity D, Girones R, Gonzalez R, Haramoto E, Harris A,
493 Holden PA, Islam MT, Jones DL, Kasprzyk-Hordern B, Kitajima M, Kotlarz N, Kumar M,
494 Kuroda K, La Rosa G, Malpei F, Mautus M, McLellan SL, Medema G, Meschke JS,
495 Mueller J, Newton RJ, Nilsson D, Noble RT, Van Nuijs A, Peccia J, Perkins TA, Pickering
496 AJ, Rose J, Sanchez G, Smith A, Stadler L, Stauber C, Thomas K, Van Der Voorn T,
497 Wigginton K, Zhu K, Bibby K. 2020. Wastewater-based epidemiology: global collaborative
498 to maximize contributions in the fight against COVID-19. *Environmental Science and*
499 *Technology*. American Chemical Society.
- 500 16. Carrasco-Hernandez R, Jácome R, López Vidal Y, Ponce de León S. 2017. Are RNA
501 viruses candidate agents for the next global pandemic? A review. *ILAR J* 58:343–358.
- 502 17. Sims N, Kasprzyk-Hordern B. 2020. Future perspectives of wastewater-based
503 epidemiology: Monitoring infectious disease spread and resistance to the community level.
504 *Environ Int* 139:105689.
- 505 18. Hellmér M, Paxéus N, Magnus L, Enache L, Arnholm B, Johansson A, Bergström T,
506 Norder H. 2014. Detection of pathogenic viruses in sewage provided early warnings of
507 hepatitis A virus and norovirus outbreaks. *Appl Environ Microbiol* 80:6771–6781.
- 508 19. Manor Y, Handsch R, Halmut T, Neuman M, Bobrov A, Rudich H, Vonsover A, Shulman
509 L, Kew O, Mendelson E. 1999. Detection of poliovirus circulation by environmental

- surveillance in the absence of clinical cases in Israel and the Palestinian Authority. *J Clin Microbiol* 37:1670–1675.
20. Blomqvist S, El Bassioni L, El Maamoon Nasr EM, Paananen A, Kaijalainen S, Asghar H, de Gourville E, Roivainen M. 2012. Detection of imported wild polioviruses and of vaccine-derived polioviruses by environmental surveillance in Egypt. *Appl Environ Microbiol* 78:5406–5409.
21. Heijnen L, Medema G. 2011. Surveillance of influenza A and the pandemic influenza A (H1N1) 2009 in sewage and surface water in the Netherlands. *J Water Health* 9:434–442.
22. Wang XW, Li J, Guo T, Zhen B, Kong Q, Yi B, Li Z, Song N, Jin M, Xiao W, Zhu X, Gu C, Yin J, Wei W, Yao W, Liu C, Li J, Ou G, Wang M, Fang T, Wang G, Qiu Y, Wu H, Chao F, Li J. 2005. Concentration and detection of SARS coronavirus in sewage from Xiao Tang Shan Hospital and the 309th Hospital of the Chinese People’s Liberation Army. *Water Sci Technol* 52:213–221.
23. Wu F, Zhang J, Xiao A, Gu X, Lee WL, Armas F, Kauffman K, Hanage W, Matus M, Ghaeli N, Endo N, Duvallet C, Poyet M, Moniz K, Washburne AD, Erickson TB, Chai PR, Thompson J, Alm EJ. 2020. SARS-CoV-2 titers in wastewater are higher than expected from clinically confirmed cases. *mSystems* 5:e00614-20.
24. Karthikeyan S, Ronquillo N, Belda-Ferre P, Alvarado D, Javidi T, Longhurst CA, Knight R. 2021. High-throughput wastewater SARS-CoV-2 detection enables forecasting of community infection dynamics in San Diego County. *mSystems* 6.
25. Peccia J, Zulli A, Brackney DE, Grubaugh ND, Kaplan EH, Casanovas-Massana A, Ko AI, Malik AA, Wang D, Wang M, Warren JL, Weinberger DM, Arnold W, Omer SB. 2020. Measurement of SARS-CoV-2 RNA in wastewater tracks community infection dynamics. *Nat Biotechnol* 38:1164–1167.
26. Kopel E, Kaliner E, Grotto I. 2014. Lessons from a public health emergency--importation of wild poliovirus to Israel. *N Engl J Med* 371:981–983.
27. Gibbons CL, Mangen M-JJ, Plass D, Havelaar AH, Brooke RJ, Kramarz P, Peterson KL, Stuurman AL, Cassini A, Fèvre EM, Kretzschmar MEE. 2014. Measuring underreporting and under-ascertainment in infectious disease datasets: a comparison of methods. *BMC Public Health* 14:147.
28. Silverman JD, Hupert N, Washburne AD. 2020. Using influenza surveillance networks to estimate state-specific prevalence of SARS-CoV-2 in the United States. *Sci Transl Med* 12.
29. Medema G, Heijnen L, Elsinga G, Italiaander R, Brouwer A. 2020. Presence of SARS-Coronavirus-2 RNA in sewage and correlation with reported COVID-19 prevalence in the early stage of the epidemic in The Netherlands. *Environ Sci Technol Lett* 7:511–516.

30. Wigginton KR, Ye Y, Ellenberg RM. 2015. Emerging investigators series: The source and fate of pandemic viruses in the urban water cycle. *Environmental Science: Water Research and Technology* 1:735–746.
31. Dong E, Du H, Gardner L. 2020. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect Dis* 20:533–534.
32. Chan JF-W, Kok K-H, Zhu Z, Chu H, To KK-W, Yuan S, Yuen K-Y. 2020. Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerg Microbes Infect* 9:221–236.
33. 2020. WHO Statement regarding cluster of pneumonia cases in Wuhan, China.
34. Rothman JA, Loveless TB, Griffith ML, Steele JA, Griffith JF, Whiteson KL. 2020. Metagenomics of wastewater influent from Southern California wastewater treatment facilities in the era of COVID-19. *Microbiology Resource Announcements* 9:19–21.
35. Prado T, Fumian TM, Mannarino CF, Resende PC, Motta FC, Eppinghaus ALF, Chagas do Vale VH, Braz RMS, de Andrade J da SR, Maranhão AG, Miagostovich MP. 2021. Wastewater-based epidemiology as a useful tool to track SARS-CoV-2 and support public health policies at municipal level in Brazil. *Water Res* 191:116810.
36. Wu Y, Guo C, Tang L, Hong Z, Zhou J, Dong X, Yin H, Xiao Q, Tang Y, Qu X, Kuang L, Fang X, Mishra N, Lu J, Shan H, Jiang G, Huang X. 2020. Prolonged presence of SARS-CoV-2 viral RNA in faecal samples. *Lancet Gastroenterol Hepatol* 5:434–435.
37. Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, Hengartner N, Giorgi EE, Bhattacharya T, Foley B, Hastie KM, Parker MD, Partridge DG, Evans CM, Freeman TM, de Silva TI, Sheffield COVID-19 Genomics Group, McDanal C, Perez LG, Tang H, Moon-Walker A, Whelan SP, LaBranche CC, Saphire EO, Montefiori DC. 2020. Tracking changes in SARS-CoV-2 spike: evidence that D614G increases infectivity of the COVID-19 virus. *Cell* 182:812–827.e19.
38. Darby AC, Hiscox JA. 2021. Covid-19: variants and vaccination. *BMJ* 372.
39. Elbe S, Buckland-Merrett G. 2017. Data, disease and diplomacy: GISAID’s innovative contribution to global health. *Glob Chall* 1:33–46.
40. Badua CLDC, Baldo KAT, Medina PMB. 2021. Genomic and proteomic mutation landscapes of SARS-CoV-2. *J Med Virol* 93:1702–1721.
41. Fang S, Li K, Shen J, Liu S, Liu J, Yang L, Hu C-D, Wan J. 2021. GESS: a database of global evaluation of SARS-CoV-2/hCoV-19 sequences. *Nucleic Acids Res* 49:D706–D714.
42. Rouchka EC, Chariker JH, Chung D. 2020. Variant analysis of 1,040 SARS-CoV-2 genomes. *PLoS One* 15:e0241535.

43. Leite H, Lindsay C, Kumar M. 2020. COVID-19 outbreak: implications on healthcare operations. *The TQM Journal* 39:88.
44. Reiner RC, Barber RM, Collins JK, Zheng P, Adolph C, Albright J, Antony CM, Aravkin AY, Bachmeier SD, Bang-Jensen B, Bannick MS, Bloom S, Carter A, Castro E, Causey K, Chakrabarti S, Charlson FJ, Cogen RM, Combs E, Dai X, Dangel WJ, Earl L, Ewald SB, Ezalarab M, Ferrari AJ, Flaxman A, Frostad JJ, Fullman N, Gakidou E, Gallagher J, Glenn SD, Goosmann EA, He J, Henry NJ, Hulland EN, Hurst B, Johanns C, Kendrick PJ, Khemani A, Larson SL, Lazzar-Atwood A, LeGrand KE, Lescinsky H, Lindstrom A, Linebarger E, Lozano R, Ma R, Månsson J, Magistro B, Herrera AMM, Marczak LB, Miller-Petrie MK, Mokdad AH, Morgan JD, Naik P, Odell CM, O'Halloran JK, Osgood-Zimmerman AE, Ostroff SM, Pasovic M, Penberthy L, Phipps G, Pigott DM, Pollock I, Ramshaw RE, Redford SB, Reinke G, Rolfe S, Santomauro DF, Shackleton JR, Shaw DH, Sheena BS, Sholokhov A, Sorensen RJD, Sparks G, Spurlock EE, Subart ML, Syailendrawati R, Torre AE, Troeger CE, Vos T, Watson A, Watson S, Wiens KE, Woyczynski L, Xu L, Zhang J, Hay SI, Lim SS, Murray CJL, IHME COVID-19 Forecasting Team. 2021. Modeling COVID-19 scenarios for the United States. *Nat Med* 27:94–105.
45. Nemudryi A, Nemudraia A, Wiegand T, Surya K, Buyukyoruk M, Cicha C, Vanderwood KK, Wilkinson R, Wiedenheft B. 2020. Temporal detection and phylogenetic assessment of SARS-CoV-2 in municipal wastewater. *Cell Rep Med* 1:100098.
46. Crits-Christoph A, Kantor RS, Olm MR, Whitney ON, Al-Shayeb B, Lou YC, Flamholz A, Kennedy LC, Greenwald H, Hinkle A, Hetzel J, Spitzer S, Koble J, Tan A, Hyde F, Schroth G, Kuersten S, Banfield JF, Nelson KL. 2021. Genome sequencing of sewage detects regionally prevalent SARS-CoV-2 variants. *MBio* 12.
47. Shi M, Lin X-D, Tian J-H, Chen L-J, Chen X, Li C-X, Qin X-C, Li J, Cao J-P, Eden J-S, Buchmann J, Wang W, Xu J, Holmes EC, Zhang Y-Z. 2016. Redefining the invertebrate RNA virosphere. *Nature* 540:539–543.
48. Fontenele RS, Kraberger S, Hadfield J, Driver EM, Bowes D, Holland LA, Faleye TOC, Adhikari S, Kumar R, Inchausti R, Holmes WK, Deitrick S, Brown P, Duty D, Smith T, Bhatnagar A, Yeager RA, Holm RH, Hoogesteijn von Reitzenstein N, Wheeler E, Dixon K, Constantine T, Wilson MA, Lim ES, Jiang X, Halden RU, Scotch M, Varsani A. 2021. High-throughput sequencing of SARS-CoV-2 in wastewater provides insights into circulating variants. *medRxiv* <https://doi.org/10.1101/2021.01.22.21250320>.
49. Kazama S, Masago Y, Tohma K, Souma N, Imagawa T, Suzuki A, Liu X, Saito M, Oshitani H, Omura T. 2016. Temporal dynamics of norovirus determined through monitoring of municipal wastewater by pyrosequencing and virological surveillance of gastroenteritis cases. *Water Res* 92:244–253.
50. Bačnik K, Kutnjak D, Pecman A, Mehle N, Tušek Žnidarič M, Gutiérrez Aguirre I, Ravnika M. 2020. Viromics and infectivity analysis reveal the release of infective plant viruses from wastewater into the environment. *Water Res* 177:115628.

- 619 51. Brinkman NE, Fout GS, Keely SP. 2017. Retrospective surveillance of wastewater to
620 examine seasonal dynamics of enterovirus infections. *mSphere* 2.
- 621 52. Polo D, Quintela-Baluja M, Corbishley A, Jones DL, Singer AC, Graham DW, Romalde JL.
622 2020. Making waves: Wastewater-based epidemiology for COVID-19 - approaches and
623 challenges for surveillance and prediction. *Water Res* 186:116404.
- 624 53. Philo SE, Keim EK, Swanstrom R, Ong AQW, Burnor EA, Kossik AL, Harrison JC,
625 Demeke BA, Zhou NA, Beck NK, Shirai JH, Meschke JS. 2021. A comparison of SARS-
626 CoV-2 wastewater concentration methods for environmental surveillance. *Sci Total Environ*
627 760:144215.
- 628 54. Wu F, Xiao A, Zhang J, Moniz K, Endo N, Armas F, Bonneau R, Brown MA, Bushman M,
629 Chai PR, Duvallet C, Erickson TB, Foppe K, Ghaeli N, Gu X, Hanage WP, Huang KH, Lee
630 WL, Matus M, McElroy KA, Nagler J, Rhode SF, Santillana M, Tucker JA, Wuertz S, Zhao
631 S, Thompson J, Alm EJ. 2020. SARS-CoV-2 titers in wastewater foreshadow dynamics and
632 clinical presentation of new COVID-19 cases. *medRxiv* 2020.06.15.20117747.
- 633 55. Bivins A, Greaves J, Fischer R, Yinda KC, Ahmed W, Kitajima M, Munster VJ, Bibby K.
634 2020. Persistence of SARS-CoV-2 in water and wastewater. *Environ Sci Technol Lett*
635 7:937–942.
- 636 56. Bivins A, North D, Wu Z, Shaffer M, Ahmed W, Bibby K. 2021. Within-day variability of
637 SARS-CoV-2 RNA in municipal wastewater influent during periods of varying COVID-19
638 prevalence and positivity. *bioRxiv. medRxiv*.
- 639 57. Alm E, Broberg EK, Connor T, Hodcroft EB, Komissarov AB, Maurer-Stroh S, Melidou A,
640 Neher RA, O'Toole Á, Pereyaslov D, WHO European Region sequencing laboratories and
641 GISAID EpiCoV group, WHO European Region sequencing laboratories and GISAID
642 EpiCoV group*. 2020. Geographical and temporal distribution of SARS-CoV-2 clades in
643 the WHO European Region, January to June 2020. *Euro Surveill* 25.
- 644 58. Wang R, Chen J, Gao K, Hozumi Y, Yin C, Wei G-W. 2021. Analysis of SARS-CoV-2
645 mutations in the United States suggests presence of four substrains and novel variants.
646 *Commun Biol* 4:228.
- 647 59. Gupta A, Sabarinathan R, Bala P, Donipadi V, Vashisht D, Katika MR, Kandakatla M,
648 Mitra D, Dalal A, Bashyam MD. 2020. Mutational landscape and dominant lineages in the
649 SARS-CoV-2 infections in the state of Telangana, India. *bioRxiv. medRxiv*.
- 650 60. Rambaut A, Holmes EC, O'Toole Á, Hill V, McCrone JT, Ruis C, du Plessis L, Pybus OG.
651 2020. A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic
652 epidemiology. *Nat Microbiol* 5:1403–1407.
- 653 61. Amoah ID, Kumari S, Bux F. 2020. Coronaviruses in wastewater processes: Source, fate
654 and potential risks. *Environ Int* 143:105962.

62. Liu T, Chen Z, Chen W, Chen X, Hosseini M, Yang Z, Li J, Ho D, Turay D, Gheorghe C, Jones W, Wang C. 2020. A benchmarking study of SARS-CoV-2 whole-genome sequencing protocols using COVID-19 patient samples. *bioRxiv*.
63. Steele JA, Zimmer-Faust AG, Griffith JF, Weisberg SB. 2021. Sources of variability in methods for processing, storing, and concentrating SARS-CoV-2 in influent from urban wastewater treatment plants. *bioRxiv*. *medRxiv*.
64. Lu X, Wang L, Sakthivel SK, Whitaker B, Murray J, Kamili S, Lynch B, Malapati L, Burke SA, Harcourt J, Tamin A, Thornburg NJ, Villanueva JM, Lindstrom S. 2020. US CDC Real-Time Reverse Transcription PCR Panel for Detection of Severe Acute Respiratory Syndrome Coronavirus 2. *Emerg Infect Dis* 26.
65. Decaro N, Elia G, Campolo M, Desario C, Mari V, Radogna A, Colaianni ML, Cirone F, Tempesta M, Buonavoglia C. 2008. Detection of bovine coronavirus using a TaqMan-based real-time RT-PCR assay. *J Virol Methods* 151:167–171.
66. Cao Y, Raith MR, Griffith JF. 2015. Droplet digital PCR for simultaneous quantification of general and human-associated fecal indicators for water quality assessment. *Water Res* 70:337–349.
67. Zeileis A, Grothendieck G. 2005. zoo: S3 Infrastructure for Regular and Irregular Time Series. *Journal of Statistical Software* 14:1–27.
68. Harrell FE. 2019. Hmisc: Harrell Miscellaneous.
69. Bushnell B. 2014. BBTools software package.
70. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359.
71. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The sequence alignment/map format and SAMtools. *Bioinformatics* 25:2078–2079.
72. Olm MR, Crits-Christoph A, Bouma-Gregson K, Firek BA, Morowitz MJ, Banfield JF. 2021. inStrain profiles population microdiversity from metagenomic data and sensitively detects shared microbial strains. *Nat Biotechnol* 39:727–736.
73. Pedersen BS, Quinlan AR. 2018. Mosdepth: quick coverage calculation for genomes and exomes. *Bioinformatics* 34:867–868.
74. Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* 26:841–842.
75. R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

- 689 76. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR,
690 O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H. 2017. vegan:
691 Community Ecology Package.
- 692 77. Mandal S, Van Treuren W, White RA, Eggesbø M, Knight R, Peddada SD. 2015. Analysis
693 of composition of microbiomes: a novel method for studying microbial composition.
694 Microb Ecol Health Dis 26.
- 695 78. Kuznetsova A, Brockhoff PB, Christensen RHB. 2017. {lmerTest} Package: Tests in Linear
696 Mixed Effects Models. J Stat Softw 82:1–26.
- 697 79. Wickham H. 2009. ggplot2: Elegant graphics for data analysis. Springer-Verlag New York.
- 698 80. Wilkins D. 2020. gggenes: Draw gene arrow maps in “ggplot2.”
- 699 81. Kolde R. 2019. pheatmap: Pretty Heatmaps.