

# **Longitudinal analysis of antibody responses to the Pfizer BNT162b2 vaccine in Patients Undergoing Maintenance Hemodialysis**

André Weigert, Marie-Louise Bergman, Ligia Gonçalves, Iolanda Godinho, Nádia Duarte, Rita Abrantes, Patrícia Borges, Ana Brennand, Vanessa Malheiro, Paula Matoso, Onome Akpogheneta, Lindsay Kosack, Pedro Cruz, Estela Nogueira, Magda Pereira, Ana Ferreira, Marco Marques, Telmo Nunes, João Viana, Jocelyne Demengeot, Carlos Penha-Gonçalves

## **Supplemental Material**

Table of contents:

**Supplemental Table 1** - Anti-Spike IgG, IgM and IgA positivity for data

presented in Figure 2

**Supplemental Table 2** - Median and Inter quartile range [IQR] for semi-quantitative IgG, IgM and IgA data presented in Figure 2


**Supplemental Figure 1** - Correlation analysis of anti-Spike IgG at t2 with relevant pathology indicators.

**Supplemental Table 1.** Anti-Spike IgG, IgM and IgA positivity for data presented in Figure 2

		Positivity; n (%; 95% CI)			
	Isotype	Age	n	t1	t2
Patients	IgG	All	143	42 (29.4; 22.5-37.3)	130 (90.9; 85.1-94.6)
		27-70	66	22 (33.3; 23.2-45.3)	63 (95.5; 87.5-98.4)
		71-93	77	20 (26.0; 17.5-36.7)	67 (87.0; 77.7-92.8)
	IgM	27-93	143	17 (11.9; 7.6-18.2)	42 (29.4; 22.5-37.3)
	IgA	27-93	143	59 (41.3; 33.5-49.5)	120 (83.9; 77.0-89.0)
Controls	IgG	All	143	71 (49.7; 41.6-57.7)	136 (95.1; 90.2-97.6)
		30-70	66	48 (72.7; 61.0-82.0)	65 (98.5; 91.9-99.7)
		71-96	77	23 (29.9; 20.8-40.8)	71 (92.2; 84.0-96.4)
	IgM	30-96	143	22 (15.4; 10.4-22.2)	36 (25.2; 18.8-32.9)
	IgA	30-96	143	72 (50.4; 42.2-58.4)	113 (79.0; 71.6-84.9)

**Supplemental Table 2.** Median and Inter quartile range [IQR] for semi-quantitative IgG, IgM and IgA data presented in Figure 2

Median [IQR]						
	Isotype	Age	n	t0	t1	t2
Patients	IgG	All	143	0.22 [0.16- 0.32]	0.63 [0.32- 1.08]	2.05 [1.67- 2.23]
		27-70	66	0.22 [0.15- 0.32]	0.68 [0.45- 1.21]	2.11 [1.87- 2.28]
		71-93	77	0.22 [0.17- 0.30]	0.47 [0.28- 1.00]	1.87 [1.57- 2.18]
	IgM	27-93	143	0.33 [0.24- 0.56]	0.49 [0.32- 0.75]	0.66 [0.45- 1.07]
	IgA	27-93	143	0.53 [0.35- 0.73]	0.85 [0.63- 1.10]	1.22 [1.10- 1.63]
Controls	IgG	All	143	0.19 [0.15- 0.24]	0.96 [0.46- 1.39]	1.82 [1.70- 1.90]
		30-70	66	0.17 [0.14- 0.21]	1.27 [0.93- 1.49]	1.81 [1.72- 1.89]
		71-96	77	0.21 [0.16- 0.26]	0.52 [0.38- 1.21]	1.83 [1.66- 1.92]
	IgM	30-96	143	0.49 [0.34- 0.66]	0.49 [0.35- 0.75]	0.56 [0.35- 1.01]
	IgA	30-96	143	0.56 [0.38- 0.80]	1.00 [0.69- 1.16]	1.15 [1.04- 1.35]

**Supplementary Figure 1**

**Supplementary Figure 1:** Correlation analysis of anti-Spike IgG at t2 with relevant pathology indicators. Correlation with time in hemodialysis (A); with Hemoglobin levels (B); with 25 hydroxi-cholecalciferol (C); with Ferritin (D); with Epoetin/Kg (E) and Epoetin/U per week (F); with C-reactive protein (G) and normalized Protein Catabolic Rate (nPCR) (H).