Influence of Novel Coronavirus COVID – 19 and HIV: A Scoping Review of Hospital Course and Symptomatology

Mona Sheikh, MD*, Shavy Nagpal, MD**, Madiha Zaidi, MD. Rupalakshmi Vijayan, MD. Wanessa Matos, MD. Nguemadji Nagardig Ngaba, MD. Lordstrong Akano, MD. Samia Jahan, MD. Shazia Q. Shah, MD. Camille Celeste Go, MD., Sindhu Thevuthasan, George Michel MD

Conflicts of interest - None of the authors have reported any conflicts of interest.

Funding - There is no financial disclosure related to this study.

Acknowledgments - Krunal Pandav, MD; Sumedh Keul, statistician, Larkin Research Department, Miami, Florida

Keywords: HIV, COVID-19, Mortality, SARS-CoV-2, Coronavirus

** corresponding - Department of Surgery, The Research Institute of St. Joe's Hamilton, ON, Canada, shav.nagpal@gmail.com

*arztinmona@gmail.com – Department of Medicine, Larkin Community Hospital South Miami, Florida, US

Abbreviations: **HIV** – Human immunodeficiency virus , **ART** – Antiretroviral therapy, **ICU** – Intensive care unit, **PLWH** – People living with HIV, **+ssRNA** – positive single stranded RNA, **SOB** – shortness of breath, **HTN** -hypertension, **DM** - diabetes mellitus, **CKD** - chronic kidney disease, **COPD** - chronic obstructive pulmonary disease, **CVD** - cardiovascular disease, **HLD** - hyperlipidemia, **HBV** - hepatitis B virus, **HCV** - hepatitis C virus, **CAD** - coronary artery disease, **ESRD** - end stage renal disease, **NASH** - non-alcoholic steatohepatitis, **OSA** - obstructive sleep apnea, **CHF** - congestive heart failure, **TB** - tuberculosis.

Abstract:

Background: An outbreak of novel coronavirus (SARS- CoV-2) was observed on December 2019 in Wuhan, China which led to a global pandemic declared in March 2020. As a consequence, it imposed delirious consequences in patients with underlying co – morbid conditions that make them immunocompromised. The purpose of this paper is to provide an in - depth review of influence of COVID – 19 in patients with underlying HIV in terms of mortality and hospitalization.

Authors also aim to provide a thorough risk analysis of hospitalization, ICU admission and mortality of PLWH and COVID-19. The secondary objective was to analyze the CD4+ count variations and outcome of COVID - 19 and to correlate if ART provided a protective role. Authors also aim to provide an evaluation of typical clinical presentation of COVID-19 in PLWH. ART is found to show activity against SARS-CoV-2 in vitro, and there is some similarity in the structure of HIV-1 gp41 and S2 proteins of SARS-CoV since they both belong to +ssRNA type.

Methods: We conducted a literature review using search engines namely, Cochrane, PubMed and Google Scholar. The following keywords were targeted: “COVID-19,” “SARS-CoV-2,” and “HIV.” We included case reports, case series, and cohort (retrospective and prospective) studies. We excluded clinical trials and review articles. We came across 23 articles that met the inclusion criteria. PRISMA guidelines were followed for study acquisition (Fig. 9).

Results: From the 23 studies, we found a total of 651 PLWH with confirmed COVID-19 (549, 91, and 11 in cohorts, case series, and case reports, respectively). The overall risk of hospital admission from pooled data of the 23 reviewed articles was 69.13% (450/651), ICU admission was 12.90% (84/651) in total infected patients, and 18.67% (84/450) among hospitalized patients. The overall case fatality rate from the 23 reviewed articles was 11.21 (73/651). A weak positive correlation was found between CD4+ counts and hospital admissions in case series and case reports, while the weak negative correlation was found in cohorts. For mortality, there was a negative weak association in the cohorts and in case series, while a weak positive was seen in case reports (Fig. 7). We assessed the presenting symptoms of PLWH with COVID-19, and our review demonstrated this group does not greatly differ from the rest of the population, as their common presenting symptoms were cough, fever, and SOB.

Conclusion: Our results indicated that there was a high rate of hospitalization, ICU admission, and mortality among patients living with HIV and COVID-19. PLWH needs to be noted as a high-risk group for COVID-19 complications and severity. We recommend that PLWH be closely monitored by their physicians and strictly adhere to antiretroviral therapy and standard universal COVID-19 precautions.

Fig.7 Results: As scoping review of HIV and COVID-19

Introduction:

SARS-CoV-2 is a member of β -CoVs (Fig. 8) and is a global concern at present. What interests Authors were the overlapping similarities between the Human Immunodeficiency Virus (HIV) and SARS-COV-2 virus in terms of same RNA single-stranded family and protein structural motifs of HIV-1 gp41 and SARS-CoV-S2 [9]. Having said that, it was proposed in clinical trials to implement ART as a treatment modality for COVID – 19. However, it is not recommended by the Treatment Guidelines Panel for COVID-19 yet [10].

The prognosis of COVID-19 in common comorbidities such as HTN, DM and CVD is well known but its prognosis or outcome of patients living with HIV is not well understood so far (Ssentongo et al, 2020) [11]. This review article focuses on the people living with HIV which are confirmed COVID-19 positive cases.

The main focus of our review is to assess the risk of hospitalization, intensive care unit (ICU) admission, invasive mechanical ventilation (IMV) requirement and mortality in persons living with HIV (PLWH) who were COVID-19 positive. The secondary objective is to assess whether any correlation exists between CD4+ count and COVID-19 in PLWH with the third objective to evaluate the typical clinical presentation of COVID-19 in such patients.

Fig. 8 SARS-CoV-2 (Ref. 2- 10)

Methods:

Literature search

A literature search was conducted with PubMed, Cochrane, and Google scholars using the following keywords: “COVID-19”, “HIV,” “SARS-CoV-2”, and “AIDS” from January 2020 until August 31st, 2020. Only papers in English language were considered. PRISMA guidelines were followed (Fig. 9).

Inclusion & Exclusion criteria and study selection

The case reports, case series, retrospective cohort studies and prospective cohort studies published between January 1, 2020, and August 31, 2020 with data of PLWH and confirmed COVID-19 by RT-PCR were included in our review. Review articles, clinical trials, and preprint articles were excluded. Articles that did not have patient data and those limited to specific comorbidities and organ dysfunctions were also excluded to avoid selection bias.

We narrowed it down to 48 articles that were screened based on the relevancy of the titles and abstracts. We excluded 32 articles, of which 20 did not meet the inclusion criteria, 2 required access, and 3 were not peer-reviewed. We included 23 studies in this paper. Selected articles were independently reviewed by two authors. All disagreements were resolved with a discussion between the two authors or with input from a third independent reviewer and mutually agreed upon by the authors.

Data Extraction

The authors designed a predefined data extraction list to be used to extract the required data from the included studies. The following data were extracted: First author’s name, publication date, study design,

country, the total number of PLWH with confirmed COVID-19 by RT-PCR or antigen test, age, gender, comorbidities, ART, CD4+ cell counts, viral load, number of hospitalized patients (age, gender), duration of hospitalization, number admitted to ICU, a number that needed IMV and number of deaths (age, gender, comorbidity, CD4+ counts, ART list).

Data collection and Analysis

The data was collected on excel sheets and Google documents. We ran a qualitative review based on our primary and secondary outcomes. The studies used have been listed in Table 1-3. The data were tabulated using Microsoft Excel.

The Pearson correlation coefficient was used to assess the correlation between CD4+ counts and hospitalization and mortality, and the association of COVID-19 infection with hospital admission. A P-value <0.05 was considered significant and was calculated by using R factor, Z score and standard deviation.

Ethical approval and funding

We did not seek IRB approval, as our study is based on data from published articles and patients were not directly involved. No funding was obtained for this review.

Fig.9 PRISMA guidelines for study acquisition

Results:

This review included a total of 23 articles after excluding those that did not meet the inclusion criteria (20), not peer reviewed (3), and required access (2). The 23 articles were 7 cohort studies, 7 case series, and 9 case reports; (Fig. 10) and they were from different countries: USA (9), China (4), Spain (3), Italy (2), the UK (1), Germany (1), Turkey (1), India (1), Japan (1) and Uganda (1) (Fig. 11)

Our review was categorized into three groups based on the similarity of study type: which are cohort group (Ch), case series group (CS), and case report group (CR). The number of PLWH and confirmed COVID-19 were 549 from cohort studies (Ch), 91 from case series (CS), and 11 from case reports (CR). The mean age (in years) were 54 in Ch, 45 in CS, and 36 in CR. The gender distribution in the Ch was 440 male, 106 female, and 3 transgender; in CS was 62 male, 25 female, and 3 transgender; and in CR was 8 males and 3 females. [Tables 1-3]

Fig. 10 Types of included studies

Fig. 11 Geographical location of included studies

Risk of hospitalization of PLWH + confirmed COVID-19

The total number of hospitalized patients were 387 in Ch, 52 in CS, and 11 in CR. The risk of hospitalization in these 3 groups were 70.49% (387/549) in Ch, 57.14% (52/91) in CS, and 100% in CR. [Table 1-3] The overall risk of hospital admission from pooled data of the 23 reviewed articles was 69.13% (450/651).

Risk of ICU admission among hospitalized patients

The number of patients that needed critical care were 72 in Ch, 9 in CS, and 3 in CR. The calculated risk among hospitalized patients was 18.61% (72/387) in Ch, 17.31% (9/52), and 27.27% (3/11) in CR. [Table 1-3] The overall risk of ICU admission from pooled data of the 23 reviewed articles was 12.90% (84/651) in total infected patients and 18.67% (84/450) among hospitalized patients.

The risk of IMV requirement in hospitalized and ICU patients

Of those patients who were admitted to the ICU: 48 in Ch, 9 in CS, and 2 patients in CR needed invasive mechanical ventilation (IMV). The risk among hospitalized and ICU patients was (12.7% , 66.67%) in Ch, (13.46%, 77.78%) in CS, and (18.18%, 66.67%) in CR. [Table 1-3]

The mortality rate of PLWH + confirmed COVID-19

The total number of deaths in the three categories was 66 in Ch, 6 in CS, and 1 CR, respectively. The case fatality rate was 12.02% (66/549) in Ch, 6.59% (6/91) in CS, and 9.09% (1/11) in CR [Tables 1-3]. The overall case fatality rate from the 23 reviewed articles was 11.21 (73/651)

Correlation of CD4+ counts with hospitalization and mortality

The Pearson Correlation coefficient was used to assess the association of CD4+ counts with hospitalization and mortality in PLWH+COVID-19. A weak positive correlation was found between CD4+ counts and hospital admissions in case series ($r = 0.2622$, $p\text{-value} = 0.089403$) (Fig. 1) and case reports ($r = 0.2269$, $p\text{-value} = 0.5284$) (Fig. 2), while the weak negative correlation was found in cohorts ($r = -0.114$, $p\text{-value} = 0.043866$) (Fig. 3). For the mortality correlation, there was a negative weak association in the cohorts ($r = -0.1218$, $p\text{-value} = 0.031$) (Fig. 4) and in case series ($r = -0.011$, $p\text{-value} = 0.986$) (Fig. 5), while a weak positive was seen in case reports ($r = 0.0667$) (Fig. 6).

Figure 1. correlation of CD4+ counts & hospitalization in cohorts patients

Figure 2. Correlation of CD4+ counts % hospitalization in case series

Figure 3: Correlation of CD4+ counts & hospitalization in Case Reports

Figure 4. Correlation of CD4+ counts & Mortality in cohorts

Figure 5. Correlation of CD4+ counts & Mortality in Case series

Figure 6. Correlation of CD4+ counts & Mortality in case reports

Clinical presentations in PLWH + COVID-19

The clinical presentation from the pooled data of the 23 articles was available for 319 patients. The most common symptoms among these patients were cough (57.4%, no= 183), fever (55.5%, no= 177), SOB (21.9%, no= 70), myalgia (13.2%, no= 42), fatigue (11.9%, no= 38), and headache/malaise (11%, no= 35). And least common with less than 10% were dyspnea (9.7%, no= 31), gastrointestinal symptoms (nausea/vomiting/diarrhea) (6.6%, no= 21), sore throat (6.3%, no= 20), anosmia/ageusia (4.4 %, no= 14), and altered mental status (4.4%, no= 14) [Table 5].

Co-morbidities in COVID positive PLWH groups

The comorbidities from the 23 articles were reported for 417 out of 649 patients; of which the most common chronic medical conditions found were hypertension (37.4%, no=156), diabetes mellitus (21.8%, no= 91), chronic kidney disease (12.2%, no= 51), COPD/asthma/lung disease (12.2%, no= 51), cancer (8.9%, no= 37), obesity (6.7%, no= 28), cardiovascular disease (6.2%,no= 26), chronic liver disease (5.5%, no= 23), and hyperlipidemia (5.8%, no= 24) [Table 5]

Discussion:

We conducted a systematic quantitative review to evaluate certain aspects of COVID-19 in patients living with human immunodeficiency virus (PLWH). The outcomes were categorized into risk of hospitalization, ICU admission, IMV requirement and mortality. The review of 23 articles indicated that there was a consistent high risk of hospitalization, ICU admission, IMV requirement and mortality in PLWH + confirmed COVID-19-positive in the cohort studies, case series and case reports. The rate of hospitalization from the pooled data of the 3 groups (Ch, CS, Cr) was 69.13% (n=450); 70.49% in Ch [12-18], 57.14% in CS [19-25], and 100% in CR [26-34]. The rate of ICU admission was 12.9% (n=84) for the 3 groups; 18.61% in Ch [12-18], 17.31% in CS [19-25], and 27.27% in CR [26-34]. And the overall case fatality rate of all reviewed articles was 11.21% (n=73).

Our results indicate that PLWH were a high risk group with greater than 50% needed hospitalization while 1/5 of hospitalized patients needed critical care. This finding aligns with the argument that predicts a bad outcome of COVID-19 in PLWH due to immunocompromised status [14, 20] and does not support the argument predicting a better prognosis for PLWH. [15, 17, 18, 19] The better prognosis could be explained by a lower risk for cytokine storm as a result of immunodeficiency, as there are low CD4 T lymphocytes which are responsible for activating innate immune cells like B-lymphocytes, cytotoxic T cells, and non-immune cells. There is also some suppression of the immune reaction through subtype T helper cells. [35]

Correlation analysis of CD4+ counts with hospitalization and mortality were not consistent in all reviewed studies as it showed weak positive in case series and case reports while weak negative in cohorts [12, 14-18]. And in mortality, there was a weak negative correlation in cohorts and case series [19-25] while weak positive in case reports [26-34]. More studies needed to focus on this to determine if CD4+ counts strongly correlate with the outcome of COVID-19.

We found that PLWH and confirmed COVID-19 do not present differently than the rest of the population as the common clinical presentations are cough (28.2%), fever (27.3%), and shortness of breath (10.8%), which is consistent with the finding of cohorts [12, 14, 16-18] and case series [19-25].

Analyzing the available data for deceased patients, we found the age range to be between 47 and 82 with a male to female ratio of 3:1, this correlates with the general population as data reveal that males had a mortality rate more than females. [37] The comorbidity data of deaths were available for 26 deceased individuals: HTN (no=9), DM (no=6), COPD (no=3), CKD (no=8), ESRD (no=1), CVD/CHF (no=2), organ transplant (no=1), prior stroke (no=2), cancer (no=2), liver cirrhosis (no=1), and obesity (no=3). The CD4+ counts ranged from below 50 to 800. The average CD4+ counts for Ho et al. [14], Meyerowitz et al. [16], and Harter et al. [18] of the deaths the cohort studies were 686, 426, and 389, respectively, and mortality rates in these cohorts were 20.5%, 5.6%, and 9.1%, respectively.

Nearly 98% of the cohort and 95% of the case series patients were being treated with ART at the time of diagnosis. According to Del Amo et al. [13], HIV-positive patients who were receiving TDF/FTC had a lower risk of contracting COVID-19 and ultimately being hospitalized, concluding that further investigation is necessary. However, Kate Child et al [20] concluded that patients on suppressive ART were not protected from moderate or severe COVID-19. We could not find any correlation between a specific type of ART and the risk of hospitalization, ICU admission, or mortality.

Despite the fact that this review was done following a standardized methodology, it presents with specific limitations. Cohort studies that we reviewed, such as Del Amo et al. and Meyerowitz et al. did not report an HIV viral load which makes it difficult to assess in relation to the other cohorts. Del Amo et al. also did not disclose a CD4+ count. Among the case reports, the HIV viral load and the total number of patients on IMV were not reported by Maomao et al. Lastly, Byrd et al. did not present the total number of patients admitted in the ICU and the number of patients on IMV. We did not run a study assessment because these limitations create barriers which make it hard to assess and compare the studies with one another.

Conclusion:

In conclusion, our results indicate that PLWH with COVID-19 are at increased risk for hospitalization, ICU admission, IMV requirement and mortality. We recommend classifying this group as high risk and they should adhere strictly to COVID-19 precautions. Authors conclude that larger sample size and study

volume is needed to fully evaluate the outcome of this novel virus and confirm the association of CD4+cell counts with severe forms of the disease and mortality.

References:

1. Listings of WHO's response to COVID-19. Who.int. <https://www.who.int/news-room/detail/29-06-2020-covidtimeline>. Accessed June 20, 2020.
2. Zhu N, Zhang D, Wang W, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med*. 2020;382(8):727-733. doi:10.1056/NEJMoa2001017
3. Zhou P, Yang XL, Wang XG, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 2020;579(7798):270-273. doi:10.1038/s41586-020-2012-7
4. Kankyokansen.org.
http://www.kankyokansen.org/uploads/uploads/files/jsipc/protocol_V7.pdf. Accessed September 15, 2020
5. Carrillo-Larco RM, Altez-Fernandez C. Anosmia and dysgeusia in COVID-19: A systematic review. *Wellcome Open Res*. 2020;5:94. Published 2020 May 13. doi:10.12688/wellcomeopenres.15917.1
6. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China [published correction appears in *Lancet*. 2020 Jan 30;:]. *Lancet*. 2020;395(10223):497-506. doi:10.1016/S0140-6736(20)30183-5
7. Cheng Y, Luo R, Wang K, et al. Kidney disease is associated with in-hospital death of patients with COVID-19. *Kidney Int*. 2020;97(5):829-838. doi:10.1016/j.kint.2020.03.005
8. Lauer SA, Grantz KH, Bi Q, et al. The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application. *Ann Intern Med*. 2020;172(9):577-582. doi:10.7326/M20-0504
9. Wu Zhang X, Leng Yap Y. Structural similarity between HIV-1 gp41 and SARS-CoV S2 proteins suggests an analogous membrane fusion mechanism. *Theochem*. 2004;677(1):73-76. doi:10.1016/j.theochem.2004.02.018
10. Lopinavir/Ritonavir and Other HIV Protease Inhibitors | Coronavirus Disease COVID-19. Nih.gov. <https://www.covid19treatmentguidelines.nih.gov/antiviral-therapy/lopinavir-ritonavir-and-other-hiv-protease-inhibitors/>. Accessed July 17, 2020.
11. Ssentongo P, Ssentongo AE, Heilbrunn ES, Ba DM, Chinchilli VM. Association of cardiovascular disease and 10 other pre-existing comorbidities with COVID-19 mortality: A systematic review and meta-analysis. *PLoS One*. 2020 Aug 26;15(8):e0238215. doi: 10.1371/journal.pone.0238215. PMID: 32845926; PMCID: PMC7449476.
12. Vizcarra P, Pérez-Elías MJ, Quereda C, et al. Description of COVID-19 in HIV-infected individuals: a single-centre, prospective cohort. *Lancet HIV*. 2020;7(8):e554-e564. doi:10.1016/S2352-3018(20)30164-8
13. Del Amo J, Polo R, Moreno S, et al. Incidence and Severity of COVID-19 in HIV-Positive Persons Receiving Antiretroviral Therapy: A Cohort Study [published online ahead of print, 2020 Jun 26]. *Ann Intern Med*. 2020;M20-3689. doi:10.7326/M20-3689
14. Ho HE, Peluso MJ, Margus C, et al. Clinical outcomes and immunologic characteristics of Covid-19 in people with HIV [published online ahead of print, 2020 Jun 30]. *J Infect Dis*. 2020;jiaa380. doi:10.1093/infdis/jiaa380
15. Sigel K, Swartz T, Golden E, et al. Covid-19 and People with HIV Infection: Outcomes for Hospitalized Patients in New York City [published online ahead of print, 2020 Jun 28]. *Clin Infect Dis*. 2020;ciaa880. doi:10.1093/cid/ciaa880
16. Meyerowitz EA, Kim AY, Ard KL, et al. Disproportionate burden of coronavirus disease 2019 among racial minorities and those in congregate settings among a large cohort of people with HIV. *AIDS*. 2020;34(12):1781-1787. doi:10.1097/QAD.0000000000002607

17. Gervasoni C, Meraviglia P, Riva A, et al. Clinical features and outcomes of HIV patients with coronavirus disease 2019 [published online ahead of print, 2020 May 14]. *Clin Infect Dis*. 2020;ciaa579. doi:10.1093/cid/ciaa579
18. Härter G, Spinner CD, Roeder J, et al. COVID-19 in people living with human immunodeficiency virus: a case series of 33 patients [published online ahead of print, 2020 May 11]. *Infection*. 2020;1-6. doi:10.1007/s15010-020-01438-z
19. Blanco JL, Ambrosioni J, Garcia F, et al. COVID-19 in patients with HIV: clinical case series. *Lancet HIV*. 2020;7(5):e314-e316. doi:10.1016/S2352-3018(20)30111-9
20. Childs K, Post FA, Norcross C, et al. Hospitalized patients with COVID-19 and HIV: a case series [published online ahead of print, 2020 May 27]. *Clin Infect Dis*. 2020;ciaa657. doi:10.1093/cid/ciaa657
21. Ridgway JP, Farley B, Benoit JL, et al. A Case Series of Five People Living with HIV Hospitalized with COVID-19 in Chicago, Illinois. *AIDS Patient Care STDS*. 2020;34(8):331-335. doi:10.1089/apc.2020.0103
22. Byrd KM, Beckwith CG, Garland JM, et al. SARS-CoV-2 and HIV coinfection: clinical experience from Rhode Island, United States. *J Int AIDS Soc*. 2020;23(7):e25573. doi:10.1002/jia2.25573
23. Altuntas Aydin O, Kumbasar Karaosmanoglu H, Kart Yasar K. HIV/SARS-CoV-2 coinfecting patients in Istanbul, Turkey [published online ahead of print, 2020 Apr 29]. *J Med Virol*. 2020;10.1002/jmv.25955. doi:10.1002/jmv.25955
24. Marimuthu J, Kumar BS, Aravind Gandhi P. HIV and SARS CoV-2 co-infection: A retrospective, record based, case series from South India [published online ahead of print, 2020 Jul 7]. *J Med Virol*. 2020;10.1002/jmv.26271. doi:10.1002/jmv.26271
25. Calza L, Bon I, Tadolini M, et al. COVID-19 in patients with HIV-1 infection: a single-centre experience in northern Italy [published online ahead of print, 2020 Aug 3]. *Infection*. 2020;1-5. doi:10.1007/s15010-020-01492-7
26. Zhu F, Cao Y, Xu S, Zhou M. Co-infection of SARS-CoV-2 and HIV in a patient in Wuhan city, China. *J Med Virol*. 2020;92(6):529-530. doi:10.1002/jmv.25732
27. Zhao J, Liao X, Wang H, et al. Early virus clearance and delayed antibody response in a case of COVID-19 with a history of co-infection with HIV-1 and HCV [published online ahead of print, 2020 Apr 9]. *Clin Infect Dis*. 2020;ciaa408. doi:10.1093/cid/ciaa408
28. Wang M, Luo L, Bu H, Xia H. One case of coronavirus disease 2019 (COVID-19) in a patient co-infected by HIV with a low CD4+ T-cell count. *Int J Infect Dis*. 2020;96:148-150. doi:10.1016/j.ijid.2020.04.060
29. Mahmood K, Rashed ER, Oliveros E, et al. Predisposition or Protection?: COVID-19 in a Patient on LVAD Support With HIV/AIDS. *JACC Case Rep*. 2020;2(9):1337-1341. doi:10.1016/j.jaccas.2020.05.015
30. Toombs JM, Van den Abbeele K, Democratis J, Merricks R, Mandal AKJ, Missouriis CG. COVID-19 in three people living with HIV in the United Kingdom [published online ahead of print, 2020 Jun 15]. *J Med Virol*. 2020;10.1002/jmv.26178. doi:10.1002/jmv.26178
31. Baluku JB, Mwebaza S, Ingabire G, Nsereko C, Muwanga M. HIV and SARS-CoV-2 coinfection: A case report from Uganda [published online ahead of print, 2020 May 21]. *J Med Virol*. 2020;10.1002/jmv.26044. doi:10.1002/jmv.26044
32. Nakamoto T, Kutsuna S, Yanagawa Y, et al. A case of SARS-CoV-2 infection in an untreated HIV patient in Tokyo, Japan [published online ahead of print, 2020 Jun 3]. *J Med Virol*. 2020;10.1002/jmv.26102. doi:10.1002/jmv.26102
33. Haddad S, Tayyar R, Risch L, et al. Encephalopathy and seizure activity in a COVID-19 well controlled HIV patient. *IDCases*. 2020;21:e00814. Published 2020 May 16. doi:10.1016/j.idcr.2020.e00814

34. Menghua W, Xin Z, Jianwei L, Yu Z, Qinwei Y. Case report: one case of coronavirus disease 2019 (COVID-19) in a patient co-infected by HIV with a normal CD4+ T cell count. *AIDS Res Ther.* 2020;17(1):46. Published 2020 Jul 23. doi:10.1186/s12981-020-00301-3
35. Mascolo, S., Romanelli, A., Carleo, M.A. and Esposito, V. (2020), Could HIV infection alter the clinical course of SARS-CoV-2 infection? When less is better. *J Med Virol*, 92: 1777-1778. doi:[10.1002/jmv.25881](https://doi.org/10.1002/jmv.25881)
36. Luckheeram RV, Zhou R, Verma AD, Xia B. CD4⁺T cells: differentiation and functions. *Clin Dev Immunol.* 2012;2012:925135. doi:10.1155/2012/925135
37. Zheng Z, Peng F, Xu B, Zhao J, Liu H, Peng J, Li Q, Jiang C, Zhou Y, Liu S, Ye C, Zhang P, Xing Y, Guo H, Tang W. Risk factors of critical & mortal COVID-19 cases: A systematic literature review and meta-analysis. *J Infect.* 2020 Aug;81(2):e16-e25. doi: 10.1016/j.jinf.2020.04.021. Epub 2020 Apr 23. PMID: 32335169; PMCID: PMC7177098

Author/Study type	Location	Total no. infect. pts	Total no. hosp. pts	Rate of hosp.	Total no. ICU pt.	ICU rate (total pt., hosp.)	Total no. IMV pt.	Rate of IMV (total pt., hosp., ICU)
Viscarra et al/ Prospective	Spain	35	28	80%	6	21.4%, 17.1%	5	14.3%, 17.9%
Del Amo et al/ Prospective	Spain	236	151	64%	15	9.9%, 6.4%	NA	N/A
Ho et al/ Retrospective	USA	93	72	77.40%	19	20.4%, 26.4%	15	16.1%, 20.8%
Sigel et al / Retrospective	USA	88	88	100%	18	20.5%, 205%	18	20.5%, 20.5%
Meyerowitz et al / Retrospective	USA	36	21	58%	6	28.6%, 16.6%	4	11.1%, 19.1%
Gervasoni et al / Retrospective	Italy	28	13	50%	2	15.4%, 7.1%	2	4.3%, 15.4%

Harter et al / Retrospective	Germany	33	14	42%	6	42.9%, 18.2%	4	12.1%, 28.6%
Total		549	387	70.49%	72	13.11%, 18.61%	48	8.74%, 12.70%, 66.67%

Table 1. Outcome of PLWH with COVID-19: Cohort Studies

Author/Study type	Location	Total no. infect pts.	Total no. hosp. pts	Rate of hosp	Total no. ICU pts	ICU rate (total pt., hosp.)	Total no. IMV pts.	Rate of IMV (total pt., hosp., ICU)
Blanco et al / Case Series	Spain	5	5	100%	2	40%, 40%	1	20%, 20%
Childs et al / Case Series	USA	18	18	100%	5	27.7%, 27.7%	5	27.78%, 27.8%
Ridgeway et al / Case Series	USA	5	5	100%	1	20%, 20%	0	0
Byrd et al / Case Series	USA	27	9	33.30%	0	0	0	0
Altuntas Aydin et al / Case Series	Turkey	4	4	100%	1	25%, 25%	1	25%, 25%
Marimuthu et al / Case Series	India	6	6	100%	0	0	0	0

Calza et al / Case Series	Italy	26	5	19.20%	0	0	0	0
Total		91	52	57.14%	9	9.89%, 17.31%	7	7.69%, 13.46%, 77.78%

Table 2. Outcome of PLWH with COVID-19: Case Series

Author/Study type	Location	Total no. infected pts	Total no. hosp. pts	Rate of hosp.	Total no. ICU pt.	ICU rate (total pt., hosp.)	Total no. IMV pt.	Rate of IMV (total pt., hosp., ICU)
Zhu et al / Case Report	China	1	1	100%	0	0	0	0
Zhao et al / Case Report	USA	1	1	100%	0	0	0	0
Wang et al / Case Report	China	1	1	100%	1	100%, 100%	0	0
Kiran Rashed et al / Case Report	USA	1	1	100%	0	0	0	0
Toombs et al / Case Report	UK	3	3	100%	1	33.3%, 33.3%	1	33.3%, 33.3%, 100%
Baluku et al / Case Report	Uganda	1	1	100%	0	0	0	0
Nakamoto et al / Case Report	Japan	1	1	100%	0	0	0	0
Hadded et al / Case Report	USA	1	1	100%	1	100%, 100%	1	100%, 100%

Menghua et al / Case Report	China	1	1	100%	0	0	0	0
Total		11	11	100%	3	27.27% both	2	18.18%, 18.18%, 66.67%

Table 3. Outcome of PLWH with COVID-19: Case Reports

Table 4. Mortality data from all reviewed articles

Author (no. death)	Age (mean) Gender	Comorbidities	ART
Viscarr et al (no= 2)	N/A	N/A	N/A
Del Amo et al (no= 20)	N/A 16 m, 4 f	N/A	-10 pt. on TAF/FTC -8 pt. on ABC/3TC
Ho et al (no= 19)	62 (55-68) 13 m, 6 f	13 obesity	-10 pt. on TDF -3 pt. on PI
Sigel et al (no= 18)	62 (57-67) 13 m, 5 f	4 DM, 6 HTN, 2 COPD, 6 CKD, 3 organ transplant, 1 liver cirrhosis, 1 obesity	-13 pt. on Integrase -5 pt. on PI -16 pt. on NRT
Meyerowitz et al (no= 2)	59, 63 1 m, 1 f	1st pt.: CHF, CKD, DM, HLD, HTN, prior stroke, 40 BMI 2nd pt.: HTN, prior stroke, CKD	-1st pt.: CTG+ABC/ETG+DRT/r+ETR -2nd pt.: DTG+TAF/FTC
Gervasoni et al (no= 2)	47, N/A 2 m	-1st pt.: obesity -2nd pt.: CVD, lung cancer	N/A

Harter et al (no= 3)	59, 55, 82 3 m	-1 pt.: HTN, COPD, DM -other 2: N/A	-1st pt.: DOR/TDF/FTC -2nd pt.: BIC/TAF/FTC -3rd pt.: DRV/RTV/RGV
Byrd et al (no= 1)	59 m	ESRD, cancer	EFV, ABC

Author (no= T. PLWH + COVID)	Comorbidities	Clinical Presentation	Viral Load	CD4+ Counts	ART, Y or N/%
Viscarra et al (no= 35)	15 (42.9%) HTN, 10 (%) CVD, 6 (17.1%) DM, 3 (8.6%) CKD, 16 (45.7%) chronic liver disease, 8 (22.9%) chronic respiratory disease, 9 (25.7%) cancer	25 fever, 7 sore throat, 24 cough, 21 dyspnea, 4 anosmia/ageusia, 13 myalgia, 8 headache, 21 fatigue, 10 diarrhea, 3 N/V	34 pt.: <50	502 (295 – 771)	Y/100%
Del Amo et al (no= 236)	N/A	N/A	N/A	N/A	100%
Ho et al (no= 93)	49 (52.7%) HTN, 32 (34.4%) DM, 25 (26.8%) COPD/Asthma, 16 (17.2%) CKD, 8 (8.6%) cancer, 7 (7.5%) ESRD, 4 (4.3%) autoimmune disease	61 fever, 71 cough, 57 SOB, 10 altered mental status	57 pt.: <50	554 (339-752)	Y/95.7%
Sigel et al (no= 88)	33 (37.5%) HTN, 24 (27.3%) DM, 5 (5.7%) liver cirrhosis, 6 (6.8%) CAD, 19 (21.6%) CKD, 15 (17.1%) cancer, 4 (4.6%) organ transplant, 9 (10.2%) obesity, 48 (54.6%) smoking	N/A	-66 pt.: <50 -16 pt.: >50	<50 - >500	Y/100%
Meyerowitz et al (no= 36)	11 (30.6%) HTN, 8 (22.2%) DM, 4 (11.1%) CHF, 8 (22.2%) HLD, 6 (16.7%) CKD, 2 (5.6%) ESRD, 5 (13.9%) stroke, 3 (8.3%) COPD, 3 (8.3%) smoking, 1 (2.8%) NASH, 1 (2.8%) NAFLD, 1 (2.8%) asthma, 1 (2.8%) dementia, 2 (5.6%) lymphomas (in remission)	22 fever, 21 cough, 6 sore throat, 8 myalgia, 5 fatigue, 3 dyspnea, 13 SOB, 10 myalgia, 4 headache, 3 altered mental status, 4 anosmia/ageusia, 7 GI (diarrhea, appetite loss, N/V), 3 congestion	N/A	691 (16 – 1485)	Y/97.2%
Gervasoni et al (no= 28)	14 (50%) HTN, 3 (10.7%) DM, 15 (53.6%) HLD, 5 (17.9%) HCV/HBV co-infection, 4 (14.3%) renal disease, 2 (7.1%) epilepsy, 2 (7.1%) CVD, 3 (10.7%) neoplasm, 2 (7.1%) COPD, 1 (3.6%) organ transplant	41 fever, 23 cough, 10 dyspnea, 7 diarrhea, 4 myalgia, 3 headaches	<50	636 (346 – 926)	Y/100%
Harter et al (no= 33)	10 (30.3%) HTN, 4 (12.1%) DM, 6 (18.2%) COPD, 2 (6.1%) CVD, 2 (6.1%) renal disease, 5 (15.2%) HBV, 3 (9.1%) CVD, 1 (3%) HCV	25 cough, 22 fever, 7 myalgia/arthritis, 7 headache, 7 sore throat, 6 anosmia	-30 pt.: <50 -920 -824	670 (69-1715)	Y/100%
Blanco et al (no= 5)	1 (20%) hypothyroidism, 1 (20%) asthma, 3 none	5 fever, 5 cough, 4 malaise/headache, 1 dyspnea	4 pts <50; 1pt. 45,500	563.7	Y/80%
Childs et al (no= 18)	6 (33.3%) HTN, 4 (22.2%) DM, 5 (27.8%) CKD, 10 (55.6%) obesity	11 Fever, 13 cough, 12 SOB	<50	395	Y/100%
Ridgeway et al (no= 5)	2 (40%) HTN, 1 (20%) DM, 1 (20%) HLD, 3 (60%) obesity, 1 (20%) OSA, 1 (20%) TB, 1 (20%) Addison's dis, 1 CHF, 1 COPD, 1 PE	3 fever, 4 cough, 3 SOB; 3 diarrhea, 2 myalgia, 2 headache, 2 n/V, 1 abd pain, 1 chest pain, 1 dyspnea	20	331	Y/100%

Byrd et al (no= 27)	N/A	cough, dyspnea, lethargy, fever, headache, sore throats, chest pain, myalgia, anosmia	<200	87-1441	Y/n/a
Altuntas aydin et al (no= 4)	1 (25%) DM, 1 (25%) COPD, 1 (25%) HTN, 1 (25%) HBV, 1 (25%) bipolar disorder, 1 (25%) obesity (35.5 BMI)	Dyspnea, cough, fever, diarrhea	N/A	627	Y/100%
Marimuthu et al (no= 6)	2 (33.3%) HTN	5 fever, 2 cough, 1 sore throat	N/A	535	Y/83.3%
Calza et al (no= 26)	11 (42.3%) HTN, 4 (15.4%) DM, 4 (15.4%) obesity, 3 (11.5%) asthma / chronic respiratory disease, 3 (11.5%) hypothyroidisms	19 Fever, 14 cough, 11 fatigue, 11 myalgia, 3 dyspnea, 4 tachypnea	<50	350	Y/n/a
Zhu et al (no= 1)	DM2, smoker	Fever, cough, SOB	N/A	N/A	N
Zhao et al (no= 1)	None	Fever, myalgia	<500	216	Y/100%
Wang et al (no= 1)	None	Fever, cough, chest pain	N/A	34	N/A
Kiran Rashed et al (no= 1)	CAD, DM	Fever, cough, dyspnea, myalgia	Und.	266	Y/100%
Toombs et al (no= 3)	2 (66.7%) DM, 2 (66.7%) HTN, 2 (66.7%) smoker, 1 (33.3%) TB, 1 (33.3%) obesity, 1 (33.3%) Graves' disease	2 Fever, 3 cough, 2 dyspnea, 1 headache, 1 anorexia	-2 pt.: und. -1 pt.: 1,000,000	373	Y/100%
Baluku et al (no= 1)	N/A	Headache, chest pain, myalgia, anorexia	<1000	965	Y/100%
Nakamoto et al (no= 1)	HBV, syphilis	Cough	100	194	N/A
Hadded et al (no= 1)	None	Fever, cough, altered mental status, GI (vomiting. Abd. pain)	Und.	604	Y/100%
Menghua et al (no= 1)	h/o syphilis (cured)	Fatigue	Und.	224	Y/100%

Table 5. Patient Characteristics of all reviewed articles (comorbidity, CD4 counts, viral load)