Variations in Non-Pharmaceutical Interventions by State Correlate with COVID-19 Disease Outcomes

Annika J. Avery¹, Jiayi Wang², Xinyu Ma², Qingkai Pan², Elizabeth E. McGrady¹, Zongyuan Yuan², Yuqing Liang¹, Rebecca Nugent^{2*}, Seema S. Lakdawala^{1,3*}

1 – Department of Microbiology and Molecular Genetics, University of Pittsburgh, School of Medicine, Pittsburgh, PA 15219

2 – Department of Statistics and Data Science, Carnegie Mellon University, Pittsburgh PA, 15213

3 – Center for Vaccine Research, University of Pittsburgh, School of Medicine, Pittsburgh, PA 15219

*send correspondence: Lakdawala@pitt.edu and rnugent@stat.cmu.edu

Abstract:

The COVID-19 pandemic highlighted the lack of understanding around effective public health interventions to curtail the spread of an emerging respiratory virus. Here, we examined the public health approaches implemented by each state to limit the spread and burden of COVID-19. Our analysis revealed that stronger statewide interventions positively correlated with fewer COVID-19 deaths, but some neighboring states with distinct intervention strategies had similar SARS-CoV-2 case trajectories. Additionally, more than two weeks is needed to observe an impact on SARS-CoV-2 cases after an intervention is implemented. These data provide a critical framework to inform future interventions during emerging pandemics.

Introduction:

Starting in December of 2019, there were reports of a novel coronavirus (CoV) emerging in China; on March 11, 2020, the World Health Organization (WHO) declared COVID-19 a pandemic. In response, many countries around the world began implementing non-pharmaceutical interventions (NPI) and providing guidance to keep their citizens safe, with several countries (such as China, South Korea, Singapore, Japan and in Europe) doing so in late February and March of 2020 (1-3). In contrast, the United States federal government did not implement and oversee any federal NPIs to restrict transmission of COVID-19 during all of 2020. Instead, individual states carried out the responsibility of implementing restrictions and creating reopening plans with their respective state-level resources (4).

This necessary action from US state and local governments resulted in a variegated approach to managing an emerging pandemic and over 600,000 deaths in the US as of July 2021 (<https://coronavirus.jhu.edu/>). In the face of this unfathomable loss, there exists an opportunity to understand how specific NPIs may affect transmission of COVID-19 in the United States. We aimed to characterize the effectiveness of NPIs implemented at the state level at reducing COVID-19 transmission as well as compare the overall effectiveness of NPI implementation strategies across states.

To explore the dynamics of NPI implementation and changes in daily COVID-19 cases, we created an interactive dashboard (www.phightcovid.org) that displays state-specific time series with normalized daily COVID-19 cases to compare the potential impact of NPIs across states. This dashboard is periodically updated to incorporate more recent case counts and NPIs. Clustering the

estimated basis-spline models underlying the state time series revealed geographical groupings, indicating that similar SARS-CoV-2 case trajectories may be based on state location in addition to NPIs. Finally, we observed a statistically significant negative correlation between COVID-19 deaths and state NPI score, solidifying the importance of implementation of public health interventions.

Heterogeneity in State NPI restrictions

To capture the heterogeneity in protective measures implemented by each state, including the District of Columbia (DC), we collected state mandated COVID-19 NPI orders from each governor website for one year (March 11, 2020-March 31, 2021). NPI orders for the following five categories were included in this study: stay-at-home orders, non-essential business restrictions, indoor gathering limitations, restaurant/bar restrictions, and mask/face covering mandates. Daily cumulative NPI scores were calculated for each state based on the stringency of each intervention in these five categories (see methods for score rubric). Fig. 1 highlights the heterogeneity in the NPIs implemented by each state, at three distinct times during the pandemic. A time-lapse movie of the NPI map with daily COVID-19 cases is provided in movie S1. Interestingly, implementation strategies varied by state with some states directly placing orders on all residents, while others delegated the responsibility for ordering NPIs to county or city governments (fig. S1A). In some cases, states initially placed statewide orders on all residents, but as the pandemic progressed, implementation of interventions was moved to a county or city level. Fig. S1 provides a snapshot of states that implemented statewide, county-level, or city-level NPIs, note that the various levels of NPI implementation were not exclusive. In some states, statewide NPIs were implemented for all residents, but individual counties would extend or modify the NPIs (fig. S1). To explore the accessibility of NPI guidelines per state, we additionally measured the time it took for a new user to find NPI information for each state (fig. S1B). We observed that information for the majority of states could be found in under five minutes (fig. S1B).

All states, except for South Dakota where there weren't any mandatory statewide NPIs, implemented a different combination of interventions statewide during the pandemic (www.phightcovid.org). Initially, 43 states implemented stay-at-home orders, of which 93% were lifted by June 15, 2020. Restaurant/bar closures were implemented in 49 states (including DC)

between March 15 and April 3, 2020. These states, except Missouri, eased restaurant/bar restrictions for the first time between April 24 and June 15, 2020, with the largest proportion doing so between May 9 and May 19, 2020. Missouri did not ease this NPI, rather the closure was completely lifted on June 16, 2020. Between June 1, 2020, and January 2, 2021, 41 states and DC reissued restaurant/bar restrictions for some duration of time before again easing them. A similar scenario of alternating between easing and re-issuing mandates occurred with indoor gathering limitations and non-essential business restrictions. Temporal restrictions by state are available in an interactive format on www.phightcovid.org/graphs.

SARS-CoV-2 case trajectories cluster by geographical region

Fig 2: SARS-CoV-2 cases and non-pharmaceutical interventions are distinct between states. Time series graphs for Maryland (A) and Tennessee (B) from March 11, 2020, through March 31, 2021. Daily SARS-CoV-2 cases were normalized to state population represented per 500,000 people (black circles), and the seven-day rolling average is indicated by the grey line. State NPI score per day is indicated by the colored bar along the x-axis. The NPIs are labeled with the category abbreviation, and colored as magenta if restricting, or teal if easing. The duration of the stay-at-home order is indicated by the grey band; black vertical lines represent implementation of statewide mask mandate, labor-day, and each vaccine FDA approval dates.

To compare the implementation of various NPIs per state over the last year to the trajectory of COVID-19 cases, we created time series graphs, (including a seven-day rolling average) for each state as displayed in Fig. 2 and fig. S2. In Fig. 2, Maryland and Tennessee are highlighted as representative states that implemented strong interventions throughout the past year (Maryland) versus states that had implemented fewer interventions (Tennessee) (table S1). The implementation dates of the statewide government NPIs are indicated on the graphs. As NPIs were implemented and eased, the NPI score changed based on the heuristic NPI scoring rubric, indicated by the colored horizontal bar (table S2). While most states, including Maryland and Tennessee, implemented NPIs early in the pandemic, Tennessee was far less stringent with NPIs in the fall of 2020 when cases began to rise. At the beginning of October, Tennessee lifted all NPIs (NPI score = 0), and in December the state experienced record new daily SARS-CoV-2 cases during the fall/winter wave. Tennessee briefly reissued indoor gathering limitations of ten people from December 21, 2020 to January 19, 2021 to curtail the spread of COVID-19 (table S1). Many states, like Maryland, eased restrictions during the summer, but reissued multiple restrictions in the fall as cases began to increase. For example, Maryland reissued restrictions in November that limited indoor gatherings to 25 people as well as restrictions on restaurant/bar occupancy, alcohol curfews,

and occupancy limits at non-essential business (table S1). Like most states, Maryland and Tennessee experienced record new daily COVID-19 cases in winter months (December 2020 to January 2021), although the amplitude of the peaks differed. The COVID-19 peak of new daily COVID-19 cases was 271% greater in Tennessee compared to Maryland. Across all 50 states there was variation in SARS-CoV-2 case trajectories, but states with consistent NPI restrictions throughout the year had smaller fall/winter peaks (www.phightcovid.org/graphs). This observation would suggest states may have similar epidemic curves related to the strength of the NPI restrictions implemented in each state.

To identify similarities between state SARS-CoV-2 case curves, we estimated a basis spline (B-spline) model for every state (5). Each estimated B-spline is a weighted piecewise combination of 15 polynomials, connected at “knots”. Examples of the estimated B-splines and their SARS-CoV-2 case curves for three states (Texas, Georgia, and Maine) are included in fig. S3A-F; these panels show how the different polynomials are combined into the overall estimated spline. Estimated splines and weight coefficients are closely related in states with similar underlying case curves. We compared the estimated state splines by using the K-means algorithm to cluster similar sets of weight coefficients, identifying groups of states with similar SARS-CoV-2 case trajectories. Returning to our three example states, we see that the coefficients for two states in the same cluster (Texas, Georgia) are more similar than those in a different cluster (Maine) (fig. S3D&E compared to F).

Standard best-fit criteria for K-means and repeated simulation identified a set of seven clusters of states, each with a distinct case curve shape shared across each group. The most common clustering is depicted in (Fig. 3A). Interestingly, the clusters demonstrate that states grouped geographically even though no geographic or NPI information was included when identifying the clusters, indicating that neighboring states tended to have similar case curve trajectories even if their NPI interventions differ (Fig. 3A).

An assessment of state median NPI score from March 2020 to March 2021 is depicted in Fig. 3B; surprisingly, a similar clustering pattern was not observed between these two maps. For example, Minnesota and Wisconsin have distinct state median NPI scores but fall within the same

K-means clusters (Fig. 3A and B), suggesting that implementation of various NPIs may not solely dictate the trajectory of SARS-CoV-2 cases. Aside from geographical location, environmental conditions such as humidity and temperature are thought to contribute to the seasonality of influenza viruses (6). Recently, humidity and temperature have been implicated in SARS-CoV-2 spread (7). However, the identified clusters may not correlate directly to environmental conditions, rather it could be driven by interstate travel patterns, similarities in state demographics, or when specific interventions were implemented(8, 9). Mobility and social behavior have also been linked to transmission of COVID-19 and may also influence the spread between states(10-12). These potential confounders all require further investigation.

Impact of NPI on case change occurs at greater than 2 weeks

Figure 4: More than two-weeks is needed before a NPI has an impact on case change. Case change between the date a restrictive NPI intervention was issued two (A), four (B) and five (C) weeks later for Michigan. Interventions are represented by the points, with easing shown in red and restricting in blue. The circled areas show where the points will be located if there is a considerably greater (red), or less (blue) number of cases at 2, 4 and 5 weeks later. The line of identity, where $x=y$ is the diagonal dashed line in the center of the graph. (D) Overlapping time series graphs showing the number of days since the first time a restaurant/bar easing was issued after May 2020, represented by day 0 on the x-axis. NPIs that were eased are represented by circles. NPIs that were restricted are represented by triangles. The Fourth of July is marked with an X for reference.

Effective implementation of NPI requires an understanding of the impact of NPIs on the trajectory of SARS-CoV-2 transmission. During the COVID-19 pandemic, discussion around the impact of interventions focused on case changes less than two weeks after an intervention. This included both easing or a restricting of an NPI (13, 14). However, that may not be a realistic time lag between NPI and case change. To examine this relationship, we graphed the number of cases at the time of an intervention versus the number of cases some point in the future, either 2-, 4-, and 5-weeks after an intervention (Fig. 4). Michigan was chosen as a representative of states that implemented many NPI over the course of the last year. By examining the NPIs within Michigan, we found that cases 2-weeks after either an easing or restricting NPI were similar to cases when this NPI was implemented, as the spots are still along the line of identity ($x=y$). The farther away from the line of identity the greater the case change; we anticipate that as cases raise after lifting of restrictions the cases will increase and move toward the red circle. In contrast as cases reduce after the implementation of an NPI, cases will decrease and move toward the blue circle. Fig. 4B and 4C examine case change after NPIs at 4- and 5- weeks later. Cases 4 weeks after a NPI is eased have increased such that the red easing restrictions are in the red circle (Fig. 4B). In contrast, cases 4 weeks after a restrictive NPI is implemented have lower cases, but they continue to decrease 5 weeks after the restrictive intervention (Fig. 4C). The observations made in Michigan pose interesting implications regarding the duration of time an NPI should be left in place until it becomes effective or when NPIs should be implemented. This analysis did not distinguish between the type of NPI and impact on cases, which could impact the effect on cases and requires further exploration.

To assess the impact of a specific type of NPI, we overlaid the SARS-CoV-2 case trajectories from multiple states on the day a specific intervention was implemented. Using states from multiple different clusters as shown earlier (Fig. 3A), we graphed the lag in days from the first-time restaurant/bar restrictions were eased after May 2020. Two weeks after the first restaurant and bar easing in May all states were relatively similar in that all experienced little case change. However, between two and four weeks, and even more so between four and five weeks, cases increased. This analysis further supports the observation that a lag of more than 2-weeks is needed before a change in cases after an NPI is eased or implemented (Fig. 4D).

COVID-19 disease outcome correlates with NPI score

Figure 5: Higher state NPI scores are associated with lower COVID-19 cases and deaths. (A) Linear regression of state median score and SARS-CoV-2 case rate March 2020-April 2021. State cumulative cases are normalized to 100,000 by state population. Dashed lines show the 95% confidence interval. (B) Linear regression of state median score and SARS-CoV-2 mortality rate June 1, 2020- April 13, 2021. State mortality rate is normalized to 100,000 by state population. Dashed lines show the 95% confidence interval. (C) Scatter plot of all state median scores, US mortality rate and population density. Dashed line along the y-axis highlights median score of 2.5. States are colored as blue if their COVID-19 mortality rate was at or above the US COVID-19 mortality rate. State population density is the number of people per square mile and is represented by the circle size. States with a large circle size have a high population density. We grouped population density as follows, extra-small: 1-50 people/sq. mile; small: 50-100 people/sq. mile; medium: 100-200 people/sq. mile; large: 200-350 people/sq. mile; extra-large: 350+ people/sq. mile.

To determine whether the implementation of various NPIs and their stringency impacted COVID-19 outcomes in each state we compared the median state NPI score for each state to SARS-CoV-2 cases and COVID-19 deaths. A linear regression of cumulative SARS-CoV-2 cases normalized to state population and state median NPI score revealed a statistically significant negative relationship (Fig. 5A, $p < 0.0001$). This suggested that greater NPI restrictions led to lower cases, similar to what was reported by other recent modeling approaches (15, 16). Since case reporting has been found to vary by state (17, 18), we compared state median NPI score to cumulative COVID-19 deaths, normalized to state population, from June 2020 - March 31, 2021 to remove the potential confounder of variable case reporting (Fig. 5B). Linear regression indicated that median state scores were negatively correlated with cumulative COVID-19 deaths ($p = 0.02$), indicating that states with more interventions had reduced mortality rates in addition to fewer cases.

While the trendlines are clearly significant between NPI score and SARS-CoV-2 cases or COVID-19 deaths, there are a number of states that reside outside of the 95% confidence interval (Fig. 5A and B). In particular, we were surprised by the states that had high NPI scores, but a similar case or cumulative death value as states with much lower NPI scores, suggesting that there may be an optimal NPI restriction threshold. To define the ‘goldilocks’ or ‘just right’, of NPI combinations, we compared the states with cumulative cases below the national cumulative case or death average for common interventions, including mask orders, some level of restaurant/bar restrictions, and gathering limitations. Using our rubric, states with all three intervention combinations at some stringency level for a prolonged period of time during the pandemic would have a median state NPI score of ~2.0-3.0, and a greater likelihood of lower mortality and fewer cases than states without these interventions. For example, Maryland, shown in Fig. 2A, had a median score of 2.55 and had the 12th lowest mortality rate out of all states.

Overall, 45% of states had a median NPI score above 2.5 and of these only 8 out of 23 had a mortality rate above the national average. In contrast, in states that had a median score below 2.5, 18 out of 25 had a mortality rate higher than the US average mortality rate (Fig. 5C). It is interesting to note that in states with the highest population density, 14 of 16, had a median score above 2.5, indicating stronger restrictions. In contrast, only 5 out of 24 states with the smallest population densities had median NPI scores above 2.5. To compare how population density may contribute to COVID-19 mortality in the context of NPI orders, a regression between average cumulative deaths per state by population density was performed for states with median NPI. Surprisingly, we did not find a statistically significant relationship between population density and cumulative mortality rate from June 2020 - March 31, 2021, for state median NPI scores either above or below a median score of 2.5 (fig. S4).

Discussion

During the course of the COVID-19 pandemic a number of studies have attempted to examine the effectiveness of NPIs on SARS-CoV-2 cases around the world and in the US. These studies focused on NPIs implemented early in the pandemic (prior to June 2020) and found that NPIs reduced SARS-CoV-2 cases and mobility with both real and modeled data (1, 2, 10, 15, 18-21). Comparison of 190 countries from January 23, 2020 to April 13, 2020 found that countries with NPIs (specifically: face mask, quarantines, social distancing, and travel restrictions) may have had a lower R_t value, suggesting a decrease in secondary SARS-CoV-2 transmission events in these spaces (1). However, this broad view does not integrate social behavioral variations or climate impacts that could also influence SARS-CoV-2 transmission. In other studies focused on U.S. counties or states, it was found that NPIs reduced mobility of the population and increased the doubling time of SARS-CoV-2 infections (18, 19). These studies again examined NPIs implemented early in the pandemic and could not account for issues with SARS-CoV-2 testing or the multiple rounds of issuing and easing interventions throughout the course of the past year. In contrast, our study includes data from all 51 U.S. states (including DC) and encompasses the NPIs implemented from March 2020 to April of 2021, to incorporate multiple waves of SARS-CoV-2 infections in states.

Taken together our results indicate that state implemented NPIs can lead to less COVID-19 cases and mortality. While it was previously thought that more NPIs led to a better outcome of disease burden (15), we found that even a moderate amount of restrictions can have a substantial impact

on lowering COVID-19 transmission. In addition, we observed a clear geographical impact on SARS-CoV-2 trajectories, and it is likely that consideration of geographical neighbors should be considered when designing future pandemic NPI plans. Finally, additional analysis into the impact of interventions in each state is needed to account for the numerous confounders that limit this type of analysis. A more refined analysis examining the impact of interventions, when they were implemented, the state of the disease at time of implementation, the strength of interventions, and implementation of similar interventions in neighboring states, is needed.

Methods

Data Collection

COVID-19 case and death data were obtained from the COVID-19 Data Repository by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University. For JHU CSSE COVID-19 Data see <https://github.com/CSSEGISandData/COVID-19>. Using this data, we calculated the new daily cases/500,000 people for every state by normalizing a state's new confirmed cases to the state population then multiplying by 500,000.

NPI intervention data was collected from a variety of sources and evolved over time. Data was collected manually from state government and governor websites. Data obtained on these websites was found in press releases, FAQ pages, reading through executive orders, or in the most accessible format, through an informative interactive portal detailing current restrictions and any changes from past restrictions. Data obtained through various news sources or governor's twitter page, was validated through a search for an official state government or governor announcement or order.

Implementation of NPIs varied between statewide or county specific order, which will impact the ability for residents to obtain this information and their awareness of the NPI in their region. To address the accessibility of the NPI orders in place for each state, we measured the time it took to 1) find state issued COVID-19 restrictions and 2) determine if state government orders apply statewide or county-specific. Additionally, we identified if there were county government issued NPIs and the specific stringency of the orders. The search process was consistent for all 50 states and took place Feb 17, 2021. Starting at google.com the keywords "[state of interest] covid restrictions" was queried. The most relevant (state government page or governor page) first result was chosen. After entering the website, a process of looking for the above-mentioned questions was carried out. The time was recorded upon finding each of the questions of interest. Furthermore, if the site was exceptionally easy to navigate and the current restrictions in place were found in less than a minute without downloading and opening additional files, the state was noted as being outstanding in user accessibility (n=16, CA, CO, CT, DE, DC, ID, IL, MI, MN, NJ, NM, OH, OR, PA, RI, and WA). Although states without any NPI orders or no detail on the NPI orders state were also easy to navigate. In situations such as this the state was not marked as being outstanding in user accessibility due to their lack of information displayed at all (AR, FL, and SD). Details of the time, sources, and notes per state for this analysis is available as table S3.

Scoring System

In order to compare the stringency of interventions implemented in each state, we developed a detailed scoring system. The system scores on the following five categories: Stay at Home Orders, Non-Essential Business Restrictions, Indoor Gathering Bans, Restaurant/Bar Restrictions and Mask/Face Covering Mandates. The highest score possible for each of the five

categories is 1.00, with the collective highest score possible being 5.00. The scoring criteria are presented in table S2 and briefly described here:

- Stay at home orders had a score impact of 1.0 when issued and -1.0 when removed.
- Non-essential business sectors that we impacted the score on are as follows: retail (+/- 0.25), hair salons/barbershops (+/- 0.10), personal care services (+/- 0.10), gyms/fitness centers (+/- 0.2), indoor entertainment (+/- 0.2). Each of these business sectors could have between 0 and their max score depending on the level of the restrictions in place. For example, if hair salons were open but only at 50% occupancy, the score impact would be 0.05).
- Indoor gathering bans primarily applied to indoor private social gatherings, however, as states eased restrictions, indoor gathering sizes described were more closely associated with larger private events. See table S2, category 3 for the scores associated with gathering sizes.
- Restaurants and bars being completely closed resulted in a score impact of 1.0. The breakdown of the 1.0 is as follows: restaurant outdoor dining (+/- 0.1), restaurant indoor dining (+/- 0.45), bar outdoor service (+/- 0.1), bar indoor service (+/- 0.25), restaurant and/or bar having any distancing or sanitation practices (+/- 0.05) and restaurant and/or bar having a service or alcohol curfew (+/- 0.05). See table S2, category 4, for further details.
- Mask and facial covering restrictions had a score impact of 1.0 if they were required in both indoor and outdoor spaces and a score impact of 0.8 if they were only required indoors.

Scoring of states that adopted a state government issued county risk-level metric (CA, CO, HI, IL, IN, NM, ND, OR, UT, WA) was more complicated than just scoring for the state as by itself. In these states the NPIs scored on were whatever they were for the majority risk level out of all counties. For example, if Oregon had 16 counties at low, 6 at moderate and 2 at high risk levels, then the entire state would be considered as being at a low risk level. The low risk level NPIs would be recorded and the state as a whole scored accordingly.

Time Series

Time series graphs were generated for each state to visualize an individual state's precise interventions (either restrictive or easing), NPI score (calculated based on the system described above), new daily SARS-CoV-2 cases, and seven day rolling average line. Daily SARS-CoV-2 cases were normalized to state population (based on the U.S. census bureau 2017 population estimates) and presented per 500,000 people. Interactive time series graphs are displayed on www.phightcovid.org, where the user can hover over each intervention and see the description of the type of intervention. The code to generate each of the time series is available on the phightcovid Github repository (see code availability section).

Lags

Lag graphs were created for gaining a better initial understanding of the time needed between when an NPI is implemented and when cases begin to change as a result. Using the programming language R we plotted today's new daily cases per 500,000 people on the x-axis and the new daily cases per 500,000 people at a lag of 2-, 4, and 5- weeks later, on the y-axis. We then use larger circles to highlight where it was expected to see the points if there was a considerable case change. R code for the lag graphs can be found in our Github repository

Spline model and clustering: Basis spline models were generated with 15 knots per state. K-mean clustering of the 15 B-spline coefficients (adapted from Rory Michelen, <https://github.com/RoryMichelen/Medium-Articles/blob/master/>) was used to compare the SARS-CoV-2 case trajectories between states. Code for generation of splines and clustering can be found at the Github repository.

Code and data accessibility: The data analyzed in this manuscript, including the NPI scores per state, all code developed to generate time series graphs, lag graphs, and spline models as well as a collection of many pdfs and screenshots of the raw data sources organized by state are available on Github at https://github.com/Lakdawala-Lab/PHIGHTCOVID_StNPI_Publ2021. Case data for time series plots was obtained every other month from March 2020 to April of 2021 from the COVID-19 Data Repository by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University, available on Github at <https://github.com/CSSEGISandData/COVID-19> (17).

Acknowledgments: We thank Andrea J. French, Gabriella H. Padovani, Logan Hellinger, Roma Kerby and Shreya Jyotishi for their help with data collection during the summer of 2020 and Arhat Pradhan for technical assistance with creation of supplemental movies. Additionally, we thank Joe Schmid from Symphony RM, for discussion to automate data curation from governor's websites. Finally, all members of the Lakdawala lab for their critical review of this manuscript.

Funding statement: This work was funded with internal startup funds from the University of Pittsburgh.

References

1. Y. Bo *et al.*, Effectiveness of non-pharmaceutical interventions on COVID-19 transmission in 190 countries from 23 January to 13 April 2020. *Int J Infect Dis* **102**, 247-253 (2021).
2. B. J. Cowling *et al.*, Impact assessment of non-pharmaceutical interventions against coronavirus disease 2019 and influenza in Hong Kong: an observational study. *Lancet Public Health* **5**, e279-e288 (2020).
3. N. Imai *et al.*, Adoption and impact of non-pharmaceutical interventions for COVID-19. *Wellcome Open Res* **5**, 59 (2020).
4. C. Adolph, K. Amano, B. Bang-Jensen, N. Fullman, J. Wilkerson, Pandemic Politics: Timing State-Level Social Distancing Responses to COVID-19. *J Health Polit Policy Law* **46**, 211-233 (2021).
5. G. D. Knott, *Interpolating cubic splines*. Progress in computer science and applied logic (Birkhauser, Boston, 2000), pp. xii, 244 p.
6. J. Shaman, V. E. Pitzer, C. Viboud, B. T. Grenfell, M. Lipsitch, Absolute humidity and the seasonal onset of influenza in the continental United States. *PLoS Biol* **8**, e1000316 (2010).
7. K. Chennakesavulu, G. R. Reddy, The effect of latitude and PM2.5 on spreading of SARS-CoV-2 in tropical and temperate zone countries. *Environmental Pollution* **266**, 115176 (2020).
8. M. Chinazzi *et al.*, The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. *Science* **368**, 395-400 (2020).

9. W. K. Pan *et al.*, COVID-19: Effectiveness of Non-Pharmaceutical Interventions in the United States before Phased Removal of Social Distancing Protections Varies by Region. *medRxiv*, 2020.2008.2018.20177600 (2020).
10. R. Abouk, B. Heydari, The Immediate Effect of COVID-19 Policies on Social-Distancing Behavior in the United States. *Public Health Rep* **136**, 245-252 (2021).
11. B. M. Althouse *et al.*, The unintended consequences of inconsistent pandemic control policies. *medRxiv*, (2020).
12. G. A. Wellenius *et al.*, Impacts of social distancing policies on mobility and COVID-19 case growth in the US. *Nat Commun* **12**, 3118 (2021).
13. S. Lai *et al.*, Effect of non-pharmaceutical interventions to contain COVID-19 in China. *Nature* **585**, 410-413 (2020).
14. L. Pellis *et al.*, Challenges in control of COVID-19: short doubling time and long delay to effect of interventions. *Philos Trans R Soc Lond B Biol Sci* **376**, 20200264 (2021).
15. D. H. Morris, F. W. Rossine, J. B. Plotkin, S. A. Levin, Optimal, near-optimal, and robust epidemic control. *Communications Physics* **4**, 78 (2021).
16. P. D. Wibbens, W. W. Koo, A. M. McGahan, Which COVID policies are most effective? A Bayesian analysis of COVID-19 by jurisdiction. *PLoS One* **15**, e0244177 (2020).
17. E. Dong, H. Du, L. Gardner, An interactive web-based dashboard to track COVID-19 in real time. *The Lancet Infectious Diseases* **20**, 533-534 (2020).
18. E. R. White, L. Hebert-Dufresne, State-level variation of initial COVID-19 dynamics in the United States. *PLoS One* **15**, e0240648 (2020).
19. S. Singh, M. Shaikh, K. Hauck, M. Miraldo, Impacts of introducing and lifting nonpharmaceutical interventions on COVID-19 daily growth rate and compliance in the United States. *Proc Natl Acad Sci U S A* **118**, (2021).
20. J. M. Brauner *et al.*, Inferring the effectiveness of government interventions against COVID-19. *Science* **371**, (2021).
21. J. Dehning *et al.*, Inferring change points in the spread of COVID-19 reveals the effectiveness of interventions. *Science* **369**, (2020).

Table S1. List of NPIs from Maryland and Tennessee by Date

Maryland				Tennessee			
Date	NPI Type	Score	NPI Description	Date	NPI Type	Score	NPI Description
03/12/2020	GL-R	0.90	Restricting: Indoor Gathering Limit, 10 people	03/22/2020	GL-R	1.90	Restricting: Indoor Gathering Limit, 10 people
03/16/2020	B-R	2.40	Closing: Entertainment	03/22/2020	RB-R	1.90	Closing: Restaurants and Bars
03/16/2020	RB-R	2.40	Closing: Restaurants and Bars	04/01/2020	B-R	3.90	Closing: All Non-Essential Businesses
03/23/2020	B-R	2.90	Closing: All Non-Essential Businesses	04/27/2020	RB-E	3.50	Easing: Restaurant Indoor Service (50%)
05/15/2020	B-E	2.70	Easing: Non-Essential Businesses, Retail; Personal Care (50%)	04/29/2020	B-E	3.35	Easing: Non-Essential Businesses, Retail (50%)
05/29/2020	RB-E	2.60	Easing: Restaurant and Bar Outdoor Service Opens	05/01/2020	B-E	2.25	Easing: Non-Essential Businesses, Gyms (50%)
06/05/2020	B-E	2.55	Easing: Non-Essential Businesses, Hair Salons	05/06/2020	B-E	2.15	Easing: Non-Essential Businesses, Hair and Personal Care (50%)
06/12/2020	B-E	1.90	Easing: Non-Essential Businesses, Entertainment	05/22/2020	B-E	1.30	Easing: Non-Essential Businesses, Entertainment (100%)
06/12/2020	RB-E	1.90	Easing: Restaurant and Bar Indoor Service (50%)	05/22/2020	GL-E	1.30	Easing: Indoor Gathering Limit, 50 people
06/19/2020	B-E	1.80	Easing: Non-Essential Businesses, Gyms (50%)	05/22/2020	RB-E	1.30	Easing: Bar Indoor Service (50%)
09/04/2020	B-E	2.70	Easing: Non-Essential Businesses, Retail (75%); Entertainment	09/29/2020	B-E	0.00	Lifting: All Non-Essential Businesses Restrictions
09/04/2020	GL-E	1.80	Lifting: Indoor Gathering Limit, 10 People	09/29/2020	GL-E	0.00	Lifting: Indoor Gathering Limit, 50 People
09/21/2020	RB-E	1.80	Easing: Restaurant and Bar Indoor Service (75%)	09/29/2020	RB-E	0.00	Lifting: Restaurant and Bar Restrictions
11/10/2020	GL-R	2.55	Restricting: Indoor Gathering Limit, 25 people	12/21/2020	GL-R	0.90	Restricting: Indoor Gathering Limit, 10 people
11/11/2020	RB-R	2.55	Restricting: Restaurant Indoor Service (50%)	01/19/2021	GL-E	0.00	Lifting: Indoor Gathering Limit, 10 People
11/20/2020	B-R	2.65	Restricting: Non-Essential Businesses, Retail (50%)				
11/20/2020	RB-R	2.65	Restricting: Restaurants and Bars, Alcohol Curfew				
03/12/2021	B-E	2.00	Easing: Non-Essential Businesses, Retail; Hair, Personal Care; Gyms; Entertainment (100%)				
03/12/2021	RB-E	2.00	Easing: Restaurant and Bar Indoor and Outdoor Service (100%)				

Table S2: Heuristic NPI Scoring Rubric

Category	Condition	Score Given (+ for issuing, - for easing/lifting condition)	If Curbside Pickup (Retail Only)	If 25%	If 50%	If 75%	If 100%
Category 1: Stay at home order	Stay at home order implemented	1					
	Stay at home order lifted	-1					
Category 2: Non-essential business closures	Non-essential business closures	1					
	Hair Salon/Barbershop	0.1		0.025	0.05	0.075	
	Entertainment (movies, theaters, museums, etc)	0.2		0.05	0.1	0.15	
	Gyms, fitness centers	0.2		0.05	0.1	0.15	
	Personal care services	0.1		0.025	0.05	0.075	
	Retail	0.25		0.05	0.1	0.15	0.2
	Open Businesses with distancing and sanitation practices	0.15	0.05	0.1	0.15	0.2	
Category 3: Indoor gathering bans	Size = Own Household Members Only	1					
	Size ≤ 5	0.95					
	Size ≤ 10	0.9					
	Size ≤ 15	0.85					
	Size ≤ 20	0.8					
	Size ≤ 25	0.75					
	Size ≤ 45	0.55					
	Size ≤ 50	0.5					
	Size ≤ 75	0.4					
	Size ≤ 100	0.25					
	Size ≤ 250	0.1					
	Size ≤ 500	0.05					
	Any gathering size restriction	0.05					
Category 4: Restaurant and bar limitations/restrictions	Restaurant and bar closures	1					
	Restaurant: outdoor dining closed	0.1					
	Restaurant: outdoor dining open with restrictions	0.05					
	Restaurant: outdoor dining open at 100% capacity	0					
	Restaurant: indoor dining closed	0.45					
	Restaurant: indoor dining open with 0-49% capacity restrictions	0.2					
	Restaurant: indoor dining open with 50-99% capacity restrictions	0.1					
	Restaurant: indoor dining open at 100% capacity	0					
	Bar: outdoor service closed	0.1					
	Bar: outdoor service open with restrictions	0.05					
	Bar: outdoor service open at 100% capacity	0					
	Bar: indoor service closed	0.25					
	Bar: indoor service open with 0-49% capacity and seating only restrictions	0.15					
	Bar: indoor service open with 50-99% capacity restrictions	0.1					
	Bar: indoor service open with 50-99% capacity and seating only restrictions	0.1					
	Bar: indoor service open with 50-99% capacity restrictions	0.05					
	Bar: indoor service open at 100% capacity and seating only restrictions	0.05					
	Bar: indoor service open at 100% capacity	0					
	Restaurant and Bar: Any distancing and sanitation practices	0.05					
	Restaurant and/or Bar: Service or alcohol sale curfew	0.05					
Category 5: Mask Mandate	If masks/face coverings must be worn both indoors and outdoors:	1					
	If masks/face coverings must be worn indoors only:	0.8					
	If masks/face coverings must be worn indoors only and only if individuals are within 6ft of each other for 15 minutes or longer (only in Iowa as of Feb. 2021):	0.2

Fig S1: Implementation strategies and accessibility of NPI in each state (A) State governments issuing NPI on all residents either with entire state orders or county-based orders. Indicated with a star are state governments that adopted a county risk-level metric and regularly changed specific county NPIs depending on the current COVID-19 risk level determined by the state (green). State governments delegating NPI responsibility to counties (red). County governments either delegating NPI responsibilities, or city and health departments deciding to implement NPI on their own (brown and blue respectively). Given the complexity of differentiating between health department and county level implementation, we were unable to capture all the states that implemented NPI through local health departments (B) To quantify accessibility of state COVID-19 NPI information, we timed how long we had to spend searching for NPI information on each state COVID-19 website. Data is from Feb 2021. Some states may have changed the NPI implementation strategy and website for ease.

Fig S2: COVID-19 deaths and non-pharmaceutical interventions between states. Time series graphs for Maryland (A) and Tennessee (B) from March 11, 2020 through March 31, 2021. Daily COVID deaths were normalized to state population represented per 1,00,000 people (black circles) and the 7-day rolling average is indicated by the grey line. State NPI score per day is indicated by the colored bar along the x-axis. The NPIs are labeled with the category abbreviation, and colored as magenta if restricting, or teal if easing. Additionally, the duration of the stay-at-home order is highlighted by the grey band and black lines indicate implementation of statewide mask mandate, labor-day, and each vaccine FDA approval dates.

Fig S3: Generation of Basis Spline for Each State. Basis spline models were created for the SARS-CoV-2 cases in each state using R package XX, with 15 different knots. (A-C) represents the spline model (red) for three different states (Texas, Georgia, and Maine) on top of each daily case increase. (D-F) Breakdown of the 15 different polynomials that created the spline model and list of the specific coefficient for each spline model.

426

428

430

432

434

436

438