Title page

Title: Shocked, relaxed, tired: Mental health during the three waves of the Covid-19

Authors & affiliations:

Jan S. Novotný, PhD¹, Juan P. Gonzalez-Rivas, MD^{2,3}, Šárka Kunzová, MD, PhD², Mária Skladaná, MD^{2,4}, Anna Pospíšilová, BA², Anna Polcrová, MA^{2,5}, Maria Vassilaki, MD PhD⁶, Jose R. Medina-Inojosa, MD⁷, Francisco Lopez-Jimenez, MD⁷, Yonas E. Geda, MD⁸, Gorazd B. Stokin, MD, PhD^{1,9,10}

¹Translational Neuroscience and Aging Program, Centre for Translational Medicine, and

²Kardiovize study, International Clinical Research Centre, St. Anne's University Hospital, Brno, Czech Republic

³Department of Global Health and Population, Harvard TH Chan School of Public Health, Harvard University, Boston, MA, USA

⁴2nd Department of Internal Medicine, St. Anne's University Hospital in Brno and Faculty of Medicine, Masaryk University Brno, Czech Republic

⁵Research Centre for Toxic Compounds in the Environment (RECETOX), Masaryk University, Brno, Czech Republic

⁶Division of Epidemiology, Department of Quantitative Health Sciences, ⁷Division of Preventive Cardiology, Department of Cardiovascular Medicine and ⁹Translational Neuroscience and Aging Program, Mayo Clinic, Rochester, MN, USA

⁸Department of Neurology, Barrow Neurological Institute, Phoenix, AZ, USA

¹⁰Division of Neurology, University Medical Centre, Ljubljana, Slovenia

Corresponding author:

Gorazd B. Stokin, gbstokin@alumni.ucsd.edu, Translational Neuroscience and Aging Program, Centre for Translational Medicine, International Clinical Research Centre, St.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.
Anne's University Hospital, Pekarska 53, 656 91 Brno, Czech Republic

Abstract

Background: Several recent studies reported that the Covid-19 pandemic induces significant mental distress. However, it remains unknown whether mental distress persists or subsides in response to consecutive Covid-19 waves. Therefore, we here examine mental distress before and across the three Covid-19 waves.

Methods: We measured stress and depressive symptoms prior to and across the three Covid-19 waves in a well-established population-based sample. A total of 330 participants completed e-questionnaires prior to and during the Covid-19 waves. Friedman's repeated-measures and Wilcoxon rank-sum pairwise test with Bonferroni correction were used to investigate repeated-measure differences, while the Kruskal-Wallis test was used to assess the effect of behavioral and physiological parameters on mental health.

Results: We found significantly increased mental distress during each of the Covid-19 waves compared with the pre-Covid-19 period. Increased mental distress was more common in women and most pronounced during the 1st and in particular, the 3rd Covid-19 wave. The stress and depressive symptoms increased during the pandemic compared with the pre-Covid-19 period by a maximum of 1.58 and 3.63 times, respectively. Negative emotions and loneliness were identified as the most significant risk factors of Covid-19-induced mental distress, while resilience was found to be protective.

Conclusions: Mental distress surged during all the Covid-19 waves without showing extinction. This pattern is mechanistically reminiscent of the post-traumatic stress disorder. Our results suggest that the Covid-19 pandemic underlies the development of a novel chronic mental health disorder and do not exclude possible further deterioration of mental health in the case of future Covid-19 waves.

Keywords: Covid-19, longitudinal, stress levels, depressive symptoms, chronic mental health disorder, post-traumatic stress disorder

Introduction

The Covid-19 pandemic, which hit the world in early 2020, affected most people's lives in many areas ranging from health to their basic needs, including social interaction and free movement. Although the effects of epidemics on mental health were described previously¹⁻⁴, the magnitude of their impact on mental health only emerged during the Covid-19 pandemic⁵⁻⁹. This impact is documented in a string of recently reported studies¹⁰⁻¹⁴, which all demonstrate a significant surge in mental distress during the Covid-19 pandemic. These studies, however, often lack pre-pandemic data and do not address the effects of individual Covid-19 waves and, in general, the long-term consequences of the Covid-19 pandemic on mental health. It remains, therefore, unknown whether consecutive Covid-19 waves lead to eventual extinction¹⁵ of the Covid-19-induced mental distress similarly to other fear behaviours^{16,17}. As a result, it also remains unclear whether the observed mental distress is a temporary short-term response to the Covid-19-pandemic or a matter of a novel pandemic-driven chronic mental health disorder sustained by the consecutive Covid-19 waves. Considering the protracted and continuously evolving course of the Covid-19 pandemic, including the emergence of novel SARS-CoV-2 variants, knowledge of the dynamics of the Covid-19-induced mental distress is imperative also to inform about mental health during probable future Covid-19 waves. We here investigate longitudinal changes in mental health before and across the three waves of the Covid-19 pandemic in a well-characterized population-based sample.

Methods

Study design and study population

Kardiovize study is a prospective longitudinal epidemiological cohort investigating cardiovascular, neuropsychiatric, and other health-related topics in Central Europe. A summary of the baseline examination protocol and general characteristics of the population-based sample have been published previously¹⁸. The inclusion criteria for the Covid-19 add-on study were all the participants of the Kardiovize study with available data on stress and

depressive symptoms prior to the onset of the Covid-19 pandemic. Overview of the recruitment process for the Covid-19 add-on study has been published previously⁹.

Procedure

The Covid-19 add-on study was conducted across the three Covid-19 waves. The first wave took place from April 23 to May 27, 2020, during the first nationwide lockdown. The second wave took place from August 31 to October 23, 2020, following the restriction-free summer. The third wave took place from March 4 to April 7, 2021, and coincided with several months of severe restrictions. Total of 1823 Kardiovize study participants were invited to join the Covid-19 add-on study, of whom 715 participated in the first wave of the study. 330 out of 715 participants who originally enrolled into the Covid-19 add-on study (46%) completed e-questionnaire prior to and during all of the three Covid-19 waves. E-questionnaire was completed through an online survey module using validated RedCap software (Research Electronic Data Capture) tool¹⁹.

E-questionnaire

The e-questionnaire consisted of several sections that surveyed demographic characteristics, mental health (levels of stress and depressive symptoms), lifestyle changes due to the government's measures to contain the Covid-19 transmission and select other factors that may influence mental health. The primary outcomes were stress level and severity of depressive symptoms. Stress level was measured using the Percieved Stress Scale (PSS) ranging from 0-40 and categorized as "low" (score of 0-13), "moderate" (score of 14-26), and "high" (score of 27-40) stress²⁰. Severity of depressive symptoms was measured using the Patient Health Questionnaire (PHQ) ranging 0-6 and the presence of depressive symptoms defined as a score ≥ 3 ^{21,22}. A detailed description of the other tools was published previously⁹.

Statistical analysis

Descriptive statistics were conducted for the socio-demographic variables. Missing values amounting to 1.8% of the total sample were identified in the pre-Covid-19 stress (N = 6) and depressive symptoms (N = 6) scales. No missing value imputations were performed. The missing data were completely at random and non-overlapping in cases. The prevalence of moderate-to-high stress and the presence of depressive symptoms was estimated as the percentage of participants meeting the cut-off values of each test. Fold change in the prevalence of mental distress described how many times the presence of elevated stress and depressive symptoms increased/decreased between study timepoints (e.g., Wave 3 prevalence divided by pre-Covid prevalence). Due to violation of the normality, we used Friedman's repeated-measures test to evaluate overall changes in stress and depressive symptoms and the Wilcoxon rank-sum pairwise test with Bonferroni correction to identify differences between study timepoints. Kruskal-Wallis test with eta-square effect size was used to examine the impact of different variables on mental health. Data analysis and visualization were performed in the R (v.4.1.0) environment with rstatix, sjmisc, ggplot2, and pheatmap packages.

Results

The sample consisted of 330 participants who completed the e-questionnaire prior to and during all three Covid-19 waves; mean age($\pm$ SD) were 45.9 ($\pm$ 10.7) years and 149 (45.2%) were males. Participants were largely well-educated, considering the majority of them held university degrees (Table 1). The Covid-19 add-on study sample characteristics were similar to the baseline Kardiovize study population-based sample, maintaining its representativeness. We observed significantly increased mean stress levels ($\chi^2=33.622$, $P<0.001$) during the 1st and 3rd Covid-19 waves compared with the pre-Covid-19 period as well as during the 3rd compared with the 2nd Covid-19 wave (Figure 1, eTable 1 in Supplementary Appendix). Prevalence of moderate-to-high stress increased 1.58 times during the 3rd Covid-19 wave compared with the pre-Covid-19 period (eTable 2 in Supplementary Appendix). Mean stress

levels and the prevalence of moderate-to-high stress were on average 1.2 times and 1.48 times higher in women compared to men (eTables 1 and 2 in Supplementary Appendix).

Similarly, we observed significantly increased severity of depressive symptoms ($\chi^2=38.961$, $P<0.001$) between all three Covid-19 waves and the pre-Covid-19 period as well as between the 3rd compared with the 2nd and 1st Covid-19 waves (Figure 2, eTable 1 in Supplementary Appendix). The prevalence of significant depressive symptoms increased 3.63 times during the 3rd Covid-19 wave compared with the pre-Covid-19 period (eTable 2 in Supplementary Appendix). As with stress, mean severity of depressive symptoms and the prevalence of significant depressive symptoms were on average 1.28 times and 1.55 times higher in women compared to men (eTables 1 and 2 in Supplementary Appendix).

We next searched for lifestyle and psychosocial variables that show significant association with mental distress developed during the Covid-19 pandemic. We identified negative emotional perception of the Covid-19 illness and feelings of loneliness as the most significant risk factors for developing mental distress during the Covid-19 pandemic. In contrast, higher level of resilience was found to be protective of Covid-19-induced mental distress (eFigure 1 in Supplementary Appendix).

Discussion

We examined longitudinally changes in mental health in response to the Covid-19 pandemic by measuring stress levels and severity of depressive symptoms prior to and during all three of the Covid-19 waves in a representative population-based sample. Consistent with previous findings, we find significantly increased stress levels and severity of depressive symptoms during the Covid-19 pandemic compared with the pre-Covid-19 period^{8,10,11,23–25} and higher mean scores and prevalence of mental distress in response to Covid-19 in women compared with men^{26–30}.

We extend these findings by showing that the observed mental distress matched the severity of the measures undertaken during specific Covid-19 waves. More specifically, during the 1st Covid-19 wave, there were not many individuals afflicted by the SARS-CoV-2 infection in the

Czech population, and people focused largely on adapting to the severely restrictive measures of the complete nationwide lockdown imposed by the government. These restrictive measures led to significant surge in mental distress^{7,31–34}. The 2nd Covid-19 wave followed the summer vacation period with the population well acquainted with then already relaxed restrictive measures, which further increased mental distress. During the 3rd wave, however, there was a significant increase in SARS-CoV-2 infections, hospitalizations and deaths, reintroduction of severe restrictive measures, and prolonged uncertainty about “returning to the normal life” with increasing fatigue of the population, which translated into significantly increased mental distress^{10,35–38}.

Intriguingly, consecutive Covid-19 waves did not alleviate but rather sustained mental distress. This suggests that Covid-19-induced mental distress is not a temporary short-term condition but rather a yet to be defined chronic mental health disorder. Considering fear extinction plays an important role in the pathogenesis of anxiety and the post-traumatic stress disorder^{17,39,40}, the observed lack of decrease in mental distress following consecutive Covid-19 waves suggests that similar delayed fear extinction mechanisms may play a role also in the development of the Covid-19-induced mental distress.

We also screened for risk factors of Covid-19-induced mental distress across all the Covid-19 waves. The finding of negative emotional perception of the Covid-19 illness and feelings of loneliness as the major risk factor of Covid-19-induced mental distress is consistent with concerns about mental health, emotional experience, and in particular negative emotional perceptions of illness, as well as feelings of loneliness, both considered general risk factors of mental distress^{41–47}. Similarly, the finding of resilience protecting against Covid-19-induced mental distress is consistent with its very definition, the ability to overcome adversity without compromising its own mental and emotional stability^{23,48–52}.

Conclusions and recommendations

Our results demonstrate a significant increase in stress levels and severity of depressive symptoms in all three Covid-19 waves compared with the pre-Covid-19 period. This sustained

state of mental distress suggests that Covid-19-induced a novel yet to be characterized chronic mental health disorder. Lack of extinction of mental distress during the consecutive Covid-19 waves further suggests that the mechanisms underlying Covid-19-related chronic mental health disorder are similar to the ones involved in the development of the post-traumatic stress syndrome. In agreement with established mental health disorders, Covid-19-induced mental distress is related primarily to the negative emotional experience and the feeling of loneliness. Further research is needed to characterize and define Covid-19-related chronic mental health disorder and decipher mechanisms underlying its pathogenesis.

Importantly, the ongoing Covid-19 pandemic with the situation in many countries now similar to the 2nd Covid-19 wave described in this study (relaxation of restrictive measures), together with delta-variant-related resurgence of Covid-19 infections, indicate high probability of further increases in mental distress. Based on the observations provided in our study, we can be concerned that future increases in mental distress might cause even greater mental health issues. Several measures can be undertaken to prevent and improve Covid-19-induced mental distress. First, more intense and better organised approaches to mental health care must be incorporated into public health policies to protect mental health. Second, individuals at high risk of developing mental distress need to be identified in a timely manner and offered appropriate interventions (increased social contacts, programs to build resilience, cognitive-behavioural techniques, etc.)^{53–55}. And third, the development and use of online technologies and web-based prevention and intervention tools should be incorporated into the mental health care process^{56,57}.

Statements

Supported by the European Regional Development Fund - Project ENOCH 750 (No. CZ.02.1.01/0.0/0.0/16_019/0000868).

The research protocol was approved by St. Anne's University Hospital ethics committee and the Internal Review Board. All participants were acquainted with the research and provided written informed consent.

Maria Vassilaki has received research funding from Roche and Biogen in the past; she currently consults for Roche, receives research funding from NIH/NIA and EU/ St. Anne's University Hospital Brno (Czech Republic), and has equity ownership in Abbott Laboratories, Johnson and Johnson, Medtronic, and Amgen. In addition, she is currently an Associate Editor for the Journal of Alzheimer's Disease and a Guest Editor for the Frontiers Research Topic collection "Multimorbidity in the Context of Neurodegenerative Disorders" (participating journals: Frontiers in Neuroscience-Neurodegeneration and Frontiers in Aging Neuroscience). No other disclosures were reported.

The data used in this study are available on request immediately following the publication to anyone who submits the online request that will be approved by the St. Anne's University Hospital International Clinical Research Centre internal board. The researchers have to provide their research intentions and goals, and specify, and justify requested variables. The data will be provided for a limited and well-defined time via cloud service or e-mail in csv format. After defined period the data should be returned, and all other copies should be destroyed.

The authors contributed to this article as follows: GBS and JPGR conceived the idea of this study, SK, MS, APos, and APol contributed to the discussion and battery creation, JSN and Apos verified the data, JSN made the statistical analysis, JSN and GBS wrote the draft of the manuscript, MS, YEG, JRMI, FLJ, MV, GBS contributed to the discussion and reviewed.

We thank the participants of the Kardiovize study for their long-term participation in this Covid-19 add-on study.

References

- 1 Hawryluck L, Gold WL, Robinson S, Pogorski S, Galea S, Styra R. SARS Control and Psychological Effects of Quarantine, Toronto, Canada. *Emerg Infect Dis* 2004; **10**: 1206–12.
- 2 Jalloh MF, Li W, Bunnell RE, *et al.* Impact of Ebola experiences and risk perceptions on mental health in Sierra Leone, July 2015. *BMJ Glob Heal* 2018; **3**: e000471.
- 3 Maunder RG. Was SARS a mental health catastrophe? *Gen Hosp Psychiatry* 2009; **31**: 316–7.
- 4 Cénat JM, Felix N, Blais-Rochette C, *et al.* Prevalence of mental health problems in populations affected by the Ebola virus disease: A systematic review and meta-analysis. *Psychiatry Res* 2020; **289**: 113033.
- 5 Knolle F, Ronan L, Murray GK. The impact of the COVID-19 pandemic on mental health in the general population: a comparison between Germany and the UK. *BMC Psychol* 2021; **9**: 60.
- 6 Ren Y, Qian W, Li Z, *et al.* Public mental health under the long-term influence of COVID-19 in China: Geographical and temporal distribution. *J Affect Disord* 2020; **277**: 893–900.
- 7 McGinty EE, Presskreischer R, Han H, Barry CL. Psychological Distress and Loneliness Reported by US Adults in 2018 and April 2020. *JAMA* 2020; **324**: 93.
- 8 Pierce M, Hope H, Ford T, *et al.* Mental health before and during the COVID-19 pandemic: a longitudinal probability sample survey of the UK population. *The Lancet Psychiatry* 2020; published online July. DOI:10.1016/S2215-0366(20)30308-4.
- 9 Novotný JS, Gonzalez-Rivas JP, Kunzová Š, *et al.* Risk Factors Underlying COVID-19 Lockdown-Induced Mental Distress. *Front Psychiatry* 2020; **11**. DOI:10.3389/fpsy.2020.603014.
- 10 Hamama-Raz Y, Goodwin R, Leshem E, Ben-Ezra M. The toll of a second lockdown: A longitudinal study. *J Affect Disord* 2021; **294**: 60–2.
- 11 Ramiz L, Contrand B, Rojas Castro MY, *et al.* A longitudinal study of mental health

- before and during COVID-19 lockdown in the French population. *Global Health* 2021; **17**: 29.
- 12 Petersen MW, Dantoft TM, Jensen JS, *et al.* The impact of the Covid-19 pandemic on mental and physical health in Denmark – a longitudinal population-based study before and during the first wave. *BMC Public Health* 2021; **21**: 1418.
 - 13 Stroud I, Gutman LM. Longitudinal changes in the mental health of UK young male and female adults during the COVID-19 pandemic. *Psychiatry Res* 2021; **303**: 114074.
 - 14 Ribeiro F, Schröder VE, Krüger R, Leist AK. The evolution and social determinants of mental health during the first wave of the COVID-19 outbreak in Luxembourg. *Psychiatry Res* 2021; **303**: 114090.
 - 15 Phelps EA, Delgado MR, Nearing KI, LeDoux JE. Extinction Learning in Humans. *Neuron* 2004; **43**: 897–905.
 - 16 Ahrenholtz R, Hiser J, Ross MC, *et al.* Unique neurocircuitry activation profiles during fear conditioning and extinction among women with posttraumatic stress disorder. *J Psychiatr Res* 2021; **141**: 257–66.
 - 17 Linton SR, Levita L. Potentiated perceptual neural responses to learned threat during Pavlovian fear acquisition and extinction in adolescents. *Dev Sci* 2021; published online April 4. DOI:10.1111/desc.13107.
 - 18 Movsisyan NK, Vinciguerra M, Lopez-Jimenez F, *et al.* Kardiovize Brno 2030, a prospective cardiovascular health study in Central Europe: Methods, baseline findings and future directions. *Eur J Prev Cardiol* 2018; **25**: 54–64.
 - 19 Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009; **42**: 377–81.
 - 20 Cohen S, Williamson GM. Perceived stress in a probability sample of the United States. In: Spacapan S, Oskamp S, eds. *The Claremont Symposium on Applied Social Psychology. The social psychology of health.* Newbury Park, CA: Sage Publications,

- Inc., 1988: 31–67.
- 21 Löwe B, Wahl I, Rose M, *et al.* A 4-item measure of depression and anxiety: Validation and standardization of the Patient Health Questionnaire-4 (PHQ-4) in the general population. *J Affect Disord* 2010; **122**: 86–95.
 - 22 Kroenke K, Spitzer RL, Williams JBW. The PHQ-9. *J Gen Intern Med* 2001; **16**: 606–13.
 - 23 Olff M, Primasari I, Qing Y, *et al.* Mental health responses to COVID-19 around the world. *Eur J Psychotraumatol* 2021; **12**: 1929754.
 - 24 Wang Z, Luo S, Xu J, *et al.* Well-Being Reduces COVID-19 Anxiety: A Three-Wave Longitudinal Study in China. *J Happiness Stud* 2021; published online March 27. DOI:10.1007/s10902-021-00385-2.
 - 25 Daly M, Sutin AR, Robinson E. Longitudinal changes in mental health and the COVID-19 pandemic: evidence from the UK Household Longitudinal Study. *Psychol Med* 2020; : 1–10.
 - 26 Lahav Y. Psychological Distress Related to COVID-19 – The Contribution of Continuous Traumatic Stress. *J Affect Disord* 2020; published online Aug. DOI:10.1016/j.jad.2020.07.141.
 - 27 Xiong J, Lipsitz O, Nasri F, *et al.* Impact of COVID-19 pandemic on mental health in the general population: A systematic review. *J. Affect. Disord.* 2020; **277**: 55–64.
 - 28 Pieh C, Budimir S, Delgadillo J, Barkham M, Fontaine JRJ, Probst T. Mental Health During COVID-19 Lockdown in the United Kingdom. *Psychosom Med* 2021; **83**: 328–37.
 - 29 Albert P. Why is depression more prevalent in women? *J Psychiatry Neurosci* 2015; **40**: 219–21.
 - 30 Salk RH, Hyde JS, Abramson LY. Gender differences in depression in representative national samples: Meta-analyses of diagnoses and symptoms. *Psychol Bull* 2017; **143**: 783–822.
 - 31 Mertens G, Gerritsen L, Duijndam S, Salemink E, Engelhard IM. Fear of the

- coronavirus (COVID-19): Predictors in an online study conducted in March 2020. *J Anxiety Disord* 2020; **74**: 102258.
- 32 Satici B, Saricali M, Satici SA, Griffiths MD. Intolerance of Uncertainty and Mental Wellbeing: Serial Mediation by Rumination and Fear of COVID-19. *Int J Ment Health Addict* 2020; published online May 15. DOI:10.1007/s11469-020-00305-0.
 - 33 Marroquín B, Vine V, Morgan R. Mental health during the COVID-19 pandemic: Effects of stay-at-home policies, social distancing behavior, and social resources. *Psychiatry Res* 2020; **293**: 113419.
 - 34 Ladapo JA, Rothwell JT, Ramirez CM. Association of COVID-19 Risk Misperceptions with Household Isolation in the United States (Preprint). *JMIR Form Res* 2021; published online May 3. DOI:10.2196/30164.
 - 35 Pietrabissa G, Simpson SG. Psychological Consequences of Social Isolation During COVID-19 Outbreak. *Front Psychol* 2020; **11**. DOI:10.3389/fpsyg.2020.02201.
 - 36 Brooks SK, Webster RK, Smith LE, *et al*. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 2020; **395**: 912–20.
 - 37 Rettie H, Daniels J. Coping and tolerance of uncertainty: Predictors and mediators of mental health during the COVID-19 pandemic. *Am Psychol* 2021; **76**: 427–37.
 - 38 Moradian S, Bäuerle A, Schweda A, *et al*. Differences and similarities between the impact of the first and the second COVID-19-lockdown on mental health and safety behaviour in Germany. *J Public Health (Bangkok)* 2021; published online Feb 26. DOI:10.1093/pubmed/fdab037.
 - 39 Nees F, Pohlack ST, Grimm O, Winkelmann T, Zidda F, Flor H. White matter correlates of contextual pavlovian fear extinction and the role of anxiety in healthy humans. *Cortex* 2019; **121**: 179–88.
 - 40 Norrholm SD, Jovanovic T. Fear Processing, Psychophysiology, and PTSD. *Harv Rev Psychiatry* 2018; **26**: 129–41.
 - 41 Charles ST, Piazza JR, Mogle J, Sliwinski MJ, Almeida DM. The Wear and Tear of Daily Stressors on Mental Health. *Psychol Sci* 2013; **24**: 733–41.

- 42 Hankin BL. Personality and Depressive Symptoms: Stress Generation and Cognitive Vulnerabilities to Depression in a Prospective Daily Diary Study. *J Soc Clin Psychol* 2010; **29**: 369–401.
- 43 Costa EC V., Vale S, Sobral M, Graça Pereira M. Illness perceptions are the main predictors of depression and anxiety symptoms in patients with chronic pain. *Psychol Health Med* 2016; **21**: 483–95.
- 44 Muscat P, Weinman J, Farrugia E, Callus R, Chilcot J. Illness perceptions predict distress in patients with chronic kidney disease. *BMC Psychol* 2021; **9**: 75.
- 45 Skapinakis P, Bellos S, Oikonomou A, *et al*. Depression and Its Relationship with Coping Strategies and Illness Perceptions during the COVID-19 Lockdown in Greece: A Cross-Sectional Survey of the Population. *Depress Res Treat* 2020; **2020**: 1–11.
- 46 Killgore WDS, Cloonan SA, Taylor EC, Dailey NS. Loneliness: A signature mental health concern in the era of COVID-19. *Psychiatry Res* 2020; **290**: 113117.
- 47 Rosenberg M, Luetke M, Hensel D, Kianersi S, Herbenick D. Depression and loneliness during COVID-19 restrictions in the United States, and their associations with frequency of social and sexual connections. *medRxiv* 2020; : 2020.05.18.20101840.
- 48 Smith BW, Tooley EM, Christopher PJ, Kay VS. Resilience as the ability to bounce back from stress: A neglected personal resource? *J Posit Psychol* 2010; **5**: 166–76.
- 49 Petzold MB, Bendau A, Plag J, *et al*. Risk, resilience, psychological distress, and anxiety at the beginning of the COVID-19 pandemic in Germany. *Brain Behav* 2020; published online July 7. DOI:10.1002/brb3.1745.
- 50 Killgore WDS, Taylor EC, Cloonan SA, Dailey NS. Psychological resilience during the COVID-19 lockdown. *Psychiatry Res* 2020; **291**: 113216.
- 51 Peker A, Cengiz S. Covid-19 fear, happiness and stress in adults: the mediating role of psychological resilience and coping with stress. *Int J Psychiatry Clin Pract* 2021; : 1–9.
- 52 Friesen KA, Weiss JA, Howe SJ, Kerns CM, McMorris CA. Mental Health and

Resilient Coping in Caregivers of Autistic Individuals during the COVID-19 Pandemic: Findings from the Families Facing COVID Study. *J Autism Dev Disord* 2021; published online July 8. DOI:10.1007/s10803-021-05177-4.

- 53 Masi CM, Chen H-Y, Hawkley LC, Cacioppo JT. A Meta-Analysis of Interventions to Reduce Loneliness. *Personal Soc Psychol Rev* 2011; **15**: 219–66.
- 54 Bavel JJV, Baicker K, Boggio PS, *et al.* Using social and behavioural science to support COVID-19 pandemic response. *Nat. Hum. Behav.* 2020; **4**: 460–71.
- 55 Wu AW, Connors C, Everly GS. COVID-19: Peer Support and Crisis Communication Strategies to Promote Institutional Resilience. *Ann Intern Med* 2020; **172**: 822–3.
- 56 Renton T, Tang H, Ennis N, *et al.* Web-Based Intervention Programs for Depression: A Scoping Review and Evaluation. *J Med Internet Res* 2014; **16**: e209.
- 57 Fiol-DeRoque MA, Serrano-Ripoll MJ, Jiménez R, *et al.* A Mobile Phone–Based Intervention to Reduce Mental Health Problems in Health Care Workers During the COVID-19 Pandemic (PsyCovidApp): Randomized Controlled Trial. *JMIR mHealth uHealth* 2021; **9**: e27039.

	No (%)
N	330
Age[†]	45.88 ±10.72
Sex	
Males	149 (45.2)
Females	181 (54.8)
Education[‡]	
Without GCSE	37 (11.2)
With GCSE	114 (34.5)
University	179 (54.2)

Table 1. The basic characteristics of the Covid-19 population-based sample.

[†]Presented as mean ±SD.

[‡]GCSE - General Certificate of Secondary Education, University education includes higher vocational school, bachelor, master, and doctoral degrees.

Figure 1. Changes in stress levels during the Covid-19 pandemic. Boxplots show changes in stress level scores (median, IRQ) during the individual waves of the Covid-19 pandemic. Associated bar plots show presence of moderate-to-high stress (PSS score ≥ 14), filled part denotes high stress prevalence (PSS score ≥ 27). Dashed lines show changes in mean scores for males and females separately, diamonds indicate mean scores of the sample. Horizontal upper bars represent differences between individual timepoints with indicated levels of significance (* $P < .05$, *** $P < .001$).

Figure 2. Changes in the severity of depressive symptoms during the Covid-19 pandemic. Boxplots show changes in the depressive symptoms score (median, IQR) during the individual waves of the Covid-19 pandemic. Associated bar plots show presence of significant depressive symptoms (PHQ score ≥ 3). Dashed lines show changes in mean scores for males and females separately, diamonds indicate mean scores of the sample. Horizontal upper bars represent differences between individual timepoints with indicated levels of significance (* $P < .05$, ** $P < .01$, *** $P < .001$).