

Supplementary Appendix for „Shocked, relaxed, tired: Mental health during the three waves of the Covid-19“

**Table of contents**

eTable 1. Mean scores (SD) of stress and depression in study timepoints (N=330)? ..... 2

eTable 2. Prevalence of significant mental distress in study timepoints (% (no of cases)) ..... 2

eFigure 1. Effect of selected ..... 3

**eTable 1.** Mean scores (SD) of stress and depression in study timepoints (N=330)

	Sample	Males	Females	P-value <sup>†</sup>
<b>Stress<sup>‡</sup></b>				
Pre-Covid	11.7 ±6.0	10.7 ±5.1	12.5 ±6.5	.005
Wave 1	13.4 ±7.0	11.8 ±6.8	14.6 ±7.0	<.001
Wave 2	12.3 ±6.8	11.3 ±6.3	13.1 ±7.1	.015
Wave 3	13.9 ±7.4	12.3 ±7.0	15.1 ±7.5	<.001
<b>Depression<sup>§</sup></b>				
Pre-Covid	0.6 ±1.1	0.6 ±1.0	0.6 ±1.1	.76
Wave 1	1.1 ±1.3	0.9 ±1.4	1.3 ±1.3	.007
Wave 2	0.9 ±1.3	0.7 ±1.1	1.0 ±1.4	.06
Wave 3	1.5 ±1.5	1.3 ±1.3	1.6 ±1.6	.03

<sup>†</sup>Sex differences estimated using t-test.

<sup>‡</sup>Mean scores of PSS.


<sup>§</sup>Mean scores of PHQ.

**eTable 2.** Prevalence of significant mental distress in study timepoints (% (no of cases)) (N=330)

	Sample	Males	Females
<b>Moderate-to-high stress<sup>†</sup></b>			
Pre-Covid	33.0 (109)	24.2 (36)	40.3 (73)
Wave 1	45.2 (149)	35.6 (53)	53.0 (96)
Wave 2	40.9 (135)	33.6 (50)	47.0 (85)
Wave 3	52.1 (172)	43.6 (65)	59.1 (107)
<b>Significant depressive symptoms<sup>‡</sup></b>			
Pre-Covid	5.8 (19)	4.0 (6)	7.2 (13)
Wave 1	15.5 (51)	13.4 (20)	17.1 (31)
Wave 2	12.1 (40)	8.7 (13)	14.9 (27)
Wave 3	20.9 (69)	16.8 (25)	24.3 (44)

<sup>†</sup>Defined as PSS score of ≥ 14.

<sup>‡</sup>Defined as PHQ score ≥ 3.



**eFigure 1. Effect of selected factors on mental health in study timepoints.** Heatmaps show eta-squared effect sizes (with indicated levels of significance) of selected factors on stress (pane a) and depression (pane c) based on Kruskal-Wallis test. Line plots indicate trends in mean scores of stress (pane b) and depression (pane d) in individual levels of selected factors. Y-axis ranges of line plots are set to [8,28] for stress and [0,4] for depression. Factors levels are (from left): age [25-40, 41-55, 56-66 yrs], sex [males, females], education [Without GCSE, With GCSE, University], loneliness [absent, present], resilience [low, medium, high], cognitive and emotional Covid-19 illness perc. [1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> tertile], Undergone Covid-19 [no, yes], Impact on finances [none, moderate, extreme], changes in sleep, nutrition and physical activity [improved, no change, worsen], freq. of searching Covid-19-related info [max 1-2x/week, every 1-3 days, several times/day].