Adverse events following COVID-19 virus vaccination in Japanese young population: The first cross-sectional study conducted by a questionnaire survey after the first-time-injection

Marie Suehiro¹, Shinya Okubo², Kensuke Nakajima², Kosuke Kanda³, Masanobu Hayakawa³, Shigeru Oiso², Tsutomu Kabashima⁴, Hideaki Fujita⁵, Yukio Ando⁶, Takahiro Muro¹.

¹ Division of Medical Informatics, ²Division of Pharmaceutical Health Care and Science, ³Division of Clinical pharmaceuticals, ⁴Division of Pharmaceutical Technology, ⁵Section of Functional Morphology, ⁶Division of Amyloidosis Research, Department of Pharmacy, Faculty of Pharmaceutical Sciences, Nagasaki International University

Abstract

A cross-sectional study was conducted to clarify the adverse events of COVID-19 vaccines in Japanese young population.

The proportion of participants with adverse events at the vaccination site (immediately or within 30 minutes after vaccination) was 0.5%, and anaphylaxis occurred in one female student (0.03%).

We analyzed 1,877 data obtained from a questionnaire survey of 1,993 vaccinated individuals. Eighty-two percent of participants complained of local adverse events. Injection site pain was the most common local adverse event (71%). Systemic adverse events occurred in 48% of participants. The most common adverse event was myalgia (34%).

A multivariable logistic regression model was used to determine risk factors.

Local adverse events were associated with sex (female) and allergy history, with odds ratios (ORs) (95% confidence interval [CI]) of 2.15 (1.69-2.73) and 1.73 (1.10-2.74), respectively. Systemic adverse events were associated with sex (female), age (<20 years old), allergy history, and history of adverse events with previous medications, with ORs (95% CI) of 2.49 (2.03-3.06), 1.80 (1.44-2.24), 1.39 (1.03-1.89), and 1.53 (1.02-2.29).

The results of this study clarified for the first time that age less than 20 years is a risk factor for systemic adverse events from the COVID-19 Vaccine Moderna Intramuscular Injection. This information will give impacts on considering adverse events and its mechanisms in mRNA vaccination.

Introduction

The coronavirus disease 2019 (COVID-19) crisis has been spreading worldwide. The COVID-19 vaccine is expected to be an effective preventive measure.

In Japan, vaccinations for priority groups, such as medical staff and elderly people (older than 65 years old), were being conducted first. Vaccinations at workplaces began on June 21, 2021 in limited universities and companies, and our university employees and students were subject to vaccination.

Currently, in Japan, the COMIRNATY intramuscular injection is used for priority vaccinations, and COVID-19 Vaccine Moderna Intramuscular Injection is used for vaccination in workplaces.

The adverse events of the coronavirus vaccine in elderly people have been studied; however, those of the coronavirus vaccine in young people have not been studied in Japan.

The reported adverse events were following vaccination using the COMIRNATY intramuscular injection, and there have been no reports of adverse events following the COVID-19 Vaccine Moderna Intramuscular Injection in Japanese population ^{1,2)}.

In addition, racial differences in the adverse events following coronavirus vaccines have not been sufficiently studied.

Therefore, to clarify the adverse events following coronavirus vaccination in young population, we completed a questionnaire survey after the first dose for students, faculty, and staff who belong to an educational foundation, Kyushu Bunka Gakuen.

Study population and method

Design

This study used a cross-sectional design with a questionnaire survey.

Study population

The study population included people who had a first time vaccination using COVID-19 Vaccine Moderna Intramuscular Injection on workplace vaccination of an educational foundation, Kyushu Bunka Gakuen.

This workplace vaccination included students attending educational facilities run by the educational foundation Kyushu Bunka Gakuen, universities, junior colleges, training facilities for cooking, training facilities for dental hygienists, and high schools.

In addition, this workplace vaccination included those employed by local companies related to the educational foundation of Kyushu Bunka Gakuen.

Data collection

Information regarding adverse events at the vaccination site was obtained from medical records.

To obtain information on that adverse events occurred after leaving the site, we conducted a questionnaire survey on the website.

An explanation of this study was provided on the website, and answers to questions regarding consent to research were obtained one by one.

Following adverse events related to vaccination, we obtained the following in a questionnaire: type of adverse event, date of occurrence, period of occurrence, and medication/care.

In addition, the attributes (age, sex, allergy history, and history of adverse events with previous medication) of the respondents were also obtained. Fever was defined as an increase of more than 1° C above the usual body temperature.

Statistical analysis

Patients' demographics and answers to the questionnaire are summarized with frequencies and percentages for categorical data and median plus range for continuous data.

We compared the patients who experienced adverse events (AE) and those who did not experience adverse events (No-AE) using the Wilcoxon rank sum test for continuous variables and Fisher's exact test for dichotomous variables. Differences were considered significant at $p < 0.05$.

To estimate the risk factors of adverse events, we first evaluated the odds ratio (OR), 95% confidence intervals (95% CI), and p-values of each potential risk factor using unadjusted logistic regression models.

Second, we determined the risk factors using a multivariable logistic regression model.

Statistical significance was determined by 95% CIs not including 1.00 in logistic analyses.

Candidate predictors were selected according to a literature review and clinical expertise.

We selected four variables, including patient demographics (age, sex, allergy history, and history of adverse events for past medication). Age, a continuous variable, was categorized as minor (<20 years old) or adult.

All statistical analyses were performed using JMP Pro 16 (SAS Institute Inc., Cary, NC, USA).

Ethics statement

This study was approved by the Nagasaki International University Ethics Committee (Approval number 50, July 2, 2021).

Results

Adverse events at the vaccination site

In this workplace vaccination program, 3,998 people were vaccinated.

Adverse events occurring at the vaccination site (immediately or within 30 min after vaccination) are shown in Table 1.

The proportion of individuals with adverse events at the vaccination site was 0.5%, and anaphylaxis occurred in one female student who had anaphylaxis at influenza vaccination once (0.03%).

**Table 1. Adverse events occurring at the vaccination site
(immediately or within 30 min after vaccination)**

Adverse events	Number of people	(%)	sex	
			female	male
Vagal syncope	5	0.13	4	1
Hyperventilation syndrome	4	0.10	4	
Feeling sick	4	0.10	3	1
Suffocation	3	0.08	3	
Nausea	1	0.03	1	
Anaphylaxis	1	0.03	1	
Hyperventilation with involuntary movement	1	0.03	1	
Severe urticaria	1	0.03	1	
Total	20	0.5	18	2

Demographic characteristics of participants

Table 2 shows the demographic characteristics of participants in the questionnaire survey.

We obtained data from 1,993 subjects who responded to the questionnaire survey (response rate 49.8%). A total of 1,877 subjects who agreed to participate in the present study were enrolled in the analysis. Female participants accounted for 66% of the study population, with a median age of 22 years. Twenty-four percent of the study population was younger than 20 years old.

Table 2. Demographic Characteristics of participants

Characteristic	Number of people (N=1877)	(%) ^{a)}
Female	1242	66
Age, median (range), y	22	(18-70)
Age (minor*)	444	24
Allergy history	219	12
Adverse events history for past medication	118	6

a) Data are expressed as No.(%) unless otherwise indicated.

*minor : <20 years old

Adverse events after leaving the vaccination site

The incidence of each adverse event is presented in Table 3. Local and systemic adverse events were generally mild. Eighty-two percent of participants complained of local adverse events. Injection site pain was the most common local adverse event (71%). Systemic adverse events occurred in 48% of participants. The most common adverse event was myalgia (34%), followed by general fatigue (31%).

Table 3. The incidence of each adverse event

Adverse events	Number of people	(%)
Local adverse events		
Any	1533	82
Pain	1334	71
Swelling	757	40
thermal sense	641	34
Redness	476	25
Itching	370	20
Induration	312	17
Systemic adverse events		
Any	894	48
Myalgia	632	34
Fatigue	581	31
Fever	416	22
Headache	386	21
Joint pain	119	6
Chills	86	5

Figure 1 shows the time of onset of adverse events, Figure 2 disclosed the duration of adverse events, and the type of treatment for adverse events are demonstrated in Figure 3. Injection site pain occurred from day 0 to day 1 and continued for a couple of days. However, almost all the symptoms improved without any treatments. The percentage of participants with itching and local redness after day 7 were 27% and 19%, respectively.

The onset of myalgia was often on the day and the day after injection (98%), and symptoms lasted 2-3 days, which was a little longer than other systemic adverse reactions (73%, 31% for 3 days). Of those who developed symptoms, 75% did not undergo any treatments, and 17% were administered acetaminophen.

General fatigue often occurred the day of injection (94%) and continued for 2-3 days thereafter (66%).

Fig. 1 The time of onset of adverse events (days after vaccination)

Fig. 2 The duration of adverse events

Fig.3 The type of treatment for adverse events

Comparison of the AE and No-AE groups

The characteristics of the AE and No-AE groups are shown in Table 4. Regarding local adverse events, the number of female participants in the AE group was significantly larger than that in the No-AE group ($p < 0.001$). Similarly, the number of people with an allergy history of AE was greater than that of No-AE ($p = 0.003$). Regarding systemic adverse events, the number of female and minors in the AE group was significantly higher than the number in the No-AE group ($p < 0.001$). The number of people with an allergy history in the AE group was greater than that in the No-AE group ($p = 0.004$).

Table 4. The characteristics of the AE and No-AE groups^{a)}

Local adverse events

Characteristic	AE ^{b)} (N=1,533)	No-AE ^{b)} (N=344)	P-Value ^{c)}
sex (female)	1068 (70%)	174 (51%)	<0.001
age, median,(range),y	22 (18-70)	26 (18-64)	<0.001
age (minor) ^{d)}	374 (24%)	70 (20%)	0.139
allergy history	194 (17%)	25 (7%)	0.003
adverse events history for past medication	103 (7%)	15 (4%)	0.006

Systemic adverse events

Characteristic	AE ^{b)} (N=894)	No-AE ^{b)} (N=983)	P-Value ^{c)}
sex (female)	691 (77%)	551 (56%)	<0.001
age,median,(range),y	21 (18-70)	27 (18-64)	<0.001
age (minor) ^{d)}	267 (30%)	177 (18%)	<0.001
allergy history	125 (14%)	94 (10 %)	0.004
adverse events history for past medication	71 (8%)	47 (5%)	0.111

a) Data are expressed as No.(%) unless otherwise indicated.

b)Abbreviations:AE, the patients who experienced adverse events; No-AE, the patients who did not experience adverse events

c) Wilcoxon's rank sum test was performed for continuous variables, and Fisher's exact test was performed for categorical variables.

d) minor : <20 years old

Multivariate analysis of AE and No-AE groups

Results of the multivariable analysis of the AE and No-AE groups are shown in Table 5.

Local adverse events were associated with sex (female) and allergy history, with ORs (95% CI) of 2.15 (1.69-2.73) and 1.73 (1.10-2.74), respectively. Looking in detail at individual adverse events, in local adverse events, redness and thermal sense showed similar tendencies. Induration and itching were associated with age (<20 years old) (OR [95% CI]: 0.56 [0.40-0.78] and 0.62 [0.46-0.83], respectively), and injection site pain was associated with a history of adverse events with prior medication OR [95% CI]: 3.11 [1.12-8.65]).

Systemic adverse events were associated with sex (female), age (<20 years old), allergy history, and history of adverse events with past medications, with ORs (95% CI) of 2.49 (2.03-3.06), 1.80 (1.44-2.24), 1.39 (1.03-1.89), and 1.53 (1.02-2.29), respectively.

Table 5. Results of the multivariable analysis of the AE and No-AE groups

Local adverse events						
Adverse events	Characteristic		Unadjusted OR(95%CI)	P-Value	Adjusted OR(95%CI)	P-Value
Any	sex	male	1 [Reference]		1 [Reference]	
		female	2.24(1.77-2.85)	<0.001	2.15 (1.69-2.73)	<0.001
	age(a minor* or an adult.)	adult	1 [Reference]		1 [Reference]	
		minor	1.25(0.94-1.66)	0.130	1.15(0.86-1.54)	0.354
	allergy history	NO	1 [Reference]		1 [Reference]	
		Yes	1.85(1.20-2.85)	0.006	1.73 (1.10-2.74)	0.018
	adverse events history for past medication	NO	1 [Reference]		1 [Reference]	
		Yes	1.58(0.91-2.75)	0.106	1.27(0.72-2.26)	0.413
Redness	sex	male	1 [Reference]		1 [Reference]	
		female	2.17(1.68-2.81)	<0.001	2.18 (1.68-2.83)	<0.001
	age(a minor* or an adult.)	adult	1 [Reference]		1 [Reference]	
		minor	0.84(0.65-1.08)	0.178	0.77 (0.60-1.00)	0.054
	allergy history	NO	1 [Reference]		1 [Reference]	
		Yes	1.47(1.08-2.01)	0.016	1.40 (1.01-1.94)	0.042
	adverse events history for past medication	NO	1 [Reference]		1 [Reference]	
		Yes	1.32(0.87-2.00)	0.191	1.14(0.74-1.76)	0.545
Swelling	sex	male	1 [Reference]		1 [Reference]	
		female	2.06(1.65-2.58)	<0.001	2.00 (1.60-2.51)	<0.001
	age(a minor* or an adult.)	adult	1 [Reference]		1 [Reference]	
		minor	1.29(1.02-1.63)	0.036	1.21 (0.95-1.53)	0.125
	allergy history	NO	1 [Reference]		1 [Reference]	
		Yes	1.57(1.15-2.13)	0.004	1.48 (1.08-2.05)	0.016
	adverse events history for past medication	NO	1 [Reference]		1 [Reference]	
		Yes	1.33(0.89-1.99)	0.162	1.15 (0.76-1.75)	0.516
Induration	sex	male	1 [Reference]		1 [Reference]	
		female	0.90(0.69-1.18)	0.456	0.93 (0.71-1.21)	0.575
	age(a minor* or an adult.)	adult	1 [Reference]		1 [Reference]	
		minor	0.55(0.40-0.77)	<0.001	0.56 (0.40-0.78)	<0.001
	allergy history	NO	1 [Reference]		1 [Reference]	
		Yes	1.61(1.14-2.26)	0.007	1.56 (1.09-2.23)	0.015
	adverse events history for past medication	NO	1 [Reference]		1 [Reference]	
		Yes	1.50(0.95-2.36)	0.080	1.28 (0.80-2.06)	0.302
Pain	sex	male	1 [Reference]		1 [Reference]	
		female	1.60(1.17-2.18)	0.003	1.60 (1.17-2.20)	0.004
	age(a minor* or an adult.)	adult	1 [Reference]		1 [Reference]	
		minor	0.77(0.55-1.08)	0.127	0.73 (0.52-1.03)	0.071
	allergy history	NO	1 [Reference]		1 [Reference]	
		Yes	1.96(1.11-3.45)	0.020	1.78 (0.98-3.21)	0.058
	adverse events history for past medication	NO	1 [Reference]		1 [Reference]	
		Yes	3.73(1.36-10.25)	0.011	3.11 (1.12-8.65)	0.030
Thermal sense	sex	male	1 [Reference]		1 [Reference]	
		female	3.16(2.47-4.03)	<0.001	3.13 (2.44-4.01)	<0.001
	age(a minor* or an adult.)	adult	1 [Reference]		1 [Reference]	
		minor	1.08(0.85-1.37)	0.517	0.97 (0.76-1.24)	0.798
	allergy history	NO	1 [Reference]		1 [Reference]	
		Yes	1.60(1.18-2.16)	0.002	1.43 (1.04-1.97)	0.030
	adverse events history for past medication	NO	1 [Reference]		1 [Reference]	
		Yes	1.71(1.14-2.56)	0.009	1.50 (0.98-2.30)	0.061
Itching	sex	male	1 [Reference]		1 [Reference]	
		female	2.84(2.09-3.84)	<0.001	2.96 (2.18-4.02)	<0.001
	age(a minor* or an adult.)	adult	1 [Reference]		1 [Reference]	
		minor	0.70(0.52-0.93)	0.014	0.62 (0.46-0.83)	0.002
	allergy history	NO	1 [Reference]		1 [Reference]	
		Yes	0.98 (0.69-1.40)	0.916	0.91 (0.63-1.32)	0.625
	adverse events history for past medication	NO	1 [Reference]		1 [Reference]	
		Yes	1.18(0.75-1.85)	0.474	1.14(0.71-1.83)	0.587

Systemic adverse events

Adverse events	Characteristic		Unadjusted OR(95%CI)	P-Value	Adjusted OR(95%CI)	P-Value
Any	sex	male	1 [Reference]		1 [Reference]	
		female	2.67 (2.18-3.26)	<0.001	2.49 (2.03-3.06)	<0.001
	age(a minor* or an adult.)	adult	1 [Reference]		1 [Reference]	
		minor	1.91 (1.54-2.37)	<0.001	1.80 (1.44-2.24)	<0.001
	allergy history	NO	1 [Reference]		1 [Reference]	
		Yes	1.54 (1.16-2.04)	0.003	1.39 (1.03-1.89)	0.033
	adverse events history for past medication	NO	1 [Reference]		1 [Reference]	
		Yes	1.71(1.17-2.51)	0.005	1.53 (1.02-2.29)	0.039
Fever	sex	male	1 [Reference]		1 [Reference]	
		female	1.25(0.99-1.58)	0.065	1.22 (0.96-1.54)	0.106
	age(a minor* or an adult.)	adult	1 [Reference]		1 [Reference]	
		minor	1.37(1.07-1.75)	0.013	1.34 (1.05-1.72)	0.020
	allergy history	NO	1 [Reference]		1 [Reference]	
		Yes	0.87(0.61-1.23)	0.433	0.82 (0.57-1.17)	0.274
	adverse events history for past medication	NO	1 [Reference]		1 [Reference]	
		Yes	1.33(0.87-2.03)	0.182	1.41 (0.91-2.18)	0.123
Headache	sex	male	1 [Reference]		1 [Reference]	
		female	1.96(1.40-2.73)	<0.001	1.88 (1.35-2.63)	<0.001
	age(a minor* or an adult.)	adult	1 [Reference]		1 [Reference]	
		minor	1.57(1.18-2.10)	0.002	1.52 (1.13-2.03)	0.005
	allergy history	NO	1 [Reference]		1 [Reference]	
		Yes	1.35(0.92-1.96)	0.123	1.31 (0.88-1.94)	0.189
	adverse events history for past medication	NO	1 [Reference]		1 [Reference]	
		Yes	1.25(0.77-2.02)	0.376	1.20(0.72-2.01)	0.478
Fatigue	sex	male	1 [Reference]		1 [Reference]	
		female	1.64(1.19-2.25)	0.002	1.64(1.19-2.27)	0.002
	age(a minor* or an adult.)	adult	1 [Reference]		1 [Reference]	
		minor	0.97(0.72-1.30)	0.822	0.93 (0.69-1.26)	0.636
	allergy history	NO	1 [Reference]		1 [Reference]	
		Yes	1.47(0.97-2.23)	0.072	1.47 (0.95-2.28)	0.086
	adverse events history for past medication	NO	1 [Reference]		1 [Reference]	
		Yes	1.14(0.68-1.92)	0.612	1.01 (0.59-1.73)	0.972
Chills	sex	male	1 [Reference]		1 [Reference]	
		female	1.33(0.75-2.34)	0.328	1.32(0.74-2.34)	0.345
	age(a minor* or an adult.)	adult	1 [Reference]		1 [Reference]	
		minor	0.97(0.60-1.58)	0.908	0.97 (0.59-1.60)	0.912
	allergy history	NO	1 [Reference]		1 [Reference]	
		Yes	2.34(1.38-3.96)	0.002	2.17 (1.25-3.78)	0.006
	adverse events history for past medication	NO	1 [Reference]		1 [Reference]	
		Yes	1.84(0.93-3.65)	0.082	1.39 (0.67-2.88)	0.371
Myalgia	sex	male	1 [Reference]		1 [Reference]	
		female	1.46(0.49-0.96)	0.026	1.47(1.05-2.06)	0.023
	age(a minor* or an adult.)	adult	1 [Reference]		1 [Reference]	
		minor	1.01(0.74-1.39)	0.945	0.98 (0.71-1.35)	0.897
	allergy history	NO	1 [Reference]		1 [Reference]	
		Yes	0.77(0.52-1.15)	0.208	0.71 (0.47-1.08)	0.113
	adverse events history for past medication	NO	1 [Reference]		1 [Reference]	
		Yes	1.13(0.65-1.95)	0.668	1.30 (0.73-2.30)	0.377
Joint pain	sex	male	1 [Reference]		1 [Reference]	
		female	1.13(0.70-1.81)	0.613	1.14(0.71-1.84)	0.582
	age(a minor* or an adult.)	adult	1 [Reference]		1 [Reference]	
		minor	0.91(0.59-1.40)	0.667	0.91(0.59-1.40)	0.663
	allergy history	NO	1 [Reference]		1 [Reference]	
		Yes	1.28(0.76-2.15)	0.362	1.21 (0.70-2.08)	0.500
	adverse events history for past medication	NO	1 [Reference]		1 [Reference]	
		Yes	1.39(0.72-2.67)	0.328	1.29 (0.65-2.55)	0.470

*minor : < 20 years old

Abbreviations:OR, Odds Ratio; CI, Confidence Interval

Discussion

The results of this study clarified for the first time that minors aged <20 years old are at greater risk of systemic adverse events from the COVID-19 Vaccine Moderna Intramuscular Injection.

According to a report by the Japanese Self-Defense Forces, the incidences tended to be higher in younger people; the factors analysed in this study confirmed this statistically¹⁾.

Second, the results of this study showed that female sex is a major risk factor for many adverse events.

According to Ministry of Health, Labour, and Welfare, Japan and the CDC of the United States, women are more likely to have adverse events from vaccination with this vaccine, which was supported by the results of this study.

The incidence of adverse events in this study was higher than that reported in previous reports of adverse events from the COVID-19 Vaccine Moderna Intramuscular Injection in Japan and other countries^{1,3,4)}.

We considered that this reason is because this study group included more younger women than those in previous studies.

In reports performed in foreign countries, headache and fatigue were the most frequently reported compared to other systemic events³⁾. However, in this study, many cases of myalgia and fatigue were also observed. In this study, we found that female sex was a risk factor for myalgia. This study is characterized by a higher proportion of younger people and women than those of other reports, which was also reflected in the incidence of adverse events³⁾.

In this study, to obtain information of adverse events that occurred after leaving the site, we conducted a questionnaire survey on a website. Therefore, the existence of a reporting bias cannot be completely ruled out. This is one of the limitations of the present study.

Information on adverse events that did not require treatment was not available, except for voluntary reports. This is a general limitation in adverse event investigations.

The COVID-19 Vaccine Moderna Intramuscular Injection efficacy after second doses was 94.1% in preventing symptomatic, laboratory-confirmed COVID-19 among persons without evidence of previous SARS-CoV-2 infection^{3,5)}. This vaccine requires two times injections to be fully effective, with the second dose having a higher frequency of adverse events than the first time injection¹⁻⁴⁾. Although there are no detailed data on Japanese who received the COVID-19 Vaccine Moderna Intramuscular Injection, data on Japanese healthcare workers who received the COMIRNATY intramuscular injection show that female and young people are more likely to increase of adverse events at the time of the second dose²⁾. Similar adverse events are predicted with the COVID-19 Vaccine Moderna Intramuscular Injection. In addition, there is a high possibility that this vaccine will be administered in younger people.

Age is often a risk factor for adverse events and should be pursued with caution. However, the adverse events caused by vaccination with this vaccine are rarely serious, and the benefits of vaccination should be emphasized⁶⁾. For peace of mind at the time of vaccination, those with many risk factors or those who had adverse events at the time of the first inoculation should prepare antipyretic analgesics, such as acetaminophen, before the second inoculation in anticipation of any adverse events. We will also investigate, analyze, and report adverse events after the second vaccination. This information will give impacts on considering adverse events and its mechanisms in mRNA vaccination.

Conflict of Interest

The authors have no conflicts of interest to declare regarding this study.

Reference

- 1) Japan Ministry of Health, Labor and Welfare. web site
https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/vaccine_kenkoujoukyoutyousa.html,
(View Date: 2021/07/17)
- 2) Takehiro Izumo, Naoyuki Kuse, Nobuyasu Awano, Mari Tone, Keita Sakamoto, Kohei Takada, Yutaka Muto, Kazushi Fujimoto, Ayae Saiki, Yu Ito, Haruko Matsumoto, Minoru Inomata, Side effects and antibody titer transition of the BNT162b2 messenger ribonucleic acid coronavirus disease 2019 vaccine in Japan, Respiratory Investigation . 2021 Jun 19; 2212-5345(21)00111-8. doi: 10.1016/j.resinv.2021.06.003.
- 3) Lindsey R. Baden, Hana M. El Sahly, Brandon Essink, Karen Kotloff, Sharon Frey, Rick Novak, David Diemert, Stephen A. Spector, Nadine Rouphael, C.Buddy Creech, John McGettigan, Shishir Khetan, et al., for the COVE Study Group, Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine, N Engl J Med. 2021; 384(5): 403-416. doi:10.1056/NEJMoa2035389. Epub 2020 Dec 30.
- 4) Johanna Chapin-Bardales, Julianne Gee, Tanya Myers, Reactogenicity Following Receipt of mRNA-Based COVID-19 Vaccines, JAMA. 2021; 325(21): 2201-2202. doi:10.1001/jama.2021.5374
- 5) Sara E. Oliver, Julia W. Gargano, Mona Marin, Megan Wallace, Kathryn G. Curran, Mary

Chamberland, Nancy McClung, Doug Campos-Outcalt, Rebecca L. Morgan, Sarah Mbaeyi, José R. Romero, H. Keipp Talbot, Grace M. Lee, Beth P. Bell, Kathleen Dooling, The Advisory Committee on Immunization Practices' Interim Recommendation for Use of Moderna COVID-19 Vaccine — United States, December 2020, Morbidity and Mortality Weekly Report (MMWR), January 1, 2021; 69(5152): 1653-1656

6) CDC - Centers for Disease Control and Prevention web site
Clinical Considerations: Myocarditis and Pericarditis after Receipt of mRNA COVID-19 Vaccines Among Adolescents and Young Adults, <https://www.cdc.gov/vaccines/covid-19/clinical-considerations/myocarditis.html> (View Date: 2021/07/23)