

Online resource table S4: Conservation of mutation sites. Evidence was obtained from Clinvar.

Variant	Clinical significance	Evidence
c.325C>T (p.Arg109Trp)	Pathogenic	This sequence change replaces arginine with tryptophan at codon 109 of the <i>RYR1</i> protein (p.Arg109Trp). The arginine residue is highly conserved and there is a moderate physicochemical difference between arginine and tryptophan.
c.6487C>T (p.Arg2163Cys)	Pathogenic	The R2163C variant is a non-conservative amino acid substitution, which occurs at a position that is conserved across species.
c.7268T>A (p.Met2423Lys)	Pathogenic	In 3 sibs with minicore myopathy with external ophthalmoplegia (255320) originally reported by Swash and Schwartz (1981), Jungbluth et al. (2005) identified a 7268T-A transversion in exon 45 the <i>RYR1</i> gene, resulting in a met2423-to-lys substitution in a highly conserved region.
c.14111C>T (p.Thr4704Met)	Pathogenic	This sequence change replaces threonine with methionine at codon 4709 of the <i>RYR1</i> protein (p.Thr4709Met). The threonine residue is highly conserved and there is a moderate physicochemical difference between threonine and methionine.
c.742G>T (p.Gly248Trp)	Likely pathogenic	No evidence about the conservation of the mutation site.
c.1186G>T (p.Glu396X)	Likely pathogenic	No evidence about the conservation of the mutation site.
c.2654G>A (p.Arg885His)	Likely pathogenic	This sequence change replaces arginine with histidine at codon 885 of the <i>RYR1</i> protein (p.Arg885His). The arginine residue is highly conserved and there is a small physicochemical difference between arginine and histidine.
c.8554C>T (p.Arg2852X)	Likely pathogenic	No evidence about the conservation of the mutation site.

