

Self-Report, Hair Toxicology, and Acute Toxicology

All ABCD participants completed a substance use interview, including a 12-month Timeline Follow-Back if they reported any substance use. In addition, all participants were asked to give a sample of their hair for analysis. Ten percent of participants were randomly selected to undergo some or all of the following acute toxicology tests: breathalyzer test, urinary cotinine screen, and/or oral fluid testing. Finally, participants who reported any past year substance use were given all three acute toxicology assessments. Data reported in Table S2 show self-reported substance use, detailed hair toxicology results, and acute toxicology results in participants who had positive hair results.

Table S2. Positive other toxicology results and self-reported use endorsements for participants with confirmed positive hair results, by drug class.

Positive result per self-report or biospecimen:	Cocaine (n=13)	Opiates (n=2)	Amphetamines (n=63)	THCCOOH (n=26)	Natural Cannabinoids (n=21)	Alcohol (n=10)	Nicotine (n=25)
Self-Reported use:							
Alcohol sips	38%	50%	38%	46%	5%	60%	52%
Alcohol full drink	0%	0%	0	4%	5%	0%	0%
Nicotine puffs	15%	50%	5%	15%	19%	10%	24%
Nicotine more than puff	8%	0%	0	8%	10%	10%	12%
Cannabis puff	8%	50%	0	12%	10%	10%	8%
Cannabis more than puff	8%	50%	0	4%	15%	10%	4%
Other drugs	0%	0%	0%	0%	0%	0%	0%
Hair analyte confirmation ^b :							
Cocaine ^d	13/13						
Benzoyllecgonine	12/12						
Norcocaine	2/11						
Oxymorphone ^c		2/2					
Amphetamines ^e			52/62				
Methamphetamine			9/51				
THCCOOH				25/26	9/10		
THC				10/11	21/21		
CBN				3/6	5/7		

CBD				5/8	7/10		
THCV				0/3	0/4		
ETG						9/10	
Cotinine							18/25
3-hydroxycotinine							6/11
Norcotinine							0/0
Nornicotine							8/8
Draeger oral fluid test ^f :							
Amphetamine	1/1 ^a	0/0	33/33 ^a	0/3	2/3 ^a	0/0	2/4 ^a
NicAlert ^g	0/4	0/0	0/2	1/3	1/3	0/0	1/1

Notes: 11-nor-9-carboxy-tetrahydrocannabinol = THCCOOH, Δ9-tetrahydrocannabinol = THC, cannabidiol = CBD, cannabinol = CBN, Δ9-tetrahydrocannabivarin = THCv; ETG=ethyl glucuronide

^a Positive results are consistent with the youth's prescription medication

^b Variation in analyte n due to insufficient quantity of hair (e.g., norcocaine; THCv)

^c Other opioids tested for (fentanyl, codeine, morphine, 6-acetylmorphine, hydrocodone, hydromorphone, and oxycodone) were negative for all.

^d Other cocaine metabolites tested for (cocaethylene, methylenedioxycocaine, orthohydroxycocaine, and parahydroxycocaine) were all negative.

^e Other amphetamines tested for (MDA, MDEA, and MDMA) were all negative.

^f Seven drug classes (cocaine, opiates, cannabis, benzodiazepines, amphetamine, methamphetamine, and methadone) were tested as described in the primary manuscript. Only drug classes with a positive are displayed here.

^gNicAlert is a urinary cotinine screen. Positive results on NicAlert here indicate urinary cotinine results at a level which indicates recent, personal nicotine exposure.