The fomite contribution to the transmission of COVID-19 in the UK: an evolutionary population estimate

A. Meiksin

*School of Physics and Astronomy
University of Edinburgh
James Clerk Maxwell Building
Peter Guthrie Tait Road
Edinburgh
EH9 3FD
meiksin@ed.ac.uk*

Abstract

A SEIR model with an added fomite term is used to constrain the contribution of fomites to the spread of COVID-19 under the Spring 2020 lockdown in the UK. Assuming uniform priors on the reproduction number in lockdown and the fomite transmission rate, an upper limit is found on the fomite transmission rate of less than 1 contaminated object in 7 per day per infectious person (95% CL). Basing the prior on the reproduction rate during lockdown instead on the CoMix study results for the reduction in social contacts under lockdown, and assuming the reproduction number scales with the number of social contacts, provides a much more restrictive upper limit on the transmission rate by contaminated objects of fewer than 1 in 30 per day per infectious person (95% CL). Applied to postal deliveries and groceries, the upper limit on the fomite transmission rate corresponds to a probability below 1 in 70 (95% CL) that a contaminated object transmits the infection. Fewer than about half (95% CL) of the total number of deaths during the lockdown are found to arise from fomites, and most likely fewer than a quarter. These findings apply only to fomites with a transmission rate that is unaffected by a lockdown.

Keywords: SARS-CoV-2; infectious disease epidemiology; mathematical modelling

Preprint submitted to Elsevier

June 30, 2021

1. Introduction

COVID-19, a respiratory infection caused by the SARS-CoV-2 virus, is believed to be transmitted primarily through viral-loaded respiratory droplets and contact with an infectious person, along with a suspected contribution from fomites (WHO, 2020). Respiratory aerosols may also be a contributing factor (Zuo et al., 2020). Establishing the role of fomites and respiratory aerosols is difficult (Lewis, 2021; Morawska and Milton, 2020). Whilst viable amounts of virus survive under laboratory conditions on contaminated surfaces (van Doremalen et al., 2020), and articles in proximity to an infectious patient can show traces of the virus RNA (Jiang et al., 2020; Ong et al., 2020), viable viruses may not survive in a natural environment in sufficient concentration to transmit the infection. On the other hand, prolonged infectivity has been measured on organic surfaces such as skin (Harbourt et al., 2020) and in a protein-rich environment (Pastorino et al., 2020), although fomites are an unlikely source of contamination in the presence of a vigorous cleaning regimen (Colaneri et al., 2020). The demonstration of indirect means of transmission in a natural environment is hampered by the expected rarity of fomites (Meiksin, 2020), so that they are difficult to detect except where an outbreak has already been identified, in which case the direction of contamination is uncertain (WHO, 2020). Indirect evidence supports the possibility of fomite transmission. Comparison between groups with and without regular hand hygiene suggests good hand hygiene reduces the transmission (Al-Ansary et al., 2020; Lio et al., 2021).

In the first half of 2020, many countries instituted national lockdowns in an effort to contain the pandemic (Flaxman et al., 2020). The decline in COVID-19 deaths following the lockdowns provides evidence that forms of non-pharmaceutical intervention that interrupt social contact successfully suppress the spread of the virus. Since a moderately high proportion of the infectious population may be pre-symptomatic or asymptomatic (WHO, 2020), even under lockdown conditions they may unwittingly contribute to the spread of COVID-19 through the accidental contamination of objects within popular circulation, such as packaged food items or mail parcels. Transmission in homes through commonly touched objects, or even through respiratory aerosols built up in poorly ventilated rooms, may also be a contributory factor.

The possibility of constraining the amount of transmission by fomites from modelling the population evolution under lockdown is explored here. It

is shown that, whilst the inclusion of fomites modifies the evolution of the populations, the modification is too small to improve constraints on the rate of indirect transmission. Including the results of surveys of social contacts, however, allows more stringent constraints to be placed. On 23 March 2020, the UK Government imposed a partial lockdown, asking the public not to leave home except for essential work, outdoor exercise once per day and to buy essential items such as food and medicine. The day after the partial lockdown was imposed, the CoMix survey initiated polling of the UK adult population on their social contacts. Participants in a statistically representative sample were asked to self-report their behaviour over a 16-week period (Jarvis et al., 2020). The responses were compared to responses from a similar survey that had been conducted by the UK arm of the POLYMOD study 15 years earlier (Mossong et al., 2008). Assuming direct proportionality between the reproduction number and number of social contacts, as well as similarity in behaviour before lockdown and during the POLYMOD survey, the CoMix study inferred a reproduction number of 0.62 (95% CI: 0.37-0.89) in the UK during lockdown. Significantly, virtually all the social contacts reported were in the home, raising the question of how the epidemic, though mitigated, was able to survive well beyond the durations of the exposure and infectious periods when transmission within a household would have ceased. Evidently, transmission of the virus between households was still occurring throughout the lockdown period, possibly through non-adherence to the partial lockdown or through inter-site circulation of fomites.

In this paper, population dynamics are approximated by the SEIR model extended to include fomites to constrain the role of fomites in the epidemic in the UK. It is shown that the additional information provided by the CoMix study significantly reduces the upper limit on the fomite contribution that may be placed compared with a uniform prior on the reproduction number in lockdown, allowing for conservative modelling assumptions.

2. Material and methods

2.1. SEIR equations with a fomite term

The evolution of the pandemic is modelled using the standard set of SEIR differential equations, augmented by a fomite term (Meiksin, 2020). The equations follow the dynamics of four sub-populations: the fraction s of the population susceptible to infection, the fraction e exposed to infection, the fraction i of infectious individuals, and the fraction r of removed (recovered

or perished) individuals. It is assumed here that no removed individual becomes susceptible again. Sub-populations s and i are coupled through a term $R_t si/D_i$ where R_t , the (time-dependent) reproduction number, is the average number of people an infectious person infects. The infectious period is taken to last for a duration D_i . The duration of an exposed individual before becoming infectious is D_e .

A fomite term f represents the number of contaminated objects per capita. If C_f is the average number of potentially contaminatable articles a person comes into contact with per day, then $C_f i$ is the per capita number of objects contaminated per day. (The possibility of inter-article contamination is not included.) For simplicity, an article that comes into close proximity to an infectious carrier is considered contaminated, and the average effectiveness of the contaminated article to transmit the infection is quantified through the transmittivity T_f , representing the average number of members of the susceptible population a contaminated object infects (or, equivalently, the fraction of contaminated objects that transmit the infection to a susceptible person). The coupling term between the susceptible population and fomites is then $T_f s f/D_f$. This represents the transmission rate per capita to an average T_f members of the susceptible population per capita by a number f of contaminated objects per capita for the duration D_f viruses survive on a contaminated object. The susceptible, exposed and infectious fractions depend only on the product $N_f = C_f T_f$. The epidemic is initiated by the introduction of exposed and infectious carriers at the respective rates c_e and c_i per capita (of the initial population).

The model equations are

$$\begin{aligned}\frac{ds}{dt} &= -\left(\frac{R_t}{D_i}i + \frac{N_f}{D_f}f_*\right)s, \\ \frac{de}{dt} &= \left(\frac{R_t}{D_i}i + \frac{N_f}{D_f}f_*\right)s - \frac{e}{D_e} + c_e, \\ \frac{di}{dt} &= \frac{e}{D_e} - \frac{i}{D_i} + c_i, \\ \frac{df_*}{dt} &= i - \frac{f_*}{D_f}.\end{aligned}\tag{1}$$

Here, the variable $f_* = f/C_f$ has been introduced to explicitly show the dependence of the infectious population only on N_f . Initially, $R_t = R_0$,

where R_0 is the basic reproduction number when the epidemic starts. The solution to the fomite equation may be expressed in terms of the infectious population as

$$f_*(t) = f_*(t_0)e^{-(t-t_0)/D_f} + e^{-(t-t_0)/D_f} \int_{t_0}^t dt' i(t')e^{(t'-t_0)/D_f}, \quad (2)$$

where $f_*(t_0)$ is the value of f_* at an initial time t_0 . This form shows explicitly that the number of fomites arises from the cumulative contributions from the infectious population. In the limit $D_f \ll (t - t_0)$, the fomite term is directly proportional to the instantaneous infectious population fraction when $i(t)$ varies slowly over the time interval D_f , $f_*(t) \rightarrow D_f i(t)$. In this limit, substitution into Eqs. (1) shows that the fomites act simply to rescale the reproduction number to $R_t + N_f D_i$, so that direct and indirect transmission may not be distinguished through the evolution of the populations.

The daily death rate per capita depends on the susceptible population through

$$\frac{dn_d(t)}{dt} = p_d \left[\frac{R_t(t - t_{\text{lag}})}{D_i} i(t - t_{\text{lag}}) + \frac{N_f}{D_f} f_*(t - t_{\text{lag}}) \right] s(t - t_{\text{lag}}), \quad (3)$$

where $n_d(t)$ is the total number of deaths per capita allowing for a lag time t_{lag} from exposure to death and p_d is the fraction of infected individuals who die.

Estimates for values of the SEIR parameter are taken from (Davies et al., 2020) for COVID-19 in the UK. The initial reproduction number without intervention is estimated at $R_0 = 2.68 \pm 0.57$. The average time from exposed to infectious state is taken to be $D_e = 4$ days, and the characteristic time during which an individual is infectious is taken to be $D_i = 5$ days. A mean infected fatality ratio 0.0050 is adopted (Meiksin, 2020; PHAS, 2020). Estimates for the mean lag time from onset of the infection to death range from 19 to 22 days (Davies et al., 2020; Flaxman et al., 2020). A time of 3 weeks is adopted here.

This paper concentrates on the lockdown period. Although R_t will not have changed to a new fixed value instantaneously after lockdown, for simplicity lockdown conditions are modelled by taking $R_t = R_0$ before the lockdown and $R_{t,\text{ld}}$ after. Estimates in the literature for both reproduction numbers generally do not distinguish direct transference of the virus from indirect transference through fomites.

Parameter	Description	Value	Reference
R_0	initial reproduction number	$2.1 < R_0 < 3.2$	(a)
R_{ld}	post-lockdown reproduction number	$0 \leq R_{t,ld} \leq 1$	(b)
T_f	fomite transmittivity	$0 \leq T_f \leq 0.3$	Assumed
D_e	duration of exposed period	4 days	(a)
D_i	duration of infectious period	5 days	(a)
D_f	duration of fomite infectious period	$0.2 \leq D_f \leq 4$ day	(c)
c_0	peak source rate per capita	10^{-6} day^{-1}	(d)
t_{c0}	time of source peak	day 77	(d)
FWHM	source distribution full width at half maximum	8 days	(d)
$\bar{f}_{ld}$	mean reduction factor f_{ld} in social contacts	0.27	(e)
$\sigma_{f_{ld}}$	standard deviation in f_{ld}	0.086	(e)

Table 1: Model parameters.

(a) Davies et al. (2020); (b) Russell et al. (2020); (c) van Doremalen et al. (2020); Pastorino et al. (2020); (d) du Plessis et al. (2020); (e) Jarvis et al. (2020)

The fomite contribution may in principle be identified through its effect on the evolution of the populations for finite fomite durations ($D_f > 0$). Laboratory estimates for the lifetime of fomites on various substances range
135 from under an hour on copper to several hours on plastic (van Doremalen et al., 2020). When embedded in protein-rich material designed to reproduce natural concentrations, the infectivity of the virus is prolonged to as long as 4 days and possibly longer (Pastorino et al., 2020). The virus was similarly
140 found to last at least 4 days on skin at room temperature, although its survival is severely shortened in a hot environment (Harbourt et al., 2020). Four representative duration times are considered here, 0.21 d and 0.41 d, typical of cardboard (0.14–0.30 d, 95% CI) and plastic (0.34–0.49 d, 95% CI), respectively (van Doremalen et al., 2020), 1 d and 4 d. A summary of the assumed model parameters is provided in Table 1.

145 An alternative means of constraining the fomite contribution is through an independent measurement of the change in the number of social contacts. Whilst a fomite contamination rate that scales in direct proportion to the number of social contacts cannot be constrained through this means, a fomite rate that is independent of, or only weakly dependent on, the mean number
150 of social contacts may be. For example, postal deliveries and food and medical purchases continued under the lockdown, and so may have served as an indirect means of transmitting the virus between households. By contrast, indirect transmission that occurred primarily in the workplace would have been severely curtailed by the lockdown.

155 An estimate of the change in R_t before and after lockdown may be determined using the results of the CoMix survey of social contacts during lockdown (Jarvis et al., 2020). The mean number of social contacts returned from the survey was 2.9 (with an inter-quartile range of $IQR = 1-4$), a substantial reduction compared with the earlier POLYMOD survey result of
160 10.8 ($IQR = 6-14$). This reduction factor is adopted to provide a normal-distributed prior on R_t during the lockdown, taking into account a possible contribution of fomites. Here, N_f is assumed to apply to a fomite component that was unchanged by the lockdown, and so is held constant before and after the lockdown. If additional fomites were present, then they would
165 have contributed even more to the overall death rate, including before the lockdown.

2.2. Power-series approximate solutions

Insight into the dynamical role played by fomites is provided by an approximate power-series solution to Eqs. (1). The first two equations show that the role of the fomites may be absorbed into an effective reproduction number

$$\tilde{R}_t = R_t + N_f \frac{D_i}{D_f} \frac{f_*}{i}. \quad (4)$$

This is a formal expression, in that the evolution of both f_* and i depend on R_t . Evaluating $\tilde{R}_t$ at its value near lockdown, however, provides an accurate description of the evolution of the populations afterwards. This may be demonstrated through power-series solutions to Eqs. (1). When the removed fraction r is small, it is convenient to use r as the independent variable rather than t . Power-series solutions around $r = r_0 = r(t_0)$ after a time t_0 are sought in the form:

$$s(r) = s_0 + \sum_{n=1}^{\infty} s_n (r - r_0)^n, \quad e(r) = e_0 + \sum_{n=1}^{\infty} e_n (r - r_0)^n, \quad i(r) = i_0 + \sum_{n=1}^{\infty} i_n (r - r_0)^n. \quad (5)$$

The solution to second order is described in the Appendix. The first order coefficients are

$$s_1 = -\tilde{R}_{t_0} s_0, \quad e_1 = \tilde{R}_{t_0} - \frac{D_i}{D_e} \frac{e_0}{i_0}, \quad i_1 = \frac{D_i}{D_e} \frac{e_0}{i_0} - 1, \quad (6)$$

where the effective reproduction number $\tilde{R}_{t_0} = R_t + (D_i/D_f)N_f f_*(r_0)/i_0$ has been defined. The role of the fomites, other than in rescaling R_t , arises only at second order (and not until even higher order for i), so that the presence of fomites may not be inferred from the evolution of the populations unless the fomite contribution is large and with a long duration.

The time development of the death rate, Eq. (3), is affected by fomites only at the second and higher order contributions from fomites. The role of fomites may then in principle be detected either through their second order contributions, or from independent knowledge of the reproduction number R_t . Each of these possibilities is considered in turn.

2.3. Statistical quantification

Eq. (3) is used to model the mortality rates from COVID-19, seeking maximum likelihood estimates for R_t and N_f . Following Flaxman et al.

(2020), the number of deaths is drawn from a negative binomial distribution with mean N_d and variance $N_d + N_d^2/\psi$, where N_d is the model prediction for the mean number of weekly deaths and ψ is a free parameter. In the limit $\psi \rightarrow \infty$, the distribution becomes Poisson. Following Flaxman et al. (2020), ψ is drawn from a half normal distribution with mode 0 and variance 5.

The model parameters are fit using the post-lockdown data and before lockdown is eased. The general decline in mortality rate is well-modelled using a constant reproduction rate (Flaxman et al., 2020). To seek a fomite contribution, a two-step procedure is followed. First maximum likelihood values for R_0 and $R_{t,ld}$ are found assuming no fomites contribute ($N_f = 0$); then, for the value of R_0 obtained, the joint likelihood for $R_{t,ld}$ and $N_f(\geq 0)$ is computed to determine whether adding a fomite term increases the likelihood. This procedure is followed because R_0 is made uncertain by limitations in assessing the death rate during the rapid rise of the infections. This uncertainty could produce possibly spurious higher likelihoods into lockdown when a fomite term is added if pre-lockdown deaths are included in the likelihood. The limits obtained are conservative in that the procedure minimizes the fomite signal arising from terms that may not be absorbed into a re-scaling of $R_{t,ld}$. Since any signal found may in principle be attributable to evolution in $R_{t,ld}$ during lockdown, the signal is regarded as an upper limit to the fomite contribution.

2.4. Reported mortality rates

The COVID-19 mortality rates compiled by the European Centre for Disease Control¹ (ECDC) are used for fitting the model parameters. The ECDC compiles data from up to 500 sources in each country each week from national and regional competent authorities.

3. Results

3.1. Uniform priors

A maximum likelihood fit to the death rates from the start of lockdown on 23 March 2020 until its easing in July 2020 gives a basic reproduction number $R_0 \simeq 3.107$ and $R_{t,ld} = 0.760$ during the lockdown period. (Although not used in this analysis, after the lockdown is eased the maximum-likelihood model for the death rates until the end of August corresponds to

¹<https://www.ecdc.europa.eu/en/covid-19/data-collection>

Figure 1: *Upper panel:* Weekly death rates in the UK. Black data points: the rates reported by the ECDC. Solid blue lines: Mean weekly deaths predicted for $D_f = 0.41$ d and assuming uniform priors for N_f and $R_{t,ld}$ after lockdown. Also shown is the 68% confidence range of uncertainty in the mean predicted number of deaths during lockdown (shaded region). Broken cyan line: Maximum likelihood fit without a fomite contribution. *Bottom panel:* Solid blue line: Mean predicted excess deaths per week arising from fomites along with the 95% confidence upper limit (shaded region), assuming uniform priors for N_f and $R_{t,ld}$.

Figure 2: *Upper panel:* Posterior probability density for $R_{t,ld}$, marginalised over N_f , both for a uniform prior for $R_{t,ld}$ (solid black line) and a prior based on the CoMix study (dashed blue line). Shown for $D_f = 0.41$ d. *Lower panel:* Probability density for excess number of deaths from fomites for $D_f = 0.41$ d over lockdown period, both for uniform and the CoMix prior on $R_{t,ld}$.

Figure 3: Relative difference between the effective reproduction number $\tilde{R}_{t,ld}$ for models with $N_f \simeq 0.14$, $R_{t,ld} \simeq 0.04$, corresponding to the 95% confidence upper limits on N_f , that would be inferred from the death rates assuming a counterfactual model with $N_f = 0$, compared with the maximum likelihood value for $R_{t,ld}$ for a model with $N_f = 0$. Shown for various values of D_f .

D_f	N_f (95% CL)	$R_{t,ld}$ (M-L)	$R_{t,ld}$ (95% UL)	N_f (M-L)
d	d ⁻¹			d ⁻¹
0.21	0.14	0.04	0.72	0.013
0.41	0.14	0.04	0.72	0.013
1	0.14	0.04	0.72	0.013
4	0.13	0.05	0.73	0.009

Table 2: Model results for uniform priors in N_f and $R_{t,ld}$. The columns are: (1) fomite duration (days); (2) 95% confidence upper limit on N_f (per infectious person per day) and (3) the corresponding maximum likelihood value for $R_{t,ld}$; (4) 95% confidence upper limit on $R_{t,ld}$ and (5) the corresponding maximum likelihood value for N_f (per infectious person per day).

$R_{t,ld} = 0.867$.) The solutions with fomites show only small departures in the death rates from a model with constant reproduction rate after lockdown, in accordance with the power-series approximation (see Appendix). The magnitude of the departure depends on the duration D_f of the fomites. The average numbers of weekly deaths predicted are shown in the upper panel of Fig. 1, along with the 68% confidence range in uncertainty during lockdown.

The joint likelihood function for $R_{t,ld}$ and N_f during lockdown is very flat, with the highest values corresponding to a ridge at $R_{t,ld} + D_f N_f \approx 0.76$ for $0.21 \leq D_f \leq 4$ d. The probability distribution for $R_{t,ld}$, marginalised over N_f , is shown in the upper panel of Fig. 2 for $D_f = 0.41$. The results are nearly independent of D_f for $0.21 \leq D_f \leq 4$ d. The marginalised 95% confidence upper limits on N_f are provided in Table 2. The maximum likelihood values for $R_{t,ld}$ corresponding to the 95% confidence upper limit values for N_f are in the next column. Also shown are the marginalised 95% confidence upper limits on $R_{t,ld}$ and the corresponding maximum likelihood values for N_f . The upper limits on N_f and $R_{t,ld}$ are not very sensitive to the fomite duration D_f , and correspond to the pandemic's being spread almost entirely by either fomites or direct transmission, respectively. The overall maximum likelihood solution corresponds to $N_f = 0$ and $R_{t,ld} = 0.76$ for all values of $0.21 \leq D_f \leq 4$ d.

The power-series solutions in Sec. 2.2 show fomites affect the death rates in second order. The magnitude of the departure from the prediction for a constant reproduction number may be seen from the effective reproduction number that would be inferred from the death rates assuming a counterfactual model with no fomites. Taking the exposure rate to be proportional to $\tilde{R}_{t,ld}(D_f)is/D_i$, it follows from Eqs. (1) that $\tilde{R}_{t,ld}(D_f) = R_{t,ld} + (D_i/D_f)N_f f_*/i$. The effective reproduction numbers $\tilde{R}_{t,ld}$ are computed from the solutions to models corresponding to the 95% confidence upper limits on N_f and their corresponding maximum likelihood values $R_{t,ld}$ in Table 2 for the various values of fomite duration D_f . Fig. 3 shows the fractional differences $\tilde{R}_{t,ld}(D_f)/R_{t,ld} - 1$, where $R_{t,ld}$ is the maximum likelihood value for the reproduction number for a model with $N_f = 0$. The effective reproduction numbers become time-dependent, increasing with time by an amount that depends on D_f . The fractional differences, however, are very small, exceeding 1% only when D_f is comparable to D_i . The effect of fomites is too small to be discernable from the population statistics without additional information.

Similarly, the mean excess numbers of weekly deaths for $N_f > 0$ compared with a counterfactual model having $N_f = 0$ are shown in the bottom panel of Fig. 1, along with the 68% confidence interval. The expected number of excess deaths over the lockdown period ranges from about 13000–16000 (with fewer excess deaths for large D_f), corresponding to a fraction 0.6–0.7 of all deaths in this period. The probability distribution in the number of excess deaths, however, is very broad, as shown in the bottom panel of Fig. 2. No strong statement on the number of excess deaths resulting from fomite transmission may be made: any number up to all cannot be excluded at the 95% confidence level.

3.2. CoMix study prior

The CoMix study provides new information on the reduction in the number of social contacts per person during lockdown that limits the allowed range in the reproduction number in lockdown, breaking the degeneracy (co-linearity) between R_t and N_f in the effective reproduction number $\tilde{R}_t(N_f, D_f)$. Allowing for fomites for which the transmission rate is constant, unchanged under lockdown conditions, the CoMix result on the reduction of social contacts before and after lockdown may be used to estimate a prior probability distribution on the reproduction number in lockdown, $R_{t,ld}$, as follows. The basic reproduction number found before lockdown is re-interpreted as an effective basic reproduction number $\tilde{R}_0 = R_0 + (D_i/D_f)N_f f_*/i$. For

Figure 4: *Upper panel:* Weekly death rates in the UK. Black data points: the rates reported by the ECDC. Solid blue lines: Mean weekly deaths predicted for $D_f = 0.41$ d and assuming a uniform prior for N_f and a prior for $R_{t,ld}$ after lockdown based on the CoMix study. Also shown is the 68% range of uncertainty in the mean predicted number of deaths during lockdown (shaded region). Red dotted line: Power series approximate solution during the lockdown period (see Appendix). *Bottom panel:* Blue solid line: Mean predicted excess deaths per week arising from fomites, along with the 95% upper limit (shaded region).

D_f	N_f (95% ul)	$R_{t,ld}$ (ml)	$R_{t,ld}$ (95% ul)	N_f (ml)
d	d ⁻¹			d ⁻¹
0.21	0.034	0.59	0.77	0
0.41	0.035	0.59	0.77	0
1	0.034	0.58	0.77	0
4	0.028	0.61	0.77	0

Table 3: Model results for a uniform prior for N_f and a prior for $R_{t,ld}$ based on the CoMix study. The columns are: (1) fomite duration (days); (2) 95% upper limit on N_f (per infectious person per day) and (3) the corresponding maximum likelihood value for $R_{t,ld}$; (4) 95% upper limit on $R_{t,ld}$ and (5) the corresponding maximum likelihood value for N_f (per infectious person per day).

any given values of N_f and D_f , the value of $\tilde{R}_0$ at the moment of lockdown, at time t_{ld} , is used to provide an estimate for the actual basic reproduction number in the model, $R_0(N_f, D_f) = \tilde{R}_0 - (D_i/D_f)N_f f_*(t_{ld})/i(t_{ld})$. Under the assumption that the reproduction numbers are directly proportional to the number of social contacts, for any given reproduction number $R_{t,ld}$ during lockdown, the likelihood for the model is multiplied by the likelihood of the ratio $f_{ld} = R_{t,ld}/R_0(N_f, D_f)$, taken to be proportional to a normal distribution with mean $\bar{f}_{ld} = 0.27$ and standard deviation $\sigma_{f_{ld}} = 0.086$, as inferred from Jarvis et al. (2020). (It is noted in practice it is sufficient to use the small D_f limit for obtaining $R_0(N_f) = \tilde{R}_0 - D_i N_f$, as the difference in estimators is small compared with R_0 .)

The probability distribution for $R_{t,ld}$, marginalised over N_f , is shown in Fig. 2 for $D_f = 0.41$ d. The results are nearly independent of D_f over $0.21 \leq D_f \leq 4$ d. In contrast to models with a uniform prior on $R_{t,ld}$, the CoMix study result on the reduction of social contacts in lockdown corresponds to a maximum likelihood value for the reproduction number during lockdown of $R_{t,ld} = 0.74^{+0.03}_{-0.07}$ (68% CI), after marginalising over N_f . This may be compared with the estimate for a model with $N_f = 0$ of $R_{t,ld} = 0.76 \pm 0.02$ (68% CI), so that allowing for fomites little affects the expected value but broadens the confidence interval.

By contrast, the marginalised probability distribution for N_f continues
 305 to be flat, peaking at $N_f = 0$. The marginalised 95% upper limit on
 N_f is $N_f \lesssim 0.035 \text{ d}^{-1} \text{ infectious person}^{-1}$ for $0.21 \leq D_f \leq 1 \text{ d}$ and $N_f <$
 $0.028 \text{ d}^{-1} \text{ infectious person}^{-1}$ for $D_f = 4 \text{ d}$. Corresponding to these limits on
 N_f , the maximum likelihood value for the reproduction number during lock-
 down is $R_{t,\text{ld}} = 0.6$. Marginalised over N_f , $R_{t,\text{ld}} < 0.77$ at 95% confidence.
 310 The corresponding maximum likelihood values for N_f vanish for all values
 of D_f . The overall maximum likelihood solution corresponds to $N_f = 0$ and
 $R_{t,\text{ld}} = 0.76$ for all values of $0.21 \leq D_f \leq 4 \text{ d}$. Consequently, conservatively
 no fomite signal is detected, but a much more restrictive upper limit on the
 fomite rate may be set compared with the case of a uniform prior on $R_{t,\text{ld}}$.
 315 The mean weekly death rate for $D_f = 0.41 \text{ d}$ is shown in the upper
 panel of Fig. 4, along with the 68% confidence interval. The lower panel
 shows the mean excess numbers of weekly deaths for $N_f > 0$ compared
 with a counterfactual model having $N_f = 0$, along with the 95% upper
 limit. Summing over the shown lockdown period gives an expectation of
 320 about 4960 excess deaths, compared with about 22900 total deaths. This
 corresponds to an expected fractional death excess of 0.22. The probability
 density for the total number of excess deaths over the lockdown period is
 broad, but peaks at a lower number of excess deaths compared with a uniform
 prior on $R_{t,\text{ld}}$, as shown in the lower panel of Fig. 2. The 95% upper limit
 325 on the number of excess deaths is 12000, corresponding to a fraction 0.52
 of all deaths. This is typified by a model for $D_f = 0.41 \text{ d}$ with $N_f =$
 $0.035 \text{ d}^{-1} \text{ infectious person}^{-1}$ and $R_{t,\text{ld}} = 0.59$ (see Table 3), for which the
 fraction of excess deaths from fomites during lockdown is 0.51. Similar results
 are found for $D_f = 0.21$ and 1 d . The number of excess deaths for $D_f = 4 \text{ d}$,
 330 however, is somewhat smaller. The expected number of excess deaths is
 about 4200 over the lockdown period, corresponding to a fraction 0.18 of all
 deaths during this time, with an upper limit of 10300 excess deaths (95%
 CL), corresponding to a fraction 0.45 of all deaths during lockdown.

4. Discussion

335 4.1. Parameter constraints

Fomites are recognised as potential contributors to the transmission of the
 novel coronavirus SARS-CoV-2, however statistically quantifying the contri-
 bution is rendered difficult by the rarity of fomites in the general environ-
 ment. During a long lockdown period, the reproduction number from direct

transmission may be nearly constant, simplifying the evolution of the infection. Population statistics are examined as a possible means to constrain the fomite transmission rate N_f during a lockdown. Fomite transmission alters the evolution in the rate of deaths compared with a non-fomite model with a constant reproduction number, although the difference is small. Adding a fomite contribution is not found to improve the likelihood during the lockdown phase allowing for uniform priors on the fomite transmission rate N_f and reproduction number for direct transmission during lockdown $R_{t,ld}$. Upper limits are placed of $N_f < (0.13-0.14) \text{ d}^{-1} \text{ infectious person}^{-1}$ (decreasing with D_f) and $R_{t,ld} < 0.72-0.73$ (increasing with D_f) (95% CL). The joint likelihood function for both N_f and $R_{t,ld}$ has a sharp ridge at $R_{t,ld} + D_i N_f \approx 0.76$, expressing their near statistical co-linearity. Whilst the formal maximum likelihood model corresponds to $N_f = 0$, the probability distribution for the number of excess deaths is broad. The expected number of excess deaths arising from fomites is 0.6–0.7 of the total during lockdown (decreasing with D_f), and the possibility that all deaths arose from fomites may not be excluded at better than 95–98% confidence (increasing with D_f).

Including the CoMix study results on the reduction in social contacts between people in the UK following the near complete lockdown imposed in March 2020 modifies the prior distribution on the reproduction number, allowing a tighter constraint to be placed on the fomite transmission rate of $N_f \lesssim 0.03 \text{ d}^{-1} \text{ infectious person}^{-1}$ (95% CL) for $0.21 < D_f < 4 \text{ d}$. The reproduction number during lockdown is found to be $R_{t,ld} = 0.74^{+0.03}_{-0.07}$ (68% CI), after marginalising over N_f . The probability distribution for $R_{t,ld}$ is little sensitive to the addition of fomites, so that models without fomites should provide reliable estimates even if fomites are active. The value found here is consistent with other estimates, but on the lower end of the range (DHSC/SAGE, 2021).

To give the limits on N_f some context, the mean number C_f of potentially contaminatable objects a person may come into contact with per day must be specified. An estimate for the post is $C_f \simeq 0.57 \text{ d}^{-1} \text{ person}^{-1}$ (Meiksin, 2020). For food items, the Office for National Statistics estimates an average spend of £63.80 per week for an average household (ONS, 2020) of 2.4 members (ONS, 2021). For an average product value of £2, this corresponds to $C_f \sim 2.5 \text{ d}^{-1} \text{ person}^{-1}$ for post and food items combined. The upper limit allowing for the CoMix prior then corresponds to a fomite transmissivity upper limit of $T_f < 0.014$ (95% CL), or at most about 1 in 70 objects that comes into contact with an infectious person transmits the infection to a susceptible

person.

Quantifying the impact of fomites in terms of the excess deaths compared with a counterfactual non-fomite model, the expected number of excess deaths during lockdown is found to be 22% of all deaths during this period. The probability distribution on the number of excess deaths is found to be broad, however, so that the 95% upper limit on the number of excess deaths is 52% of all deaths during lockdown for $0.2 \leq D_f \leq 1$ d. For fomites with a duration time comparable to the infectious period, the limits are somewhat more restrictive. For $D_f = 4$ d, the expected number of excess deaths is 0.18 of all deaths during lockdown, with a 95% upper limit of 0.45. Regardless of fomite duration time, allowing for the CoMix study prior on the reproduction number during lockdown shows that it is highly unlikely that most deaths from COVID-19 were caused by transmission through lockdown-independent fomites.

4.2. Limitations of study

Placing constraints on the contribution of fomites during lockdown requires some key assumptions. In this context, fomites may be divided into two types, those that scale like the number of social contacts and those independent of the number of social contacts. Only the latter, such as fomites arising from essential services that continued through the lockdown like post deliveries and food and medicine purchases, may be constrained using the change in the evolution of the population statistics following lockdown. Fomite transmission rates that scale with the number of social contacts, such as may occur for fomites in office spaces, likely scale with the reproduction number for direct transmission, and so their effects on population evolution statistics are indistinguishable from those of direct transmission. It is not possible to rule out from population statistics the possibility that most of the transmission prior to lockdown arose from workplace fomites. Although contamination through household fomites may have then been expected to increase during lockdown, an increase in fomite transmission in a domestic setting may have been masked by the limited number of additional family members that could have been infected. Increased hand-washing in the home could in principle also be a factor.

The strongest constraints on fomite transmission in this study rely on the measured change in the number of social contacts following lockdown from the CoMix study in the UK. It is assumed the change in the reproduction number is proportional to the change in the number of social contacts, and

415 that the proportionality factor is independent of age. The measured reduction may be subject to several biases (Jarvis et al., 2020): there is a possibility of recall bias, as the study requested information about the previous day; the sample may be subject to selection bias if preferably people observing the lockdown replied to the survey; also children were not interviewed, so that
420 child-child contacts were inferred from the POLYMOD survey.

The maximum likelihood models with fomites were found to result in a small increase in the transmission with time compared with the maximum likelihood models without fomites. This difference accounts for the more stringent upper limits on the fomite contribution for longer fomite duration
425 times. The increase may be interpreted as evidence for a slightly increasing reproduction number with time, in which case the fomite contribution would be even smaller. The alternative of a decreasing reproduction number with a larger fomite contribution than the upper limits found here cannot be ruled out, but it would seem unlikely the reproduction number would decrease in
430 anticipation of an easing of the lockdown, and in any case this possibility would have only a small effect on the limits found.

4.3. Suggestions for further progress

CoMix is part of a European partnership involving 17 countries². As more results on the change in social contacts are published, similar analyses
435 to this may be performed to tighten constraints on fomite transmission of the SARS-CoV-2 virus within Europe.

The high availability of tests for infections may now allow more direct estimates of transmission through post and grocery deliveries by correlating illnesses between recipients and deliverers, including sorters and stockers, who
440 may have been in an asymptomatic infectious phase when handling items. Such correlations may be made possible with the assistance of data from the track-and-trace smartphone app system ³ in conjunction with appropriate data protection measures.

5. Conclusions

445 Because of their longevity on surfaces, fomites alter the time evolution of populations in an epidemic compared with direct transmission alone. The

²<https://www.uhasselt.be/UH/71795-start/The-CoMix-study>

³<https://www.nhs.uk/apps-library/nhs-covid-19/>; <https://protect.scot>

differences, however, are too small to detect when the fomite duration is shorter than the duration of the infectious phase. For uniform priors on the reproduction rate and fomite transmission rate, only coarse upper limits may
 450 be set on the fomite transmission rate. Allowing for a measurement of the change in the number of social contacts when a lockdown is implemented, as provided by the CoMix study, and under the assumption that the reproduction number scales with the number of social contacts, more restrictive constraints may be placed on the role of fomites. Using data for the UK
 455 from the lockdown in March to July 2020, it is found that fomites that act independently of a lockdown, such as delivered post or grocery items, contributed to fewer than about half of the total deaths from COVID-19 during the lockdown (95% CL), and likely fewer than a quarter.

Declaration of Competing Interests: None.

460 Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Appendix A.

Power-series solutions of Eqs. (1) in r are constructed by first dividing
 465 the equations by $dr/dt = i/D_i$ to transform them to

$$\begin{aligned}\frac{ds}{dr} &= -R_t s - \frac{D_i}{D_f} N_f \frac{s}{i} f_*, \\ \frac{de}{dr} &= R_t s + \frac{D_i}{D_f} N_f \frac{s}{i} f_* - \frac{D_i}{D_e} \frac{e}{i} + c_e \frac{D_i}{i}, \\ \frac{di}{dr} &= \frac{D_i}{D_e} \frac{e}{i} - 1 + c_i \frac{D_i}{i}, \\ \frac{df_*}{dr} &= D_i - \frac{D_i}{D_f} \frac{f_*}{i}.\end{aligned}\tag{A.1}$$

Power-series solutions around $r = r_0 = r(t_0)$ after a time t_0 are assumed:

$$s(r) = s_0 + \sum_{n=1}^{\infty} s_n (r-r_0)^n, \quad e(r) = e_0 + \sum_{n=1}^{\infty} e_n (r-r_0)^n, \quad i(r) = i_0 + \sum_{n=1}^{\infty} i_n (r-r_0)^n. \quad (\text{A.2})$$

It is numerically convenient to use the integrated solution for the fomite term:

$$\begin{aligned} f_*(r) &= f_*(r_0) \exp \left[- \int_{r_0}^r dr' \frac{D_i}{D_f} \frac{1}{i(r')} \right] \\ &+ D_i \exp \left[- \int_{r_0}^r dr' \frac{D_i}{D_f} \frac{1}{i(r')} \right] \int_{r_0}^r dr'' \exp \left[\int_{r_0}^{r'} dr'' \frac{D_i}{D_f} \frac{1}{i(r'')} \right] \end{aligned} \quad (\text{A.3})$$

To second order, the coefficients are

$$s_1 = -\tilde{R}_{t0} s_0, \quad s_2 = \frac{1}{2} \tilde{R}_{t0}^2 s_0 - \frac{1}{2} \frac{D_i}{D_f} N_f \left(1 - \frac{D_i}{D_e} \frac{e_0}{i_0} - \frac{D_i}{D_f} \right) \frac{f_*(r_0) s_0}{i_0^2} - \frac{1}{2} \frac{D_i^2}{D_f} N_f \frac{s_0}{i_0}, \quad (\text{A.4})$$

$$e_1 = \tilde{R}_{t0} - \frac{D_i}{D_e} \frac{e_0}{i_0}, \quad (\text{A.5})$$

$$\begin{aligned} e_2 &= -\frac{1}{2} \tilde{R}_{t0}^2 s_0 - \frac{D_i}{2 D_e i_0} \left\{ \tilde{R}_{t0} s_0 + \frac{e_0}{i_0} \left[1 - \frac{D_i}{D_e} \left(1 + \frac{e_0}{i_0} \right) \right] \right\} \\ &+ \frac{1}{2} \frac{D_i}{D_f} N_f \frac{s_0}{i_0} \left\{ D_i + \frac{f_*(r_0)}{i_0} \left[1 - \frac{D_i}{D_f} \left(1 + \frac{D_f e_0}{D_e i_0} \right) \right] \right\}, \end{aligned} \quad (\text{A.6})$$

and

$$i_1 = \frac{D_i}{D_e} \frac{e_0}{i_0} - 1, \quad i_2 = \frac{D_i}{2 D_e i_0} \left\{ \tilde{R}_{t0} s_0 + \frac{e_0}{i_0} \left[1 - \frac{D_i}{D_e} \left(1 + \frac{e_0}{i_0} \right) \right] \right\}, \quad (\text{A.7})$$

470 where the effective reproduction number $\tilde{R}_{t0} = R_t + (D_i/D_f) N_f f_*(r_0)/i_0$ has been defined.

The forms of the coefficients show that to first order, the fomite term does not affect the evolution of the populations other than through a re-scaling of R_t to $\tilde{R}_{t0}$. At second order, the evolution of the susceptible and exposed

475 populations show additional contributions from the fomites that are not eliminated by a re-scaling, whilst the evolution of the infectious population, in contrast, may still be re-scaled to second order. Using the expression for i to second order, Eq. (A.3) becomes

$$f_*(r) \simeq f_*(r_0) \left[\frac{y_+(r)}{y_+(r_0)} \frac{y_-(r_0)}{y_-(r)} \right]^\delta + \frac{D_i}{4i_2\Delta(1+\delta)} y_+(r) y_-(r) \left\{ \left[\frac{y_+(r)}{y_+(r_0)} \frac{y_-(r_0)}{y_-(r)} \right]^{1+\delta} - 1 \right\}, \quad (\text{A.8})$$

where $y_\pm(r) = \Delta \pm [i_1 + 2i_2(r - r_0)]$, $\Delta = (i_1^2 - 4i_0i_2)^{1/2}$ and $\delta = D_i/(D_f\Delta)$.
480 (For applications here, $i_1^2 - 4i_0i_2 > 0$.)

The evolution of the populations may be expressed as a function of t through

$$t = t_0 + D_i \int_0^{r-r_0} \frac{dx}{i_0 + i_1x + i_2x^2} = \frac{D_i}{\Delta} \log \left[\frac{y_-(r)}{y_+(r)} \right]. \quad (\text{A.9})$$

An example solution is illustrated in the upper panel of Fig. 4 for a solution at the 98% upper limit of $N_f = 0.042$ allowing for the CoMix prior,
485 and the corresponding maximum likelihood value $R_{t,\text{ld}} = 0.55$. The predicted number of deaths agrees with the numerical solution for these parameter values to within 1%.

References

- Al-Ansary, L., Bawazeer, G., Beller, E., *et al.*, 2020. Physical interventions
490 to interrupt or reduce the spread of respiratory viruses. part 2 - hand hygiene and other hygiene measures: systematic review and meta-analysis doi:10.1101/2020.04.14.20065250. (Last accessed 14 June 2021).
- Colaneri, M., Seminari, E., Novati, S., *et al.*, 2020. Severe acute respiratory
495 syndrome coronavirus 2 rna contamination of inanimate surfaces and virus viability in a health care emergency unit. Clin Microbiol Infect. 26, 1094.e1–1094.e5. doi:10.1016/j.cmi.2020.05.009.
- Davies, N., Kucharski, A., Eggo, R., Gimma, A., Edmunds, W., 2020. Effects of non-pharmaceutical interventions on covid-19 cases, deaths, and demand

- 500 for hospital services in the uk: a modelling study. *Lancet Public Health* 5, e375–e385. doi:10.1016/S2468-2667(20)30133-X.
- DHSC/SAGE, 2021. The r value and growth rate. Department of Health and Social Care (DHSC) and Scientific Advisory Group for Emergencies (SAGE). URL: <https://www.gov.uk/guidance/the-r-value-and-growth-rate>.
505 (Last accessed 15 June 2021).
- du Plessis, L., McCrone, J., Zarebski, A., *et al.*, 2020. Establishment & lineage dynamics of the sars-cov-2 epidemic in the uk doi:10.1101/2020.10.23.20218446.
510 (Last accessed 14 June 2021).
- Flaxman, S., Mishra, S., Gandy, H., *et al.*, 2020. Estimating the effects of non-pharmaceutical interventions on covid-19 in europe. *Nature* 584, 257–261. doi:10.1038/s41586-020-2405-7.
- Harbourt, D., Haddow, A., Piper, A., *et al.*, 2020. Modeling the stability
515 of severe acute respiratory syndrome coronavirus 2 (sars-cov-2) on skin, currency, and clothing. *PLoS Negl Trop Dis* 14, e0008831. doi:10.1371/journal.pntd.0008831.
- Jarvis, C., Van Zandvoort, K., Gimma, A., *et al.*, 2020. Quantifying the impact of physical distance measures on the transmission of covid-19 in
520 the uk. *BMC Med* 18. doi:10.1186/s12916-020-01597-8.
- Jiang, F., Jiang, X.L., Wang, Z.G., Meng, Z.H., Shao, S.F., Anderson, B., Ma, M.J., 2020. Detection of severe acute respiratory syndrome coronavirus 2 rna on surfaces in quarantine rooms. *Emerg Infect Dis.* 26, 2162–2164. doi:10.3201/eid2609.201435.
- 525 Lewis, D., 2021. Covid-19 rarely spreads through surfaces. so why are we still deep cleaning? *Nature* 590, 26–28. doi:10.1038/d41586-021-00251-4.
- Lio, C., Cheon, H., Lei, C., *et al.*, 2021. Effectiveness of personal protective health behaviour against covid-19. *BMC Public Health* 21, 827. doi:10.1186/s12889-021-10680-5.

- 530 Meiksin, A., 2020. Dynamics of covid-19 transmission including indirect transmission mechanisms: a mathematical analysis. *Epidemiology and Infection* 148, 1–7. doi:10.1017/S0950268820002563.
- Morawska, L., Milton, D., 2020. It is time to address airborne transmission of coronavirus disease 2019 (covid-19). *Clin Infect Dis.* 71, 2311–2313. doi:10.1093/cid/ciaa939.
- 535 Mossong, J., Hens, N., Jit, M., *et al.*, 2008. Social contacts and mixing patterns relevant to the spread of infectious diseases. *PLoS Med.* 5, e74. doi:10.1371/journal.pmed.0050074.
- Ong, S., Tan, Y., Chia, P., *et al.*, 2020. Air, surface environmental, and personal protective equipment contamination by severe acute respiratory syndrome coronavirus 2 (sars-cov-2) from a symptomatic patient. *JAMA* 323, 1610–1612. doi:10.1001/jama.2020.3227.
- ONS, 2020. Expenditure on food and non-alcoholic drinks by place of purchase: table a2. Office for National Statistics (ONS). URL: <https://www.ons.gov.uk>.
- 545 (Last accessed 11 June 2021).
- ONS, 2021. Families and households in the uk: 2020. Statistical bulletin. Office for National Statistics (ONS). URL: <https://www.ons.gov.uk>. (Last accessed 11 June 2021).
- 550 Pastorino, B., Touret, F., Gilles, M., de Lamballerie, X., Charrel, R., 2020. Prolonged infectivity of sars-cov-2 in fomites. *Emerg Infect Dis.* 26, 2256–2257. doi:10.3201/eid2609.201788.
- PHAS, 2020. The infection fatality rate of COVID-19 in Stockholm. Technical Report. Folkhälsomyndigheten: Public Health Agency of Sweden (PHAS). URL: <https://www.folkhalsomyndigheten.se/publicerat-material/>.
- 555 (Last accessed 28 May 2021).
- Russell, T., Hellewell, J., Jarvis, C., *et al.*, 2020. Estimating the infection and case fatality ratio for coronavirus diseases (covid-19) using age-adjusted data from the outbreak on the diamond princess cruised ship, february 2020. *Euro Surveill* 25. doi:10.2807/1560-7917.ES.2020.25.12.2000256.
- 560

- van Doremalen, N., Bushmaker, T., Morris, D., *et al.*, 2020. Aerosol and surface stability of sars-cov-2 as compared with sars-cov-1. N Engl J Med 382, 1564–1567. doi:10.1056/NEJMc2004973.
- WHO, 2020. Transmission of sars-cov-2: implications for infection prevention precautions. Scientific Brief. World Health Organization (WHO). URL: <https://www.who.int/news-room/commentaries/detail/transmission-of-sars-cov-2-implications-for-infection-prevention-precautions>. (Last accessed 25 May 2021).
- Zuo, Y., Uspal, W., Wei, T., 2020. Airborne transmission of covid-19: Aerosol dispersion, lung deposition, and virus-receptor interactions. ACS Nano 14, 16502–16524. doi:10.1021/acsnano.0c08484.