The relationship between sleep disturbance and cognitive impairment in mood disorders: A systematic review

Oliver Pearson^{1,2}, Nora Uglik-Marucha¹, Kamilla W. Miskowiak³, Scott A. Cairney^{4,5}, Ivana Rosenzweig^{6,7}, Allan H. Young^{2,8}, Paul R.A. Stokes^{2,8}

¹Psychometrics and Measurement Lab, Biostatistics and Health Informatics
Department, Institute of Psychiatry, Psychology and Neuroscience, King's College
London, SE5 8AF, London, UK

²Centre for Affective Disorders, Department of Psychological Medicine, Institute of
Psychiatry & Psychology and Neuroscience, King's College London

³Copenhagen Affective Disorder Research Centre (CADIC), Psychiatric Centre
Copenhagen, Copenhagen University Hospital, Rigshospitalet, Copenhagen,
Denmark.

⁴Department of Psychology, University of York, York, YO10 5DD

⁵York Biomedical Research Institute, University of York, York, YO10 5DD

⁶Sleep and Brain Plasticity Centre, Department of Neuroimaging, IoPPN, KCL

⁷Sleep Disorders Centre, Guy's and St Thomas' Hospital, GSTT NHS, London

⁸South London and Maudsley NHS Foundation Trust, Bethlem Royal Hospital,
Monks Orchard Road, Beckenham, Kent, BR3 3BX, United Kingdom

Corresponding Author:

Oliver Pearson

Email: oliver.pearson@kcl.ac.uk

Research assistant

Psychometrics and Measurement Lab

Department of Biostatistics and Health Informatics

Institute of Psychiatry, Psychology & Neuroscience, King's College London, 16 De
Crespigny Park, London, SE5 8AF, UK

Abstract word count: 256, Text word count: 3436,

Number of Tables: 2, Number of Figures: 1

Abstract

Background: Cognitive impairment experienced by people with bipolar disorders (BD) or major depressive disorder (MDD) is associated with impaired psychosocial function and poorer quality of life. Sleep disturbance is another core symptom of mood disorders which may be associated with, and perhaps worsen, cognitive impairments. The aim of this systematic review was to critically assess the relationship between sleep disturbance and cognitive impairment in mood disorders.

Methods: In this systematic review, relevant studies which examined sleep disturbance and cognitive function in people with mood disorders were identified using electronic database searches of PsychINFO, MEDLINE, Embase and Web of Science.

Findings: Fourteen studies were included; eight investigated people with BD, five investigated people with MDD, and one included both people with MDD and people with BD. One study was an intervention for sleep disturbance and the remaining thirteen studies used either a longitudinal or cross-sectional observational design. Ten studies reported a significant association between subjectively measured sleep disturbance and cognitive impairment in people with MDD or BD after adjusting for demographic and clinical covariates, whereas no such association was found in healthy participants. Two studies reported a significant association between objectively measured sleep abnormalities and cognitive impairment in mood disorders. One study of cognitive behavioural therapy for insomnia modified for BD (CBTI-BD) found an association between improvements in sleep and cognitive performance in BD. **Interpretation:** There is preliminary evidence to suggest a significant association between sleep disturbance and cognitive impairment in mood disorders. These findings suggest that identifying and treating sleep disturbance may be important when addressing cognitive impairment in MDD and BD. **Funding:** none.

Key words: sleep disturbance, sleep disorder, insomnia, hypersomnia, cognitive impairment, bipolar disorder, major depressive disorder

Introduction

Cognitive impairment in people with bipolar disorders (BD) or major depressive disorder (MDD) is associated with reduced functional capacity,^{1,2} poor illness prognosis,^{3,4} and poor quality of life.^{5,6} Meta-analyses have reported significant deficits in executive function, memory and attention in people with MDD currently experiencing a depressive episode, with effect sizes ranging from $d = -0.32$ to -0.97 .^{7,8} In BD, a meta-analysis has reported moderate effect size deficits in executive function in people experiencing depressive episodes ($d = -0.55$) and large effect sizes in people experiencing manic episodes ($d = -0.72$)⁹. However, cognitive impairment is not confined to mood episodes, but persists during euthymia in MDD and BD. For example, Rock et al.⁷ found moderate effect size deficits in executive function and attention in people with remitted MDD, while Semkovska et al.¹⁰ identified deficits in working memory and long-term memory. In BD, medium to large effect size deficits in executive function, verbal memory, sustained attention and psychomotor speed have been identified by a meta-analysis.¹¹ Nevertheless, there is considerable heterogeneity in cognitive impairments across people with mood disorders and approximately 35% of people with MDD¹² and 20-40% of people with BD^{13,14} experience moderate to severe impairment in several cognitive domains.

The aetiology of cognitive impairment in mood disorders is not fully understood,¹⁵ and determining how cognitive impairment is mediated in MDD and BD is a key priority. Sleep disturbance may be one contributing factor. Sleep disturbances are core features of manic episodes, hypomanic episodes and major depressive episodes manifesting as insomnia or hypersomnia and fatigue in major depressive episodes and a decreased need for sleep in manic or hypomanic episodes.¹⁶ There is also growing evidence to suggest that sleep disturbance persists during euthymia in a substantial proportion of people with mood disorders. Clinically significant sleep disturbances are reported by 70% of euthymic people with BD¹⁷ and 44-60% of euthymic people with MDD¹⁸⁻²⁰. Moreover, sleep disturbance is associated with higher rates of relapse and suicide attempts in both BD²¹ and MDD²². Sleep disturbances escalate just before manic and depressive relapse in BD, and worsen during these episodes.²³

There is strong evidence detailing the negative effects of sleep disturbance on many aspects of cognition in healthy individuals²⁴, including fragmented memory loss²⁵ and impaired suppression of unwanted thoughts²⁶. Recently, evidence has emerged to suggest an association between sleep disturbance and cognitive impairment in mood disorders. For example, Bradley et al.²⁷ found that people with BD and abnormal sleep patterns, as measured by sleep actigraphy, performed significantly poorer than healthy participants in cognitive performance, whereas people with BD and normal sleep patterns did not significantly differ from healthy participants. As yet, there has not been a systematic review of the relationship between sleep disturbance and cognitive impairment in mood disorders.

The aim of this systematic review was to critically analyse the relationship between sleep disturbance and cognitive impairment in MDD and BD. In particular, the association between sleep disturbance and cognitive impairment, and relevant factors moderating, mediating, and confounding the relationship will be examined.

Methods

Literature search

Two reviewers (OP and NUM) independently performed the literature search. Relevant studies were identified using electronic database searches of PsychINFO, MEDLINE, Embase and Web of Science. The following search string was used: [(sleep disturb*) OR (sleep disorder*) OR (insomn*) OR (hypersomn*)] AND [(cognitive impair*)] AND [(bipolar*) OR (depress*)]. Papers were extracted from inception until February 2021. Only studies written in English were considered. The relevance of the article was initially verified by title and abstract review and then by a further review of each manuscript to verify whether they met the inclusion and exclusion criteria. This systematic review followed PRISMA 2020²⁸ guidelines, although a study protocol was not registered.

Inclusion and exclusion criteria

All databases were screened following the same protocol. Included studies were required to include participants with a diagnosis of MDD, BD-I or BD-II, meeting criteria for ICD-9²⁹ or subsequent editions, or DSM-III³⁰ or subsequent editions. Included studies were required to measure both sleep disturbance and cognition.

Studies were excluded if they: did not primarily include people with a diagnosis of MDD, BD-I or BD-II; included only participants with dysthymia or cyclothymia or with comorbid neurological disorders (e.g. dementia, epilepsy or stroke) or mild cognitive impairment (MCI); included primarily older participants; were follow-up studies or case reports; were not published by a peer-reviewed scientific journal.

Data extraction

The following data were extracted: study name and year; participant sample size, age range, disorder and mood state of participants, measures of sleep and cognition, a summary of main findings and study limitations.

Quality assessment

Two reviewers (OP and NUM) independently assessed the quality of included studies using the Quality Assessment Tool for Quantitative Studies, developed by the Effective Public Health Practice Project (EPHPP).³¹ The following criteria of each study were rated as either strong, moderate or weak: selection bias, design, confounders, blinding, data collection methods, and withdrawals/drop-outs (if applicable). These ratings were aggregated to form a global rating: studies were rated globally as strong if they received no weak ratings; moderate if they received one weak rating; and weak if they received two or more weak ratings.

Results

A total of 3094 studies were identified, which was reduced to 2208 after duplicates were removed. Following primary assessment of titles and abstracts, this was reduced to 30 studies. Based on secondary assessment of full texts, we included 14 studies that fulfilled the eligibility criteria (please see Figure 1 for further details).

INSERT FIGURE 1 HERE PLEASE

Studies

Of the 14 included studies, eight included people with BD^{27,32–37}, five included people with MDD^{38–42}, and one included both people with MDD and people with BD⁴³ (see *Table 1*). These studies were published between years 2012–2020. A total of 2353 people with mood disorders were included in these studies, 1083 people with MDD

and 1270 people with BD. The mood state of participants varied across studies. Of the eight BD studies, six included only euthymic participants^{32,33,35,35,37,44}, whereas two included participants in any mood state.^{27,34} Of the five MDD studies, four only included people currently experiencing a depressive episode; however, Wilckens et al. included both people with current depression and people with remitted depression. Mood state was not described in the one study of people with MDD or BD⁴³.

INSERT TABLE 1 HERE PLEASE

BD results

Six of the eight BD studies reported associations between sleep disturbance and cognitive impairment. Kanady et al.³³ found that greater sleep time variability predicted poorer working memory ($p < 0.01$) and verbal learning performance ($p < 0.01$) in BD. Furthermore, following treatment with cognitive behaviour therapy for insomnia modified for bipolar disorder (CBTI-BD⁴⁵), a reduction in total wake time predicted improved working memory ($p < 0.05$) and a reduction in sleep time variability predicted improved verbal learning ($p < 0.05$). Laskemoen et al.³⁴ found significant associations between insomnia and processing speed ($p < 0.001$), insomnia and inhibition ($p < 0.01$), hypersomnia and processing speed ($p < 0.01$), and hypersomnia and inhibition ($p < 0.05$) in people with BD, whereas no association between sleep disturbances and cognition was found in healthy participants. Samalin³⁵ et al. found that sleep disturbances were indirectly associated with cognitive function potentially mediated via residual depressive symptoms and perceived cognitive performance ($p = 0.001$). Another study by Samalin et al.⁴⁶ reported that participants with impaired cognitive functioning had significantly higher scores ($p < 0.02$), and thus more severe sleep disruptions, on PSQI than participants with satisfactory cognitive functioning. Volkert et al.³⁷ reported that people with BD experiencing persistent sleep disturbances show more severe cognitive dysfunctions ($p < 0.05$). Bradley et al.²⁷ found that people with BD with normal sleep patterns were not significantly different to healthy participants in cognitive functions, but people with BD and sleep disturbance (including objective measures) performed significantly worse in cognitive tests than healthy participants ($p < 0.001$). Russo et al.⁴⁴ found that sleep disturbance was significantly negatively associated with social cognition (r

= -0.273, $p = 0.004$), but only associated at a trend level with working memory ($r = -0.179$, $p = 0.061$) and visual learning ($r = -0.161$, $p = 0.094$). Boland et al.³² reported no significant associations between sleep disturbance (including objective measures) and cognition, except for a significant negative correlation between daytime dysfunction due to sleep disturbance and working memory ($r = -0.57$, $p = 0.03$).

MDD results

In MDD, Cabanel et al.³⁸ reported that patients with severely impaired sleep quality (PSQI > 10) had significantly poorer performance on the TMT-A (visuomotor processing speed; $p = 0.011$) and the TMT-B (executive function; $p = 0.015$) than people with MDD with normal or moderately impaired sleep quality (PSQI < 10). Cha et al.⁴¹ found that sleep quality predicted impairments in subjective cognitive performance ($p < 0.001$) and objective cognitive impairments ($p = 0.017$). Depression severity was a partial mediator of the relationship between subjective cognitive function and sleep quality ($B = -0.43$; 95% CI, -0.56, -0.33; $B = -0.18$; 95% CI, -0.30, -0.07 after controlling for depression severity). Wilckens et al.⁴² reported that higher wake after sleep onset (i.e., longer periods of wakefulness occurring after defined sleep onset), and shorter total sleep time, were linearly associated with slower response time across single-task ($p = 0.10$) and switching blocks ($p = 0.005$) considered a proxy for condition-independent psychomotor slowing⁴⁷; however, these associations were also observed in healthy controls and therefore not specific to MDD. Müller et al.³⁹ found that 25-40% of participants with MDD reported 'almost always' daytime sleepiness, non-restorative sleep, attention deficits or memory complaints, with significant correlations between these variables ($p < 0.05$), as well as curvilinear associations between sleep duration and sleep quality (quadratic contrast, $p < 0.05$). However, Peng et al.⁴⁰ reported no significant association between cognition and sleep disturbance ($r = 0.029$, $p = 0.219$) in adolescents with first episode MDD.

Results in mixed MDD & BD groups

Soehner and Harvey⁴³ (N = 639 MDD participants; N = 138 BD participants) reported that participants with a mood disorder and insomnia in the past year experienced

significantly greater cognitive impairment in the past 30 days than participants with a mood disorder and no insomnia symptoms in the past year ($p = 0.004$).

Methodological quality assessment

Methodological quality assessment of studies using the EPHPP Quality Assessment Tool for Quantitative Studies³¹ are displayed in *Supplementary Table 1*. Selection bias was generally rated as moderate because in most studies participants were sampled from either outpatient clinics or inpatient settings. Study design was generally rated as weak because most were cross-sectional. Most studies controlled for at least 80% of relevant confounding variables, such as age, coffee consumption etc. Blinding was rated as moderate if it was not described, as was the case in almost all studies. Most studies used validated and reliable measures for sleep disturbance (e.g. Pittsburgh Sleep Quality Index, PSQI⁴⁸) and cognition (e.g. MATRICS Consensus Cognitive Battery, MCCB⁴⁹), and were therefore rated strong. Where applicable, withdrawals were reported satisfactorily by most studies. There was no association between global rating and positivity of findings.

Discussion

The purpose of this systematic review was to critically analyse the published literature on the relationship between sleep disturbance and cognitive impairment in BD and MDD. Fourteen studies were included in this review, for which quality assessment ranged from weak to strong.

Kanady et al.³³ was the only RCT included in this review. This study, in which CBTI-BD⁴⁵ was administered to participants with BD, found that a reduction in total wake time predicted improved working memory. The study also found that a reduction in sleep time variability predicted improved verbal learning. These findings suggest that sleep disturbance may contribute to cognitive impairment in BD, and that CBTI-BD may be an effective treatment. The remaining studies included in our review used either a longitudinal or cross-sectional design, and therefore did not allow for causal inferences to be made.

Ten of the twelve studies that measured sleep disturbance using subjective measures in participants with either MDD or BD found a significant association between sleep disturbance and cognitive impairment when adjusting for demographic and clinical covariates. In contrast, no significant association was reported in healthy participants, which suggests that this relationship may be specific to mood disorders. Given the strong evidence on the effect of objective sleep disturbance on cognitive impairment in healthy individuals²⁴, this may indicate that subjective experience of sleep impairment may also increase risks for cognitive impairment in people with mood disorders. Some studies also found a curvilinear association, indicating that both extremely low (<5 hours) and extremely high (>11 hours) self-reported sleep durations were associated with cognitive impairment; this was observed in participants with MDD (e.g. Müller et al.³⁹) and in BD (e.g. Laskemoen et al.³⁴). However, there was also evidence to suggest that insomnia may have more deleterious effects on cognitive impairment than hypersomnia. For example, Laskemoen et al.³⁴ found a larger effect size between insomnia and processing speed ($F = 15.43$, $p < 0.001$, $\eta^2 = 0.019$) than hypersomnia ($F = 6.87$, $p < 0.01$, $\eta^2 = 0.009$) in people with BD.

The main disadvantage of using subjective measures to assess sleep disturbance in people with mood disorders is that there is a considerable risk of bias. Euthymic people with either BD or MDD tend to significantly underestimate sleep duration and quality.^{17,50} Indeed, Boland et al.³² found that people with BD had significantly higher insomnia severity index (ISI) scores than healthy participants (indicating worse sleep quality), yet found no significant difference between BD and healthy participants in objective sleep parameters, as measured by actigraphy. This effect may be due to maladaptive sleep-related beliefs developed during mood episodes, in which sleep disturbance is particularly severe, which are then carried over into euthymia.⁵⁰ Nevertheless, subjective measures provide valuable information about the perception of sleep disturbance and sleep-related cognitions, and interventions of maladaptive sleep beliefs may be a useful way of treating sleep disturbance in mood disorders⁵¹. Although antidepressant treatment produces no change in maladaptive sleep beliefs in MDD⁵⁰, Cognitive Behavioural Treatment for Insomnia (CBT-I) has been shown to significantly improve maladaptive sleep beliefs in people with insomnia. CBT-I has been found to significantly improve both subjective sleep

measures, measured by sleep diaries, and objective sleep measures, measured by polysomnography (PSG).⁵¹ Therefore, sleep-related cognitions may be an important target for treating sleep disturbance and potentially cognitive impairment in people with mood disorders.

Only three studies included in this review measured sleep disturbance using objective measures: two examined sleep disturbance in participants with BD using actigraphy^{27,32} and one investigated sleep disturbance in participants with MDD using polysomnography⁴². The MDD study that used polysomnography found that longer periods of wakefulness occurring after defined sleep onset were associated with psychomotor slowing, however these associations were not specific to MDD⁴². One advantage of polysomnography is that it is more accurate than actigraphy in recognising sleep and wake times, and is therefore the ‘gold standard’ of objective sleep measurement. Another key advantage of polysomnography is the ability to measure sleep architecture, however, the Wilckens et al MDD study did not report sleep architecture measurements.

Mood disorders are associated with several sleep architecture abnormalities, particularly altered distribution of rapid eye movement (REM) sleep in MDD⁵² and BD⁵³. Components of sleep architecture have been linked to aspects of cognitive impairment in healthy individuals. For example, time in slow wave sleep contributes to processing speed in healthy individuals.⁵⁴ Abnormal sleep architecture may be associated with cognitive impairment in mood disorders, so sleep architecture may be a highly relevant variable to measure. Moreover, knowledge of the specific components of sleep architecture associated with improvements in mental health or cognition could be targeted in interventions for treatment or prevention. For example, enhancing slow oscillation activity via non-invasive auditory stimulation has been shown to improve memory in healthy individuals⁵⁵. Thus, we suggest that future studies of the role of sleep in cognition in mood disorders use polysomnography where possible.

Interestingly, the two BD studies which used actigraphy had opposite findings. Bradley et al.²⁷ found a significant association between objective sleep disturbance and cognitive impairment in participants with BD ($p < 0.001$). Specifically,

participants with BD and abnormal sleep patterns were found to experience deficits in attention and executive function but normal verbal memory. In contrast, Boland et al.³² found no significant association between objective sleep disturbance and cognitive impairment in participants with BD. However, this study included a relatively small sample size of participants with BD (n=24) which implies low statistical power and risk of type-II error. Nevertheless, taken together, these two BD studies are inconclusive as to the association between objective sleep disturbance and cognitive impairment in people with BD and highlights the need for future research using larger sample sizes and PSG.

Limitations

This review reflects the limited number of studies in the field and so the relatively small number of participants in which the associations between sleep disturbance and cognitive impairment have been investigated, particularly regarding MDD, limits our power to draw wider conclusions. The studies included in our review generally had small sample sizes ($N = 48 - 4555$), which limited statistical power and increased risks of type-II error. A meta-analysis was not performed because there was significant heterogeneity between studies in the age and mood state of participants, as well as the measures of sleep and cognition used.

In studies examining sleep disturbance and cognitive impairment in mood disorders, it is normally best practice to include only euthymic participants because mood state may affect both sleep disturbance and cognitive impairment, and therefore may be a confound.⁵⁶ The MDD studies were particularly weak in this regard, as four of the five MDD studies only included people currently experiencing a depressive episode. The BD studies were stronger, but still varied: six studies only included participants that were euthymic,^{32,33,35,37,44,46} and two included participants in any mood state.^{27,34}

With the exception of Wilckens et al.⁴², none of the included studies adjusted for the use of psychotropic medications in the relationship between sleep disturbance and cognitive impairment. Psychotropic medication may be a confounding variable, as they can affect both sleep and cognition in people with mood disorders.⁵⁷ Indeed, Volkert et al.³⁷ reported that antipsychotic treatment and polypharmacy were related to cognitive impairment. Furthermore, many antipsychotics affect REM sleep, which

has been linked to emotional memory processing⁵⁸. Without adjusting for the use of psychotropic medications, it is unclear which medications alter the relationship between sleep disturbance and cognitive impairment in mood disorders, and to what extent.

Suggestions for future research

Theoretically, sleep disturbance may cause, predispose or exacerbate cognitive impairment in mood disorders but this link has yet to be adequately evaluated. Further research is clearly needed in this area in participants with mood disorders. Based on the evaluation of current evidence, we would suggest that future studies follow five recommendations. First, studies should include only euthymic participants, to remove the confounding effect of mood state. Second, they should use objective measures of sleep, ideally polysomnography, to evaluate the role of sleep disturbance in mediating cognitive impairment, and to determine which aspects of sleep architecture are linked to cognition in these individuals. Third, all studies should use a comprehensive cognitive battery such as the BACS-A⁵⁹ or ideally the ISBD-BANC⁶⁰. Fourth, analyses should adjust for the use of psychotropic medications, as these may also have a confounding effect. Finally, we would suggest that studies use a prospective longitudinal design, and, if possible, use a randomised clinical trial design for intervention studies.

Conclusion

In conclusion, our systematic review provides preliminary evidence for an association between sleep disturbance and cognitive impairment in mood disorders. The studies identified by this systematic review found increased rates of both subjective sleep disturbances and cognitive impairment in people with MDD and BD, compared to healthy participants. Ten of the twelve studies that measured sleep disturbance using *subjective* measures of sleep found a significant association between sleep disturbance and cognitive impairment across both diagnoses when adjusting for demographic and clinical covariates, whereas no significant association was found in healthy participants. In contrast, the evidence was scarcer and more conflicting with regards to whether *objectively* measured sleep abnormalities are associated with cognitive impairment in people with mood disorders: two out of three studies reported significant associations. Only one study³³ conducted an

intervention, and reported that improved sleep predicted an improvement in cognition in BD. As cognitive impairment has significant implications for psychosocial functioning in people with mood disorders, these findings highlight the importance of effectively identifying and treating sleep disturbances in people with MDD and BD.

Contributors

OP wrote the report with input from NUM, KWM, SAC, IR, AHY and PRAS. OP and NUM independently performed the literature search and quality assessment. All authors had full access to all the papers in the study and had final responsibility for the decision to submit for publication.

Declaration of interests

OP, NUM, SAC and IR declare no competing interests.

KWM has received consultancy fees from Lundbeck and Janssen-Cilag in the past three years.

PRAS reports non-financial support from Janssen Research and Development LLC, personal fees and non-financial support from Frontiers in Psychiatry, personal fees from Allergan, a grant from H Lundbeck, grants and non-financial support from Corcept Therapeutics, outside the submitted work.

AHY declares:

Employed by King's College London; Honorary Consultant SLaM (NHS UK)

Deputy Editor, BJPsych Open

Paid lectures and advisory boards for the following companies with drugs used in affective and related disorders:

Astrazenaca, Eli Lilly, Lundbeck, Sunovion, Servier, Livanova, Janssen, Allegan, Bionomics, Sumitomo Dainippon Pharma, COMPASS

Consultant to Johnson & Johnson

Consultant to Livanova

Received honoraria for attending advisory boards and presenting talks at meetings organised by LivaNova. Principal Investigator in the Restore-Life VNS registry study funded by LivaNova.

Principal Investigator on ESKETINTRD3004: "An Open-label, Long-term, Safety and Efficacy Study of Intranasal Esketamine in Treatment-resistant Depression."

Principal Investigator on "The Effects of Psilocybin on Cognitive Function in Healthy Participants"

Principal Investigator on "The Safety and Efficacy of Psilocybin in Participants with Treatment-Resistant Depression (P-TRD)"

UK Chief Investigator for Novartis MDD study MIJ821A12201

Grant funding (past and present): NIMH (USA); CIHR (Canada); NARSAD (USA); Stanley Medical Research Institute (USA); MRC (UK); Wellcome Trust (UK); Royal

College of Physicians (Edin); BMA (UK); UBC-VGH Foundation (Canada); WEDC (Canada); CCS Depression Research Fund (Canada); MSFHR (Canada); NIHR (UK). Janssen (UK)

No shareholdings in pharmaceutical companies

References

1. McIntyre RS, Cha DS, Soczynska JK, et al. Cognitive deficits and functional outcomes in major depressive disorder: determinants, substrates, and treatment interventions. *Depress Anxiety*. 2013;30(6):515-527. doi:10.1002/da.22063
2. O'Donnell LA, Deldin PJ, Grogan-Kaylor A, et al. Depression and executive functioning deficits predict poor occupational functioning in a large longitudinal sample with bipolar disorder. *J Affect Disord*. 2017;215:135-142. doi:10.1016/j.jad.2017.03.015
3. Martino DJ, Igoa A, Scápola M, Marengo E, Samamé C, Strejilevich SA. Functional Outcome in the Middle Course of Bipolar Disorder: A Longitudinal Study. *J Nerv Ment Dis*. 2017;205(3):203—206. doi:10.1097/nmd.0000000000000583
4. Zhang B-H, Feng L, Feng Y, et al. The Effect of Cognitive Impairment on the Prognosis of Major Depressive Disorder. *J Nerv Ment Dis*. 2020;208(9). https://journals.lww.com/jonmd/Fulltext/2020/09000/The_Effect_of_Cognitive_Impairment_on_the.8.aspx
5. Depp CA, Mausbach BT, Harmell AL, et al. Meta-analysis of the association between cognitive abilities and everyday functioning in bipolar disorder. *Bipolar Disord*. 2012;14(3):217-226.
6. Cambridge OR, Knight MJ, Mills N, Baune BT. The clinical relationship between cognitive impairment and psychosocial functioning in major depressive disorder: A systematic review. *Psychiatry Res*. 2018;269:157-171. doi:10.1016/j.psychres.2018.08.033
7. Rock P, Roiser J, Riedel W, Blackwell A. Cognitive impairment in depression: a systematic review and meta-analysis. *Psychol Med*. 2014;44(10):2029.
8. Snyder HR. Major depressive disorder is associated with broad impairments on neuropsychological measures of executive function: A meta-analysis and review. *Psychol Bull*. 2013;139(1):81-132. doi:10.1037/a0028727
9. Kurtz MM, Gerraty RT. A meta-analytic investigation of neurocognitive deficits in bipolar illness: Profile and effects of clinical state. *Neuropsychology*. 2009;23(5):551-562. doi:10.1037/a0016277
10. Semkovska M, Quinlivan L, O'Grady T, et al. Cognitive function following a major depressive episode: a systematic review and meta-analysis. *Lancet Psychiatry*. 2019;6(10):851-861. doi:10.1016/S2215-0366(19)30291-3
11. Bora E, Yucel M, Pantelis C. Cognitive endophenotypes of bipolar disorder: A meta-analysis of neuropsychological deficits in euthymic patients and their first-degree relatives. *J Affect Disord*. 2009;113(1-2):1-20. doi:10.1016/j.jad.2008.06.009
12. Pu S, Noda T, Setoyama S, Nakagome K. Empirical evidence for discrete neurocognitive subgroups in patients with non-psychotic major depressive disorder: clinical implications. *Psychol Med*. 2018;48(16):2717-2729. doi:10.1017/S003329171800034X

13. Burdick KE, Russo M, Frangou S, et al. Empirical evidence for discrete neurocognitive subgroups in bipolar disorder: clinical implications. *Psychol Med*. 2014;44(14):3083-3096. doi:10.1017/S0033291714000439
14. Jensen JH, Knorr U, Vinberg M, Kessing LV, Miskowiak KW. Discrete neurocognitive subgroups in fully or partially remitted bipolar disorder: Associations with functional abilities. *J Affect Disord*. 2016;205:378-386.
15. Depp CA, Dev S, Eyler LT. Bipolar Depression and Cognitive Impairment. *Psychiatr Clin North Am*. 2016;39(1):95-109. doi:10.1016/j.psc.2015.09.004
16. American Psychiatric Association, American Psychiatric Association, eds. *Diagnostic and Statistical Manual of Mental Disorders: DSM-5*. 5th ed. American Psychiatric Association; 2013.
17. Harvey AG, Schmidt DA, Scarnà A, Semler CN, Goodwin GM. Sleep-Related Functioning in Euthymic Patients With Bipolar Disorder, Patients With Insomnia, and Subjects Without Sleep Problems. *Am J Psychiatry*. 2005;162(1):50-57. doi:10.1176/appi.ajp.162.1.50
18. Iovieno N, van Nieuwenhuizen A, Clain A, Baer L, Nierenberg AA. Residual symptoms after remission of major depressive disorder with fluoxetine and risk of relapse. *Depress Anxiety*. 2011;28(2):137-144.
19. Nierenberg AA, Keefe BR, Leslie VC, et al. Residual symptoms in depressed patients who respond acutely to fluoxetine. *J Clin Psychiatry*. 1999;60(4):221-225.
20. Carney CE, Segal ZV, Edinger JD, Krystal AD. A comparison of rates of residual insomnia symptoms following pharmacotherapy or cognitive-behavioral therapy for major depressive disorder. *J Clin Psychiatry*. Published online 2007.
21. Sylvia LG, Dupuy JM, Ostacher MJ, et al. Sleep disturbance in euthymic bipolar patients. *J Psychopharmacol (Oxf)*. 2012;26(8):1108-1112. doi:10.1177/0269881111421973
22. Li SX, Lam SP, Chan JWY, Yu MWM, Wing Y-K. Residual Sleep Disturbances in Patients Remitted From Major Depressive Disorder: A 4-Year Naturalistic Follow-up Study. *Sleep*. 2012;35(8):1153-1161. doi:10.5665/sleep.2008
23. Jackson A, Cavanagh J, Scott J. A systematic review of manic and depressive prodromes. *J Affect Disord*. 2003;74(3):209-217.
24. Krause AJ, Simon EB, Mander BA, et al. The sleep-deprived human brain. *Nat Rev Neurosci*. 2017;18(7):404-418. doi:10.1038/nrn.2017.55
25. Ashton JE, Harrington MO, Langthorne D, Ngo H-VV, Cairney SA. Sleep deprivation induces fragmented memory loss. *Learn Mem*. 2020;27(4):130-135.
26. Harrington MO, Ashton JE, Sankarasubramanian S, Anderson MC, Cairney SA. Losing Control: Sleep Deprivation Impairs the Suppression of Unwanted Thoughts. Published online 2021:17.

27. Bradley AJ, Anderson KN, Gallagher P, McAllister-Williams RH. The association between sleep and cognitive abnormalities in bipolar disorder. *Psychol Med*. 2020;50(1):125-132. doi:10.1017/S0033291718004038
28. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. Published online March 29, 2021:n71. doi:10.1136/bmj.n71
29. World Health Organization. International Classification of Diseases—Ninth Revision (ICD-9). *Wkly Epidemiol Rec*. 1988;63(45):343-344.
30. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (3rd Ed.)*; 1980.
31. Thomas H, Ciliska D, Dobbins M. Quality assessment tool for quantitative studies. *Tor Eff Public Health Pract Proj McMaster Univ*. Published online 2003.
32. Boland EM, Stange JP, Molz Adams A, et al. Associations between sleep disturbance, cognitive functioning and work disability in Bipolar Disorder. *Psychiatry Res*. 2015;230(2):567-574. doi:10.1016/j.psychres.2015.09.051
33. Kanady JC, Soehner AM, Klein AB, Harvey AG. The association between insomnia-related sleep disruptions and cognitive dysfunction during the inter-episode phase of bipolar disorder. *J Psychiatr Res*. 2017;88:80-88. doi:10.1016/j.jpsychires.2017.01.001
34. Laskemoen JF, Büchmann C, Barrett EA, et al. Do sleep disturbances contribute to cognitive impairments in schizophrenia spectrum and bipolar disorders? *Eur Arch Psychiatry Clin Neurosci*. Published online October 5, 2019. doi:10.1007/s00406-019-01075-0
35. Samalin L, Boyer L, Murru A, et al. Residual depressive symptoms, sleep disturbance and perceived cognitive impairment as determinants of functioning in patients with bipolar disorder. *J Affect Disord*. 2017;210:280-286. doi:10.1016/j.jad.2016.12.054
36. Ullah MH. Effects of sleep disturbance on cognitive functioning in bipolar disorder type 1: a correlational study design. *ProQuest Diss Publ*. Published online 2017.
37. Volkert J, Kopf J, Kazmaier J, et al. Evidence for cognitive subgroups in bipolar disorder and the influence of subclinical depression and sleep disturbances. *Eur Neuropsychopharmacol*. 2015;25(2):192-202. doi:10.1016/j.euroneuro.2014.07.017
38. Cabanel N, Schmidt A-M, Fockenberg S, et al. Evening preference and poor sleep independently affect attentional-executive functions in patients with depression. *Psychiatry Res*. 2019;281:112533. doi:10.1016/j.psychres.2019.112533
39. Müller MJ, Olschinski C, Kundermann B, Cabanel N. Sleep duration of inpatients with a depressive disorder: associations with age, subjective sleep quality, and cognitive complaints. *Arch Psychiatr Nurs*. 2017;31(1):77-82. doi:10.1016/j.apnu.2016.08.008
40. Peng S, Li J, Wu Z, et al. Characteristics of and factors influencing cognitive impairment in adolescent patients with first-episode depression. *Vulnerable Child Youth Stud*. 2018;13(4):317-324. doi:10.1080/17450128.2018.1515517

41. Cha DS, Carmona N, Cha RH, et al. Perceived sleep quality predicts cognitive function in adults with major depressive disorder independent of depression severity. *Ann Clin PSYCHIATRY*. Published online 2019:10.
42. Wilckens KA, Kline CE, Bowman MA, et al. Does objectively-assessed sleep moderate the association between history of major depressive disorder and task-switching? *J Affect Disord*. 2020;265:216-223. doi:10.1016/j.jad.2020.01.003
43. Soehner AM, Harvey AG. Prevalence and Functional Consequences of Severe Insomnia Symptoms in Mood and Anxiety Disorders: Results from a Nationally Representative Sample. *Sleep*. 2012;35(10):1367-1375. doi:10.5665/sleep.2116
44. Russo M, Mahon K, Shanahan M, et al. The relationship between sleep quality and neurocognition in bipolar disorder. *J Affect Disord*. 2015;187:156-162. doi:10.1016/j.jad.2015.08.009
45. Harvey AG, Soehner AM, Kaplan KA, et al. Treating insomnia improves mood state, sleep, and functioning in bipolar disorder: A pilot randomized controlled trial. *J Consult Clin Psychol*. 2015;83(3):564-577. doi:10.1037/a0038655
46. Samalin L, de Chazeron I, Vieta E, Bellivier F, Llorca P-M. Residual symptoms and specific functional impairments in euthymic patients with bipolar disorder. *Bipolar Disord*. 2016;18(2):164-173. doi:10.1111/bdi.12376
47. White DA, Myerson J, Hale S. How cognitive is psychomotor slowing in depression? evidence from a meta-analysis. *Aging Neuropsychol Cogn*. 1997;4(3):166-174. doi:10.1080/13825589708256645
48. Buysse DJ, Reynolds III CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res*. 1989;28(2):193-213.
49. Wysokinski A, Dzienniak M, Kloszewska I. Assessment of cognitive performance using CNS vital signs after electroconvulsive treatment of schizophrenia. *J ECT*. 2014;30(1):e5-e6.
50. Carney CE, Harris AL, Friedman J, Segal ZV. Residual sleep beliefs and sleep disturbance following cognitive behavioral therapy for major depression. *Depress Anxiety*. 2011;28(6):464-470.
51. Morin C, Blais F, Savard J. Are changes in beliefs and attitudes about sleep related to sleep improvements in the treatment of insomnia? *Behav Res Ther*. 2002;40(7):741-752.
52. Hutka P, Krivosova M, Muchova Z, et al. Association of Sleep Architecture and Physiology with Depressive Disorder and Antidepressants Treatment. *Int J Mol Sci*. 2021;22(3):1333. doi:10.3390/ijms22031333
53. Eidelman P, Talbot LS, Gruber J, Hairston I, Harvey AG. Sleep architecture as correlate and predictor of symptoms and impairment in inter-episode bipolar disorder: taking on the challenge of medication effects: Sleep and inter-episode bipolar disorder. *J Sleep Res*. 2010;19(4):516-524. doi:10.1111/j.1365-2869.2010.00826.x

54. della Monica C, Johnsen S, Atzori G, Groeger JA, Dijk D-J. Rapid Eye Movement Sleep, Sleep Continuity and Slow Wave Sleep as Predictors of Cognition, Mood, and Subjective Sleep Quality in Healthy Men and Women, Aged 20–84 Years. *Front Psychiatry*. 2018;9:255. doi:10.3389/fpsyt.2018.00255
55. Ngo H-VV, Martinetz T, Born J, Mölle M. Auditory closed-loop stimulation of the sleep slow oscillation enhances memory. *Neuron*. 2013;78(3):545-553.
56. Miskowiak K, Burdick K, Martinez-Aran A, et al. Assessing and addressing cognitive impairment in bipolar disorder: the International Society for Bipolar Disorders Targeting Cognition Task Force recommendations for clinicians. *Bipolar Disord*. 2018;20(3):184-194. doi:10.1111/bdi.12595
57. Staner L. Sleep disturbances, psychiatric disorders, and psychotropic drugs. *Dialogues Clin Neurosci*. 2005;7(4):323.
58. Walker MP, van Der Helm E. Overnight therapy? The role of sleep in emotional brain processing. *Psychol Bull*. 2009;135(5):731.
59. Keefe RS, Goldberg TE, Harvey PD, Gold JM, Poe MP, Coughenour L. The Brief Assessment of Cognition in Schizophrenia: reliability, sensitivity, and comparison with a standard neurocognitive battery. *Schizophr Res*. 2004;68(2-3):283-297.
60. Yatham LN, Torres IJ, Malhi GS, et al. The international society for bipolar disorders—battery for assessment of neurocognition (ISBD□BANC). *Bipolar Disord*. 2010;12(4):351-363.
61. Fietze I, Wiesenacker D, Blau A, Penzel T. Sleep quality in a hospital and effects of noise. *Somnologie-Schlafforschung Schlafmed*. 2008;12(2):167.
62. Wechsler D. Wechsler Test of Adult Reading: WTAR: Psychological Corporation. *San Antonio Tx*. Published online 2001.
63. Wechsler D. *Wechsler Memory Scale (WMS-III)*. Vol 14. Psychological corporation San Antonio, TX; 1997.
64. Morin CM. *Insomnia: Psychological Assessment and Management*. Guilford press; 1993.
65. Rush AJ, Gullion CM, Basco MR, Jarrett RB, Trivedi MH. The inventory of depressive symptomatology (IDS): psychometric properties. *Psychol Med*. 1996;26(3):477-486.
66. Endicott J, Spitzer R. Schedule for affective disorders and schizophrenia (SADS). *Acta Psychiatr Belg*. 1987;87(4):361-516.
67. Montgomery SA, Åsberg M. A new depression scale designed to be sensitive to change. *Br J Psychiatry*. 1979;134(4):382-389.
68. Kaufman AS. Manual for the Kaufman brief intelligence test. *Circ Pines MN Am Guid Serv*. Published online 1990.

69. Delis DC, Kramer JH, Kaplan E, Holdnack J. Reliability and validity of the Delis-Kaplan Executive Function System: an update. *J Int Neuropsychol Soc JINS*. 2004;10(2):301.
70. Shallice T. Specific impairments of planning. *Philos Trans R Soc Lond B Biol Sci*. 1982;298(1089):199-209.
71. Niemann H, Sturm W, Thöne-Otto A, Wilmes K. California verbal learning test, version 21. *Mödling Schuhfried GmbH*. Published online 2011.
72. Woods SP, Delis DC, Scott JC, Kramer JH, Holdnack JA. The California Verbal Learning Test—second edition: test-retest reliability, practice effects, and reliable change indices for the standard and alternate forms. *Arch Clin Neuropsychol*. 2006;21(5):413-420.
73. Nelson HE. *National Adult Reading Test (NART): For the Assessment of Premorbid Intelligence in Patients with Dementia: Test Manual*. Nfer-Nelson; 1982.
74. Dinges DF, Powell JW. Microcomputer analyses of performance on a portable, simple visual RT task during sustained operations. *Behav Res Methods Instrum Comput*. 1985;17(6):652-655.
75. Fan J, McCandliss BD, Sommer T, Raz A, Posner MI. Testing the efficiency and independence of attentional networks. *J Cogn Neurosci*. 2002;14(3):340-347.
76. Pariante CM, Alhaj HA, Arulnathan VE, et al. Central glucocorticoid receptor-mediated effects of the antidepressant, citalopram, in humans: a study using EEG and cognitive testing. *Psychoneuroendocrinology*. 2012;37(5):618-628.
77. Samalin L, Llorca PM, Giordana B, et al. Residual symptoms and functional performance in a large sample of euthymic bipolar patients in France (the OPTHYUM study). *J Affect Disord*. 2014;159:94-102.
78. Rosa AR, Sánchez-Moreno J, Martínez-Aran A, et al. Validity and reliability of the Functioning Assessment Short Test (FAST) in bipolar disorder. *Clin Pract Epidemiol Ment Health*. 2007;3(1):5.
79. Glahn DC, Almasy L, Barguil M, et al. Neurocognitive endophenotypes for bipolar disorder identified in multiplex multigenerational families. *Arch Gen Psychiatry*. 2010;67(2):168-177.
80. Zimmermann P, Fimm B. A test battery for attentional performance. *Appl Neuropsychol Atten Theory Diagn Rehabil*. 2002;110:151.
81. Aschenbrenner S, Tucha O, Lange K. Manual zum RWT (Regensburger Wortflüssigkeits-Test; Handanweisung). Published online 2000.
82. Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *sleep*. 1991;14(6):540-545.

Tables

Table 1. Characteristics and findings of studies examining sleep disturbance and cognitive impairment in MDD

Study	Mood disorder(s) studied	Participants	Measures	Main findings	Limitations
Cabanel et al.	MDD	Total N = 63 N = 34 MDD participants (27 with single and 7 with recurrent depression; 9 with comorbid anxiety disorders), mean age 33.0 ± 8.9 N = 29 healthy participants, mean age 31.5 ± 7.8	Sleep: PSQI Cognition: TMT-A, TMT-B	Patients with severely impaired sleep quality (PSQI > 10) had significantly poorer performance on the TMT-A (visuomotor processing speed; $p = 0.011$) and the TMT-B (executive function; $p = 0.015$), and remained significant after Bonferroni correction ($\alpha = 0.025$ per tests), than patients with no or moderately impaired sleep quality (PSQI > 10).	The use of subjective measure of sleep instead of objective sleep battery. Neuropsychological assessment only evaluated a subdomain of executive functions. Particularly small sample size.

Cha et al.	MDD	Total N = 200, aged 18 to 65 N = 100 participants with recurrent MDD N = 100 sex-, age-, education-matched healthy participants	Sleep: PSQI Cognition: computerised cognitive assessment tool THINC-integrated tool that included CRT, the One Back Memory task, DSST, TMT-B, PDQ-5-D.	Sleep quality predicted impairments in subjective cognitive performance ($p < 0.001$) and objective cognitive impairments ($p = 0.017$). Depression severity was a partial mediator of the relationship between subjective cognitive function and sleep quality ($B = -0.43$; 95% CI, -0.56, -0.33). After controlling for the depression severity, the direct effect of sleep quality on subjective cognition performance remained significant ($B = -0.18$; 95% CI, -0.30, -0.07).	The use of subjective measure of sleep instead of objective sleep battery. Complete case analysis of cognitive tests resulting in low statistical power.
Wilckens et al.	MDD	Total N=53, aged 48 to 79 years of age N=17 participants with recurrent	Sleep: two nights of PSG Cognition: computer-based task-switching paradigm	Higher wake after sleep onset and shorter total sleep time were linearly associated with slower response time across single-task ($p = 0.10$) and switching blocks ($p = 0.005$); however, these associations were not specific to MDD.	Participants with history of MDD were more likely to take antidepressants at the time of the

		MDD N=36 participants with no recurrent MDD N=28 participants with no lifetime MDD N=25 participants with lifetime MDD			study, thus possibly affecting sleep and/or cognition. Did not exclude ages >65. Particularly small sample size.
Mueller et al. ²⁹	MDD	Total N = 154, aged 19-89 N = 154 MDD participants	Sleep: SQH Cognition: SQH (items pertaining to daytime attention deficits and memory	25-40% of participants with MDD reported 'almost always' daytime sleepiness, non-restorative sleep, attention deficits or memory complaints, with significant correlations between these variables ($p < 0.05$), as well as curvilinear associations between sleep duration and sleep quality (quadratic contrast, $p < 0.05$).	The use of subjective measure of sleep instead of objective sleep battery. Did not use a validated measure of

			complaints only)		cognition. No control group.
Peng et al. ³¹	MDD	Total N = 178, aged 15–30 N = 128 first-episode MDD participants N = 50 healthy participants	Sleep: PSQI Cognition: MCCB	No significant association between cognition and sleep disturbance ($r = 0.029$, $p = 0.219$) in adolescents with first episode MDD.	The use of subjective measure of sleep instead of objective sleep battery. Included participants aged <18.
Boland et al. ³²	BD	Total N = 48, aged 18-65 N = 24 euthymic BD participants N = 24 healthy participants	Sleep: ISI, UCII, PSQI, wrist actigraphy Cognition: KBIT-II, DKEFS (Stroop subtest only), Tower of London, CVLT-II, WMS-III (digit span subtest only)	No significant associations between sleep disturbance (including objective measures) and cognition, except for a significant negative correlation between daytime dysfunction due to sleep disturbance and working memory ($r = -0.57$, $p = 0.03$).	Particularly small sample size.

Bradley et al. ²⁴	BD	Total N = 88, aged 18-65 N = 46 BD participants, in any mood state N = 42 healthy control participants	Sleep: wrist actigraphy Cognitive assessment: NART, PVT, ANT, DSST, NSMT	People with BD with normal sleep patterns were not significantly different to healthy participants in cognitive functions, but people with BD and sleep disturbance (including objective measures) performed significantly worse in cognitive tests than healthy participants ($p < 0.001$).	Did not adjust for the mood state of participants. Particularly small sample size.
Kanady et al. ³³	BD	Total N = 66, aged 18-65 47 euthymic BD participants with comorbid insomnia 19 BD participants without sleep disturbance in the last six	Sleep: ISI, PSQI, DSISD, sleep diary Cognition: N-Back Task and WMS-III (digit span subtest only)	Greater sleep time variability predicted poorer working memory ($p < 0.01$) and verbal learning performance ($p < 0.01$) in BD. Furthermore, following treatment of cognitive behaviour therapy for insomnia modified for bipolar disorder (CBTI-BD ³⁸), a reduction in total wake time predicted improved working memory ($p < 0.05$) and a reduction in sleep time variability predicted improved verbal learning ($p < 0.05$).	No healthy control group. Particularly small sample size.

		months			
Laskemoen et al. ³⁴	BD	Total N = 979, aged 18-65 N = 340 BD participants, in any mood state N = 457 schizophrenia participants N = 182 healthy control participants	Sleep: IDS-C (items pertaining to sleep disturbance) Cognition: MCCB	Significant associations between insomnia and processing speed ($p < 0.001$), insomnia and inhibition ($p < 0.01$), hypersomnia and processing speed ($p < 0.01$), and hypersomnia and inhibition ($p < 0.05$) in people with BD, whereas no association between sleep disturbances and cognition was found in healthy participants.	The use of subjective measure of sleep instead of objective sleep battery. Did not adjust for the mood state of participants.
Russo et al.	BD	Total N = 117, aged 18-65 N = 117 euthymic BD participants	Sleep: ESS, PSQI Cognition: MCCB	Sleep disturbance was significantly negatively associated with social cognition ($r = -0.273$, $p = 0.004$), and approached significance with working memory ($r = -0.179$, $p = 0.061$) and visual learning ($r = -0.161$, $p = 0.094$). Path analyses indicated that social cognition and working memory were directly (negatively) predicted by sleep disruptions.	No control group. Only used subjective measures of sleep.

Samalin et al. ³⁵	BD	Total N = 468, aged >18 N = 468 euthymic BD participants	Sleep: PSQI Cognition: VAS and FAST	Sleep disturbances were indirectly associated with functioning (including cognition) via residual depressive symptoms and perceived cognitive performance ($p = 0.001$).	The use of subjective measures of cognitive functioning and sleep instead of objective neuropsychological and sleep batteries. Did not exclude ages >65. No control group.
Samalin et al.	BD	Total N = 468, mean age 47.7 ± 12.5 N=468 adult outpatients with bipolar I disorder or bipolar II disorder	Sleep: PSQI Cognition: VAS and FAST	Participants with impaired cognitive functioning had significantly higher scores ($p < 0.02$), and thus more severe sleep disruptions, on PSQI than participants with satisfactory cognitive functioning.	The use of subjective measures of cognitive functioning and sleep instead of objective neuropsychological and sleep

		in an euthymic period			batteries. Did not exclude ages >65. No control group.
Volkert et al. ³⁷	BD	Total N = 140, aged 18–65 N = 70 euthymic BD participants N = 70 healthy participants	Sleep: MADRS (item pertaining to sleep disturbance) Cognition: TAP, CVLT, TOL, RWT	People with BD suffering from persistent sleep disturbances show more severe cognitive dysfunctions ($p < 0.05$).	Did not use a validated measure of sleep disturbance.
Soehner and Harvey ⁴³	BD & MDD	Total N = 5,692, aged >18 N = 3,711 participants with no mood or anxiety disorder	Sleep: series of self-report questions Cognition: items from WHO-DAS pertaining to cognition	Participants with a mood disorder and any insomnia symptom in the past year had significantly greater 30-day cognitive impairment than participants with a mood disorder and no insomnia symptoms in the past year ($p = 0.004$).	Subjective measures of sleep and cognition.

		N = 327 participant s with mood disorders only			
		N = 1,137 participa nts with anxiety disorders only			
		N = 517 participant s with comorbid mood and anxiety disorders			

Abbreviations: TMT, Trail Making Test part A and B; CRT: Choice Reaction Time; DSST, Digit Symbol Substitution Test; PDQ-5-D, Perceived Deficits Questionnaire- 5-item for Depression; PSG, polysomnography; MCI, mild cognitive impairment; SQH, Sleep Quality in Hospital settings in German language⁶¹; PSQI, Pittsburgh Sleep Quality Index⁴⁸; WTAR, Wechsler Test of Adult Reading⁶²; WMS-III, Wechsler Memory Scale-III⁶³; MCCB, MATRICS Consensus Cognitive Battery⁴⁹; ISI, Insomnia Severity Index⁶⁴; DSISD, Duke Structured Interview for Sleep Disorder¹⁶; IDS-C, Inventory of Depressive Symptoms — Clinician rated scale⁶⁵; SADS-C, Schedule for Affective Disorders and Schizophrenia-Change⁶⁶; MADRS, Montgomery–Asperg Depression Ratings Scale⁶⁷; UCII, Unstructured Clinical Interview for Insomnia; KBIT-II, Kauffman Brief Intelligence Test Second Edition⁶⁸;

DKEFS, Delis-Kaplan Executive Functioning System⁶⁹; TOL, Tower of London⁷⁰; CVLT, California Verbal Learning Test⁷¹; CVLT-II, California Verbal Learning Test Second Edition⁷²; NART, National Adult Reading Test⁷³; PVT, Psychomotor vigilance Test⁷⁴; ANT, Attention Network Test⁷⁵; DSST, Digit Symbol Substitution Test⁶³; NSMT, Newcastle Spatial Memory Task⁷⁶; VAS, visual analogic scale⁷⁷; FAST, Functioning Assessment Short Test⁶⁰; STAN, South Texas Assessment of Neurocognition⁷⁹; TAP, Test battery of Attentional Performance⁸⁰; RWT, Regensburger Wortflüssigkeits Test⁸¹; ESS, Epworth Sleepiness Scale⁸²; WHO-DAS, World Health Organization Disability Assessment Schedule-II.

Supplementary Tables

Supplementary Table 1: Methodological evaluation of studies using the EPHPP Quality Assessment Tool for Quantitative Studies

Author and year	Mood disorder(s) studied	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdrawals and dropouts	Global rating
Soehner and Harvey, 2012	MDD and BD	Moderate	Weak	Strong	Moderate	Moderate	N/A	Moderate
Cabanel et al., 2019	MDD	Moderate	Weak	Strong	Moderate	Strong	N/A	Moderate
Cha et al., 2019	MDD	Moderate	Weak	Strong	Moderate	Strong	Strong	Moderate
Muller et al., 2017	MDD	Moderate	Weak	Strong	Moderate	Weak	N/A	Weak
Peng et al., 2018	MDD	Moderate	Weak	Strong	Moderate	Strong	N/A	Moderate
Wilckens et al., 2020	MDD	Moderate	Weak	Strong	Moderate	Strong	N/A	Moderate
Boland et al., 2015	BD	Moderate	Moderate	Strong	Strong	Strong	Weak	Moderate
Bradley et al., 2020	BD	Strong	Moderate	Strong	Moderate	Strong	Strong	Strong
Kanady et al., 2017	BD	Moderate	Strong	Strong	Moderate	Strong	Strong	Strong
Laskemoen et al., 2020	BD	Strong	Weak	Strong	Moderate	Moderate	N/A	Moderate
Russo et al., 2015	BD	Moderate	Weak	Moderate	Moderate	Strong	N/A	Moderate
Samalin et al., 2017	BD	Strong	Weak	Weak	Moderate	Strong	N/A	Weak
Samalin et al., 2016	BD	Strong	Weak	Weak	Moderate	Strong	N/A	Weak

Volkert et al., 2015	BD	Moderate	Weak	Strong	Moderate	Moderate	N/A	Moderate
----------------------	----	----------	------	--------	----------	----------	-----	-----------------

Figure 1. PRISMA 2020 Flow Diagram of study selection process