1 Prevalence of cognitive impairment following chemotherapy treatment 2 for breast cancer: A systematic review and meta-analysis

3

4 Alexandra L Whittaker ^{a,b*}, Rebecca P George ^a, Lucy O'Malley ^c

5 a School of Animal and Veterinary Sciences, The University of Adelaide, Roseworthy Campus,
6 Roseworthy, South Australia 5371, Australia

7 b Faculty of Biology, Medicine and Health, University of Manchester, Manchester, UK.

8 c Division of Dentistry, School of Medical Sciences, Manchester Academic Health Sciences
9 Centre, University of Manchester, Manchester, UK.

10 * Corresponding Author email: alexandra.whittaker@adelaide.edu.au,

11 Short Title: Prevalence of cognitive impairment in breast cancer survivors

12 MeSH key words: Chemotherapy-Related Cognitive Impairment; Psycho-Oncology; Breast
13 Neoplasms; Chemotherapy, Adjuvant

14

15 Abstract

16 Breast cancer survival rates have markedly improved. Consequently, survivorship issues
17 have received increased attention. One common sequela of treatment is chemotherapy-
18 induced cognitive impairment (CICI). CICI causes a range of impairments that can have
19 a significant negative impact on quality of life. Knowledge of the prevalence of this
20 condition is required to inform survivorship plans, and ensure adequate resource
21 allocation and support is available for sufferers.

22 Objective

23 To estimate the prevalence of cognitive impairment following chemotherapy
24 treatment for breast cancer.

25 Methods

26 Medline, Scopus, CINAHL and PSYCHInfo were searched for
27 eligible studies which included prevalence data on CICI, as ascertained through the use of
28 self-report, or neuropsychological tests. Methodological quality of included studies was
29 assessed. Findings were synthesised narratively, with meta-analyses being used to
30 calculate pooled prevalence when impairment was assessed by neuropsychological
31 tests.

32

33 **Results and discussion**

34 The review included 52 studies. Time-points considered ranged from the chemotherapy
35 treatment period to greater than 10 years after treatment cessation. Summary prevalence
36 figures (across time-points) using self-report, short cognitive screening tools and
37 neuropsychological test batteries were 44%, 16% and 21-34% respectively (very low
38 GRADE evidence).

39 **Conclusion**

40 Synthesised findings demonstrate that 1 in 3 breast cancer survivors may have
41 clinically significant cognitive impairment. Prevalence is higher when self-report based
42 on patient experience is considered. This review highlights a number of study design
43 issues that may have contributed to the low certainty rating of the evidence. Future
44 studies should take a more consistent approach to the criteria used to assess impairment.
45 Larger studies are urgently needed.

46
47 **Keywords:** breast cancer, chemobrain, chemotherapy-induced cognitive impairment,
48 prevalence, systematic review

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63 Summary of Findings Table

Outcomes	N ₂ of participants (studies) Follow up	Certainty of the evidence (GRADE)	Impact
Prevalence of cognitive impairment diagnosed by self-report measures (Self-report)	3817 (15 observational studies) ^a	⊕○○○ VERY LOW ^{b,c}	Prevalence rates for global impairment varied from 21-83% (mean 44%) across time-points ranging from during chemotherapy treatment to greater than 10 years after cessation of treatment.
Prevalence of cognitive impairment diagnosed by short cognitive screening tools (Short Cognitive Screening)	1533 (9 observational studies)	⊕○○○ VERY LOW ^{b,c}	Prevalence rates for global impairment varied from 0-50% (mean 16%) across time-points ranging from during chemotherapy treatment to 2-3 years after cessation of treatment.
Prevalence of cognitive impairment diagnosed by objective neuropsychological testing (Objective Neuropsychological Test)	1938 (35 observational studies) ^d	⊕○○○ VERY LOW ^{c,e,f}	The overall random-effects pooled prevalence (95% CIs) of cognitive impairment from time points which ranged from during chemotherapy treatment, to 2-3 years after cessation of chemotherapy (T1-T5) respectively was: 34% (24-44), 30% (20-40), 23% (16-31), 31% (14-50), and 21% (15-28). Late term prevalence at 5-10 years, and greater than 10 years after cessation of treatment, was 13% (5-32) and 8% (2-26) respectively (non-pooled data with only one study contributing to each value).

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval

GRADE Working Group grades of evidence

High certainty: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate certainty: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low certainty: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Explanations

a. 14 observational studies, 1 RCT

b. Point estimates varied widely across studies

c. Confidence intervals are wide

d. 30 observational studies, 5 RCT

e. Heterogeneity as determined by I² universally high

f. Tests for publication bias show evidence of bias at a number of time-points

64

65 Background

66 In women, breast cancer is the most frequently diagnosed cancer, with an estimated 2.3
67 million new cases globally per year. [1] Prognosis has significantly improved over time,
68 with current 5-year survival rates of around 90% and 10-year survival at 80%.[1] This
69 improvement has likely been brought about largely through the ability to characterise
70 cancer subtypes, enabling development of targeted agents, increased use of personalised
71 medicine and consequently improved treatment success. [1] Concurrently, improvements
72 in surgical resection methods have also been seen. [2] As a result, the number of breast
73 cancer patients living beyond a diagnosis has grown significantly, and survivorship
74 issues are receiving more attention, with a particular focus on quality of life issues.

75 Breast cancer treatment generally uses a multi-modal treatment approach utilising
76 combinations of surgery, chemotherapy, endocrine therapy and radiotherapy, dependant
77 on disease stage and sub-type classification. [3] However, chemotherapy combined with
78 surgery, especially in advanced stages of the disease, forms the mainstay of therapy.
79 Chemotherapy has been implicated as a significant contributor to cognitive impairment

80 observed in cancer patients, both during and after cessation of treatment. The resultant
81 condition has been termed chemotherapy-induced cognitive impairment (CICI), or
82 colloquially, chemofog or chemobrain. [4] Whilst the exact mechanisms for this are
83 unclear, proposed mechanisms include direct neurotoxic injury, a reduction in
84 neurogenesis (new neuron formation), central nervous system white matter abnormalities,
85 and neuroinflammation. [5] Most of the evidence for these theories comes from the pre-
86 clinical literature, [5-7] or neuro-imaging studies [8, 9].

87 The most common neuropsychological effects that arise in CICI are impairments to
88 visual processing, visual motor function, executive function and attention. [10] Sufferers
89 typically report side effects with a wide range of severity, from subtle to more severe
90 impairment, which may persist for up to 20 years post-chemotherapy treatment. [11]
91 Impairments typically manifest as survivors feeling ‘less-sharp’, being unable to recall
92 words, having episodes of mental confusion, and requiring more mental effort to perform
93 everyday activities. [12] [10] As a result, CICI has the potential to significantly impact
94 patient quality of life. Furthermore, it has been suggested that a lack of prior information,
95 coupled with minimal validation and understanding from family and health care
96 providers leads to patients feeling disempowered, and subsequently not receiving the
97 emotional and rehabilitative support they may need. [13]

98
99 The reported prevalence of CICI following treatment for breast cancer is somewhat
100 variable. A commonly cited range from the narrative review of Janelins et al. 2014
101 suggests that 12 – 82% of breast cancer patients will experience impairment as a result of
102 their treatment regime. [14] This variability may be attributable to a range of patient
103 factors such as age, [15] IQ, [16] menopausal status, [16] and education level. [17]
104 Alternately, study design factors such as the employment of cross-sectional, versus
105 longitudinal study designs, with the latter able to account for baseline pre-treatment
106 cognitive function, likely influences rates. [18] Furthermore, in spite of the name, it has
107 been shown that cognitive impairment is often present before chemotherapy treatment
108 has commenced, potentially arising as a result of cancer itself. [19, 20] This may occur
109 via direct effects of the tumour itself, as a result of associated co-morbidities, or due to
110 psychological factors such as worry and fatigue. [21] Time since treatment, also likely
111 contributes to the variance in prevalence reported. Neuroimaging has demonstrated
112 structural and functional changes in various patient brain regions, which likely correlate

113 with the corresponding changes seen in executive function and memory. However, these
114 alterations also show partial recovery over time, [22, 23] presumably leading to an
115 improvement in symptoms. Finally, prevalence rates may be influenced by the method
116 used to assess cognitive impairment, with the three common methods employed being the
117 ‘gold-standard’ neuropsychological testing, [24] short cognitive screening tools [25, 26]
118 and self-report. [14]

119 A preliminary database search for previous systematic reviews on prevalence of CICI
120 was conducted. One recent systematic review was sourced which reports prevalence data
121 for cognitive impairment in breast cancer patients. [3] This review is limited in scope
122 compared to the current review since the timeline of consideration was the breast cancer
123 treatment period, rather than longer-term follow up points. The review authors also only
124 included longitudinal studies, which utilized objective neuropsychological tests. Whilst
125 the reasons for these inclusions are understandable, the present review expands on this
126 knowledge base by evaluating both self-report and objective psychological testing to
127 compare prevalence rates of CICI ascertained by these different testing modalities. In the
128 current review all study designs were eligible, not just longitudinal designs. This broad
129 inclusion does not allow the teasing apart of the contribution of chemotherapy, as
130 opposed to cancer itself, on impairment. However, from the point of view of the patient
131 suffering cognitive difficulties, or health-care providers and policy makers needing
132 evidence-based information, this distinction is of little consequence; external validity of
133 the findings is assured. Understanding the burden of CICI will help inform survivorship
134 strategies, and drive funding priorities for future research and targeted support, such as
135 rehabilitation options. Furthermore, by identifying CICI incidence over the longer term
136 this will indicate the need for longer-term support to inform cancer survivorship care
137 plans.

138 *Review Question*

139 The aim of this review was to synthesise the evidence on prevalence rates of cognitive
140 impairment following chemotherapy treatment in breast cancer survivors, taking into
141 consideration factors such as age and time since cessation of treatment.

142 **Methods**

143 The Joanna Briggs Institute guidelines on conducting prevalence reviews [27] and the
144 PRISMA guidelines [28] were used to guide review performance and reporting. The

objectives, inclusion criteria and methods of analysis for this review were specified in advance and documented in a protocol registered on PROSPERO; ref CRD42021228541.

Search Strategy

The search strategy aimed to locate published studies in English. An initial search of Medline was undertaken to identify articles on the topic. Keywords used in the titles and abstracts of relevant articles, and the index terms used to describe the articles were used to develop a full search strategy for Medline via Pubmed using MeSH and free text terms. The four databases were searched in December 2020 using the developed search strategies (see supplementary material S1). Key concepts used for searching were “cancer”, “chemotherapy” and “cognitive impairment”. Hand searching of reference lists was performed to identify additional studies. Studies published from database inception were eligible for inclusion. Publications were excluded if they were conference abstracts, review articles and grey literature.

Study Selection

Following the search, all identified citations were uploaded into EndNote X8.0.1 and duplicates removed. Potentially relevant studies were retrieved in full and their citation details imported into Covidence (Veritas Health Innovation, Melbourne, Australia). Title, abstract and full text screening for assessment against the inclusion criteria for the review (see below) was performed by one reviewer (A.W), with checking of the information performed by a second reviewer (R.P.G).

Eligibility Criteria

Population

Females of any age treated with systemic chemotherapeutic agents for any form of breast cancer were considered for inclusion. Patients currently in treatment or that had ceased treatment (remission) were eligible for inclusion. Studies with patients receiving other concurrent drug therapy or with other co-morbidities were eligible for inclusion with noting (see exclusion criteria). Patients of any socioeconomic status or level of education were eligible for inclusion with noting of any differences reported. Patients that were either pre-or post-menopausal were eligible for inclusion with noting.

The following groups of patients were excluded:

- 178 • patients with previous history of traumatic brain injury, neurodegenerative
- 179 conditions such as Parkinson's or Alzheimer's or severe depressive symptoms,
- 180 since these conditions affect cognition either directly or indirectly. [29]
- 181 • patients who received palliative care (due to differences in the treatment pathways
- 182 and treatment regimens).[30]
- 183 • patients with metastatic CNS cancers or who have previously received CNS
- 184 targeted therapy for other conditions due to the risk of direct effect on cognition
- 185 as a result of the cancer/treatment [31, 32]

186

187 *Intervention*

188 Studies evaluating patients administered chemotherapy for breast cancer treatment were
 189 included. This included a range of common chemotherapy regimens using either sole or
 190 combination chemotherapy across any number of treatment cycles. Studies evaluating
 191 radiation and newer targeted therapies for cancer treatment were excluded. However,
 192 patients who received hormone therapy, such as tamoxifen, for prophylactic purposes
 193 following the treatment of their cancer, or accompanying local radiation therapy were
 194 eligible for inclusion with noting. This represents a deviation from the published protocol
 195 since it was discovered on screening that the majority of the studies have included these
 196 patients within their chemotherapy –treatment groups. Given that these treatment
 197 strategies are commonly used in breast cancer patients, this inclusion is justified in
 198 enhancing study external validity.

199

200 *Condition*

201 Studies were included if they were investigating cognitive impairment arising as a result
 202 of interventions administered for cancer as described above. This condition may be
 203 variably described as ‘chemobrain’, ‘chemofog’, chemotherapy-induced cognitive
 204 impairment (CICI), or cancer-related cognitive impairment (CRCI). Studies where
 205 cognitive decline was indicated by any one of several commonly used modalities were
 206 eligible for inclusion. These measurement modalities included self-reported measures,
 207 imaging, or objective neuropsychological testing [33].

208

209 *Context & Study Design*

210 Studies from any country or geographic region were eligible for inclusion with noting of
211 location. Patients from community or clinic settings were eligible for inclusion. All study
212 designs were eligible for inclusion. Studies that selected participants into groups based on
213 report of cognitive decline were excluded due to the risk of sampling bias limiting the
214 generalisability of the results.

216 *Data extraction*

217 Data were extracted from the included studies by one reviewer (A.W), with checking
218 performed on all studies by a second reviewer using a modified extraction template in
219 Covidence (supplementary material S2). The template was piloted on three studies to
220 confirm that all required information was being considered. Contact with study authors
221 was undertaken where necessary to clarify findings or seek further information. Key data
222 extracted included the population sampled, chemotherapy and other treatment
223 administered, timeframe since chemotherapy treatment or remission, identified
224 confounding factors, method of cognitive testing and criteria for classifying impairment.
225 Prevalence (%) was extracted or calculated from the available data. Where necessary,
226 data were extracted from figures using GetData Graph Digitiser (version 2.26, S.
227 Federov, Moscow, Russia). Prevalence estimates reported by sub-group, such as
228 cognitive domain affected, age, or treatment were also recorded.

229

230 *Assessment of Methodological Quality*

231 Articles included in the study were assessed for methodological quality by one reviewer
232 (AW), with confirmation provided by a second reviewer, using the JBI critical appraisal
233 checklist and guidance notes developed for prevalence reviews (supplementary material
234 S3). [34] A study was deemed to be of high quality if it scored greater than 70%,
235 moderate quality with scores between 50-70% and low quality when scoring below 50%,
236 as described in Dijkshoorn et al 2021 [3]. Summary graphs were created in Review
237 Manager (RevMan) ([Computer program], Version 5.3. Copenhagen: The Nordic
238 Cochrane Centre, The Cochrane Collaboration, 2014).

239 *Data synthesis*

240 An initial descriptive analysis of the studies was undertaken to identify relevant
241 subgroups where studies with similar characteristics could be combined. Synthesis was
242 undertaken narratively where there was heterogeneity in these study characteristics.
243 Studies were grouped for synthesis based on time point of cognitive assessment. Basic
244 statistical analysis was performed using Megastat Excel Add-In (version 10.3, McGraw-
245 Hill Higher Education, New York, NY).

246
247 A meta-analysis of the prevalence rates of cognitive impairment was conducted for
248 studies that utilised objective neuropsychological tests. Many of the included studies
249 contained repeated observations where the unit of interest was the individual rather than
250 observation. Consequently, these cannot be combined in a standard meta-analysis, with
251 time-point as sub-group, since they introduce a unit-of-analysis error. [35] A number of
252 approaches to perform meta-analyses in this situation have been proposed. [36] For this
253 review, an All Time-points Meta-analysis (ATM) method was selected with studies at the
254 time-points of interest being analysed separately, with a qualitative comparison being
255 made with estimates at other time-points. Analysis was performed using the MetaXL
256 (www.epigear.com) add-in for Microsoft Excel. A pooled prevalence figure was
257 calculated with 95% CI. Overall estimates were calculated with random effects models
258 employing the DerSimonian & Laird method as a between-study variance estimator.
259 Cochran's Q (verifying the presence of heterogeneity) and the I^2 statistic (amount of
260 heterogeneity) were calculated. A quality model [37] was run as part of the meta-analysis
261 where the weight of each study was calculated based on the previously calculated quality
262 score. An $I^2 > 70\%$ was considered substantial and these pooled data were subject to
263 sensitivity analyses.

264
265 In a meta-analysis of prevalence, if the estimates for a study get closer to the limits of
266 zero or one, the weight may be overestimated in the meta-analysis. Consequently, it has
267 been recommended to transform the prevalence to a variable not constrained to the 0-1
268 range with an approximately normal distribution. [38] The double arcsine transformation
269 is usually preferred [38] and was applied here. For final presentation, the pooled
270 transformed proportions and corresponding CIs were back transformed.

271

272 Publication bias was examined using doi plots and the Luis Furuya-Kanamori asymmetry
273 index (LFK index) in MetaXL. The doi plot is suggested to be more sensitive than the
274 funnel plot, particularly when small numbers of studies are involved. [39]

275 Potential influences on prevalence estimates were investigated using subgroup analyses
276 and meta-regression. The criteria for performing a meta-regression were derived from
277 Cochrane guidance. [35] Meta-regression was performed using Comprehensive Meta-
278 Analysis (Version 3, Borenstein, M., Hedges, L., Higgins, J., & Rothstein, H. Biostat,
279 Englewood, NJ 2013). Age, cognitive impairment criteria stringency, sample size,
280 methodological quality and geographical location were identified a priori as potential
281 sources of variation in the prevalence estimates.

282 **Results**

283 *Search Results*

284 A total of 3010 titles were identified from the electronic searches, with 5 identified from
285 reference lists (Figure 1). Following removal of duplicates 2194 titles remained. After
286 title screening, 127 records remained. A further 43 records were excluded after abstract
287 examination. Eighty-nine full-text articles were screened. Fifty-two of these met the
288 eligibility criteria for the review, with thirty-seven being excluded at this stage (refer to
289 supplementary material S4).

290

291

Figure 1: The PRISMA [28] flow diagram for the systematic review detailing the databases searched, the number of abstracts screened and the full texts retrieved

Included Studies

The 52 included studies contributed 84 prevalence estimates across a series of time points in relation to chemotherapy treatment. It should be noted that the majority of studies did not seek to determine prevalence as a primary aim of the study, with the data being presented as one of a number of outcomes reported. Three studies did however have a focus on prevalence as expounded in their study aims. [40-42] The majority of studies adopted an observational study design (48 studies, 92%). The remainder were randomised controlled trials (RCT). The studies that employed RCT designs were evaluations where the primary study question related to differences between different chemotherapy regimes or alternate therapeutic strategies and there was no selection into groups based on presence of cognitive impairment (an exclusion criterion for this review). Out of the studies that were observational in nature, 25 (52%) were cohort designs, 20 (42%) cross-sectional and 13 (27%) prospective longitudinal. Sample sizes were generally small with a range from 20 - 1147. Studies with larger sample sizes tended to employ self-report measures rather than objective test modalities.

312 With the exception of two studies [43, 44], data came from countries with high-income
313 economies (World Bank Classification). The vast majority of study data originated from
314 the continents of North America and Europe (42% and 44% respectively). Age of study
315 participants was generally similar across studies and to be expected given the breast
316 cancer age risk. The pooled mean across the studies was age 52 (range of included study
317 means: 45-71). Three studies [15, 45, 46] were specifically interested in older patients
318 and included only women over the age of 65.

319

320 Consistent with clinical practice, women received a range of chemotherapy drugs as part
321 of their treatment cycles across the studies. Common combinations reported were FEC
322 (5-fluorouracil, epirubicin, and cyclophosphamide), CMF (cyclophosphamide,
323 methotrexate and 5-fluorouracil), doxorubicin + paclitaxel, doxorubicin +
324 cyclophosphamide. This information has not been presented in the review due to a
325 general lack of separation of outcome data based on treatment in the primary studies,
326 necessitating assimilation as a group. In the vast majority of studies, a significant
327 percentage of patients, especially at later follow-up points were receiving hormonal
328 therapy such as tamoxifen. Hormonal therapies have been shown to influence cognition
329 [47, 48]. These agents are administered frequently and can be considered a standard part
330 of treatment. Inclusion of these data hence supports the external validity of the review's
331 findings.

332

333 The included studies generally assessed cognition at similar time-points to coincide with
334 typical phases of breast cancer treatment. Due to this similarity studies were grouped
335 based on these time points for further assimilation, labelled T1-T7 (Refer to
336 supplementary material S5). Data from baseline, pre-chemotherapy assessments (T0),
337 were not extracted as part of this review since we did not have as an inclusion criterion
338 that studies had to be longitudinal. Further, the review question relates to prevalence of
339 impairment after breast cancer treatment, not any measured decline as a result of
340 treatment.

341

342 The included studies utilized a range of methods for assessment of cognitive impairment.
343 It was considered that these differences could introduce substantial heterogeneity when
344 assimilating prevalence estimates. Hence, assimilation was undertaken based on use of

345 three different methods; self-report measures, short cognitive screening tools, and
 346 objective neuropsychological test batteries (supplementary material S6). Fifteen studies
 347 utilized self-report measures, nine employed short cognitive screening tools and a
 348 majority of 35 used neuropsychological test batteries. Seven studies utilized two of these
 349 testing modalities [15, 49-51] [52] [53] [42]. Studies using objective cognitive tests for
 350 the most part used tests which spanned the various cognitive domains of executive
 351 function, language function, motor, processing speed, verbal and visual learning and
 352 memory, visuo-spatial function and working memory. From here on, review data is
 353 presented based on these 3 groupings of outcome assessment method (refer to Table 1, 2
 354 and 3).

Table 1: Summary of included studies that used self-report measures of cognitive impairment

Study	Study Design	Location	Sample Size	Sample Age (Range, Mean)	Assessment Method	Methodological Quality	Definition of cognitive decline	Time points □	Prevalence % (95% CIs where available)
Buchanan 2015 [54]	Cross-sectional	United States	Chemotherapy = 288 Chemotherapy + hormone = 859	34 - 82	FACT-Cog	High	Responses of cognitive complaints “sometimes”, “often”, or “always”	Mixed*	74 Chemotherapy 78 Chemotherapy + hormone
Debess 2010 [50]	Cohort	Denmark	75	29–59, 47.2	Author-derived questions assessed on 7 point scale	High	Scores of 5–7	T2	Memory: 29 Concentration: 13 Mental fatigue: 33 Vigour: 43
Ganz 2013 [55]	Cohort	United States	189	51.8	PAOFI	Moderate	Classified based on PAOFI score as high range versus normal/no complaints	Mixed*	Memory: 23 High level cognition: 19
Hurria 2006 [45]	Prospective Longitudinal	United States	45	65–84, 70	Squire	Moderate	Unclear but based on score	T3	51
Janelins 2017 [56]	Prospective Longitudinal	United States	505	22–81, 53.4	FACT-Cog	High	A 1/2 standard deviation representing a minimal clinically important difference	T2, T3	T2: 45 T3: 37
Jenkins 2006 [51]	Cohort	UK	85	51.5	Broadbent	High	Self-report of change	T2, T3	T2: Memory: 83 Concentration: 80 T3:

									memory: 60 concentration: 45
Koppelmans 2012 [11]	Cohort	Netherlands	196	50-80, 64.1	Basic self- report	High	Yes response to question	T7	More problems remembering: 53 Forgetting (daily) pursuits: 43 Word-finding problems: 38
Lange 2016 [15]	Cohort	France	58	65 – 81, 70	FACT-Cog	High	A difference of more than 10% between baseline and T2 was considered clinically significant	T2	Perceived Cognitive Impairment: 34 Perceived Cognitive abilities: 49 (80 over 75 yrs)
Ng 2018 [52]	Prospective Longitudinal	Singapore	166	50.7	FACT-Cog	High	A drop of 10.6 points in the global score from baseline.	T1, T3	T1: 22 T3: 31
Rey 2012 [57]	Cohort	France	222	18-40, 37	Telephone Interview	High	Women who reported memory and/or attention troubles “very often” or “often”	T3, T4, T5	T3:37 T4: 37 T5: 42

Schagen 1999 [53]	Cohort	Netherlands	39	47.1	Interview with questions scored on a Likert scale	Moderate	A score of 2 (moderate) or more was considered a distinct complaint about the cognitive functioning in the domain concerned.	T5	Concentration problems: 31 Memory problems: 21 Thinking: 8 Language: 8
Schmidt 2016 [58]	Cross-sectional	United States	904	48.8 (across range of cancer types)	Based on Quality of Life in Adult Cancer Survivors (QLACS) with some amendment	High	An endorsement of the items was categorised as perceived cognitive decline.	Unclear	58
Shilling 2007 [59]	Cohort	UK	85	51.7	Interview	Moderate	An endorsement of the items	T2, T4	Memory T2: 83 T4: 60 Concentration: T2: 78 T3: 45
Tager 2010 [60]	Cohort	United States	31	46.95–70.96, 60.7	Participants asked to rate perceived memory abilities on a five-point Likert scale	Moderate	Ratings >2 were coded as cognitive problems	T2,T3	T2: 43 T3: 46

Van Dam 1998 [42]	RCT	Netherlands	FEC [□] :36 CTC [#] : 34	FEC [□] : 48.1 CTC [#] : 45.5	Cognitive problems in daily life interview	High	An endorsement of the items was categorised as perceived cognitive decline.	T5	Concentration: High-dose:38 Standard-dose: 31 Memory High-dose: 32 Standard-dose: 28 Thinking: High-dose: 21 Standard-dose: 11 Language: High-dose:12 Standard-dose: 11
----------------------	-----	-------------	---	--	---	------	---	----	--

* Time-points measured in study cross time-points used in this review preventing accurate assimilation

□ T1 = During chemotherapy treatment, T2 = Just after cessation of treatment, T3 = 6 months after treatment cessation, T4 = 1 year after treatment cessation, T5 = 2-3 years after treatment cessation, T6 = 5 – 10 years after treatment cessation, T7= ≥10 years after treatment cessation

□ Fluorouracil, epirubicin, cyclophosphamide (standard dose)

[#] Four cycles of FEC followed by cyclophosphamide, thiotepa, and carboplatin (high dose)

Table 2: Summary of included studies employing short cognitive screening tools to assess cognitive impairment

Study	Study Design	Location	Sample Size	Sample Age (Range, Mean)	Assessment Method	Methodological Quality	Definition of cognitive decline	Time point(s) □	Prevalence % (95% CIs where available)
Biglia 2012 [49]	Prospective Longitudinal	Italy	40	38–65, 51	MMSE	Moderate	Score under the mean	T2	31
Brezden 2000 [25]	Cohort	Canada	71 across two groups: 1) Currently receiving chemotherapy, and 2) one year after cessation	34-70, 49 26-61, 46	HSCS	High	Classification as mild, moderate or severe according to pattern of scores as described by test authors	T1, T4	T1 mild: 13 moderate: 10 severe: 12/31 = 39 T4 mild: 20 moderate: 28 severe: 9/40 = 23
Fan 2005 [26]	Cohort	Canada	81	48 (median)	HSCS	High	Scores from each item on the test were subject to an algorithm which generated bands of impairment	T2, T4, T5	T2 Mild: 35 Mod-Severe: 16 T4 Mild: 30 Mod-severe: 4 T5: Mild: 21 Mod-severe: 4
Fontes 2016 [40]	Prospective Longitudinal	Portugal	475	54.7 (median)	MoCA	High	A MoCA score at least 2.0 standard deviations below age- and education-adjusted cut-offs	T3, T5	T3: 7 T5: 8

Ng 2018 [52]	Prospective Longitudinal	Singapore	166	50.7	Headminder computerized test	High	Reliable change index (RCI) score calculated to determine cognitive decline in each cognitive domain. A RCI score of lower than -1.5 was used as criteria for decline.	T1, T3	T1: 6.1-21.6 T3: 1.2-14.6
Pillai 2019 [43]	Prospective Longitudinal	India	152	27 -72, 47	MMSE	Low	MMSE score of ≤ 24 (out of 30)	T2,T3,T4, T5	T2: 0 T3: 0 T4: 0 T5: 0
Prokasheva 2011 [61]	Cohort	Israel	20	30-57, 49.3	Doors and People Test	Moderate	Below 1SD as indicative of mild impairment and below 2SD for moderate impairment	T5	40
Ramalho 2017 [62]	Cohort	Portugal	418	27 -87	MoCA	High	MoCA score values at least 1.5 standard deviations below age- and education-adjusted cut-offs	T3	8 (5.8,11)
Tchen 2003 [63]	Cohort	Canada	110	27-60, 48	HSCS	High	Based on standard test criteria	T1	T1 Mild: 34 Moderate: 2 Severe: 14

□ T1 = During chemotherapy treatment, T2 = Just after cessation of treatment, T3 = 6 months after treatment cessation, T4 = 1 year after treatment cessation, T5 = 2-3 years after treatment cessation

Table 3: Summary of included studies employing objective neuropsychological testing to assess cognitive impairment

Study	Study Design	Location	Sample Size	Sample Age (Range, Mean)	Number of Tests in Battery	Methodological Quality	Definition of Cognitive Decline	Time point(s) □	Prevalence % (95% CIs where available)
Andryszak 2018 [64]	Cohort	Poland	31	52.4	1	Moderate	Scores lower than 2 SD compared to control group averages	T1,T2	T1:32 T2: 26
Biglia 2012 [49]	Prospective Longitudinal	Italy	40	38–65, 51	10	Moderate	“Unsatisfying score” not defined in report	T2	15
Collins 2009 [65]	Cohort	Canada	53	50-65, 57.9	18	High	SRB* scores of –2 or less on at least two tests	T2, T4	T2: 34 T4: 11
Collins 2013 [66]	Prospective longitudinal	Canada	60	52.4	17	High	Scores lower than 2 SD on at least two tests	T1 (measured at end of each cycle)	30 (mean across all cycles)
Collins 2014 [67]	Cohort	Canada	56	51.8	19	High	SRB scores of –2 or less on at least two tests	T1 (measured at end of each cycle) T4	T1: 48 T4: 22
Debess 2010 [50]	Cohort	Denmark	75	29–59, 47.2	4	High	Significant changes (between 5th and 95th percentile in controls) on at least two tests	T2	4
Hermelink 2007 [68]	RCT	Germany	101	48.6	12	High	Reliable-change index (RCI) with a probability of error set at 10%	T1	27
Hermelink	Cohort	Germany	91	27.3–64.9,	18	High	Five or more scores	T2, T4	T2: 6

2017 [69]				52.3			below 1.5 standard deviations and/or four below 2 standard deviations		T4: 18
Hurria 2006 [46]	Prospective longitudinal	United States	28	65–84, 71	13	Moderate	Scores lower than 2 SD on at least two tests	T3	39
Jansen 2011 [70]	Prospective longitudinal	United States	71	30 – 65, 50.3	4	High	Scores lower than 1 SD on at least two tests	T1, T2, T3	T1:23 T2:52 T3:20
Jenkins 2006 [51]	Cohort	UK	85	51.5	7	High	Reliable change index on at least two measurement	T2, T3	T2: 20 T3: 18
Jim 2009 [71]	Cohort	United States	97	50	9	High	Scores lower than 1.5 SD on at least two tests	T3	34
Jung 2014 [72]	Cross sectional study	Korea	32	31-61, 46	3	High	Cutoff as 5/4 on Digit Span -DS F; 4/3 on Digit Span B; 28-30/25 on Controlled Oral Word Association- COWA)	T2	DSF mild: 32 moderate: 13 DSB mild: 32 moderate: 42 COWA* (A) mild: 3 moderate: 58 COWA (B) mild: 52
Kesler 2017 [73]	Cohort	United States	31	34–65, 48.6	4	Moderate	Scores lower than 1.5 SD on at least two tests or lower than 2SD on any one test.	T4	55
Kreukels	Cohort	Netherland	63	46.6	10	High	Scores lower than 2	T3/T4	33

2008 [74]		s					SD on at least three tests		
Lange 2016 [15]	Cohort	France	58	65 – 81, 70	8	High	A significant change in a domain score	T2	64
Mehlsen 2009 [75]	Cross-sectional	Denmark	36	48.6	13	High	Decline on at least 3 of the cognitive measures.	T2	29
Mehnert 2007 [76]	RCT	Germany	23	33–65, 53	18	Moderate	Scores lower than 1.4 SD on at least four tests	T6	13
Menning 2016 [77]	Cohort	Netherlands	31	49.8	11	High	Multivariate normative comparison comparing against distribution of scores in control group.	T3	16
Reid-Arndt 2009 [78]	Cross sectional study	United States	46	53.4	11	High	Scores of below 1 SD	T2	Executive Functioning: 22 Verbal Fluency: 41
Reid-Arndt 2010 [79]	Prospective longitudinal	United States	39	53.4	7	High	Below 1SD as indicative of mild impairment and below 1.5 SD for moderate impairment	T3, T4	WMS Log Mem: T3: 6 T4: 33 RAVLT T3: 13 T4: 6 WMS Log Mem II T3: 3 T4: 33 Trails A T3: 3 T4: 9

									Trails B T3: 15 T4: 24 SCW T3: 3 T4: 3 COWA T3: 0 T4: 9 Category Fluency: T3: 0 T4: 16
Ruzich 2007[80]	Prospective longitudinal	Australia	35	30–66, 53	15	Moderate	Scores below 1 SD on at least two tests	T1,T2,T3	T1: 15 T2: 37 T3: 30
Schagen 1999 [53]	Cohort	Netherlands	39	47.1	14	Moderate	Scores below 2 SD on at least three tests	T5	28
Schagen 2002 [81]	Cross-sectional	Netherlands	Two groups of different chemotherapy regimes: FEC ^a : 23 CMF ^a : 31	50.3	13	High	Number of tests with scores below 2SD	T5	FEC: 9 CMF: 13
Schagen 2006 [82]	RCT	Netherlands	Two groups of different chemotherapy	45	10 tests (not listed)	High	Scores below 2 SD on at least three tests	T3	FEC: 10 CTC: 20

			regimes: FEC $\square$: 39 CTC $\#$: 28						
Schrauwen 2020 [41]	Cohort	Belgium	66	27 – 64, 46.7	5	High	Standardized difference score exceeding -2.5 SD on at least one test.	T2	24
Shilling 2005 [83]	Cohort	UK	50	51.1	8	High	Reliable decline on two or more tests	T2	34
Stewart 2008 [84]	Cohort	Canada	61	50-66, 57.5	18	High	Two or more SRB scores of -2.0 or less on at least two tests	T2	31
Stouten- Kemperma n 2015 [85]	RCT	Netherland s	Two groups of different chemoth erapy regimes: Conventi onal Dose =24 High Dose = 17	56.3- 59.8	8	Moderate	Score larger than 2SDs below the mean considered impaired on a test. The fifth percentile of the overall impairment score of healthy control scores was cutoff for impairment	T7	Conventional Dose: 8 High Dose: 11
Syarif 2019 [44]	Cross- sectional	Indonesia	82	43	1	Low	Not reported	Unclear	87 (mild-serious impairment)
Van Dam 1998 [42]	RCT	Netherland s	FEC $\square$: 36 CTC $\#$: 34	FEC: 48.1 CTC: 45.5	13	High	Scores below 2 SD on at least three tests	T5	FEC: 17 CTC: 32
Van Dyk 2018 [86]	Cross- sectional	United States	20	47	17	Moderate	Impaired by ICCTF guidelines	T2	45

Vearncombe 2009 [87]	Prospective longitudinal	Australia	136	49.38	10	High	Reliable change on at least 2 measures	T2	17
Wefel 2010 [88]	Prospective longitudinal	United States	42	33-65, 48,4	6	High	Scores below 2SD on one test.	T2, T4	T2:65 T4:61
Wieneke 1995[89]	Cross sectional	United States	28	28 – 54, 42	15	Moderate	Score below 2 SD on at least one test classified as moderate impairment	T2-T4 (varies across period)	75

□ T1 = During chemotherapy treatment, T2 = Just after cessation of treatment, T3 = 6 months after treatment cessation, T4 = 1 year after treatment cessation, T5 = 2-3 years after treatment cessation, T6 = 5 – 10 years after treatment cessation, T7 = ≥ 10 years after treatment cessation

*Standardised Regression Based

□ Fluorouracil, epirubicin, cyclophosphamide

^a Cyclophosphamide, methotrexate, 5-fluorouracil

[#] Four cycles of FEC followed by cyclophosphamide, thiotepa, and carboplatin,

Methodological quality

A summary of the methodological quality assessment for the included articles is provided in Figure 2 and supplementary material (S7). Thirty-five (67%) studies were considered to be of high quality, fifteen (29%) of moderate quality and only two [43, 44] (4%) studies of low quality (see Tables 1, 2 and 3). The main domains of concern were participant sampling, sample size, condition measurement and statistical analysis. Of these, the outcomes of assessment of risk in the domains of sample size and condition measurement are considered to have the most impact on the review outcomes. However, use of a quality model in meta-analysis and evaluating the impact of sample size by meta-regression minimizes this impact.

Figure 2: Risk of bias graph: review authors' judgements about each methodological quality item presented as percentages across all included studies. Study number = 52.

Prevalence estimates based on self-report measures of cognitive impairment

It was considered that there was significant diversity in the studies that used self-report measures in terms of method of assessment of cognitive impairment, timing of assessment in relation to chemotherapy treatment and form of report (based on cognitive domain rather than overall impairment). Therefore, these results are presented narratively.

Fifteen studies (Table 1) examined prevalence of cognitive impairment after breast cancer treatment contributing 22 prevalence estimates across all study time-points with the exception of T6. There were a number of studies with large sample sizes in this group [54, 56, 58],

however the median sample size for the group remained modest at 85 (range 31- 1147). Studies utilized a variety of methods to assess impairment which ranged from validated self-report instruments such as FACT-Cog or PAOFI, to semi-structured interviews or Likert responses to questions on cognitive function. Seven (47%) of studies used the former. When the studies employing validated tests were contrasted with those using non-validated methods median global impairment across the time points was 46% and 39.5% respectively, with no significant difference in prevalence rate between the two subgroups (two tailed T-test; $t(6,5) = -1.2, p = 0.27$).

Prevalence rates for global impairment varied from 21-83% (mean 44%) across all time-points. Noting the paucity of data at T6 and T7, no obvious downward trend in impairment across time is evident (Figure 3). Two studies in this sub-group focused on CICI in older women over the age of 65 [45] [15] reporting prevalence rates of 51% [45], and 34%. [15] Whilst, the former is clearly above the mean reported above of 44%, the value within Lange et al. 2016 [15] actually falls under the calculated average.

Figure 3: Boxplot of prevalence reported for cognitive impairment, via self-report methods, across the time-points included in the review (data from 12 studies in Table 1, 3 studies were excluded due to mixed or unclear timing of assessment [54, 55, 58])

Prevalence estimates based on use of short cognitive screening tools to measure cognitive impairment

There was considerably less diversity in assessment method for cognitive impairment, and time points of measure for the studies that utilised short cognitive screening measures compared to those employing self-report. However, at each time point there were fewer than five studies. Study has suggested that at least 5 studies are needed to achieve statistical power from random effects meta-analyses that are superior to the individual contributing studies. [90] Studies within this group were therefore not subjected to meta-analytic techniques and are reported narratively.

Nine studies (Table 2) utilized short cognitive screening tools as part of this grouping, contributing 17 prevalence estimates. No studies performed longer-term follow up at the T6 and T7 time-points. Sample size ranged from 20-475 (mean =170). Five different validated methods of assessing impairment were employed (Table 1). With the exception of the Doors and People test these methods assess a range of cognitive domains expected to be impacted by chemotherapy administration. In two studies [26] [25] test outcomes were reported based on a banding of scores as mild, moderate or severe. For the purposes of assimilation, only prevalence estimates for moderate-severe impairment have been considered.

In contrast to studies that used self-report none of these studies specifically examined older populations. However, three of these studies [25, 43, 62] had a wider age range with patients above 65 years of age being included. This has not however led to a substantially increased mean participant age in comparison to the other studies.

Prevalence rates across all time-points ranged from 0-50. The mean prevalence was 16%. Exclusion of the four prevalence estimates of zero provided by Pilai et al. 2019 [43] leads to a range of 4%-50%, with mean prevalence of 21%. This action may be justified given the low quality rating assigned to this study. There was a general downward trend in prevalence across time, but range at each time-point (with the exception of T3) was large (Figure 4).

Figure 4: Boxplot of prevalence reported for cognitive impairment, via short cognitive screening methods, across the time-points included in the review. Ng et al. 2018 [52] was excluded from the calculation since a range was reported, and only considering moderate-severe classification of impairment from [26] and [25].

Prevalence estimates based on use of objective neuropsychological testing to measure cognitive impairment

Thirty-five studies (Table 3) utilized objective neuropsychological tests to screen for cognitive impairment. These studies contributed 46 prevalence estimates. Sample size ranged from 20-136 (median = 53) having the lowest range and average of all the groups. Studies reported prevalence across all time points of interest, although only one estimate was reported at each of the longer-term follow up points (T6 and 7). Two studies in the group evaluated women over the age of 65. [15, 46] The neuropsychological tests used were all established, validated tests which when administered in a battery broadly encompass the major cognitive domains. Two studies only employed one test for cognitive assessment, [44, 64] and as such did not evaluate all cognitive domains.

There were differences between the studies in the criteria used to determine cognitive impairment. In general a diagnosis of impairment was made based on two factors being met:

1) the level of decline in score relative to some reference value, for example healthy control patients 2) that this decline occurred in a pre-determined number of tests. Fourteen (40%) of the studies used 2 standard deviation declines to guide level but there was heterogeneity even between these in the number of tests with declining scores required to satisfy the diagnostic criteria (see Table 3). Studies that considered a one SD decline as criteria for impairment, or were not clear on the level used, were considered less stringent (see meta-analysis). Three studies [72, 78, 79] reported test outcomes by cognitive domain or individual test, rather than global impairment rendering these data harder to assimilate with the other studies.

Meta-analysis

Pooled Prevalence

It was considered that meta-analysis was appropriate for further investigation of prevalence estimates derived from studies utilizing objective neuropsychological tests. Thirty-seven prevalence estimates from 27 articles were included in the meta-analysis, grouped based on time point of prevalence measure (from T1-T5). Only one study contributed data at each of T6 [76], and T7 [85] and consequently these time points and studies were not included. Three studies were excluded from entry since prevalence was reported based on cognitive domain or test type [72, 78, 79]. A further three studies were excluded since the timing of measure was unclear [44], or was performed across multiple time points as defined for this review. [74, 89] In two studies, prevalence estimates were reported based on chemotherapy regime. [81] [82] In order to ensure comparability with the other included studies only estimates for the standard dose chemotherapy regime from Schagen et al. 2006 [82] were included in the meta-analysis. Prevalence estimates reported by chemotherapy regimen in Schagen et al. 2002 [81] were pooled.

The overall random-effects pooled prevalence (95% CIs) of cognitive impairment from time points T1-T5 respectively was 34% (24-44), 30% (20-40), 23% (16-31), 31% (14-50%), and 21% (15-28) (Figure 5). Figure 6 synthesises these findings narratively across the time-points with the inclusion of the single individual study estimate contributed from [76] and [85] at T6 and T7. There is a general downward trend in prevalence across time, although substantial imprecision in the effect at T4 with a slightly increased prevalence than might be expected if observing the trend.

Figure 5: Forest plots of prevalence reported in studies utilising neuropsychological tests to diagnose cognitive impairment following chemotherapy treatment for breast cancer. A) Cognitive assessment undertaken during chemotherapy treatment (T1), B) Cognitive assessment undertaken just after cessation of chemotherapy treatment (T2), C) Cognitive assessment undertaken 6 months after cessation of chemotherapy treatment (T3), D) Cognitive assessment undertaken 1 year after cessation of chemotherapy treatment (T4), E) Cognitive assessment undertaken 2-3 years after cessation of chemotherapy treatment (T5).

Figure 6: Prevalence reported for cognitive impairment, via neuropsychological test methods, across the time-points included in the review. Data represent pooled prevalence estimates derived from meta-analysis, with 95% CIs. ∞ - values are not pooled figures but represent individual values from one study

Heterogeneity and Publication Bias

Heterogeneity as determined by the I^2 statistic was universally high with the exception of T5 (where there were limited estimates) at 73%, 90%, 64%, 89% and 0 (T1-5 respectively). Sensitivity analyses performed included removal of each study in turn, analysing untransformed data, considering the effect of the quality score and using a fixed effects model. In general, results were similar after these analyses (supplementary material S8). However, removal of Jansen et al. 2011 [70] at T1 had a significant effect reducing I^2 by 10% to 63% (potentially as a result of the impairment criteria used as described below). It is noteworthy that removal of the Andrysak 2018 study, [64] which only employed one cognitive test, did not impact the heterogeneity estimates, nor did removal of the studies with cross-sectional designs. [75, 86]

In order to explain the heterogeneity in the pooled effects sizes the influence of stringency of criteria for diagnosing cognitive impairment was explored via subgroup analysis. Only prevalence reports from T2 and T3 were subjected to subgroup analyses based on the number of studies contributing data at these time-points, and the number in each analysis when split

into sub-groups. At these time-points, four studies were considered to have used less stringent test methods. [70, 75, 80, 88] Heterogeneity remained high after sub-group analysis with the high stringency sub-group having a pooled prevalence of 25% (95% CIs: 16-35), $I^2 = 90\%$ at T2, and 23% (95% CIs: 13-34), $I^2 = 74\%$ at T3 (Figure 7).

Figure 7: Forest plots of prevalence sub-grouped by stringency of cognitive assessment. A) Cognitive assessment undertaken just after cessation of chemotherapy treatment (T2), B) Cognitive assessment undertaken 6 months after cessation of chemotherapy treatment (T3)

Meta-regression was performed in an attempt to account for the remaining heterogeneity. The results of five individual meta-regression analyses performed at T2 based on 1) the continuous variables of age, methodological quality and sample size, and 2) categorical variables of cognitive impairment criteria stringency and geographical location (3 subsets;

Europe (reference group), North America, Asia-Pacific) are included in Table 4. Cognitive impairment stringency explained some of the variance in prevalence, accounting for 16% of the heterogeneity at T2 ($P = 0.04$). Sample size also accounted for 16% of the heterogeneity ($P = 0.04$) with increased sample size tending to result in a lower prevalence estimate. The remainder of the covariates could not account for the variance observed.

Meta-regression was then used to examine the relationship between sample size, cognitive impairment test stringency, and effect size. A test of this model yielded a Q-value of 6.99 with 2 degrees of freedom and corresponding p-value of 0.03. The test for heterogeneity yields a Q-value of 74.7 and a corresponding p-value of 0, implying that the variation of observed effects about the regression line falls within the range that cannot be explained by sampling error alone. The model is able to explain some 23% of the variance in true effects.

The doi plots for publication bias showed some asymmetry implying the presence of bias. The LF asymmetry index confirmed there was minor asymmetry at T2, and T5 and major asymmetry at T4. There was no evidence of publication bias at T1 and T3 (see supplementary material S9).

<i>T2- meta-regression</i>	Meta-regression		
	Mean Difference (95% CIs)	P	R²
Age	0.06 (-0.01- 0.13)	0.09	0.13
Quality Score	-0.25 (-4.14 - 3.64)	0.9	0
Sample Size	-0.02 (-0.03 - 0)	0.04	0.16
Location			
Europe (ref group)			
North America	1.04 (0.11-1.98)	0.09	0.08
Asia-Pacific	0.16 (-1.14-1.46)		
Impairment criteria stringency			
High Stringency			
(ref group)		0.04	0.16
Low Stringency	0.97 (0.04-1.9)		

Table 4: Results of four individual meta-regression analyses at T2 based on age, quality score, location and cognitive impairment criterion stringency

GRADE certainty assessment and results

The evidence presented across all the studies grouped by cognitive assessment methods was assessed using the GRADE (Grading of Recommendations Assessment, Development and Evaluation) approach (GRADEpro GDT: GRADEpro Guideline Development Tool

[Software]). Results are presented in the Summary of Findings Table. The certainty of evidence was graded as very low for all three-assessment types. Inconsistency was rated as serious for all groups of studies based on the wide variation in estimates and high calculated heterogeneity (neuropsychological testing). Imprecision was similarly rated as serious for all outcomes as a result of the wide confidence intervals. Publication bias was strongly suspected for the study group that used neuropsychological tests based on the doi plots and LFK index. Due to the method of synthesis used for the self-report and short cognitive screening tool groups, publication bias was not able to be assessed and was therefore rated as undetected.

Discussion

This is the first systematic review on prevalence of CICI to have included and compared the three commonly used assessment modalities of self-report, short cognitive screen and objective neuropsychological tests. In contrast to the only other published systematic review of prevalence in CICI [3], meta-analytic techniques were applied to allow reporting of pooled effects, and studies which performed cognitive assessment beyond the breast cancer treatment period were also included. Prevalence estimates derived ranged from 0-83% across all time-points and all cognitive test methods. This crude figure is highly comparable with the commonly cited range of 12-82% based on the narrative review of Janelins et al. 2014 [14]. The current review allows greater examination of variability in these prevalence rates, through examination of the influence of factors such as test method, time since treatment and age on rates.

Impact of assessment tool choice on prevalence rates

There are striking differences in the average and maximum prevalence values reported based on method of cognitive assessment. Self-report of cognitive impairment yielded the highest values. There is substantially less variability between values obtained by short cognitive screen and neuropsychological test batteries. This difference in outcomes is to be expected for a number of reasons: 1) patient mood has been shown to have a relatively greater impact on subjective reports of impairment than actual cognitive performance. For example, aspects of mood such as anxiety, depression, poor quality of life and fatigue have all been shown to correlate with self-reported cognition complaints. [49, 91-93]; 2) expectations about treatment might influence self-report of cognitive symptoms. Study has shown that patients who were aware of the potential for cognitive side effects were more likely to report cognitive function issues than those who were unaware of the possibility; [94] 3) patients

may have been functioning above the normal range of cognitive function prior to treatment, so whilst they experience a decline, their cognitive performance falls within the normal range; [95, 96] 4) the battery of objective tests used are ineffective at diagnosing subtle impairment [95, 96], this may especially be of concern with the use of short cognitive screening tools [97]; 5) objective tests are ecologically invalid by failing to represent real-life situations where patients may experience the impairment. [95, 96] Finally, based on the studies included in this review it could be argued that some of the assessment criteria are relatively crude, for example the endorsement of an item with no grading of response [11, 58, 59] would likely lead to substantially higher reporting of issues, which may not actually differ from normal age-related memory loss experienced by healthy individuals [11, 58, 59]. In fact a cursory glance at the prevalence reports in these studies shows that reported prevalences are higher than the mean for the group. This perhaps highlights a future need to only use validated self-report instruments for cognitive assessment.

In spite of the reduced variability observed with objective tests, there was still significant statistical heterogeneity observed on meta-analysis. Approximately 16% of this heterogeneity could be explained by the use of less stringent test criteria for diagnosing impairment. Additionally, pooled prevalence was noticeably altered when subgroups based on test stringency were analysed. This finding provides strong evidence of the need for researchers to agree to, and use standardised impairment criteria, in order to make findings comparable, reproducible, and ultimately useful to practitioners.

Impact of time since treatment on prevalence rates

Prevalence rates ascertained by short screening methods or larger batteries of tests showed a general downward trend over time, with a peak rate occurring during chemotherapy treatment. This finding is supported by pre-clinical and imaging studies in CICI, which demonstrate that, over time, partial resolution of the brain structural and functional impairments occurs. [22, 23] However, it does contrast with the trend in prevalence determined by Dijkshoorn et al. 2021 [3] which only included studies using objective tests, and found elevated (although perhaps not significantly so) prevalence rates at greater than one year after treatment, compared to just after treatment with chemotherapy. Notably however, actual prevalence values calculated in the current review for objective tests are remarkably similar to those in Dijkshoorn et al. 2021 [3]. Rates of 25%, 14 % and 27% were reported in that review at time-points of just after treatment, up to 1 year following, and

greater than 1 year after treatment respectively. [3] This corresponds with 30%, 23% and 31% at equivalent time points in the current review. This similarity in rates is in spite of the differences in methodology employed in the two reviews with the current review assimilating rates via meta-analytical techniques, and the previous review only including longitudinal studies such that impairment was based on a decline in cognitive function from baseline measures.

Conversely, the current review illustrates that prevalence rates as determined by self-report remain high (often above 40%), and show little evidence of decline over time. The previously discussed suggestions explaining the general elevation in self-reported cognitive decline over measured impairment may also be at play here. Furthermore, at late follow-up stages patients will have likely returned to normal daily-life activities, such as work, and may have a greater perception of impairments as they return to navigating everyday tasks.

Whilst fewer studies have evaluated longer-term follow-up points, greater than 5 years following treatment cessation, there is evidence from both self-report and objective testing that some individuals are cognitively impaired at these time-points. This is an important finding for survivorship care and deserves dedicated research attention to fully elucidate impact.

Impact of age on prevalence rates

There is recognition of the potential for increased significance of chemobrain in older adults with cancer due to higher levels of pre-existing cognitive impairment in this age group. [98] Furthermore, older adults may be more susceptible to cancer treatment toxicity leading to heightened symptoms. [15] Whilst it might therefore be predicted that CICI prevalence increases with age, the review did not find enough evidence to either support or refute this.

When meta-regression was applied, age in general was shown to have no effect on the high heterogeneity seen at T2. This lack of effect is unsurprising given that the studies included in the meta-regression had similar mean ages of the sample population, with relatively narrow ranges. Furthermore, for objective testing cognitive score results are usually compared against age-matched controls or some other age-based correction is applied, hence age is implicitly accounted for in experimental design.

Only three studies specifically focused on patients over the age of 65 [15, 45, 46]. Two of these studies [45] [15] used self-report methods, with one reporting prevalence rates below the mean for the group [45], and the other reporting rates high than the mean. [15] The third study [46] utilized objective tests and reported a prevalence value higher than the remainder of the T3 subgroup. However, it also has the widest confidence intervals of the group suggesting the estimate to be imprecise. The study also used a low sample size, which may have influenced the estimate obtained. Noteworthy, is that the cohort study of Lange et al. 2016 also used cognitive tests, with a diagnostic criteria categorized in this review as high stringency, and reported a high prevalence of 64% at T2. Moreover, confidence intervals were of similar width to other studies within the group. [15] This finding is perhaps the most suggestive of an effect of age on CICI prevalence and would be good grounds for dedicating future research effort to this question.

Strength of Evidence on CICI prevalence

In order to account for any effect that low quality studies might have had on the review findings, the meta-analyses incorporated a quality model, and a meta-regression using quality score as a variable was performed. These tools indicated that quality had minimal effect on prevalence rates or variances in the pooled estimates. As a result, there can be confidence that study quality has not affected review outcomes, at least for those studies that utilized objective cognitive assessment tools. Whilst, it is harder to make a formal determination of the impact of quality on the prevalence reports from studies using self-report and short cognitive screening tools, there is little to indicate that the results may have been unduly influenced by poor quality studies.

In all of the meta-analyses performed, with the exception of T5 where there were limited studies, moderate to high statistical heterogeneity was found. Sensitivity testing failed to account for any meaningful heterogeneity, and the subgroup analyses failed to reduce I^2 values substantially. Meta-regression did reveal an effect of cognitive impairment test stringency and sample size on variance. These factors accounted for 23 percent of the variance in effect seen at T2 in a combined model. The remainder of the heterogeneity could possibly be accounted for by methodological diversity due to the differing test batteries used, and further considerations relating to defining impairment levels that were not considered by the sub-group analysis. Additionally, clinical heterogeneity may have arisen as a result of treatments administered, patient menopausal status, or race.

There was evidence of publication bias at a number of time-points included in the meta-analysis, which lowers certainty in the findings from this analysis. However, there are a few considerations regarding assessment of publication bias in prevalence studies. Firstly, there is no specific guidance on assessment of publication bias when proportional data are involved. The issue being that funnel plots may be imprecise at the extremes of a proportion, [99] and this may similarly apply to doi plots. It could be envisaged that the risk of publication bias should be lower in studies of prevalence since there is no favourable hypothesized outcome that may influence decision to publish. However, in the current review it is worth noting that in all of the studies the study objective was not solely around obtaining prevalence data on CICI. Taking this into consideration the possibility of publication bias likely exists, and was evidenced in this study.

Certainty of the evidence contributing to the findings of this review was assessed using the GRADE approach. It was determined that there was low certainty in the body of evidence contributing to the review. However, there is a lack of formal guidance from the GRADE working group on use of the methodology for reviews of prevalence, and the guidance criteria are not always applicable to these types of data. In spite of the lack of standardization, based on the assessment conclusions should be made tentatively.

Study Limitations

The review does have a number of limitations. First, due to the methodological diversity of the included studies, it was only possible to perform meta-analytic techniques on the studies that used neuropsychological tests. As a result, study weight has not been accounted for in the summary prevalence figures reported arising from self-report and short cognitive screening methods. This necessitates a need for extra circumspection around these values. Furthermore, even when meta-analytical techniques were used there was a resultant high statistical heterogeneity. Subgroup analyses and meta-regressions failed to account for most of this heterogeneity leaving the source(s) unknown, although some suggestions have been provided.

Second, due to the nature of the included studies with many having a repeated measures design it was not possible to use meta-analytical techniques to statistically compare prevalence across time-points; this being an aim of this review. Furthermore, in the current study using the All Time Points Meta-analysis approach assumption of independence

between time-points was violated since some subjects contributed data at more than one time-point, whilst others only contribute at one time-point. This can lead to an ecological fallacy with a trend being seen on aggregate, that is not present at an individual level. [36] As a result there can only be low confidence in the direction of trends across time reported. Alternate approaches may be preferred when trends across time are of interest; for example, the use of a trend meta-analysis which uses regression modelling, or a change-in time meta-analysis where the primary study data is used to calculate differences between successive time-points or compared to baseline. [36] These alternate methods were not employed in this review since they require certain data to be provided in the primary studies such as a slope estimate for trend, or assume a longitudinal study design.

A few points regarding the conclusions that can be taken from this review are pertinent. The focus of the review was on estimating prevalence of CICI. However, as previously described cognition deficits may be caused by a range of factors associated with the cancer experience [14]. These may include the cancer itself and associated psychological distress and fatigue, or other administered treatments such as hormonal therapies. Additionally, studies eligible for inclusion in the review were not restricted to those with a longitudinal design, thus allowing for the contrast of findings with a baseline cognitive score. Based on these two points, it is impossible to say whether the cognitive deficits reported were wholly the result of chemotherapy, as opposed to a combination of cancer and multiple treatment-related effects. This concern is academic. From the perspective of patients and health-care providers it is the nature of cognitive decline that is important, not the origin of it. Since, these combinations of factors are inherent, and common to most breast cancer patients, it is considered that the review findings are generalizable.

Conclusions

This review is the first systematic comparison of prevalence rates for cognitive impairment in breast cancer patient that has considered all methods used to ascertain impairment, and evaluates long-term prevalence. Mean prevalence rates for CICI across all time-points were 44% using self-report and 6% using short cognitive screening. Pooled prevalence rates of between 21-34% were derived for impairment diagnosed by objective neuropsychological tests. For all three assessment modalities the GRADE certainty in the evidence was rated as very low. Results therefore need to be interpreted with caution since the actual prevalence may be substantially different from this estimate.

765 *Recommendations for practice*

766 The findings of this review suggest that cognitive impairment may impact up to 1 in 3
767 patients at a level that is clinically significant. This impairment also extends beyond the
768 treatment period and is often still apparent 2-3 years post treatment cessation. Health
769 practitioners should consider the potential for cognitive impairment in breast cancer
770 survivors, and guide them and their families towards sources of information and support.
771 They may also like to discuss with patients some of the minimally invasive, non-therapeutic
772 interventions that might ameliorate symptoms, for example cognitive training or physical
773 activity [32].

774 *Recommendations for research*

775 The uncertainty around the true prevalence of cognitive impairment following cancer
776 treatment leads to under-consideration of the condition, and consequently a lack of support
777 being available for survivors. This leads to reduced patient quality of life through less
778 meaningful engagement with work, family and daily activities, as well as economic impacts
779 for both the individual and society. This review has highlighted a number of considerations
780 for future research to provide greater certainty in the effect estimates. This includes
781 increasing study sample sizes, utilising standardised cognitive test methods and batteries, and
782 applying accepted, clinically relevant criteria for diagnosing impairment. With the increasing
783 number of older cancer survivors, there is also a need for targeted research investigating this
784 group. Acquiring accurate prevalence data will better inform survivorship strategies by
785 allowing appropriate delivery of targeted resources and services, and informed resource
786 allocation for this effort. Access to accurate prevalence estimates will also inform, and
787 perhaps encourage, the development of evidence-based practice guidelines for CICI.

789 **Declaration of Competing Interest**

790 The authors declare that they have no known competing financial interests or personal
791 relationships that could have appeared to influence the work reported in this paper

793 **Funding**

794 This research did not receive any specific grant from funding agencies in the public,
795 commercial, or not-for-profit sectors. ALW was supported by a NHMRC Peter Doherty

796 Biomedical Research Fellowship (APP1140072). RPG was supported by an Australian
797 Government Research Training Program Scholarship.

798 **References**

799

- 800 [1] Nardin S, Mora E, Varughese FM, D'Avanzo F, Vachanaram AR, Rossi V, et al. Breast Cancer
801 Survivorship, Quality of Life, and Late Toxicities. *Frontiers in Oncology*. 2020;10.
- 802 [2] Mascaro A, Farina M, Gigli R, Vitelli CE, Fortunato L. Recent advances in the surgical care of breast
803 cancer patients. *World J Surg Oncol*. 2010;8:5.
- 804 [3] Dijkshoorn ABC, van Stralen HE, Sloots M, Schagen SB, Visser-Meily JMA, Schepers VPM.
805 Prevalence of cognitive impairment and change in patients with breast cancer: A systematic review
806 of longitudinal studies. *Psycho-Oncology*. n/a.
- 807 [4] Mounier NM, Abdel-Maged AE, Wahdan SA, Gad AM, Azab SS. Chemotherapy-induced cognitive
808 impairment (CICI): An overview of etiology and pathogenesis. *Life Sci*. 2020;258:118071.
- 809 [5] Seigers R, Schagen S, Van Tellingen O, Dietrich J. Chemotherapy-related cognitive dysfunction:
810 current animal studies and future directions. *Brain imaging and behavior*. 2013;7:453-9.
- 811 [6] George RP, Semendric I, Hutchinson MR, Whittaker AL. Neuroimmune reactivity marker
812 expression in rodent models of chemotherapy-induced cognitive impairment: a systematic scoping
813 review. *Brain, Behavior, and Immunity*. 2021.
- 814 [7] Nguyen LD, Ehrlich BE. Cellular mechanisms and treatments for chemobrain: insight from aging
815 and neurodegenerative diseases. *EMBO Molecular Medicine*. 2020;12:e12075.
- 816 [8] Hutchinson AD, Hosking JR, Kichenadasse G, Mattiske JK, Wilson C. Objective and subjective
817 cognitive impairment following chemotherapy for cancer: a systematic review. *Cancer treatment*
818 *reviews*. 2012;38:926-34.
- 819 [9] Scherling CS, Smith A. Opening up the window into "chemobrain": a neuroimaging review.
820 *Sensors (Basel)*. 2013;13:3169-203.
- 821 [10] Wigmore P. The effect of systemic chemotherapy on neurogenesis, plasticity and memory. *Curr*
822 *Top Behav Neurosci*. 2013;15:211-40.
- 823 [11] Koppelmans V, Breteler MMB, Boogerd W, Seynaeve C, Gundy C, Schagen SB.
824 Neuropsychological performance in survivors of breast cancer more than 20 years after adjuvant
825 chemotherapy. *Journal of Clinical Oncology*. 2012;30:1080-6.
- 826 [12] Kanaskie ML, Loeb SJ. The experience of cognitive change in women with breast cancer
827 following chemotherapy. *Journal of Cancer Survivorship*. 2015;9:375-87.
- 828 [13] Mitchell T, Turton P. 'Chemobrain': concentration and memory effects in people receiving
829 chemotherapy—a descriptive phenomenological study. *European Journal of Cancer Care*.
830 2011;20:539-48.
- 831 [14] Janelins MC, Kesler SR, Ahles TA, Morrow GR. Prevalence, mechanisms, and management of
832 cancer-related cognitive impairment. *Int Rev Psychiatry*. 2014;26:102-13.
- 833 [15] Lange M, Heutte N, Rigal O, Noal S, Kurtz JE, Lévy C, et al. Decline in cognitive function in older
834 adult with early-stage breast cancer after adjuvant treatment. *Oncologist*. 2016;21:1337-48.
- 835 [16] Greendale GA, Derby CA, Maki PM. Perimenopause and Cognition. *Obstetrics and Gynecology*
836 *Clinics of North America*. 2011;38:519-35.
- 837 [17] Vardy J. *Cognitive Function in Breast Cancer Survivors*. Springer US; 2009. p. 387-419.
- 838 [18] Wefel JS, Lenzi R, Theriault RL, Davis RN, Meyers CA. The cognitive sequelae of standard-dose
839 adjuvant chemotherapy in women with breast carcinoma: results of a prospective, randomized,
840 longitudinal trial. *Cancer: Interdisciplinary International Journal of the American Cancer Society*.
841 2004;100:2292-9.
- 842 [19] Lange M, Giffard B, Noal S, Rigal O, Kurtz J-E, Heutte N, et al. Baseline cognitive functions among
843 elderly patients with localised breast cancer. *European Journal of Cancer*. 2014;50:2181-9.

[20] Jansen CE, Dodd MJ, Miaskowski CA, Dowling GA, Kramer J. Preliminary results of a longitudinal study of changes in cognitive function in breast cancer patients undergoing chemotherapy with doxorubicin and cyclophosphamide. *Psycho-Oncology*. 2008;17:1189-95.

[21] Lange M, Joly F. How to Identify and Manage Cognitive Dysfunction After Breast Cancer Treatment. *J Oncol Pract*. 2017;13:784-90.

[22] Inagaki M, Yoshikawa E, Matsuoka Y, Sugawara Y, Nakano T, Akechi T, et al. Smaller regional volumes of brain gray and white matter demonstrated in breast cancer survivors exposed to adjuvant chemotherapy. *Cancer*. 2007;109:146-56.

[23] McDonald BC, Conroy SK, Ahles TA, West JD, Saykin AJ. Gray matter reduction associated with systemic chemotherapy for breast cancer: a prospective MRI study. *Breast cancer research and treatment*. 2010;123:819-28.

[24] Lv L, Mao S, Dong H, Hu P, Dong R. Pathogenesis, Assessments, and Management of Chemotherapy-Related Cognitive Impairment (CRCI): An Updated Literature Review. *Journal of Oncology*. 2020;2020:3942439.

[25] Brezden CB, Phillips KA, Abdoell M, Bunston T, Tannock IF. Cognitive function in breast cancer patients receiving adjuvant chemotherapy. *J Clin Oncol*. 2000;18:2695-701.

[26] Fan HGM, Houédé-Tchen N, Yi QL, Chemerynsky I, Downie FP, Sabate K, et al. Fatigue, menopausal symptoms, and cognitive function in women after adjuvant chemotherapy for breast cancer: 1- and 2-Year follow-up of a prospective controlled study. *Journal of Clinical Oncology*. 2005;23:8025-32.

[27] Munn Z MS, Lisy K, Riitano D, Tufanaru C. Chapter 5: Systematic reviews of prevalence and incidence. In: Aromataris E, Munn Z (Editors). *JBIM Manual for Evidence Synthesis*. JBI, 2020. Available from <https://synthesismanual.jbi.global>. <https://doi.org/10.46658/JBIMES-20-06>.

[28] Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71.

[29] Rodríguez Martín B, Fernández Rodríguez EJ, Rihuete Galve MI, Cruz Hernández JJ. Study of Chemotherapy-Induced Cognitive Impairment in Women with Breast Cancer. *Int J Environ Res Public Health*. 2020;17.

[30] Addington-Hall J, McCarthy M. Dying from cancer: results of a national population-based investigation. *Palliative medicine*. 1995;9:295-305.

[31] Gehring K, Aaronson NK, Taphoorn MJ, Sitskoorn MM. Interventions for cognitive deficits in patients with a brain tumor: an update. *Expert review of anticancer therapy*. 2010;10:1779-95.

[32] Treanor CJ, McMenamin UC, O'Neill RF, Cardwell CR, Clarke MJ, Cantwell M, et al. Non-pharmacological interventions for cognitive impairment due to systemic cancer treatment. *Cochrane Database of Systematic Reviews*. 2016.

[33] Craig C, Monk B, Farley J, Chase D. Cognitive impairment in gynecologic cancers: A systematic review of current approaches to diagnosis and treatment. *Supportive care in cancer : official journal of the Multinational Association of Supportive Care in Cancer*. 2013;22.

[34] Munn Z, Moola S, Lisy K, Riitano D, Tufanaru C. Methodological guidance for systematic reviews of observational epidemiological studies reporting prevalence and cumulative incidence data. *Int J Evid Based Healthc*. 2015;13:147-53.

[35] Higgins JPT TJ, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions* version 6.2 (updated February 2021). Cochrane, 2021. Available from www.training.cochrane.org/handbook.

[36] Peters JL, Mengersen KL. Meta-analysis of repeated measures study designs. *Journal of Evaluation in Clinical Practice*. 2008;14:941-50.

[37] Doi SA, Thalib L. A quality-effects model for meta-analysis. *Epidemiology*. 2008;19:94-100.

[38] Barendregt J, Doi S, Lee YY, Pacella R, Vos T. Meta-analysis of prevalence. *Journal of epidemiology and community health*. 2013;67.

[39] Furuya-Kanamori L, Barendregt JJ, Doi SAR. A new improved graphical and quantitative method for detecting bias in meta-analysis. *Int J Evid Based Healthc*. 2018;16:195-203.

895 [40] Fontes F, Pereira S, Castro-Lopes JM, Lunet N. A prospective study on the neurological
896 complications of breast cancer and its treatment: Updated analysis three years after cancer
897 diagnosis. *Breast*. 2016;29:31-8.

898 [41] Schrauwen W, Van de Cavey J, Vingerhoets G, Vanheule S, Van den Broecke R, Denys H, et al.
899 Heterogeneous response of chemotherapy-related cognitive decline in patients with breast cancer:
900 A prospective study. *Journal of the International Neuropsychological Society*. 2020;26:806-14.

901 [42] van Dam FSAM, Boogerd W, Schagen SB, Muller MJ, Droogleever Fortuyn ME, Wall Evd, et al.
902 Impairment of Cognitive Function in Women Receiving Adjuvant Treatment for High-Risk Breast
903 Cancer: High-Dose Versus Standard-Dose Chemotherapy. *JNCI: Journal of the National Cancer*
904 *Institute*. 1998;90:210-8.

905 [43] Pillai US, Kayal S, Cyriac S, Nisha Y, Dharanipragada K, Kamalanathan SK, et al. Late effects of
906 breast cancer treatment and outcome after corrective interventions. *Asian Pacific Journal of Cancer*
907 *Prevention*. 2019;20:2673-9.

908 [44] Syarif H, Waluyo A, Afianti Y, Mansyur M. Executive function in breast cancer survivors and the
909 influencing factors. *Enferm Clin*. 2019;29 Suppl 2:280-5.

910 [45] Hurria A, Goldfarb S, Rosen C, Holland J, Zuckerman E, Lachs MS, et al. Effect of adjuvant breast
911 cancer chemotherapy on cognitive function from the older patient's perspective. *Breast Cancer*
912 *Research and Treatment*. 2006;98:343-8.

913 [46] Hurria A, Rosen C, Hudis C, Zuckerman E, Panageas KS, Lachs MS, et al. Cognitive function of
914 older patients receiving adjuvant chemotherapy for breast cancer: A pilot prospective longitudinal
915 study. *Journal of the American Geriatrics Society*. 2006;54:925-31.

916 [47] Buwalda B, Schagen SB. Is basic research providing answers if adjuvant anti-estrogen treatment
917 of breast cancer can induce cognitive impairment? *Life sciences*. 2013;93:581-8.

918 [48] Shilling V, Jenkins V, Fallowfield L, Howell T. The effects of hormone therapy on cognition in
919 breast cancer. *J Steroid Biochem Mol Biol*. 2003;86:405-12.

920 [49] Biglia N, Bounous VE, Malabaila A, Palmisano D, Torta DME, D'Alonzo M, et al. Objective and
921 self-reported cognitive dysfunction in breast cancer women treated with chemotherapy: A
922 prospective study. *European Journal of Cancer Care*. 2012;21:485-92.

923 [50] Debess J, Riis JØ, Engebjerg MC, Ewertz M. Cognitive function after adjuvant treatment for early
924 breast cancer: A population-based longitudinal study. *Breast Cancer Research and Treatment*.
925 2010;121:91-100.

926 [51] Jenkins V, Shilling V, Deutsch G, Bloomfield D, Morris R, Allan S, et al. A 3-year prospective study
927 of the effects of adjuvant treatments on cognition in women with early stage breast cancer. *British*
928 *Journal of Cancer*. 2006;94:828-34.

929 [52] Ng T, Dorajoo SR, Cheung YT, Lam YC, Yeo HL, Shwe M, et al. Distinct and heterogeneous
930 trajectories of self-perceived cognitive impairment among Asian breast cancer survivors.
931 *Psychooncology*. 2018;27:1185-92.

932 [53] Schagen SB, Van Dam FSAM, Muller MJ, Boogerd W, Lindeboom J, Bruning PF. Cognitive deficits
933 after postoperative adjuvant chemotherapy for breast carcinoma. *Cancer*. 1999;85:640-50.

934 [54] Buchanan ND, Dasari S, Rodriguez JL, Lee Smith J, Hodgson ME, Weinberg CR, et al. Post-
935 treatment Neurocognition and Psychosocial Care Among Breast Cancer Survivors. *Am J Prev Med*.
936 2015;49:S498-508.

937 [55] Ganz PA, Kwan L, Castellon SA, Oppenheim A, Bower JE, Silverman DHS, et al. Cognitive
938 complaints after breast cancer treatments: Examining the relationship with neuropsychological test
939 performance. *Journal of the National Cancer Institute*. 2013;105:791-801.

940 [56] Janelsins MC, Heckler CE, Peppone LJ, Kamen C, Mustian KM, Mohile SG, et al. Cognitive
941 Complaints in Survivors of Breast Cancer After Chemotherapy Compared With Age-Matched
942 Controls: An Analysis From a Nationwide, Multicenter, Prospective Longitudinal Study. *J Clin Oncol*.
943 2017;35:506-14.

944 [57] Rey D, Bouhnik AD, Mancini J, Bendiane MK, Srr V, Viens P. Self-reported cognitive
945 impairment after breast cancer treatment in young women from the ELIPPSE40 cohort: the long-
946 term impact of chemotherapy. *Breast J.* 2012;18:406-14.

947 [58] Schmidt JE, Beckjord E, Bovbjerg DH, Low CA, Posluszny DM, Lowery AE, et al. Prevalence of
948 perceived cognitive dysfunction in survivors of a wide range of cancers: results from the 2010
949 LIVESTRONG survey. *J Cancer Surviv.* 2016;10:302-11.

950 [59] Shilling V, Jenkins V. Self-reported cognitive problems in women receiving adjuvant therapy for
951 breast cancer. *European Journal of Oncology Nursing.* 2007;11:6-15.

952 [60] Tager FA, McKinley PS, Schnabel FR, El-Tamer M, Cheung YKK, Fang Y, et al. The cognitive effects
953 of chemotherapy in post-menopausal breast cancer patients: A controlled longitudinal study. *Breast
954 Cancer Research and Treatment.* 2010;123:25-34.

955 [61] Prokasheva S, Faran Y, Cwikel J, Geffen DB. Analysis of memory deficits following chemotherapy
956 in breast cancer survivors: Evidence from the doors and people test. *Journal of Psychosocial
957 Oncology.* 2011;29:499-514.

958 [62] Ramalho M, Fontes F, Ruano L, Pereira S, Lunet N. Cognitive impairment in the first year after
959 breast cancer diagnosis: A prospective cohort study. *Breast.* 2017;32:173-8.

960 [63] Tchen N, Juffs HG, Downie FP, Yi QL, Hu H, Chemerynsky I, et al. Cognitive function, fatigue, and
961 menopausal symptoms in women receiving adjuvant chemotherapy for breast cancer. *Journal of
962 Clinical Oncology.* 2003;21:4175-83.

963 [64] Andryszak P, Wilkosc M, Zurawski B, Izdebski P, Ahles AABBB-WBBCCCCCdRdRdT-
964 MDDEFGGHHH. Verbal memory in breast cancer patients treated with chemotherapy with
965 doxorubicin and cyclophosphamide. *European Journal of Cancer Care.* 2018;27:1-11.

966 [65] Collins B, Mackenzie J, Stewart A, Bielajew C, Verma S. Cognitive effects of chemotherapy in
967 post-menopausal breast cancer patients 1 year after treatment. *Psychooncology.* 2009;18:134-43.

968 [66] Collins B, MacKenzie J, Tasca GA, Scherling C, Smith A. Cognitive effects of chemotherapy in
969 breast cancer patients: a dose-response study. *Psychooncology.* 2013;22:1517-27.

970 [67] Collins B, Mackenzie J, Tasca GA, Scherling C, Smith A. Persistent cognitive changes in breast
971 cancer patients 1 year following completion of chemotherapy. *J Int Neuropsychol Soc.* 2014;20:370-
972 9.

973 [68] Hermelink K, Untch M, Lux MP, Kreienberg R, Beck T, Bauerfeind I, et al. Cognitive function
974 during neoadjuvant chemotherapy for breast cancer: results of a prospective, multicenter,
975 longitudinal study. *Cancer.* 2007;109:1905-13.

976 [69] Hermelink K, Bhner M, Sckopke P, Neufeld F, Kaste J, Voigt V, et al. Chemotherapy and Post-
977 traumatic Stress in the Causation of Cognitive Dysfunction in Breast Cancer Patients. *J Natl Cancer
978 Inst.* 2017;109.

979 [70] Jansen CE, Cooper BA, Dodd MJ, Miaskowski CA. A prospective longitudinal study of
980 chemotherapy-induced cognitive changes in breast cancer patients. *Supportive Care in Cancer.*
981 2011;19:1647-56.

982 [71] Jim HSL, Donovan KA, Small BJ, Andrykowski MA, Munster PN, Jacobsen PB. Cognitive
983 functioning in breast cancer survivors: A controlled comparison. *Cancer.* 2009;115:1776-83.

984 [72] Jung MS, Cimprich B. Cognitive deficits in Korean women treated with chemotherapy for breast
985 cancer. *Cancer Nursing.* 2014;37:E31-E42.

986 [73] Kesler SR, Rao A, Blayney DW, Oakley-Girvan IA, Karuturi M, Palesh O. Predicting Long-Term
987 Cognitive Outcome Following Breast Cancer with Pre-Treatment Resting State fMRI and Random
988 Forest Machine Learning. *Front Hum Neurosci.* 2017;11:555.

989 [74] Kreukels BPC, Van Dam FSAM, Ridderinkhof KR, Boogerd W, Schagen SB. Persistent
990 neurocognitive problems after adjuvant chemotherapy for breast cancer. *Clinical Breast Cancer.*
991 2008;8:80-7.

992 [75] Mehlsen M, Pedersen AD, Jensen AB, Zachariae R. No indications of cognitive side-effects in a
993 prospective study of breast cancer patients receiving adjuvant chemotherapy. *Psycho-Oncology.*
994 2009;18:248-57.

995 [76] Mehnert A, Scherwath A, Schirmer L, Schleimer B, Petersen C, Schulz-Kindermann F, et al. The
996 association between neuropsychological impairment, self-perceived cognitive deficits, fatigue and
997 health related quality of life in breast cancer survivors following standard adjuvant versus high-dose
998 chemotherapy. *Patient Education and Counseling*. 2007;66:108-18.
999 [77] Menning S, de Ruiter MB, Kieffer JM, Agelink van Rentergem J, Veltman DJ, Fruijt A, et al.
1000 Cognitive Impairment in a Subset of Breast Cancer Patients After Systemic Therapy—Results From a
1001 Longitudinal Study. *Journal of Pain and Symptom Management*. 2016;52:560-9.e1.
1002 [78] Reid-Arndt SA, Yee A, Perry MC, Hsieh C. Cognitive and psychological factors associated with
1003 early posttreatment functional outcomes in breast cancer survivors. *Journal of Psychosocial*
1004 *Oncology*. 2009;27:415-34.
1005 [79] Reid-Arndt SA, Hsieh C, Perry MC. Neuropsychological functioning and quality of life during the
1006 first year after completing chemotherapy for breast cancer. *Psycho-Oncology*. 2010;19:535-44.
1007 [80] Ruzich M, Ryan B, Owen C, Delahunty A, Stuart-Harris R. Prospective evaluation of cognitive
1008 function in patients with early breast cancer receiving adjuvant chemotherapy. *Asia-Pacific Journal*
1009 *of Clinical Oncology*. 2007;3:125-33.
1010 [81] Schagen SB, Muller MJ, Boogerd W, Rosenbrand RM, Van Rhijn D, Rodenhuis S, et al. Late
1011 effects of adjuvant chemotherapy on cognitive function: A follow-up study in breast cancer patients.
1012 *Annals of Oncology*. 2002;13:1387-97.
1013 [82] Schagen SB, Muller MJ, Boogerd W, Mellenbergh GJ, van Dam FSAM. Change in cognitive
1014 function after chemotherapy: A prospective longitudinal study in breast cancer patients. *Journal of*
1015 *the National Cancer Institute*. 2006;98:1742-5.
1016 [83] Shilling V, Jenkins V, Morris R, Deutsch G, Bloomfield D. The effects of adjuvant chemotherapy
1017 on cognition in women with breast cancer--preliminary results of an observational longitudinal
1018 study. *Breast*. 2005;14:142-50.
1019 [84] Stewart A, Collins B, Mackenzie J, Tomiak E, Verma S, Bielajew C. The cognitive effects of
1020 adjuvant chemotherapy in early stage breast cancer: A prospective study. *Psycho-Oncology*.
1021 2008;17:122-30.
1022 [85] Stouten-Kemperman MM, De Ruiter MB, Boogerd W, Veltman DJ, Reneman L, Schagen SB. Very
1023 late treatment-related alterations in brain function of breast cancer survivors. *Journal of the*
1024 *International Neuropsychological Society*. 2015;21:50-61.
1025 [86] Van Dyk K, Bower JE, Crespi CM, Petersen L, Ganz PA. Cognitive function following breast cancer
1026 treatment and associations with concurrent symptoms. *npj Breast Cancer*. 2018;4.
1027 [87] Vearncombe KJ, Rolfe M, Wright M, Pachana NA, Andrew B, Beadle G. Predictors of cognitive
1028 decline after chemotherapy in breast cancer patients. *Journal of the International*
1029 *Neuropsychological Society*. 2009;15:951-62.
1030 [88] Wefel JS, Saleeba AK, Buzdar AU, Meyers CA. Acute and late onset cognitive dysfunction
1031 associated with chemotherapy in women with breast cancer. *Cancer*. 2010;116:3348-56.
1032 [89] Wieneke MH, Dienst ER. Neuropsychological assessment of cognitive functioning following
1033 chemotherapy for breast cancer. *Psycho-Oncology*. 1995;4:61-6.
1034 [90] Jackson D, Turner R. Power analysis for random-effects meta-analysis. *Research synthesis*
1035 *methods*. 2017;8.
1036 [91] Castellon SA, Ganz PA, Bower JE, Petersen L, Abraham L, Greendale GA. Neurocognitive
1037 performance in breast cancer survivors exposed to adjuvant chemotherapy and tamoxifen. *Journal*
1038 *of Clinical and Experimental Neuropsychology*. 2004;26:955-69.
1039 [92] Krolak D, Collins B, Weiss L, Harris C, Van der Jagt R. Cognitive function and its relationship to
1040 other psychosocial factors in lymphoma survivors. *Support Care Cancer*. 2017;25:905-13.
1041 [93] Boscher C, Joly F, Clarisse B, Humbert X, Grellard JM, Binarelli G, et al. Perceived Cognitive
1042 Impairment in Breast Cancer Survivors and Its Relationships with Psychological Factors. *Cancers*
1043 (Basel). 2020;12.
1044 [94] Schagen SB, Das E, Vermeulen I. Information about chemotherapy-associated cognitive
1045 problems contributes to cognitive problems in cancer patients. *Psychooncology*. 2012;21:1132-5.

1046 [95] Vardy JL, Dhillon HM, Pond GR, Rourke SB, Bekele T, Renton C, et al. Cognitive Function in
1047 Patients With Colorectal Cancer Who Do and Do Not Receive Chemotherapy: A Prospective,
1048 Longitudinal, Controlled Study. *J Clin Oncol*. 2015;33:4085-92.
1049 [96] Bray VJ, Dhillon HM, Vardy JL. Systematic review of self-reported cognitive function in cancer
1050 patients following chemotherapy treatment. *Journal of Cancer Survivorship*. 2018;12:537-59.
1051 [97] Argyriou AA, Assimakopoulos K, Iconomou G, Giannakopoulou F, Kalofonos HP. Either Called
1052 "Chemobrain" or "Chemofog," the Long-Term Chemotherapy-Induced Cognitive Decline in Cancer
1053 Survivors Is Real. *Journal of Pain and Symptom Management*. 2011;41:126-39.
1054 [98] Loh KP, Janelins MC, Mohile SG, Holmes HM, Hsu T, Inouye SK, et al. Chemotherapy-related
1055 cognitive impairment in older patients with cancer. *J Geriatr Oncol*. 2016;7:270-80.
1056 [99] Hunter JP, Saratzis A, Sutton AJ, Boucher RH, Sayers RD, Bown MJ. In meta-analyses of
1057 proportion studies, funnel plots were found to be an inaccurate method of assessing publication
1058 bias. *J Clin Epidemiol*. 2014;67:897-903.
1059