

Supplementary Tables and Figures


Supplementary Table S1: Frequency and proportion of clinical conditions among crash-involved drivers in New Jersey, ages 68 years and older, 2007-2017 (n=197,122).

Clinical conditions ^a	N (%)
Cataracts	125,865 (63.9)
Ischemic heart disease	108,452 (55.0)
Rheumatoid/Osteoarthritis	101,622 (51.6)
Diabetes	80,284 (40.7)
Glaucoma	53,272 (27.0)
Heart failure	51,814 (26.3)
Acquired hypothyroidism	51,090 (25.9)
Peripheral vascular disease	49,592 (25.2)
Chronic Obstructive Pulmonary Disease (COPD)	42,983 (21.8)
Chronic kidney disease	39,549 (20.1)
Chronic pain fatigue and fibromyalgia	38,906 (19.7)
Depression	37,135 (18.8)
Sensory deafness and hearing impairment	27,344 (13.9)
Anxiety disorders	25,373 (12.9)
Stroke or Transient Ischemic Attack (TIA)	24,287 (12.3)
Alzheimer's disease and related or senile dementia	15,361 (7.8)
Acute myocardial infarction	8,117 (4.1)
Migraine and other chronic headache	5,444 (2.8)
Mobility impairments	3,698 (1.9)
Epilepsy	2,914 (1.5)
Bipolar disorder	2,637 (1.3)
Schizophrenia and other psychotic disorders	2,043 (1.0)
Traumatic Brain Injury (TBI) & nonpsychotic mental disorders due to brain damage	1,533 (0.8)
Personality disorders	970 (0.5)
Spinal cord injury	664 (0.3)
Attention-Deficit/Hyperactivity Disorder and other conduct disorders (ADHD) ^b	638 (0.3)
Multiple sclerosis and transverse myelitis	543 (0.3)
Sensory blindness and visual impairment	306 (0.2)
Muscular dystrophy	63 (0.0)

%=Percentage

^a The frequencies and proportions for the presence of clinical conditions are displayed in the table, and those for the absence of conditions are omitted.

^b We kept the label for ADHD as assigned by the CCW although ADHD may no longer be considered a conduct disorder by many professional societies.



Supplementary Figure S1. Forest plot of differences in mean Euclidean distance from residential address to crash location and associated 95% confidence intervals (CI) among crash-involved drivers in New Jersey with and without the clinical conditions, ages 68 years and older, 2007-2017. Each difference in mean was found by subtracting the mean Euclidean distance of the group without the clinical condition from the group with it. CI=Confidence Interval, CL=Confidence Limit, TBI=Traumatic Brain Injury, ADHD=Attention-Deficit/Hyperactivity Disorder