COVID-19 OUTCOMES IN PREGNANCY: A REVIEW OF 275 SCREENED STUDIES

Rupalakshmi Vijayan MD*, Hanna Moon MD, Jasmine Joseph MD, Madiha Zaidi MD, Chhaya Kamwal MD, Andrelle Senatus MD, Shavy Nagpal MD**, Miguel Diaz MD Program Director FM Residency

*Division of Research, Larkin Community hospital, South Miami, Florida

** Corresponding author: Department of Surgery, The Research Institute of St. Joe's Hamilton, ON, Canada

rupavijayan92@gmail.com, hanna0517@gmail.com, jasminejoseph131@gmail.com,
madiha11291@gmail.com, ckamwal@gmail.com, dra509andrelle@gmail.com,
mdiaz@larkinhospital.com

ABSTRACT:

In December 2019, a novel strain of severe acute respiratory syndrome (SARS-CoV-2), was declared as a cause of respiratory illness, called coronavirus 2019 (COVID-19), characterized by fever and cough. In diagnostic imaging, the afflicted population showed pathognomonic findings of pneumonia. What started out as an epidemic in China, rapidly spread across geographical locations with a significant daily increase in the number of affected cases. According to the World Health Organization (WHO) reports, the range of worldwide mortality is 3 to 4%. Maternal adaptations and immunological changes predispose pregnant women to a prolonged and severe form of pneumonia, which results in higher rates of maternal, fetal, and neonatal morbidity and mortality. There is limited data about the consequences of COVID-19 in pregnancy, thereby limiting the prevention, counseling, and management of these patients. The objective of this literature review is to explore pregnancy and perinatal outcomes of COVID-19, complications, morbidity, and mortality in this sub-population. We conducted a literature review pertaining to COVID-19 and pregnancy in databases such as: PubMed, Google Scholar, and Science Direct. The studies we chose to focus on were systematic reviews, meta-analysis, case series, and case reports. Twenty four articles were reviewed regarding COVID-19 and pregnancy, complications and their outcomes. Due to immunological changes during pregnancy as evidenced by the flaring of auto-immune diseases; pregnant women may be at an increased risk for infection. Women (19.7%) who had underlying comorbidities such as gestational DM, HTN, hypothyroidism, and autoimmune disease, COPD, or HBV infection were considered high risk. The most common maternal outcomes were premature rupture of membranes (PROM) and pre-eclampsia. Asthma was the most common comorbidity associated with maternal mortality. The most common neonatal complications were fetal distress leading to NICU admissions and preterm birth <37 weeks. The most common laboratory changes were elevated CRP and lymphocytopenia. Most patients underwent C-section due to their underlying comorbidities. Pregnant and lactating women did not shed viral particles through their vaginal mucus and milk, as evidenced by negative nucleic-acid tests of these secretions. Neonatal infections as demonstrated by positive RT-PCR were rare, but direct evidence supporting intrauterine transmission was not confirmed. Direct evidence indicating vertical transmission of COVID-19 is not available, but risk for transmission cannot be ruled out. Pregnant women should be closely monitored due to increased risk of adverse outcomes.

KEYWORDS:

COVID-19, Maternal, Neonates, Pregnancy, Pandemic and SARS-COV-2

INTRODUCTION:

In December 2019, a novel strain of severe acute respiratory syndrome (SARS-CoV-2), was declared a cause of respiratory illness, called coronavirus 2019 (COVID-19), classically characterized by fever and cough. In diagnostic imaging, the afflicted population exhibited pathognomonic findings of pneumonia. What started out as an epidemic in China, rapidly spread across geographical locations, with a significant daily increase in the number of affected cases. According to the World Health Organization (WHO) reports, the range of worldwide mortality is 3 to 4%.

Pregnancy is a myriad of immunological states. The maternal immune system prepares itself to establish and maintain tolerance to the fetus's allergenicity, as well as retain the ability to protect the body against microbial invasion. A successful pregnancy relies on finely tuned immune adaptations, both systemically and locally. Maternal immunological states actively adapt and change with the growth and development of a fetus at different gestational stages: pro-inflammatory state (beneficial to the implantation and placentation of the embryo) in the first trimester, anti-inflammatory state (helpful for fetal growth) in the second trimester, and a second proinflammatory state (preparing for the initiation of parturition) in the third trimester [1].

Maternal adaptations and immunological changes during pregnancy tend to predispose pregnant women to a prolonged and severe form of pneumonia, which ultimately results in higher rates of maternal, fetal, and neonatal morbidity and mortality.

Systemic maternal viral infections can also play a factor in affecting the outcome of a pregnancy [2]. Previously, studies have shown that SARS infections during pregnancy can lead to high rates of spontaneous abortion, premature birth, and intrauterine growth restriction [3]. However, there has not been evidence of vertical transmission of SARS infections from the mother to the child [3]. Therefore, these pregnancy complications may be caused by the direct effect of the viruses on mothers. Although current evidence is limited, we cannot disregard the potential risks among pregnant women and their fetus. Recent literature indicates that in severe cases, COVID-19 infection is associated with a cytokine-storm, which is characterized by increased plasma concentrations of interleukins 2 (IL-2), IL-7, IL-10, granulocyte-colony stimulating factor, interferon- γ -inducible protein 10, monocyte chemoattractant protein 1, macrophage inflammatory protein 1 alpha, and tumor necrosis factor α (TNF- α) [4], which may be caused by ADE [5]. Since pregnant women in their first and third trimesters are in the pro-inflammatory state, the cytokine-storm induced by SARS-CoV-2 may induce a more severe inflammatory state. Moreover, the occurrence of maternal inflammation as a result of viral infection during pregnancy can affect several aspects of fetal brain development and may lead to a wide range of neuronal dysfunctions and behavioral phenotypes that are unmasked later in postnatal life [6].

The most common symptom at the onset of COVID-19 infection is a fever, which can be associated with an increased risk of attention-deficit/hyperactivity disorder (ADHD) in the offspring during school years.

There is limited data about the consequences of COVID-19 in pregnancy, thereby limiting the prevention, counseling, and management of these patients. The objective of this literature review is to explore pregnancy and perinatal outcomes of COVID-19, complications, morbidity, and mortality in this sub-population.

METHODS:

Search method and strategy

A literature review was conducted pertaining to COVID-19 and pregnancy in databases such as: PubMed, Google Scholar, and Science Direct from December 2019 through May 31st, 2021. We used specific keywords: “COVID-19,” “Pregnancy,” “SARS-COV-2,” “Pandemic,” “Maternal,” and “Neonates.” The studies we chose to focus on were systematic reviews, meta-analysis, case series, and case reports.

Articles that did not have patient data, and those limited to specific co-morbidities and organ dysfunctions were excluded to avoid selection bias. Therefore, we had twenty four articles for the final review. Selected articles were independently reviewed by two authors. All disagreements were resolved with a discussion between the two authors, or with input from a third independent reviewer and mutually agreed upon by the authors.

Our review included studies from various countries from across the globe. Referencing was done according to guidelines using Endnote. PRISMA guidelines were followed ([Fig. 1](#))

Ethical approval and funding

This study did not require ethical approval as data was obtained from already available databases, and patients were not directly involved. No funding was obtained for this review.

Fig. 1 PRISMA diagram for Study design

RESULTS:

Our results produced a total of 789 studies and after removing duplicates, title, and abstract screening, we chose to focus on 134 studies. A manual search was performed to look for related articles. Twenty four studies were included in the final analysis [Table 1]. Although pregnant women have an immunosuppressed state due to the physiological changes during pregnancy, most patients suffered from mild to moderate COVID-19 pneumonia with no pregnancy loss. Furthermore, this proposes a pattern like the clinical characteristics of COVID-19 pneumonia as compared to that of other adult populations.

Pregnancy being an immunocompromised state, pregnant women are more at risk of contracting COVID-19. The placenta has been shown to have ACE2 receptors on the cytotrophoblast and syncytiotrophoblast, which is the route through which COVID-19 gains entry. Patients infected with COVID-19 and those who are pregnant are known to have decreased lymphocytes and inhibitory receptors.

Due to immunological changes during pregnancy, as evidenced by autoimmune diseases' flaring, pregnant women may be at an increased risk for the infection. Most women (19.7%) had underlying comorbidities such as gestational DM, HTN, hypothyroidism, autoimmune disease, COPD, or HBV infection that put them at an increased risk. The most common maternal outcomes were PROM and pre-eclampsia. Asthma was found to be the most common comorbidity associated with maternal mortality. Fetal distress leading to NICU admissions and preterm birth <37 weeks were the neonatal complications. The most common laboratory changes were elevated CRP and lymphocytopenia. The majority of the patients underwent C-section due to their underlying comorbidities. Pregnant and lactating women did not shed viral particles through their vaginal mucus and milk, as evidenced by these secretions' negative nucleic-acid tests. As demonstrated by positive RT-PCR, neonatal infections were rare, but direct evidence supporting intrauterine transmission was not confirmed.

The findings from this systematic review show that more than 90% of hospitalized pregnant women affected by CoV-SARS infections present radiological signs suggestive for pneumonia, detected either at chest x-ray or computerized tomography (CT) and the most common symptoms are fever, cough, and lymphopenia. Pregnancies affected by CoV-SARS infections have high rates of PTB before 34 and 37 weeks. Preeclampsia and cesarean delivery are also more common than in the general population. The pooled proportion of perinatal mortality is about 10%, while the most common adverse perinatal outcome is fetal distress, with more than half of the newborns admitted in the NICU. Moreover, clinical evidence of vertical transmission was not found in any of the newborns. However, these findings should be interpreted with caution in view of the very small number of included cases and heterogeneity in clinical presentation and perinatal management among the included cases. In a systematic review, women affected by COVID-19 disease had higher rates of preterm birth, and preeclampsia, while the fetus had a 2.4% rate of stillbirth, a 2.4% rate of neonatal death, and higher rate of admission to the NICU. Severe maternal morbidity as a result of COVID-19 and perinatal deaths were reported. Vertical transmission in relation to COVID-19 could not be ruled out.

DISCUSSION:

Pregnancy is an immune compromised state due to several physiologic changes, most commonly cardiorespiratory system changes that place pregnant women even at higher risk of acquiring COVID-19 infection. COVID-19- a newly emerged virus is one of the diseases that can lead to severe consequences; hence a collaborative approach to COVID-19 positive pregnant women is required. It is vital to identify clinical features of SARS-CoV-2 infection in pregnant women as early preventive intervention can lower mortality and morbidity.

The clinical characteristics of pregnant women with COVID-19 resemble those of non-pregnant women. The most common symptoms experienced by these women are fever, cough, dyspnea, fatigue, sore throat, and myalgia [7, 10, 11]. This is supported by a systematic review with 18 studies comprising of 114 pregnant women and determined that (87.5%) of them developed fever, (53.8%) cough followed by (22.5%) having fatigue, (11.3 %) SOB, and (7.5%) sore throat, (16.3%) myalgia {5,6}. 24 studies including 136 pregnant women with confirmed COVID infection were reviewed and found that all of these affected women had similar symptoms as mentioned above [8]. However, one of the recent studies showed that pregnant women with covid-19 are less likely to manifest symptoms such as fever, dyspnea, and myalgia, and are more likely to be admitted to the intensive care unit or needing invasive ventilation and extracorporeal membrane oxygenation than non-pregnant women of reproductive age[14]. Increased maternal age, high body mass index, non-white ethnicity, and pre-existing comorbidities were found to be associated with the development of severe disease in pregnancy[14]. Moreover, a high prevalence of asymptomatic patients (42.8%) testing positive for SARS Cov 2 further emphasises the need for testing in pregnancy[15].

The most significant blood work abnormalities seen were elevated C- reactive protein in 57 % and lymphocytopenia in 50 % of pregnant women. In addition to this, 81.7 % of all the 136 chest CT scan revealed ground-glass opacity to be the most common atypical imaging findings (9,10,11). Another study done on 15 pregnant women with COVID pneumonia showed Ground- glass opacity with consolidation on Chest CT Scan [10]. There are a few studies that highlight the effects of COVID 19 in early pregnancy of which most demonstrated no significant prenatal biochemical or ultrasound markers of fetal anomalies. This was evidenced by a lack of difference in mean nuchal translucency thickness or biochemical markers (pregnancy-associated plasma protein A, alpha-fetoprotein, human chorionic gonadotropin, unconjugated estriol) between cases and controls [16,17].

Although evidence suggesting vertical transmission has not been observed in most reported cases, its possibility cannot be excluded [19]. In a meta-analysis and systemic review including 936 neonates from mothers with coronavirus disease 2019, 27 neonates had a positive result for SARS Cove 2 viral RNA test using nasopharyngeal swab, indicating a pooled proportion of 3.2% (95% confidence interval, 2.2-4.3) for vertical transmission[20].

Low incidence of maternal viremia and non-overlapping placental ACE2 and TMPRSS2 expression were found to be associated with protection against placental infection and vertical transmission in maternal COVID-19. However, reduced trans placental transfer of anti-SARS-CoV-2 antibodies may leave neonates at risk for infection [18].

For maternal related complications, the analysis found that premature rupture of membrane, fetal distress, and stillbirth were the most common complications during pregnancy [12]. In 42 studies involving 548 people who were pregnant, severe COVID-19 infection was found to be strongly associated with preeclampsia, preterm birth, gestational diabetes, and low birth weight when compared with pregnant women having mild COVID 19 infection [21].

When compared with non-pregnant women of reproductive age with covid-19, the odds of admission to the intensive care unit (odds ratio 2.13, 95% confidence interval 1.53 to 2.95; seven studies, 601 108 women) and need for invasive ventilation (2.59, 2.28 to 2.94; six studies, 601 044 women) and ECMO (2.02, 1.22 to 3.34; two studies, 461 936 women) were higher in pregnant and recently pregnant women [14].

Moreover, Pregnancies affected by CoV infections have high rates of PTB before 37 and 34 weeks. Preeclampsia and cesarean delivery are also more common than in the general population. Cesarean delivery was the most common mode of delivery in COVID 19 positive pregnant women [22]. The rate of caesarean delivery was found to be 66.4% in a study involving 533 pregnant women with COVID 19 infection [23].

COVID 19 infection during pregnancy can be associated with adverse neonatal morbidities. This is supported by a prospective cohort study involving 88 159 infants from Sweden which showed that some neonatal outcomes (procedures and morbidities) were significantly more common in infants of SARS-CoV-2-positive women (n=2323) than in infants of comparator women (n=9275): assisted ventilation at birth (6.4% vs 5.1), intubation at birth (0.6% vs 0.3), admission for neonatal care (11.7% vs 8.4), respiratory distress syndrome (1.2% vs 0.5), use of CPAP (4.9% vs 3.8), mechanical ventilation (1.6% vs 0.5), any respiratory disorder (2.8% vs 2.0%), persistent pulmonary hypertension (0.3% vs 0.1%), antibiotic therapy (2.8% vs 2.0%), and hyperbilirubinemia (3.6% vs 2.5%)[24].

The pooled proportion of perinatal mortality is about 10%, while the most common adverse perinatal outcome is fetal distress, with more than half of the newborns admitted in the NICU. Women affected by COVID-19 infection had higher rates of preterm birth and preeclampsia, while the babies had a 2.4% rate of stillbirth, 2.4% of neonatal death, and higher NICU admission [13].

CONCLUSION:

COVID-19 being relatively new to the world has made the whole spectrum of population of all ages susceptible due to lack of herd immunity. Pregnancy in general makes women more susceptible to respiratory infections and since COVID predominantly affects lungs, it makes pregnant women a high risk for SARS CoV-2 infection. Due to the pro-inflammatory immune responses in various trimesters of pregnancy by the cytokine storm of COVID-19 infection, pregnant women afflicted with COVID-19 may experience more morbidity and mortality. The maternal infection and inflammatory responses that occur as a response to COVID-19 may affect the fetus even postnatally. With the ever-expanding pandemic, more stringent efforts are required to be made to protect pregnant mothers and growing fetuses. Direct evidence indicating vertical transmission of COVID-19 is not available, but risk for transmission cannot be ruled out. Pregnant women should be closely monitored due to increased risk of adverse outcomes. More studies are required to follow-up the pregnant women having COVID-19 in the first and second trimester to determine pregnancy outcomes and postnatal development of the fetus.

REFERENCES:

1. Liu, H., Wang, L. L., Zhao, S. J., Kwak-Kim, J et al (2020). Why are pregnant women susceptible to COVID-19? An immunological viewpoint. *Journal of reproductive immunology*, 139, 103122. <https://doi.org/10.1016/j.jri.2020.103122>
2. Racicot K., Mor G. Risks associated with viral infections during pregnancy. *J. Clin. Invest.* 2017;127:1591–1599. doi: 10.1172/JCI87490. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
3. Wong S.F. Pregnancy and perinatal outcomes of women with severe acute respiratory syndrome. *Am. J. Obstet. Gynecol.* 2004;191:292–297. doi: 10.1016/j.ajog.2003.11.019. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
4. Huang C. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet.* 2020;395(10223):497–506. doi: 10.1016/S0140-6736(20)30183-5. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
5. Tetro J.A. Is COVID-19 receiving ADE from other coronaviruses? *Microbes. Infect.* 2020 doi: 10.1016/j.micinf.2020.02.006. Feb 22. pii: S1286-4579(20)30034-4. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
6. Mor G. The unique immunological and microbial aspects of pregnancy. *Nat. Rev. Immunol.* 2017;17:469–482. doi: 10.1038/nri.2017.64. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
7. Dashraath P, Wong JLI, Lim MXK, Lim LM, Li S, Biswas A et al. Coronavirus disease 2019 (COVID-19) pandemic and pregnancy. *Am J Obstet Gynecol.* 2020 Jun;222(6):521-531. doi: 10.1016/j.ajog.2020.03.021. Epub 2020 Mar 23. PMID: 32217113; PMCID: PMC7270569.
8. Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet.* 2020 Mar 7;395(10226):809-815. doi: 10.1016/S0140-6736(20)30360-3. Epub 2020 Feb 12. Erratum in: *Lancet.* 2020 Mar 28;395(10229):1038. Erratum in: *Lancet.* 2020 Mar 28;395(10229):1038. PMID: 32151335; PMCID: PMC7159281.
9. Juan J, Gil MM, Rong Z, Zhang Y, Yang H, Poon LC. Effect of coronavirus disease 2019 (COVID-19) on maternal, perinatal and neonatal outcome: systematic review. *Ultrasound Obstet Gynecol.* 2020 Jul;56(1):15-27. doi: 10.1002/uog.22088. PMID: 32430957; PMCID: PMC7276742.
10. Yang Z, Wang M, Zhu Z, Liu Y. Coronavirus disease 2019 (COVID-19) and pregnancy: a systematic review. *J Matern Fetal Neonatal Med.* 2020 Apr 30;1-4. doi: 10.1080/14767058.2020.1759541. Epub ahead of print. PMID: 32354293.
11. Matar R, Alrahmani L, Monzer N, Debiante LG, Barbari E, Fares J et al . Clinical Presentation and Outcomes of Pregnant Women With Coronavirus Disease 2019: A Systematic Review and Meta-analysis. *Clin Infect Dis.* 2021 Feb 1;72(3):521-533. doi: 10.1093/cid/ciaa828. PMID: 32575114; PMCID: PMC7337697.
12. Liu D, Li L, Wu X, Zheng D, Wang J, Yang L et al. Pregnancy and Perinatal Outcomes of Women With Coronavirus Disease (COVID-19) Pneumonia: A Preliminary Analysis. *AJR Am J Roentgenol.* 2020 Jul;215(1):127-132. doi: 10.2214/AJR.20.23072. Epub 2020 Mar 18. Erratum in: *AJR Am J Roentgenol.* 2020 Jul;215(1):262. PMID: 32186894.

13. Di Mascio D, Khalil A, Saccone G, et al. Outcome of coronavirus spectrum infections (SARS, MERS, COVID-19) during pregnancy: a systematic review and meta-analysis. *Am J Obstet Gynecol MFM*. 2020;2(2):100107. doi:10.1016/j.ajogmf.2020.100107
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7104131/> Am J Obstet Gynecol MFM. 2020 Mar 25
14. Allotey J, Stallings E, Bonet M, Yap M, Chatterjee S, Kew T et al. Clinical manifestations, risk factors, and maternal and perinatal outcomes of coronavirus disease 2019 in pregnancy: living systematic review and meta-analysis *BMJ* 2020; 370 :m3320 doi:10.1136/bmj.m3320
15. Cosma, S, Borella, F, Carosso, A, et al. The “scar” of a pandemic: Cumulative incidence of COVID-19 during the first trimester of pregnancy. *J Med Virol*. 2021; 93: 537– 540.
<https://doi.org/10.1002/jmv.26267>
16. Cosma, S.; Carosso, A.R.; Borella, F.; Cusato, J.; Bovetti, M.; Bevilacqua, et al. Prenatal Biochemical and Ultrasound Markers in COVID-19 Pregnant Patients: A Prospective Case-Control Study. *Diagnostics* 2021, 11, 398. <https://doi.org/10.3390/diagnostics11030398>
17. N la Cour Freiesleben, P Egerup, K V R Hviid, E R Severinsen, A M Kolte, D Westergaard, L Fich Olsen et al . SARS-CoV-2 in first trimester pregnancy: a cohort study, *Human Reproduction*, Volume 36, Issue 1, January 2021, Pages 40–47, <https://doi.org/10.1093/humrep/deaa311>
18. Edlow AG, Li JZ, Collier AY, et al. Assessment of Maternal and Neonatal SARS-CoV-2 Viral Load, Transplacental Antibody Transfer, and Placental Pathology in Pregnancies During the COVID-19 Pandemic. *JAMA Netw Open*. 2020;3(12):e2030455. doi:10.1001/jamanetworkopen.2020.30455
19. Wong, Y.P.; Khong, T.Y.; Tan, G.C. The Effects of COVID-19 on Placenta and Pregnancy: What Do We Know So Far? *Diagnostics* 2021, 11, 94. <https://doi.org/10.3390/diagnostics11010094>
20. Alexander M. Kotlyar, Olga Grechukhina, Alice Chen, Shota Popkhadze, Alyssa Grimshaw, Oded Tal et al. Vertical transmission of coronavirus disease 2019: a systematic review and meta-analysis, *American Journal of Obstetrics and Gynecology*,
21. Wei SQ, Bilodeau-Bertrand M, Liu S, Auger N. The impact of COVID-19 on pregnancy outcomes: a systematic review and meta-analysis. *CMAJ*. 2021 Apr 19;193(16):E540-E548. doi: 10.1503/cmaj.202604. Epub 2021 Mar 19. PMID: 33741725; PMCID: PMC8084555.
22. Papapanou, M.; Papaioannou, M.; Petta, A.; Routsis, E.; Farmaki, M.; Vlahos, N.; Siristatidis, C. Maternal and Neonatal Characteristics and Outcomes of COVID-19 in Pregnancy: An Overview of Systematic Reviews. *Int. J. Environ. Res. Public Health* 2021, 18, 596.
<https://doi.org/10.3390/ijerph18020596>
23. Sahin D, Tanacan A, Erol SA, Anuk AT, Yetiskin FDY, Keskin HL et al. Updated experience of a tertiary pandemic center on 533 pregnant women with COVID-19 infection: A prospective cohort study from Turkey. *Int J Gynaecol Obstet*. 2021 Mar;152(3):328-334. doi: 10.1002/ijgo.13460. Epub 2020 Dec 12. PMID: 33131057.
24. Norman M, Navér L, Söderling J, et al. Association of Maternal SARS-CoV-2 Infection in Pregnancy With Neonatal Outcomes. *JAMA*. Published online April 29, 2021.
doi:10.1001/jama.2021.5775

Pregnancy and COVID-19: A Systematic Review					
1	Why are pregnant women susceptible to COVID-19? An immunological viewpoint	Hong Liu et al.	Review	2020	Attributed increased risk of COVID-19 infection to changes of immune system during pregnancy and cytokine-storm characteristic of COVID-19
2	Risks associated with viral infections during pregnancy	Karen Racicot and Gil Mor	Review	2017	Discussed effects of maternal, placental, and fetal viral infection to pregnancy outcomes including maternal health and fetal development
3	Pregnancy and perinatal outcomes of women with severe acute respiratory syndrome	Shell F Wong et al.	Multi-center observational	2003	Summary of adverse pregnancy outcomes during SARS epidemic
4	Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China	Chaolin Huang et al.	Case cohort	2020	Summarized epidemiology and sequelae of COVID-19 infection
5	Is COVID-19 receiving ADE from other coronaviruses?	Jason A. Tetro		2020	Introduced the possible association between previous exposure to other forms of coronavirus and antibody-dependent enhancement during COVID-19 infection
6	The unique immunological and microbial aspects of pregnancy	Gil Mor et al.	Review	2017	Discussed the dynamics of immune responses during pregnancy
7	Coronavirus disease 2019 (COVID-19) pandemic and pregnancy	Dashraath et al.	Review	2020	Review of COVID-19 infection in pregnancy woman, and assessed the use of chloroquine as possible treatment
8	Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records	Chen et al.	Review	2020	Investigated intrauterine transmission of COVID-19 by analyzing perinatal and neonatal outcomes
9	Effects of coronavirus disease 2019 (COVID-19) on maternal, perinatal and neonatal outcomes: a systematic review	Juan et al.	Review	2020	Assessed perinatal and neonatal outcomes of COVID-19 infection in pregnancy women
10	Coronavirus disease 2019 (COVID-19) and pregnancy: a systematic review	Yang et al.	Review	2020	Reported the similarities of the clinical characteristics of pregnant women and non-pregnant population,
11	Clinical Presentation and Outcomes of Pregnant Women With Coronavirus	Matar et al.	Review & Meta-analysis	2020	Reported increased rate of preterm birth and cesarean delivery in

12	Disease 2019: A Systematic Review and Meta-analysis				pregnant COVID-19 patients
	Pregnancy and Perinatal Outcomes of Women With Coronavirus Disease (COVID-19) Pneumonia: A Preliminary Analysis	Liu et al.	Original research	2020	Attempted to provide initial evidence for treatment guidance of pregnancy women with COVID-19 pneumonia by reporting early findings of chest CT and lab abnormalities
13	Outcome of coronavirus spectrum infections (SARS, MERS, COVID-19) during pregnancy: a systematic review and meta-analysis	Mascio et al.	Review & Meta-analysis	2020	Reported increased risk of adverse perinatal outcomes in pregnant women with COVID-19 infection
14	Clinical manifestations, risk factors, and maternal and perinatal outcomes of coronavirus disease 2019 in pregnancy: living systematic review and meta-analysis	Allotey J et al	Review & Meta-analysis	2020	Reported increased ICU admission or need for invasive ventilation in pregnant women with COVID-19.
15	The “scar” of a pandemic: Cumulative incidence of COVID-19 during the first trimester of pregnancy	Cosma et al	Cohort study	2020	Confirmed high prevalence of asymptomatic COVID-19 in pregnant women.
16	Prenatal Biochemical and Ultrasound Markers in COVID-19 Pregnant Patients: A Prospective Case-Control Study	Cosma et al	Case Control study	2021	Confirmed absence of any unfavorable prenatal biochemical and ultrasound markers of fetal anomalies in COVID-19-positive patients.
17	SARS-CoV-2 in first trimester pregnancy: a cohort study.	N la Cour Freiesleben et al	Cohort study	2021	Reported no difference in Nuchal translucency thickness at the first trimester scan and no increased risk of pregnancy loss in women with SARS-CoV-2 infection in the first trimester compared to pregnant women without infection.
18	Assessment of Maternal and Neonatal SARS-CoV-2 Viral Load, Transplacental Antibody Transfer, and Placental Pathology in Pregnancies During the COVID-19 Pandemic.	Edlow AG et al	Prospective Cohort study	2020	Assessed Maternal and neonatal viral load of SARS CoV-2, Transplacental antibody transfer and Possibility of vertical transmission.
19	The Effects of COVID-19 on Placenta and Pregnancy: What	Wong et al	Review	2021	Discussed the effects of COVID 19 on placenta and

	Do We Know So Far?				the possibility of vertical transmission of SARS CoV-2.
20	Vertical transmission of coronavirus disease 2019: a systematic review and meta-analysis	Alexander M. Kotlyar et al	Systemic Review & Meta analysis	2021	Suggested the possibility of vertical transmission of SARS CoV-2
21	The impact of COVID-19 on pregnancy outcomes: a systematic review and meta-analysis	Wei SQ et al	Systemic Review & Meta analysis	2021	Discussed the impacts of COVID-19 on pregnancy and summarized the outcomes.
22	Maternal and Neonatal Characteristics and Outcomes of COVID-19 in Pregnancy: An Overview of Systematic Reviews.	Papapanou et al	Review	2021	Review of maternal and neonatal outcomes of COVID-19 in pregnancy.
23	Updated experience of a tertiary pandemic center on 533 pregnant women with COVID-19 infection: A prospective cohort study from Turkey	Sahin D et al	A prospective cohort study	2020	Reported increased rates of cesarean delivery and pregnancy complications in pregnant women with COVID-19 infection.
24	Association of Maternal SARS-CoV-2 Infection in Pregnancy With Neonatal Outcomes.	Norman M et al	Cohort study	2021	Reported association of neonatal morbidities and maternal SARS CoV-2 infection.

Table 1: Characteristics of studies