Correlation between times to SARS-CoV-2 symptom onset and secondary transmission undermines epidemic control efforts

Natalie M. Linton,^{1,2} Andrei R. Akhmetzhanov,³ Hiroshi Nishiura^{1,4*}

¹Kyoto University School of Public Health, Yoshidakonoe-cho, Sakyo-ku, Kyoto city, 606-8501, Japan

²Graduate School of Medicine, Hokkaido University, Kita 15 Jo Nishi 7 Chome, Kita-ku, Sapporo-shi, Hokkaido 060-8638, Japan

³College of Public Health, National Taiwan University, 17 Xu-Zhou Road, Taipei 10055, Taiwan

⁴Core Research for Evolutional Science and Technology (CREST), Japan Science and Technology Agency, Saitama, Japan

*Correspondence:

Hiroshi Nishiura, MD, PhD

nishiura.hiroshi.5r@kyoto-u.ac.jp; Tel.: +81-75-753-4456

Abstract

Severe acute respiratory coronavirus 2 (SARS-CoV-2) infections have been associated with substantial presymptomatic transmission, which occurs when the generation interval—the time between infection of an individual with a pathogen and transmission of the pathogen to another individual—is shorter than the incubation period—the time between infection and symptom onset. We collected a dataset of 257 SARS-CoV-2 transmission pairs in Japan and jointly estimated the mean generation interval (3.7–5.1 days) and mean incubation period (4.4–5.7 days) as well as measured their dependence (Kendall’s tau of 0.4–0.6), taking into consideration demographic and epidemiological characteristics of the pairs. The positive correlation between the two parameters demonstrates that reliance on isolation of symptomatic COVID-19 cases as a focal point of control efforts is insufficient to address the challenges posed by SARS-CoV-2 transmission dynamics. Accounting for this dependence within SARS-CoV-2 epidemic models can also improve model estimates.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Introduction

The generation interval and incubation period of an infectious disease are key epidemiological parameters used to inform outbreak response. The former describes the infectiousness of a pathogen in a host—it represents the time between when a host (an infector) is infected with a pathogen and when they transmit that pathogen to another host (an infectee). In contrast, the incubation period represents the time between infection and development of symptomatic disease. It reflects the pathogen replication rate and provides a basis for predicting prognosis. It also indicates how long an infection may remain unnoticed in an individual, and thus for emerging diseases it is used to determine quarantine periods.^{1,2} Together, the generation interval and incubation period provide insight into the intrinsic dynamics of infection and characterize the effectiveness of public health interventions on its control.

At the beginning of the coronavirus 2019 (COVID-19) pandemic, the mean incubation period of severe acute respiratory disease coronavirus 2 (SARS-CoV-2)—the pathogen causing COVID-19—was rapidly estimated.^{1–4} Estimates of the mean serial interval, which is the time between symptom onset in an infector and symptom onset in a person they infect, quickly followed.^{5–8} However, due to difficulty in ascertaining exposure times of cases and directionality of transmission between epidemiologically linked cases, few attempts were made to estimate the generation interval.^{6–10} Instead, the serial interval was often used as a proxy for the generation interval when estimating epidemiological quantities. For example, it has been used to estimate the basic reproduction number—the average number of persons infected by a single infector in a completely susceptible population—and the effective reproduction number—the average number of persons infected by a single infector in the presence of existing infections and interventions.^{11,12} However, use of the serial interval as a proxy for generation time can lead to biased estimates of the effective reproduction number due to factors such as differences in their variances, the presence of asymptomatic infections, and the ability for the serial interval to have negative values (infectee onset preceding infector onset).^{8,13–16}

Generation intervals depend on many factors, such as the ratio of infected to susceptible persons among contacts, as well as the behavior of infected persons.¹⁷ Actions such as self-isolation after symptom onset can shorten generation intervals by limiting the opportunity for infected individuals to infect others during their early symptomatic period. However, if an infected person can transmit the pathogen before symptom onset, then isolation of symptomatic persons alone is insufficient to control the spread.¹⁸ For pathogens such as SARS-CoV-2, interventions targeting nonsymptomatic cases appears vital¹⁹ due to the large fraction of presymptomatic and asymptomatic transmission.^{7,20,21}

Frequently, estimates of the generation interval of SARS-CoV-2 have been derived from the serial interval and formulated following implicit (and unsupported) assumptions that: i) there is no asymptomatic transmission,²² ii) the incubation period and generation interval are independent.²³ However, these assumptions are clearly flawed with respect to SARS-CoV-2 transmission. In the case of asymptomatic transmission, there is substantial evidence of transmission from asymptomatic infectors,^{20,24} as well as plenty

of asymptomatic cases demonstrating the existence of asymptomatic and presymptomatic infectees. Likewise, correlation between the generation interval and incubation period of SARS-CoV-2 was shown to be biologically plausible, as evidenced by the peak viral load of SARS-CoV-2 occurring around the time of symptom onset.²⁵ However, previous studies that considered such a correlation did not attempt to directly estimate it.²⁶

In this study, we provide direct evidence of their correlation and jointly estimate the generation interval and incubation period using transmission pairs identified in Japan in 2020. We assessed whether either interval or their correlation varied based on demographic and epidemiological characteristics. Accounting for correlation between the generation interval and incubation period can considerably help quantify SARS-CoV-2 transmission and improve characterization of the effectiveness of public health interventions by preventing underestimation of the proportion of presymptomatic transmission and the effect of isolation of symptomatic cases on epidemic control.

Methods

Ethics oversight

This study was approved by the Medical Ethics Board of the Graduate School of Medicine at Kyoto University (R2676). It uses data published online by public health jurisdictions in Japan.

Study population and setting

We compiled a dataset of COVID-19 transmission pairs using openly published case data from reporting jurisdictions (prefectures and cities) in Japan, focusing on detecting pairs for whom directionality of transmission could be determined with some degree of certainty. Cases were limited to those reported during the calendar year 2020. Jurisdiction reporting practices changed over time, with details generally becoming sparser over time, as concerns grew around infection-related stigmatization,⁴⁰ as well as in prefectures with large case loads.

Among the information publicly shared were links between cases and links to common exposures (e.g., a medical facility, event, or restaurant). However, clear statements as to who was the infector between linked cases or within clusters were generally not published. As well, dates of contact between cases and details of the type of link between cases were often only reported in detail if deemed to be important for public health action, limiting the number of cases for whom detailed epidemiological information related to their linkages were available. Therefore, assumptions about directionality of transmission were largely at the discretion of the authors, and in consequence we used the following bases for identifying linked cases as directional transmission pairs: 1) linkage of the infector (but not infectee) to a cluster; 2) the dates of contact, type of contact, and onset dates reported for linked cases provided some insight into directionality of transmission; 3) the infector or index case of a chain travelled to a location with increased/increasing transmission prior to onset; or 4) the infector or index case of a chain was presumed to have been infected while travelling

abroad. Households with >2 cases and links between cases in clusters where directionality of transmission and timings of contact could not be clearly identified were not selected.⁵⁵

We included pairs with infectors who had multiple possible exposures (their exposure period takes the lower and upper bounds of all possible exposures) but excluded possible pairs where potential infectees had multiple possible infectors, and it is possible that infectees with multiple potential infectors would have different contact patterns (and possibly be associated with shorter generation intervals) compared to infectees that had only one potential infector identified, as a susceptible person is likely to become infected more quickly if they are surrounded by multiple possible sources of infection.^{17,27} Further details regarding ascertainment of transmission pairs are available in the supplementary information.

Exposures were defined in relation to travel, contact with a confirmed case, or link to a cluster/common exposure. Reports of symptom onset in Japan were not restricted to any particular symptom, such as fever, but may have been reported as beginning with any of a variety of symptoms associated with SARS-CoV-2 infection such as fever, cough, fatigue, or runny nose.

Data stratification

The dataset including coarsely reported dates of exposure and contact were divided into strata to assess whether the generation interval, incubation period, or correlation between the two parameters would vary by subpopulation. Age (reported in deciles) was divided into three groups: cases under 30 years of age, cases 30–59 years of age, and cases 60+ years of age. Sex was reported as female or male. Separate age and sex strata were established for infectors and infectees. Type of contact between infector and infectee was divided into three categories: household contact, social contact-based interaction, and core community interaction. These divisions were made with public health interventions in mind. For example, social contact-based interaction includes types of contact that may not have occurred when local control measures were advised or a state of emergency was declared.⁴⁴

Generally, public health control measures in Japan promoted during 2020 focused on reducing the number of people individuals were physically in contact with in a given day, as well as reducing scenarios where the “Three C’s” — closed spaces, crowded places, and close-contact settings—were present.³⁸ Interventions in Japan included limiting the total number or proportion of people who can visit facilities and venues, limiting restaurant hours, encouraging staying at home and discouraging cross-prefecture travel, etc. Our definition of core community interaction, in contrast, focuses more on contact that occurs in schools, workplaces for general business, essential workplaces (medical facilities, care facilities, government services, etc.), and unknown sources of infection (community infection). Although these settings assigned to the core community interaction category may also be targeted by public health measures, they are perhaps less acutely affected by government decrees and social sentiment compared to settings more closely related to social contact-based interaction.⁴⁴

Japan experienced three waves of COVID-19 during 2020, with the third wave extending into 2021 (Extended Data Fig. 5). The first wave began with the first reported case, confirmed to be positive for SARS-CoV-2 on 16 January 2020. The second wave we set to begin on 1 June, which is around the center of the bottom of the trough between the peaks of the first and second waves. The third wave we set to begin on 1 October, which likewise is around the center of the bottom of the trough between the peaks of the second and third waves. Assignment to a given wave for each pair was determined by infector report date. Lastly, to check for differences given our basis for selecting transmission pairs we also stratified the dataset according to whether directionality was determined with respect to 1) importation from abroad, 2) linkage of the infector (but not infectee) to a cluster, 3) domestic travel by the infector to a location with increasing transmission, or 4) the timing and type of contact between cases in transmission chains.

Statistical analyses

Descriptive analyses and visualization were performed using R 4.1.0.⁵⁶ Bayesian parameter estimation was implemented in Stan using the cmdstanr interface to CmdStan 2.26.1.⁵⁷ Data and code are available at https://github.com/nlinton/covid19_generationinterval.

To assess correlation between the generation interval and incubation period of the infector we constructed a joint probability distribution for the generation interval and infector incubation period by use of copulas (multivariate cumulative distribution functions).^{58,59} The copulas we assessed included the Gaussian (normal), Clayton, Gumbel, and independence copulas. They are described in detail in the Supplementary Materials. Timing of pathogen transmission and symptom onset was estimated using interval censoring methods derived from Reich et al.⁶⁰ and adapted from previously published work.^{1,5} For all parameters, posterior point estimates are given by the 50th percentiles of the converged Markov chain Monte Carlo (MCMC) chains from 100,000 iterations, and the best combination of copula and parametric distributions were selected using weights from a Bayesian mixture model (see Supplementary Materials).

To consider the effect of correlation between the generation interval and incubation period on transmission, we simulated 10,000 pairs from the best-fit model of the jointly estimated generation interval and incubation period for all possible values of Kendall's tau from 0 to 1. For each simulated pair, we determined whether transmission was presymptomatic based on whether the incubation period was greater than the generation interval, and thereby calculated the proportion of presymptomatic transmission p for the 10,000 pairs for each value of Kendall's tau. We then considered that symptomatic transmission could result in a decrease in transmission as defined by the basic reproduction number R_0 —the average number of infectees generated by a single infector. We calculated the effective R_0 as $R_0^E = \varphi R_0 + (1 - \varphi)(1 - \varepsilon)R_0$ where φ is the proportion of presymptomatic transmission and ε is the percent reduction in transmission due to rapid isolation among symptomatic cases, considering $R_0 = 2.2^{4,30}$ and varied this by ± 0.7 , also considering $R_0 = 1.5$ and $R_0 = 2.9$.

Results

Characteristics of transmission pairs

Information on timing of exposure and onset for infectors as well as contact between infectors and infectees was obtained for 286 transmission pairs of confirmed cases reported in Japan during 2020, of which 257 pairs had symptom onset available for the infector. For the other 29 pairs the infectors were asymptomatic at time of report. Of the 257 pairs with symptomatic infectors, 49 (19.2%) had single dates reported for both infector exposure and contact between infector and infectee. Characteristics of the pairs in each dataset are shown in Table 1, while Extended Data Fig. 1 provides insight into the relationship between the empirical generation intervals, serial intervals, and incubation periods associated with these cases.

For the dataset of 257 pairs with symptomatic infectors, most infectors (50.2%) and infectees (44.4%) were between 30–59 years of age. There were fewer female infectors (30.7%) detected compared to female infectees (51.0%). Age and sex distributions of infectors and infectees are shown in Extended Data Fig. 2. Pairs were relatively evenly distributed between the three pandemic waves that occurred during 2020. Most pairs (53.7%) were linked to a cluster—an aggregation of cases with a common exposure, while other pairs were identified by having contact patterns indicative of directionality of transmission (21.0%), the infector had travel to another prefecture before onset (16.7%), or the infector was an imported case or otherwise linked to an imported case (8.6%). Given that it was easier to determine the directionality of pairs and obtain information on timing of exposure if the infector was linked to a cluster or had travel history, our dataset includes only a handful of infectors (1.6%) with household exposure. In contrast, nearly half (42.4%) of infectees were household/family members of their infectors. The single-date (49 pairs) and asymptomatic infector (28 pairs) datasets were similarly structured in terms of age and sex, though only one asymptomatic infector was detected during the first wave.

Joint estimates of the generation interval and incubation period

The jointly estimated mean generation interval ranged between 3.7 and 5.1 days, with the mean for the overall dataset estimated at 4.3 days (95% CrI: 4.0–4.7 days). In contrast, the estimated generation interval for the dataset of asymptomatic infectors was longer, at 4.6 days (3.9–5.5 days), resulting in a ratio of asymptomatic-to-symptomatic generation intervals of 1.1. The jointly estimated mean incubation period was consistently longer than the generation interval, ranging from 4.4–5.7 days, and estimated at 4.8 days (95% CrI: 4.4–5.1 days) for the overall dataset, providing evidence of presymptomatic transmission. The prior and posterior distributions of the generation interval and incubation period are shown in Extended Data Fig. 3. The generation interval and incubation period were positively correlated, with Kendall's tau ranging between 0.4–0.6 and estimated at 0.5 (95% CrI: 0.4–0.6) for the overall dataset (Table 2). For the dataset with single dates of reported exposure and contact, the generation interval was estimated at 4.4 days (95%

CrI: 3.9–5.0 days) while the mean incubation period was estimated at 4.9 days (95% CrI: 4.4–5.6 days). Kendall's tau was slightly higher than for the overall dataset, at 0.6 (95% CrI: 0.5–0.7).

The mean generation interval did not vary substantially between strata but was shortest for female infectors, at 3.7 days (95% CrI: 3.2–4.4 days). It was also shorter for the second wave of the epidemic (3.8 days, 95% CrI: 3.4–4.3 days) compared to the first wave (5.1 days, 95% CrI: 4.5–5.8 days).²⁷ However, the generation interval for the third wave was longer than that of the second wave—nearly as long as that of the first wave—at 4.5 days (95% CrI: 3.9–5.1 days). Estimates of the incubation period varied less between strata, although the mean incubation period for pairs linked to importation from other countries (mostly from the first wave) was a bit longer than the overall estimate, at 5.7 days (95% CrI: 4.6–6.8 days).

The Clayton copula—which emphasizes lower tail dependence—was the most frequently selected copula, although the Gumbel and Gaussian copulas were also selected for some strata. The Gumbel copula emphasizes upper tail dependence while the Gaussian copula does not consider tail dependence. The independence copula was never selected (Table 2). For the overall dataset, where the Clayton copula was selected, the lower tail dependence was 0.7 (95% CrI: 0.6–0.8), indicating that infectors with an extremely short incubation period would also be more likely to quickly transmit the virus given contact with a susceptible person (see Supplementary Materials). For the generation interval, the Weibull distribution was most often selected, although the gamma and lognormal distributions were selected for some strata. The lognormal distribution was the only distribution selected across all joint estimates of the incubation period. It is typically the best fit for infectious disease incubation period data,²⁸ including COVID-19 data.²⁹

Correlation, presymptomatic transmission, and control measures

We found that 63.2% of pairs experienced presymptomatic transmission, defined as the generation interval being shorter than the incubation period, by simulating 10,000 transmission pairs from our fitted estimates of the generation interval, incubation period, and Kendall's tau (Table 2). Using this fit, we varied Kendall's tau for the same estimates of the generation interval and incubation period and show that as Kendall's tau approached zero (independence) the proportion of presymptomatic transmission reached a lower boundary of 54.2%. Conversely, as Kendall's tau approached 1 (complete dependence), the proportion of presymptomatic transmission increased to nearly 100% (Figure 3a), and the difference between symptom onset and transmission became so small that they mostly occurred on the same day, with only a small portion of presymptomatic transmission occurring outside of one day before or after symptom onset, and no symptomatic transmission occurring.

The average difference between the generation interval and incubation period across all data subsets and strata was 0.4 days, indicating a mean time from onset of symptoms to transmission of -0.4 days. The probability density function of the time from onset of symptoms to transmission fitted with a normal distribution based on simulated data with Kendall's tau varied between 0.2, 0.5, and 0.9, is shown in Figure 3c. The mean was centered at -0.4 days, and lower correlation resulted in a larger standard deviation.

Using the same simulated dataset, Kendall's tau was varied against the fraction of transmission reduced by case isolation. Increasing Kendall's tau indicated greater difficulty in controlling transmission via isolation alone given the same level of reduction in transmission due to rapid isolation φ (Figure 4). Given a basic reproduction number of R_0 of 2.2 or 2.9,³⁰ the effective R_0 (denoted R_0^E) failed to reduce below the epidemic threshold of 1 (Figure 4b and c). Moreover, as Kendall's tau approaches 1 and the proportion of presymptomatic transmission (with a mean time from onset of symptoms to transmission of approximately 0) approaches 100%, control through isolation alone becomes impossible, and there is no difference between R_0 and R_0^E . When case isolation does not occur promptly following onset (lower φ) the dependence between the generation interval and incubation period has less impact on the reduction of R_0^E .

Discussion

The generation interval underpins many infectious disease models,^{31,32} and here we provide insight into the generation interval of COVID-19 over time and across different characteristics of transmission pairs, providing one of the most comprehensive characterizations of the generation interval of wild-type COVID-19 to date. In addition, we quantitatively measured the relationship between the generation interval and incubation period of COVID-19, the lack of which was identified as a limiting factor in previous studies.^{7,23,33} From transmission pairs identified using publicly available data reported in Japan during 2020 we found positive correlation between the generation interval and incubation period with a Kendall's tau ranging from 0.4–0.6. The mean generation interval was consistently shorter than the mean incubation period when jointly estimated, with the former ranging between 3.7–5.1 days and the latter between 4.4–5.7 days, indicating consistent presence of presymptomatic transmission.

The means of the jointly estimated generation interval and incubation period are in line with those reported elsewhere^{6–8,10} with the mean generation interval reported here—4.3 days—falling in the range of 2.8–7.5 days previously reported (see Supplementary Table 1 and Extended Data Fig. 4). The positive correlation between the generation interval and incubation period indicates that for symptomatic cases, onset is tied to infectiousness. This finding supports evidence shown in virological studies.²⁵ The estimate of the generation interval estimated with asymptomatic infectors (4.6 days, 95%: 3.8, 5.6 days) was longer than the jointly estimated generation interval using symptomatic infectors indicating that infectiousness in asymptomatic cases may be more persistent than in symptomatic infections—potentially leading to an underestimation of R_0 using estimates from symptomatic pairs.³⁴ Although evidence has indicated that asymptomatic COVID-19 cases are less infectious than symptomatic cases,²⁰ asymptomatic cases nonetheless play a notable role in epidemic dynamics.⁶

The proportion of presymptomatic transmission among symptomatic cases estimated in this study, 63.2%, is higher than estimates reported by published studies using data from early in the pandemic,^{6,25} but is similar to other estimates.^{7,8,35} Among those pairs with presymptomatic transmission, 33.5% had transmission occurring within one day of infector onset, and 49.1% within two days of infector onset.

Combining estimates of asymptomatic transmission (in the range of 18–30%)^{36,37} with the estimate of presymptomatic transmission shared here, the proportion of nonsymptomatic transmission could feasibly reach 90%. Thus, interventions such as physical distancing that do not depend on detection of potential infectors while they are not showing symptoms, enhanced surveillance to detect nonsymptomatic cases, and contact tracing to identify exposed individuals while their infected contacts are not symptomatic are crucial for COVID-19 control.^{6,18,19}

Data collection for this study focused on high certainty of directionality of transmission based on publicly announced epidemiological data. In contrast to most other countries, Japan applied backward contact tracing methods from the beginning of the pandemic in an effort to prevent large clusters of cases,³⁸ making it an apt setting for obtaining transmission pair data. This is because links between cases—and particularly those related to clusters—were more likely to have been detected compared to countries where backward contact tracing was not conducted. However, the timing of COVID-19 testing plays an important role in case ascertainment,³⁹ and infected persons who were epidemiologically linked to COVID-19 cases but did not become symptomatic after initially testing negative for SARS-CoV-2 may have been missed as cases. In addition, Japan did not promote widespread community viral testing, and this perhaps limited the number of unlinked cases that may have otherwise been detected and retrospectively linked to others during the epidemic. As well, public health jurisdiction reporting practices changed over time, with details generally becoming sparser once daily incidence became high enough to wear contact tracing capacity thin, and also towards the end of the year as concerns grew around infection-related stigmatization.⁴⁰

The shorter mean generation interval during the second and third waves of pandemic in Japan compared to the first wave may in part reflect the increase in prevalence of infection, as increased competition between infectious individuals to find susceptible contacts can lead to contraction of the generation interval.^{17,27,41} Shorter generation intervals were also noted in the United Kingdom during September–November 2020, when there was a rise in the number of new cases.³⁵ However, our results indicate that increases in incidence do not perforce lead to contraction of the generation interval. The larger value obtained for the third wave in Japan, which had a higher peak than the second wave (Extended Data Fig. 5).

During the first epidemic wave, state of emergency declarations nationwide. However, during the second wave and the 2020 half of the third wave, no such preventative measures were introduced. Conversely, campaigns intending to restart the Japanese economy following the difficulties caused by the first wave of COVID-19 were developed and implemented. In particular, the GoTo Travel campaign, which offered discounts on travel inside Japan, was a fixture of the second and third waves (Extended Data Fig. 5). The campaign began just before the peak of the second wave and was associated with an increase in COVID-19 cases reporting inter-prefecture travel.⁴² Of our pairs identified for the second wave, only 20.0% of infectors were reported before the start of the GoTo travel campaign. Although we did not find that our pairs where the infector had domestic travel experienced longer generation intervals (Figure 2), travel can

left-censor the time following infection when an infector who travelled had contact with an infectee who did not travel with them.

Most identified transmission pairs had contact in the household or in settings related to social behavior, such as eating at restaurants, visiting nightlife, singing karaoke, attending sports events, listening to live music, visiting gyms, or meeting with friends, relatives, acquaintances, etc. (Table 1). These types of social contact settings have also been associated with SARS-CoV-2 transmission in other countries.⁴³ In Japan, settings for social contact first are the first to be requested to be restricted by prefectural and local governments when control measures or a state of emergency was deemed necessary to reduce case incidence,⁴⁴ though such emergency measures were not implemented during the second or third wave portions of 2020. Similar interventions focused on limiting social contact were implemented in other parts of the world,⁴⁵ though other countries had a greater focus on reducing formalized community contact, such as by moving schools and workplaces online.^{46,47}

As SARS-CoV-2 variants begin to dominate transmission in many countries,^{48–50} it remains to be seen whether the mean and variance of the generation interval and incubation period for the new variants of concern (VOC) will be similar to estimates presented in this, or previous studies. It has been suggested that the Alpha (Pango lineage B.1.1.7) variant could have longer generation intervals,⁵¹ and may therefore be more responsive to interventions targeted towards speed, such as contact tracing.⁵² However, a study using Singapore transmission pairs did not find a large difference (>1 day) between the serial interval of wild-type and Delta (Pango lineage B.1.617.2) variant SARS-CoV-2 infections.⁵³ As the mean serial interval can approximate the mean generation interval, this finding may indicate that the generation interval of the Delta variant will not vary much from that of wild-type SARS-CoV-2. Similarly, a preprint analysis of Delta variant cases from China estimated an incubation period of 5.8 days,⁵⁴ which resembles the results from a meta-analysis on wild-type SARS-CoV-2 incubation periods.²⁹

Whether the correlation between the generation interval and incubation period would be weaker or stronger than has been presented here remains to be seen. However, this study provides detailed estimates of the generation interval, incubation period, and their correlation for different groups and time periods in Japan. In doing so, it provides basis for consideration of correlation going forward.

References

1. Linton NM, Kobayashi T, Yang Y, et al. Incubation period and other epidemiological characteristics of 2019 novel coronavirus infections with right truncation: A statistical analysis of publicly available case data. *J Clin Med*. 2020;9(538). doi:10.3390/jcm9020538
2. Lauer SA, Grantz KH, Bi Q, et al. The incubation period of coronavirus disease 2019 (COVID-19) from publicly reported confirmed cases: Estimation and application. *Ann Intern Med*. 2020;172(9):577-582. doi:10.7326/M20-0504

3. Backer JA, Klinkenberg D, Wallinga J. Incubation period of 2019 novel coronavirus (2019-nCoV) infections among travellers from Wuhan, China, 20–28 January 2020. *Eurosurveillance*. 2020;(January):1-6.
4. Li Q, Guan X, Wu P, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus–infected pneumonia. *N Engl J Med*. Published online 2020:1-9. doi:10.1056/NEJMoa2001316
5. Nishiura H, Linton NM, Akhmetzhanov AR. Serial interval of novel coronavirus (COVID-19) infections. *Int J Infect Dis*. Published online 2020:113332. doi:10.1016/j.ijid.2020.02.060
6. Ferretti L, Wymant C, Kendall M, et al. Quantifying SARS-CoV-2 transmission suggests epidemic control with digital contact tracing. *Science (80-)*. 2020;368(6491):eabb6936. doi:10.1126/science.abb6936
7. Tindale LC, Stockdale JE, Coombe M, et al. Evidence for transmission of covid-19 prior to symptom onset. *Elife*. 2020;9:1-34. doi:10.7554/eLife.57149
8. Ganyani T, Kremer C, Chen D, et al. Estimating the generation interval for coronavirus disease (COVID-19) based on symptom onset data, March 2020. *Eurosurveillance*. 2020;25(17):1-8. doi:10.2807/1560-7917.ES.2020.25.17.2000257
9. Deng Y, You C, Liu Y, Qin J, Zhou XH. Estimation of incubation period and generation time based on observed length-biased epidemic cohort with censoring for COVID-19 outbreak in China. *Biometrics*. 2020;(June):1-13. doi:10.1111/biom.13325
10. Li M, Liu K, Song Y, Wang M, Wu J. Serial interval and generation interval for imported and local infectors, respectively, estimated using reported contact-tracing data of COVID-19 in China. *Front Public Heal*. 2021;8(577431). doi:10.3389/fpubh.2020.577431
11. Jung S, Akhmetzhanov AR, Hayashi K, et al. Real-time estimation of the risk of death from novel coronavirus (COVID-19) infection: Inference using exported cases. *J Clin Med*. 2020;9(2):523. doi:10.3390/JCM9020523
12. Abbott S, Hellewell J, Thompson RN, et al. Estimating the time-varying reproduction number of SARS-CoV-2 using national and subnational case counts. *Wellcome Open Res*. 2020;5:112. doi:10.12688/wellcomeopenres.16006.1
13. Torneri A, Libin P, Scalia Tomba G, Faes C, Wood JG, Hens N. On realized serial and generation intervals given control measures: The COVID-19 pandemic case. *PLOS Comput Biol*. 2021;17(3):e1008892. doi:10.1371/journal.pcbi.1008892
14. Park SW, Sun K, Champredon D, et al. Forward-looking serial intervals correctly link epidemic growth to reproduction numbers. *Proc Natl Acad Sci*. 2021;118(2):e2011548118. doi:10.1073/pnas.2011548118
15. Knight J, Mishra S. Estimating effective reproduction number using generation time versus serial interval, with application to COVID-19 in the Greater Toronto Area, Canada. *Infect Dis Model*. 2020;5:889-896. doi:10.1016/j.idm.2020.10.009

16. Griffin J, Casey M, Collins Á, et al. Rapid review of available evidence on the serial interval and generation time of COVID-19. *BMJ Open*. 2020;10(11). doi:10.1136/bmjopen-2020-040263
17. Nishiura H. Time variations in the generation time of an infectious disease: Implications for sampling to appropriately quantify transmission potential. *Math Biosci Eng*. 2010;7(4):851-869. doi:10.3934/mbe.2010.7.851
18. Fraser C, Riley S, Anderson RM, Ferguson NM. Factors that make an infectious disease outbreak controllable. *Proc Natl Acad Sci U S A*. 2004;101(16):6146-6151. doi:10.1073/pnas.0307506101
19. Lovell-Read FA, Funk S, Obolski U, Donnelly CA, Thompson RN. Interventions targeting nonsymptomatic cases can be important to prevent local outbreaks: SARS-CoV-2 as a case-study. *J R Soc Interface*. 2021;18(20201014). doi:10.1098/rsif.2020.1014
20. Nakajo K, Nishiura H. Transmissibility of asymptomatic COVID-19: Data from Japanese clusters. *Int J Infect Dis*. 2021;105:236-238. doi:10.1016/j.ijid.2021.02.065
21. He D, Zhao S, Lin Q, et al. The relative transmissibility of asymptomatic COVID-19 infections among close contacts. *Int J Infect Dis*. 2020;94:145-147. doi:10.1016/j.ijid.2020.04.034
22. Inaba H, Nishiura H. The basic reproduction number of an infectious disease in a stable population: The impact of population growth rate on the eradication threshold. *Math Model Nat Phenom*. 2008;3(7):194-228. doi:10.1051/mmnp:2008050
23. Lehtinen S, Ashcroft P, Bonhoeffer S. On the relationship between serial interval, infectiousness profile and generation time. *J R Soc Interface*. 2021;18(174):20200756. doi:10.1098/rsif.2020.0756
24. Gandhi M, Yokoe DS, Havlir D V. Asymptomatic transmission, the achilles' heel of current strategies to control Covid-19. *N Engl J Med*. 2020;382(22):2158-2160. doi:10.1056/NEJMe2009758
25. He X, Lau EH, Wu P, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med*. 2020;26:672-675. doi:10.1101/2020.03.15.20036707
26. Bushman M, Worby C, Chang HH, Kraemer M, Hanage WP. Transmission of SARS-CoV-2 before and after symptom onset: Impact of nonpharmaceutical interventions in China. *Eur J Epidemiol*. 2021;36(4):429-439. doi:10.1101/2020.12.16.20214106
27. Kenah E, Lipsitch M, Robins JM. Generation interval contraction and epidemic data analysis. *Math Biosci*. 2008;213(1):71-79. doi:10.1016/j.mbs.2008.02.007
28. Lessler J, Reich NG, Brookmeyer R, Perl TM, Nelson KE, Cummings DAT. Incubation periods of acute respiratory viral infections: a systematic review. *Lancet Infect Dis*. 2015;9(5):291-300. doi:10.1016/S1473-3099(09)70069-6.Incubation
29. McAloon C, Collins Á, Hunt K, et al. Incubation period of COVID-19: A rapid systematic review and meta-analysis of observational research. *BMJ Open*. 2020;10(8):1-9. doi:10.1136/bmjopen-2020-039652
30. Riou J, Althaus CL. Pattern of early human-to-human transmission of Wuhan 2019 novel coronavirus (2019-nCoV), December 2019 to January 2020. *Eurosurveillance*. 2020;25(4):1-5. doi:10.2807/1560-

7917.ES.2020.25.4.2000058

31. Gostic KM, McGough L, Baskerville EB, et al. Practical considerations for measuring the effective reproductive number, Rt. *PLoS Comput Biol*. 2020;16(12):1-21. doi:10.1371/journal.pcbi.1008409
32. Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to estimate time-varying reproduction numbers during epidemics. *Am J Epidemiol*. 2013;178(9):1505-1512. doi:10.1093/aje/kwt133
33. Kremer C, Ganyani T, Chen D, et al. Authors' response: Estimating the generation interval for COVID-19 based on symptom onset data. *Eurosurveillance*. 2020;25(29):18-19. doi:10.2807/1560-7917.ES.2020.25.29.2001269
34. Park SW, Cornforth DM, Dushoff J, Weitz JS. The time scale of asymptomatic transmission affects estimates of epidemic potential in the COVID-19 outbreak. *Epidemics*. 2020;31(May):100392. doi:10.1016/j.epidem.2020.100392
35. Hart WS, Abbott S, Endo A, et al. Inference of SARS-CoV-2 generation times using UK household data. 2021;(March 2020):1-32.
36. Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020. *Eurosurveillance*. 2020;25(10):1-5. doi:10.2807/1560-7917.ES.2020.25.10.2000180
37. Nishiura H, Kobayashi T, Miyama T, et al. Estimation of the asymptomatic ratio of novel coronavirus infections (COVID-19). *Int J Infect Dis*. 2020;94(January):154-155. doi:10.1016/j.ijid.2020.03.020
38. Oshitani H, The Expert Members of the National COVID-19 Cluster Taskforce at the Ministry of Health Labour and Welfare of Japan. Cluster-based approach to coronavirus disease 2019 (COVID-19) response in Japan—February–April 2020. *Jpn J Infect Dis*. Published online 2020. doi:10.1038/pr.2014.160
39. Long DR, Gombar S, Hogan CA, et al. Occurrence and timing of subsequent severe acute respiratory syndrome coronavirus 2 reverse-transcription polymerase chain reaction positivity among initially negative patients. *Clin Infect Dis*. 2021;72(2):323-326. doi:10.1093/cid/ciaa722
40. Yoshioka T, Maeda Y. Covid-19 stigma induced by local government and media reporting in Japan: It's time to reconsider risk communication lessons from the fukushima daiichi nuclear disaster. *J Epidemiol*. 2020;30(8):372-373. doi:10.2188/jea.JE20200247
41. Ali ST, Wang L, Lau EHY, et al. Serial interval of SARS-CoV-2 was shortened over time by nonpharmaceutical interventions. *Science (80-)*. 2020;9004(July):eabc9004. doi:10.1126/science.abc9004
42. Anzai A, Nishiura H. "Go To Travel" campaign and travel-associated coronavirus disease 2019 cases: A descriptive analysis, July–August 2020. *J Clin Med*. Published online 2021.
43. Leclerc QJ, Fuller NM, Knight LE, Funk S, Knight GM. What settings have been linked to SARS-CoV-2 transmission clusters? *Wellcome Open Res*. 2020;5:83. doi:10.12688/wellcomeopenres.15889.2

44. Cabinet Secretariat of Japan. COVID-19 emergency declarations. Published 2021. Accessed May 26, 2021. <https://corona.go.jp/emergency/>
45. Imai N, Gaythorpe KAM, Abbott S, et al. Adoption and impact of non-pharmaceutical interventions for COVID-19. *Wellcome Open Res.* 2020;59(5):1-12.
46. Brauner JM, Mindermann S, Sharma M, et al. Inferring the effectiveness of government interventions against COVID-19. *Science (80-)*. 2021;802. doi:10.1126/science.abd9338
47. Liu Y, Morgenstern C, Kelly J, Lowe R, Jit M. The impact of non-pharmaceutical interventions on SARS-CoV-2 transmission across 130 countries and territories. *BMC Med.* Published online 2020:1-12. doi:10.1101/2020.08.11.20172643
48. Davies NG, Abbott S, Barnard RC, et al. Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England. *Science (80-)*. 2021;372(6538):eabg3055. doi:10.1126/science.abg3055
49. Ministry of Health Labour and Welfare (MHLW). Response to COVID-19 variants of concern (VOC). Published online 2021. <https://www.mhlw.go.jp/content/10900000/000779013.pdf>
50. Ito K, Piantham C, Nishiura H. Predicted domination of variant Delta of SARS-CoV-2 before Tokyo Olympic Games, Japan, July 2021. *Eurosurveillance.* Published online 2021:4-12.
51. Kissler SM, Fauver JR, Mack C, et al. Densely sampled viral trajectories suggest longer duration of acute infection with B.1.1.7 variant relative to non-B.1.1.7 SARS-CoV-2. *medRxiv.* Published online 2021.
52. Park SW, Bolker BM, Funk S, et al. Roles of generation-interval distributions in shaping relative epidemic strength, speed, and control of new SARS-CoV-2 variants. *medRxiv.* Published online 2021:2021.05.03.21256545. <https://doi.org/10.1101/2021.05.03.21256545>
53. Pung R, Mak TM, Kucharski AJ, Lee VJ. Serial intervals in SARS-CoV-2 B.1.617.2 variant cases. *Lancet.* 2021;6736(C):10-11. doi:10.1016/s0140-6736(21)01697-4
54. Kang M, Xin H, Yuan J, et al. Transmission dynamics and epidemiological characteristics of Delta variant infections in China. *medRxiv.* Published online 2021. doi:10.1101/2021.08.12.21261991
55. Britton T, Tomba GS. Estimation in emerging epidemics: Biases and remedies. *J R Soc Interface.* 2019;16(150). doi:10.1098/rsif.2018.0670
56. R Development Core Team. R: A Language and Environment for Statistical Computing. Published online 2019. <https://www.r-project.org/>
57. Stan Development Team. CmdStan: the command-line interface to Stan. Published online 2021. <http://mc-stan.org>
58. Sklar A. Fonctions de Répartition à n Dimensions et Leurs Marges. *Publ L'Institut Stat L'Université{é} Paris.* Published online 1959.
59. Klinkenberg D, Nishiura H. The correlation between infectivity and incubation period of measles, estimated from households with two cases. *J Theor Biol.* 2011;284(1):52–60. doi:10.1016/j.jtbi.2011.06.015

60. Reich NG, Lessler J, Cummings DAT, Brookmeyer R. Estimating incubation period distributions with coarse data. *Stat Med*. 2009;28:2769-2784. doi:10.1002/sim.3659

Figures

Figure 1 | Joint distribution of the generation interval and incubation period. Contour plots of the fitted distributions. For both a, the dataset of 49 transmission pairs with single dates of reported exposure, and b, the dataset of 257 transmission pairs that also includes pairs with more coarsely reported possible dates of exposure and contact, a Clayton copula with a Weibull marginal for the generation interval and lognormal marginal for the incubation period distribution was selected.

Figure 2 | Joint estimates of the generation interval and incubation period by stratum for COVID-19 transmission pairs from Japan. The joint distribution using best-fit Gaussian, Gumbel, or Clayton copula combined with gamma, lognormal, or Weibull distributions for the a, generation interval and b, incubation period are presented for the dataset of 257 transmission pairs that also includes pairs with more coarsely reported possible dates of exposure and contact. The points are point estimates for the means of each stratum, while the colored bars indicate

the 95% credible intervals. The grey bars show the overall point estimate and 95% CrI for all cases in the background. The estimate for asymptomatic infectors was fitted to the generation interval alone, as infector incubation period could not be estimated.

Figure 3 | Increased correlation leads to a predominance of presymptomatic transmission. From estimates made by simulating the generation interval (GI) and incubation period (IP) for 10000 pairs using the fitted Clayton copula with Weibull (GI) and lognormal (IP) marginals: **a**, the proportion of transmission that was symptomatic or presymptomatic for various values of Kendall's tau; **b**, the proportion of transmission that was symptomatic, presymptomatic, or occurred on the same day (GI-IP $\in [-1, 1]$) for various values of Kendall's tau; **c**, the time from onset of symptoms to transmission (TOST), defined as GI-IP, fitted with a normal distribution.

Figure 4 | Effect of stronger correlation between the generation interval and incubation period on effectiveness of isolation. The top figure, **a**, shows the fitted generation interval probability distribution function (4.22 days). The arrow dividing the green and blue sections indicates onset at 4.67 days (the mean incubation period). If case isolation occurs at onset this is equivalent to a 100% reduction in possible transmission for symptomatic cases. Figures **b**, **c**, and **d**, show the effective basic reproduction number (R_0^E) as a function of this reduction in transmission as well as the level of correlation between the generation interval and incubation period. The dashed line is the point estimate of Kendall's tau obtained in this study, while the shaded white rectangle shows its 95% credible interval. We assume baseline R_0 of **b**, 1.5, **c**, 2.2, and **d**, 2.9. As the generation interval and incubation period approach independence (Kendall's tau $\rightarrow 1$) case isolation will become ineffective—shown by the unchanging effective R_0 (R_0^E)—as transmission will either be presymptomatic or occur nearly at the same time as symptom onset.

Tables

Table 1. Characteristics of COVID-19 transmission pairs in Japan, 2020

Strata	Role	Characteristics	Coarse dates of exposure [†] (%)	Single dates of exposure [‡] (%)	Asymptomatic infectors (%)
		All	257 pairs	49 pairs	29 pairs
Age	Infector	Under 30 years	65 (25.3%)	11 (22.4%)	6 (20.7%)
		30-59 years	129 (50.2%)	32 (65.3%)	15 (51.7%)
		60+ years	57 (22.2%)	6 (12.2%)	7 (24.1%)
		Not reported	6 (2.3%)	-	1 (3.4%)
	Infectee	Under 30 years	69 (26.8%)	17 (34.7%)	9 (31.0%)
		30-59 years	114 (44.4%)	19 (38.8%)	10 (34.5%)
		60+ years	67 (26.1%)	13 (26.5%)	9 (31.0%)
		Not reported	7 (2.7%)	-	1 (3.4%)
Sex	Infector	Female	79 (30.7%)	17 (34.7%)	12 (41.4%)
		Male	177 (68.9%)	32 (65.3%)	16 (55.2%)
		Not reported	1 (0.4%)	-	1 (3.4%)
	Infectee	Female	131 (51.0%)	22 (44.9%)	16 (55.2%)
		Male	125 (48.6%)	27 (55.1%)	12 (41.4%)
		Not reported	1 (0.4%)	-	1 (3.4%)
Transmission setting	Infector	Household	4 (1.6%)	0 (0.0%)	0 (0.0%)
		Social	153 (59.5%)	48 (98.0%)	25 (86.2%)
		Community	100 (38.9%)	1 (2.0%)	4 (13.8%)
	Infectee	Household	109 (42.4%)	2 (4.1%)	11 (37.9%)
		Social interaction	113 (44.0%)	38 (77.6%)	15 (51.7%)
		Community	35 (13.6%)	9 (18.4%)	3 (10.3%)
Epidemic wave		Wave 1	86 (33.5%)	14 (28.6%)	2 (6.9%)
		Wave 2	95 (37.0%)	16 (32.7%)	13 (44.8%)
		Wave 3	76 (29.6%)	19 (38.8%)	14 (48.3%)
Basis for selection		Cluster	138 (53.7%)	29 (59.2%)	24 (82.8%)
		Contact pattern	54 (21.0%)	19 (38.8%)	4 (13.8%)
		Domestic travel	43 (16.7%)	1 (2.0%)	1 (3.4%)
		Import	22 (8.6%)	-	-
Region§		Hokkaido & Tohoku	35 (13.6%)	6 (12.2%)	2 (6.9%)
		Kanto	31 (12.1%)	4 (8.2%)	0 (0.0%)
		Chubu	73 (28.4%)	12 (24.5%)	11 (37.9%)
		Kinki	38 (14.8%)	6 (12.2%)	3 (10.3%)
		Chugoku & Shikoku	34 (13.2%)	6 (12.2%)	6 (20.7%)
		Kyushu & Okinawa	46 (17.9%)	15 (30.6%)	7 (24.1%)

[†]Dataset using intervals of exposure and/or contact between infector and infectee. [‡]Dataset limited to pairs where infector exposure and infector-infectee contact were limited to a single day. “Hokkaido & Tohoku” includes Hokkaido, Aomori, Iwate, Miyagi, Akita, Yamagata, and Fukushima prefectures. “Kanto” includes Ibaraki, Tochigi, Gunma, Saitama, Chiba, Tokyo, and Kanagawa prefectures. “Chubu” includes Niigata, Toyama, Ishikawa, Fukui, Yamanashi, Nagano, Gifu, Shizuoka, and Aichi prefectures. “Kinki” includes Mie, Shiga, Kyoto, Osaka, Hyogo, Nara, and Wakayama prefectures. “Chugoku & Shikoku” include Tottori, Shimane, Hiroshima, Yamaguchi, Tokushima, Kagawa, Ehime, and Kochi prefectures. “Kyushu & Okinawa” include Fukuoka, Saga, Nagasaki, Kumamoto, Oita, Miyazaki, Kagoshima, and Okinawa prefectures.

1 **Table 2.** Joint estimates of the generation interval and incubation period of COVID-19 cases by stratum

Category	Strata	N	Copula	Generation interval			Incubation period			Kendall's tau (95% CrI)
				Distribution	Mean (95% CrI)	SD (95% CrI)	Distribution	Mean (95% CrI)	SD (95% CrI)	
Exact data	All cases	49	Clayton	Weibull	4.38 (3.88, 4.98)	2.10 (1.70, 2.79)	Lognormal	4.91 (4.34, 5.60)	2.66 (2.18, 3.33)	0.61 (0.46, 0.68)
Coarse data	All cases	257	Clayton	Weibull	4.34 (3.98, 4.74)	2.31 (2.00, 2.81)	Lognormal	4.75 (4.42, 5.11)	2.57 (2.25, 2.98)	0.54 (0.44, 0.63)
Infector age	Under 30 years	65	Clayton	Lognormal	4.51 (3.95, 5.13)	2.15 (1.68, 2.89)	Lognormal	4.67 (4.06, 5.36)	2.60 (2.12, 3.27)	0.50 (0.23, 0.65)
	30-59 years	129	Gumbel	Gamma	4.43 (3.97, 4.94)	2.43 (2.01, 3.00)	Lognormal	5.10 (4.63, 5.62)	2.87 (2.45, 3.44)	0.60 (0.48, 0.68)
	60+ years	57	Gumbel	Gamma	4.58 (3.87, 5.37)	2.68 (1.98, 3.68)	Lognormal	4.59 (3.91, 5.37)	2.89 (2.31, 3.74)	0.50 (0.28, 0.65)
Infected age	Under 30 years	69	Clayton	Gamma	4.57 (4.02, 5.20)	2.39 (1.90, 3.18)	Lognormal	4.72 (4.15, 5.38)	2.62 (2.15, 3.26)	0.56 (0.37, 0.68)
	30-59 years	114	Gumbel	Weibull	4.66 (4.13, 5.22)	2.59 (2.07, 3.25)	Lognormal	5.26 (4.72, 5.85)	3.12 (2.62, 3.78)	0.58 (0.47, 0.68)
	60+ years	67	Gumbel	Lognormal	4.10 (3.48, 4.83)	2.08 (1.49, 3.01)	Lognormal	4.45 (3.84, 5.15)	2.59 (2.10, 3.32)	0.42 (0.14, 0.61)
Infector sex	Female	79	Gumbel	Gamma	3.72 (3.20, 4.35)	2.09 (1.57, 2.92)	Lognormal	4.38 (3.82, 5.02)	2.59 (2.10, 3.26)	0.54 (0.36, 0.66)
	Male	177	Clayton	Weibull	4.71 (4.29, 5.20)	2.36 (2.00, 2.90)	Lognormal	5.00 (4.60, 5.46)	2.73 (2.35, 3.22)	0.52 (0.39, 0.63)
Infected sex	Female	131	Clayton	Gamma	4.35 (3.82, 4.95)	2.50 (1.99, 3.28)	Lognormal	4.65 (4.20, 5.16)	2.60 (2.20, 3.15)	0.50 (0.33, 0.64)
	Male	125	Gumbel	Gamma	4.50 (4.05, 4.99)	2.28 (1.88, 2.87)	Lognormal	4.96 (4.48, 5.52)	2.88 (2.43, 3.48)	0.55 (0.43, 0.65)
Epidemic wave	Wave 1	109	Gumbel	Weibull	4.38 (3.74, 5.01)	1.99 (1.59, 2.58)	Lognormal	4.75 (4.21, 5.36)	2.93 (2.46, 3.63)	0.53 (0.31, 0.67)
	Wave 2	113	Clayton	Gamma	4.50 (4.04, 5.03)	2.50 (2.09, 3.13)	Lognormal	4.85 (4.39, 5.38)	2.55 (2.16, 3.07)	0.58 (0.47, 0.67)
	Wave 3	35	Gaussian	Gamma	4.52 (3.81, 5.32)	2.39 (1.74, 3.43)	Lognormal	4.98 (4.19, 5.87)	3.01 (2.38, 3.93)	0.57 (0.36, 0.68)
Transmission setting	Household	86	Gumbel	Weibull	5.09 (4.46, 5.77)	2.82 (2.25, 3.59)	Lognormal	5.13 (4.53, 5.79)	2.87 (2.38, 3.55)	0.55 (0.38, 0.67)
	Social contact	95	Gumbel	Weibull	3.84 (3.42, 4.33)	1.74 (1.38, 2.32)	Lognormal	4.53 (4.02, 5.12)	2.68 (2.22, 3.34)	0.55 (0.40, 0.66)
	Community	76	Clayton	Gamma	4.45 (3.87, 5.11)	2.36 (1.85, 3.20)	Lognormal	5.00 (4.40, 5.69)	2.93 (2.42, 3.68)	0.48 (0.29, 0.63)
Basis for selection	Cluster	138	Gumbel	Gamma	4.08 (3.62, 4.60)	2.17 (1.73, 2.81)	Lognormal	4.55 (4.14, 5.02)	2.50 (2.13, 2.99)	0.46 (0.29, 0.60)
	Contact pattern	54	Clayton	Weibull	4.62 (4.03, 5.30)	2.31 (1.82, 3.11)	Lognormal	5.12 (4.45, 5.90)	3.07 (2.48, 3.92)	0.59 (0.42, 0.68)
	Domestic travel	43	Clayton	Gamma	4.84 (4.12, 5.64)	2.64 (2.01, 3.62)	Lognormal	5.01 (4.22, 5.92)	3.02 (2.40, 3.93)	0.47 (0.23, 0.64)
	Import	22	Gaussian	Weibull	4.83 (3.91, 5.78)	2.33 (1.62, 3.50)	Lognormal	5.65 (4.60, 6.77)	3.49 (2.73, 4.55)	0.64 (0.37, 0.69)
Asymptomatic infectors		29	-	Gamma	4.62 (3.78, 5.61)	2.45 (1.78, 3.59)	-	-	-	-

2 CrI: credible interval. Wave 1 began 16 January 2020. Wave 2 is assumed to have begun 1 June 2020, and wave 3 is assumed to have begun 1 October 2020.

3

4

5 Extended Data figures

6

7

8 **Extended Data Fig. 1 | Correlation of empirical generation and serial intervals with the incubation period of**
9 **COVID-19 cases in Japan.** Scatterplots showing the generation and serial intervals plotted against the incubation
10 period for transmission pairs with single-date ($n=49$) and coarsely observed ($n=257$) exposures. For **a**, data where single
11 dates of exposure for the infector and contact between infector and infectee were reported, Kendall's tau was 0.58
12 ($p<0.001$), -0.13 ($p=0.32$), and -0.23 ($p=0.07$) for the generation interval, serial interval with only positives, and serial
13 interval including negatives, respectively. For **b**, data with coarse dates of exposure and contact, the plotted value
14 represents the midpoint of the possible exposure/contact period, and Kendall's tau was 0.53 ($p<0.001$), -0.02 ($p=0.66$),
15 and -0.04 ($p=0.39$).

16

Extended Data Fig. 2 | Age and sex distribution of transmission pairs. a, Matrix of infector and infectee age groups for 257 transmission pairs. **b**, Proportion of cases by sex, wave, and infector-infectee status. **c**, Age group and sex distribution of infectors and infectees.

Extended Data Fig. 3 | Prior and posterior densities of the mean. **a**, densities for the dataset where single dates of exposure for the infector and contact between infector and infectee were reported (n=49). **b** densities for the dataset with coarsely reported dates of exposure and contact (n=257). The priors are the same for **a** and **b**; for the generation interval the prior was obtained from the values of the serial interval reported by Nishiura et al., while the prior for the incubation period was obtained from the values of the incubation period reported by Linton et al.—both studies used cases reported worldwide during the early stages of the pandemic (early 2020).

Extended Data Fig. 4 | Generation intervals reported across COVID-19 studies. The studies by Ganyani et al., Bushman et al., Li et al., and Zhao et al. are plotted using gamma distributions. The result from Hart et al. was plotted using a lognormal distribution. Results from this study and the study by Ferretti et al. plotted using Weibull distributions.

Extended Data Fig. 5 | Epidemic curve of COVID-19 cases in Japan by date of laboratory confirmation.

Here, the second epidemic wave is shown as beginning 1 June (first dashed line), while the third epidemic wave starts 1 October (second dashed line). Hokkaido declared a local state of emergency 28 February–19 March (first two vertical dotted lines). A national state of emergency was declared for key urban prefectures (Tokyo, Saitama, Chiba, Kanagawa, Osaka, Hyogo, and Fukuoka) on 7 April (third dotted line), with the state of emergency extending nationwide on 16 April (fourth dotted line). The end of the state of emergency varied between prefectures, with most ending 14 May while some continued until 25 May (fifth dotted line). The GoTo travel campaign, offering large discounts on travel inside Japan with the intention of restarting the Japanese economy following the damage caused by COVID-19 related public health and social measures, began on 22 July (sixth dotted line) for all prefectures except Tokyo, which was added to the campaign on 1 October (start date of third wave, second dashed line). The campaign continued through the end of 2020 and into 2021.