Systematic review

Colorectal cancer and microbiota modulation for clinical use. A systematic review.

Julio Madrigal-Matute and Sara Bañón Escandell*

Biomedical Institute for Nutrition and Health (IBIONS).

**Corresponding author:*

Julio Madrigal-Matute, PhD

Biomedical Institute for Nutrition and Health (IBIONS).

Phone: +34606008917;

Madrigal-matute@ibions.com

Abstract

Colorectal cancer is one of the top contributors to the global burden of cancer incidence and mortality with genetic and environmental factors contributing to its etiology. Modifiable or environmental factors can be the cause of up to 60% of the risk of developing colorectal cancer. Hence, there is a growing interest in specifically defining what can be improved in our lifestyle to reduce this risk, improve the effectiveness of treatments, reduce side effects, and decrease the risk of recurrence. One of the elements directly related to lifestyle is gut microbiota. The microbial ecosystem has a vital role in colorectal cancer prevention and antitumoral response through modulation of the immune system and production of short-chain fatty acids. Numerous approaches have been used to identify healthy microbiota that can reduce the risk of cancer development, improve treatment efficacy, and reduce side effects.

Scientific literature in this subject is growing exponentially and, therefore, systematic reviews and meta-analysis are required to ensure that appropriate recommendations are given to patients.

This work aimed to perform a systematic analysis of the published literature to elucidate whether microbiota modulation through pre-, pro-, symbiotic treatment and/or nutritional intervention can be beneficial for patients diagnosed with colorectal cancer.

Detailed analysis of published studies shows that some prebiotics, such as inulin and resistant starch, probiotics such as lactic strains producers of short-chain fatty acids, and consumption of unprocessed plant products, can be effective recommendations for patients diagnosed with colorectal cancer. This advice should always be individually tailored and followed up by a healthcare professional with expertise in the field.

Keywords: cancer, colorectal, nutrition, diet, probiotic, prebiotic, symbiotic, immunotherapy, immune system.

Abbreviations:

CRC, colorectal cancer; ROS, reactive oxygen species; RNS, reactive nitrogen species; MSI, Microsatellite instability; SCFA, Short chain fatty acids; LPS, Lipopolysaccharide; dMMR, deficient DNA mismatch repair; AOM, Azoxymethane; DSS, Dextran sodium sulfate; TNF- α , tumor necrosis factor alpha; MPO, Myeloperoxidase; PCNA, Proliferation cellular nuclear antigen; SLC5A8, Solute Carrier Family 5 Member 8; GPR43, G protein-coupled receptor 43; CEA, Carcinoembryonic antigen; APC, Adenomatous polyposis coli; DHM, 1,2-dimethyl hydrazine; TGF- β , Transforming growth factor beta; TLR2, toll-like receptor 2; DC, dendritic cells; PD-1, Programmed cell death protein 1; PDL-1, Programmed cell death protein ligand 1; CTLA-4, Cytotoxic T-Lymphocyte Associated Protein 4; CYP, Cytochrome P450 Family; GSTM1, Glutathione S-Transferase Mu 1; SULT1, Sulfotransferase Family-1; PYY, Peptide tyrosine; PPAR γ , peroxisome proliferator activated receptor gamma; 6,BR, 6-bromoisatin; NE, marine extract; MD, Mediterranean diet; LFD, low fat diet; HFD, high fat diet; EPA, Eicosapentanoic acid; PUFA, polyunsaturated fatty acids; CRP, C reactive protein; SFRP2, Secreted Frizzled Related Protein 2; DNMT1, DNA Methyltransferase 1; COX-2, Cyclooxygenase-2; PGE₂, Prostaglandin E2; Cxcl, C-X-C Motif Chemokine Ligand; Bik, BCL2 Interacting Killer; KC, Keratinocytes-derived chemokine; Gr-1, Glucocorticoid Receptor-1; CTNNB1, Beta catenin 1; EGFR, Epidermal growth factor receptor; TYMS, Thymidylate Synthetase; LPE, Lisophosphatidiletanolamine; LPC, Lisophosphatidylcholine; G-CSF, Granulocyte Colony-Stimulating Factor; GM-CSF, Granulocyte Macrophage Colony-Stimulating Factor; HDAC, Histone Deacetylase; JAM1, F11 Receptor; 5-FU, 5-florouracil; EGCG, epigallocatechin-3-gallate;

1. Introduction

1.1. Colorectal cancer

Colorectal cancer (CRC) is a disease based on an important genetic component, coupled with key influence from our lifestyle habits. Approximately 10-20% of diagnoses of CRC have a family history with the disease,¹ and through family and twin studies, heritability is estimated to range from 12-35%.^{2, 3} Although numerous genome-wide association studies have been performed identifying genes that increase susceptibility to develop CRC,⁴ many risk factors remain unknown. However, only 5-7% of patients' colon cancer has a clear genetic origin.⁵ Hereditary CRC syndromes can be subdivided into those associated with polyposis and those without polyposis (Lynch syndrome and familial CRC).⁶ Risk factors also include the development of inflammatory bowel disease, including Chron's disease and ulcerative colitis. These pathologies significantly increase the chances of developing CRC, being highly dependent on environmental and nutritional factors.⁷ Specifically, geographical differences in incidence of CRC are influenced by genetic, and mainly environmental and nutritional risks.⁸

It is not surprising that environmental factors have such a strong influence on the risk of developing colon cancer, as the intestinal epithelium is one of the tissues with the highest turnover rate in our body and is in continuous direct contact with nutrients, food toxins, and microbiota.⁹⁻¹⁴

Etiology

Among the metabolic changes that precede the development of cancer are chronic cell damage due to a pathological increase in reactive oxygen and nitrogen species (ROS and RNS), DNA damage, and chronic inflammation.⁶ Inflammation triggers the release of pro-inflammatory cytokines into the gut to regenerate damaged tissue. In certain situations, inflammation becomes chronic, forming a positive feedback loop that facilitates tumor development.¹⁵ Colon tumors generally begin with the transformation of stem and epithelial cells that reside at the base of the colonic crypts. These cells undergo chronic damage by inflammation, oxidative stress, and/or genetic mutations that evolve into a premalignant lesion or polyp.¹⁶ These polyps follow two main routes of differentiation to malignancy: the *adenoma-carcinoma* type (70-90%) and the *serrated* (10-20%) type. In addition, another route not associated

with polyposis has been described with a lower incidence, characterized by *microsatellite instability* (MSI; 2-7%).¹⁷ The development of a malignant tumor is estimated to take 10-15 years on average.⁶

1.2. Gut microbiota

The microbiome includes all types of microorganisms associated with the host, interacting in a complex way and usually found in the epithelial barrier zones.¹⁸ It is acquired at birth by vertical transmission and is modulated throughout our lives by environmental factors.¹⁹ The microbiota is a dynamic niche that is involved in the maintenance of health and in the origin and development of numerous diseases by modulating the homeostasis of the immune system.²⁰ Although the mechanisms may be varied, one of the most notably is the stimulation of the T-cell response, through the expression of bacterial epitopes or their cross-reactivity with tumor epitopes. Additionally, gut microbiota also modulates the response to tumors by inhibiting specific receptors that promote inflammatory and immune responses. Finally, another important mechanism carried out by the microbiota is the synthesis of metabolites, such as short-chain fatty acids (SCFA), which elicit a local and systemic response in the host organism.¹⁹

But what is the ideal microbiota composition for health? This is a question that is not easy to answer, although numerous studies have attempted.^{21, 22} Analyzing a metabolite panel comparing stool from healthy versus CRC patients, it was found that the best marker to discern between a healthy patient and a CRC patient is primarily a decrease in the CRC patient's levels of certain SCFA, in particular, acetate and butyrate.²³ The protective effect of SCFA against CRC may be due to their ability to decrease inflammation and modulate the immune response.²⁴ In addition, SCFA inhibit proliferation and promote apoptosis in neoplastic colonocytes.²⁵

SCFA play a key role in maintaining intestinal health and preventing the development of numerous pathologies.²⁶ They are produced by many bacteria in the intestinal tract, in particular, lactic acid bacteria and bifidobacteria. SCFA transporters and their receptors are the communication bridges between the microbiota and the cells of the gut.²⁷ Butyrate has been shown to decrease intestinal inflammation by regulating epithelial occlusive junctions,²⁸ which are essential for maintaining the integrity of the intestinal barrier and mucosa by aiding in the homeostasis of ions, nutrients and water.²⁹

Among the SCFA-producing bacteria, and specifically butyrate, are those of the genus *Butyricoccus*, which could protect against colitis in patients with irritable bowel syndrome, as these bacteria are reduced in affected patients.³⁰ Likewise, other bacteria belonging to the phylum *Firmicutes* such as *Eubacterium rectale*, *Ruminococcus albus*, *R. callidus*, *R. bromii* and *F. prausnitzii* are 5 to 10 times more abundant in healthy patients than in patients with Crohn's disease.³¹ The decrease in species such as *Bacteroides fragilis* and *B. vulgatus* is associated with reduced levels of butyric and propionic acid and increased inflammatory markers.³² Additionally, *Akkermansia muciniphila*, one of the most abundant components of the intestinal microbiota, is found to be decreased in obese patients,^{33, 34} those with ulcerative colitis,³⁵ and those with CRC.³⁶ *A. muciniphila* ferments mainly mucins, as well as other mono- and polysaccharides, giving rise to various SCFA.³⁷

Taken together, it appears that SCFA-producing bacteria, such as acetate and butyrate are often associated with healthier microbiota profiles. For these reasons, SCFA are being tested as a therapy for CRC.³⁸

Dysbiosis

Pathological changes in the composition of the microbiota is known as dysbiosis.³⁹ This unbalance in the homeostatic microbiome is related to the origin of various types of tumors due, in part, to the modulation of tumor immunity,⁴⁰ with CRC being prominent among these tumor types.⁴¹ Patients with CRC are often characterized by a decrease in beneficial microbiota and an increase in pathogenic opportunistic microorganisms.²² This pathogenic imbalance can have a variety of origins. A frequent and often underestimated cause of dysbiosis stems from intensive antibiotic use.⁴²

Chronic intestinal inflammation and epithelial disruption, two of the most important risk factors for CRC development, are related to intestinal dysbiosis.^{21, 43} Many of the pathogenic species identified in different studies associated with the development of CRC belong to the group of hydrogen sulfide (H₂S)-producing bacteria. These bacteria can oxidize SCFA, decreasing its levels locally.⁴⁴ H₂S is highly toxic to colonocytes, causes damage to the intestinal epithelium, promotes inflammation, produces metabolic changes, especially through oxidation of butyrate, and can result in the development of CRC.^{45, 46} H₂S is also rapidly absorbed by the mucosa,⁴⁷ which can disrupt the balance between proliferation and apoptosis^{48, 49} and lead to DNA

damage.⁵⁰ Furthermore, feces of people at high risk of CRC recurrence have higher concentrations of H₂S.⁵¹ Other studies corroborate these results by finding an increase in *Fusobacterium spp.*, and more specifically *Fusobacterium nucleatum*, an H₂S-producing species, in colonic tumor tissue.^{52, 53} *F. nucleatum* also promotes tumor growth, cell proliferation and the acquisition of chemoresistance in CRC, favoring inflammation and autophagy as a cytoprotective mechanism of the tumor.^{54, 55}

At the phylum level, CRC patients show increased abundance of *Proteobacteria* and *Verrucomicrobia*.⁵⁶ Species associated with CRC development include *Bacteroides fragilis*, *Escherichia coli*, *Peptostreptococcus spp*, *Enterococcus faecalis*, *Streptococcus bovis*, or *Porphyromonas spp*.^{22, 53, 57-59} For example, *B. fragilis* and *E. coli* play a key role in the development of familial CRC. In patients with adenomatous polyposis, areas arise in the colonic mucosa where *B. fragilis* and *E. coli* accumulate, leading to a local increase in oncotoxins such as *B. fragilis* toxin and colibactin.⁶⁰ These oncotoxins, when transferred to the colon of mice, increase colonic tumor growth and mortality.⁶⁰ In addition, *B. fragilis* levels are elevated in patients with diarrhea, with chronic diarrhea being an important risk factor for the development of CRC.⁶¹

Nonetheless, it may not be so simple to discern which bacteria are beneficial. In general, it appears that it may be context-dependent and that one of the key characteristics of a healthy microbiota is its beneficial/harmful microbiota ratio and/or its diversity.

1.3. Food and gut microbiota

Due to the importance of the microbiota in health and in the origin of diseases such as CRC, there is much interest in knowing how to modify the flora in order to have a better homeostatic capacity. The main way in which we can interact directly with our microbiota is through the food we eat.⁶² A study of monozygotic and dizygotic twins, and their mothers, shows that while there is a common core microbiota, pathological conditions, such as malnutrition leading to obesity, produce a significant change in the microbiota. These changes result in reduced diversity and modifications in gene expression and metabolic pathways.⁶² Along these lines, it has been shown that a "Western" diet (high fat and sugar) can alter the microbiota in periods as short as 1-3 days, and that these changes can be reversed by following a healthier diet (high

plant).^{63, 64} These results show that a healthy diet, mainly a plant-based, can improve our microbiota faster than probably any other therapeutic intervention. In fact, adherence to a Mediterranean diet has been shown to have a beneficial outcome in modulating the microbiota. Participants following this type of diet had higher levels of SCFA, *Prevotella* and *Firmicutes* that degrade plant fibers, in addition to a higher proportion of bifidobacteria/*E. coli* and better gastrointestinal habits.⁶⁵ Therefore, it is likely that part of the health benefits of this dietary pattern are due to its effect on our commensal flora.

In fact, the health benefits of certain dietary patterns, such as fiber consumption, have been evaluated for decades. However, the mechanisms by which fiber decreases the incidence of CRC are not fully understood. Two mechanisms that may be complementary are generally postulated: i) fiber may facilitate gastrointestinal transit decreasing the contact between procancerous substances and ii) the intestinal epithelium, or gut bacteria may ferment fiber, producing SCFA.⁶⁶ In an attempt to dissect these mechanisms, a study was conducted using a mouse model in which a representative human microbiota (gnobiotic) was "seeded". It was found that a low-fiber diet leads to an overgrowth of bacterial species that cause degradation of the mucus lining the intestinal barrier. This makes the barrier more permeable to pathogenic organisms, facilitating the development of colitis.⁶⁷ In this mouse model, with a low-fiber diet, overgrowth of *B. caccae* and *A. muciniphila* was promoted, with these two species being responsible for the degradation of intestinal mucus.

The importance of nutrition is often underestimated in most scientific and medical fields; however, available data show that it is a key tool in modifying the microbiota.

1.4. Probiotics

Probiotics are defined as "live microorganisms that confer a health benefit when administered in adequate amounts".⁶⁸ The benefits are varied and range from the maintenance of intestinal homeostasis to the prevention of diarrhea, irritable bowel syndrome, and constipation.⁶⁹ In addition, numerous *in vitro*,⁷⁰ preclinical⁷¹⁻⁷³ and clinical studies⁷⁴ have shown that probiotics may have beneficial effects, specifically in cases of CRC.

Probiotics exert their beneficial action by modifying the gut microbiota, improving the gut barrier and its microenvironment, secreting anti-cancer metabolites, and reducing inflammation.⁷⁵ In addition, probiotics have an intimate relationship with the

immune system, particularly through homeostasis of epithelial cell junctions and modulation of T-lymphocytes and dendritic cells activation.^{36, 76} As an example, *Butyricicoccus pullicaecorum* reduces necrotic enteritis and pathogen abundance in the caecum and ileum in broiler chickens.⁷⁷ Furthermore, its administration promotes the proper functioning of the intestinal epithelial barrier and reduces colitis in preclinical models.³⁰ One of the most widely used probiotics is *A. muciniphila*, which has been shown in numerous studies to have a beneficial role in health maintenance and a therapeutic effect on various diseases such as obesity,⁷⁸ or epithelial tumors.^{79, 80} In this regard, it has been used in preclinical models of chronic colitis, where it improved inflammatory parameters and damage in the colon.⁸¹ Additionally, lactic acid bacteria have been frequently used for their health benefits and their immunomodulatory effect,⁸² contributing to inhibiting the development of pathogens.⁸⁹ Treatment with *L. salivary* and *L. fermentum* in combination with *L. acidophilus* can reduce CRC cell proliferation in experimental models.^{83, 84} Furthermore, oral administration of 21 different strains of *Lachnospiraceae* attenuated colorectal inflammation and improved the composition of the microbiota in preclinical models.⁸⁵

However, a meta-analysis evaluating the role of probiotics in cancer patients failed to find a clear overall beneficial outcome, except in reducing the degree of diarrhea, bowel movement, and the need for antidiarrheal drugs.⁸⁶

1.5. Prebiotics

Prebiotics are defined as "any substance that is selectively utilized by the recipient organism and confers a health benefit".⁸⁷ One of the most studied prebiotics is inulin. Inulin is part of a group of dietary fibers known as inulin-type fructans. These compounds are found in numerous foods such as garlic, artichoke, onion and asparagus.⁸⁸ Inulin-type fructans resist digestion in the small intestine and are fermented in the colon, where they are transformed into lactic acid and SCFA, improving intestinal health.^{89, 90} Inulin has been shown to increase levels of CRC-protective bacteria such as *Bifidobacteria*,⁹¹ *Bacteroides*,⁹² and *Akkermansia muciniphila*⁷⁸ in murine models.

The role of resistant starch as a prebiotic has also been examined in numerous pathologies and its benefits have also been suggested in the field of cancer, particularly in CRC.⁹³ Resistant starch is a group of carbohydrates that resist digestion by pancreatic amylase in the small intestine. This type of starch then

reaches the colon where it is fermented by the microbiota, producing numerous protective effects against tumor development, including production of SCFA (butyrate), regulation of bowel movements and the reduction of colonic pH.^{93, 94} Many foods contain resistant starch, such as cereals, vegetables, legumes, seeds, and some nuts.^{93, 95}

Recent studies are using prebiotics as a basis for the synthesis of therapeutic targets for cancer treatment because of their promising role in this field.⁹⁶

1.6. Symbiotics

Symbiotics are defined as "a mixture including live microorganisms and substrates, which used selectively by the host organism confers a health benefit".⁹⁷ A distinction is made between *cooperative* symbiotics, in which probiotic and prebiotic are administered together and act independently contributing to the health benefit, and *synergistic* symbiotics, where the probiotic uses the co-administered prebiotic as a substrate.⁹⁷ The use of symbiotics is becoming increasingly common in the biomedical literature as a method of enhancing the effects of pro- and prebiotics used individually. For example, the combination of *Bifidobacterium animalis subsp. lactis* with a resistant starch led to a decrease in the number and incidence of colon neoplasms. Interestingly, the probiotic alone did not produce a similar effect. It was the combination that resulted in protection against cancer development, in part through increased production of SCFA.⁹⁸

Therefore, it seems very promising that symbiotics may play a more effective role than pre- or probiotics alone.

1.7. Oncological treatments and microbiota in CRC

Technological advances and new therapies have dramatically increased the life expectancy of CRC patients.⁶ However, CRC patients often show particularly low responses to hospital treatments.⁹⁹ For this reason, numerous lines of research are being devoted to better understanding the mechanisms underlying this lack of response to treatments, such as immunotherapy.

The first evidence showing the key role that the microbiota plays in modulating the body's immune response to cancer treatment came from a pioneering study showing that microbial flora, and in particular lipopolysaccharide (LPS) from Gram-negative bacteria, increased the activation of dendritic cells and TCD8+ lymphocytes. In this

way, the microbiota promoted melanoma regression after radiotherapy treatment. In contrast, the use of antibiotics, inhibition of LPS and its signaling, significantly diminished the effect of radiotherapy treatment.¹⁰⁰ Numerous studies have corroborated these data, showing that therapies decrease their efficacy in axenic (without microbial flora) or antibiotic-treated mice.^{80, 101-103} A key study in this field showed that genetically identical mice from two different suppliers responded unequally to the same cancer treatment, due to differences in microbiota. Not only that, but cohousing or fecal transfer eliminated these differences, clearly pointing towards a key role of microbiota in modulating the efficacy of cancer treatment.¹⁰⁴

Although the role of the microbiota has been studied in different cancer treatments such as radiotherapy,¹⁰⁰ chemotherapy,¹⁰² and surgery,¹⁰⁵ most studies have focused in immunotherapy.^{80, 103} This may be due to both the microbiota's promising role in cancer treatment and its key role in modulating the immune response.

Immunotherapy

Unfortunately, a large proportion of CRC patients do not respond well to immunotherapy. It is known that the effectiveness of some treatments depends, to a large extent, on the individual's genetics and/or cancer phenotype⁹⁹; however, for a majority of CRC patients, the cause of their lack of response is unknown.

Immunotherapy has promising results in numerous tumor types by inducing immunity by T lymphocytes.¹⁰⁶ Thus, the anti-PD-1 drugs pembrolizumab and nivolumab are effective in metastatic patients whose tumors are deficient in DNA mismatch repair (dMMR) enzymes and have high microsatellite instability (MSI-H).⁹⁹ For the treatment of refractory dMMR-MSI-H tumors, pembrolizumab and nivolumab combined with an anti-CTLA4, ipilimumab, are used. However, in tumors with active DNA repair enzymes and low microsatellite instability, which account for 85% all tumors, these therapies tend to be ineffective.⁹⁹ Therefore, there is great interest in finding methods to improve the efficacy of these therapies in CRC.

Knowing the ideal composition of microbiota to improve the efficacy of immunotherapy treatments is no easy task. It is likely, as mentioned above, that the key lies in increasing the ratio of bacteria that are beneficial to health and decreasing those that are harmful.¹⁰⁷ In this line, mice transplanted with microbiota from CRC patients show, after treatment with anti-PD-1, a decrease in the anti-tumor response in CRC and lower survival, compared to those transplanted with microbiota from

healthy controls.¹¹⁷ Among the bacteria that may be beneficial to improve immunotherapy are: *Clostridiales*, *Ruminococcaceae*, *Enterococci*, *Collinsella*, *Alistipes*, *Faecalibacterium spp*, *A. muciniphila*, *B. fragilis*, *Eubacterium linosum* or *Bifidobacteria*, although most studies have been performed on epithelial type tumors.^{19, 79, 80}

Therefore, the study of the microbiota in the treatment of cancer patients, especially in CRC patients whose treatments do not reach the effectiveness of other types of cancer, is essential.

Effect of surgery on the microbiota

Colorectal surgery in humans aims for total tumor removal plus the addition of a safety perimeter. When this clean-border surgery is achieved, the likelihood of tumor recurrence decreases considerably.¹⁰⁸ However, local recurrences are common, up to 15%, the origin of which appear to be linked to the biology of the primary tumor and/or seeding of the tumor bed during surgery.¹⁰⁹⁻¹¹¹ Most tumors recur at the site of the anastomosis due to incomplete wound closure, the inherent porosity of the junctional area, improper healing, and/or the presence of collagenolytic organisms that increase porosity.^{112, 113} In this regard, a high-fat diet increases the production of colorectal tumors in preclinical models of anastomosis.¹⁰⁵ This mechanism is linked to the overproduction of collagenolytic organisms at the suture site. Pre-operative fecal analysis shows that obese patients respond by increasing collagenolytic microorganisms after taking antibiotics. In contrast, non-obese patients have decreased levels of collagenolytic microorganisms.¹⁰⁵ Therefore, a healthy microbiota low in collagenolytic microorganisms is key to increasing the chances of successful CRC surgery and avoiding recurrences.

1.8. Study aims

The overall aim of this work is to review published studies on the modification of the microbiota in CRC.

The specific objectives of this work are to evaluate the impact of modulating the microbiota in CRC patients through; 1) probiotics, 2) prebiotics, 3) symbiotics and 4) nutritional changes. We aim to analyze the potential use of these approaches to modulate the microbiota in CRC patients.

2. Methods

2.1. Systematic search

The search strategy was defined between January 15th and April 15th, 2021. The search and filtering in the Pubmed database took place on April 24th, 2021. The language used for the search and filtering of the publications was English. The detailed analysis of the selected publications was carried out from April 25th to May 31st, 2021.

The search strategy included six types of terms: (1) target disease of the study; (2) type of cancer; (3) microbiota and dysbiosis; (4) oncology treatment, (5) use of nutrition and/or supplementation with pro-, pre- and symbiotics and (6) original articles.

To broaden the search terms, asterisks were added to the end of the words. Keywords were combined using the Boolean operators "AND", "OR" and "NOT". The search strategy is detailed in Table 1.

The results of the different levels of the search can be found in the attached supplementary files.

Inclusion criteria

Only original articles were selected. Biographies, commentaries, duplicate publications, editorials, guides, interviews, news items, personal narratives, errata, retractions, reviews, systematic reviews and meta-analyses were excluded. In the next level of screening, only articles published in English that had mechanistic results and outcomes, at least *in vivo*, were selected by analysis of the abstract.

Publications that met the criteria above were downloaded and reviewed.

3. Results

3.1. Literature search

Figure 1 shows the flow diagram of the search and selection process. This diagram has been made by adapting the work to the guidelines for systematic searches recommended by Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA).¹¹⁴ The Pubmed search yielded 202 results. The articles were screened for eligibility criteria. As a result of the first screening, a total of 108 articles

were selected. After applying the next filtering on the type of results, a total of 62 articles remained for analysis. Of these 62 studies, 15 included human clinical trials (Fig. 1).

3.2. Characteristics of included results

Table 2 shows the characteristics of the included studies. The type of intervention performed, type of cancer, type of study, impact on the microbiota, the mechanism proposed by the authors, and its impact on cancer development are shown. Finally, the last column includes a more subjective criterion, namely the quality of the data analyzed on a scale ranging from very low = 1, low = 2, medium = 3, high = 4, very high = 5. This criterion has been assessed according to the experimental design, statistical analysis and strength of the mechanisms demonstrated. Figure 2 shows the distribution of the included studies.

4. Discussion

A patient's microbiota plays a key role in modulating the immune system as a natural defense against tumor development and during cancer treatment, especially in CRC.²⁰ However, there is no defined therapeutic strategy for the microbiota in a CRC patient. To make the transition from clinical trials to patient treatment, it is necessary to review all published data on microbiota-related therapeutic approaches in CRC in a systematic way. The therapeutic modulators of the microbiome include primarily probiotics, prebiotics, symbiotics and nutrition.

Probiotics

One of the most frequent probiotic combinations for the study of CRC is that of *Lactobacillus* and *Bifidobacterium*. In several studies, pre-inoculation with *Lactobacillus acidophilus* or a cocktail approved for commercial use in China, called Bifico, which consists of *Bifidobacterium longum*, *Lactobacillus acidophilus*, and *Enterococcus faecalis*, reduced colon tumor growth in preclinical models.^{118, 169} Bifico treatment also caused a decrease in *Desulfovibrio*, *Mucispirillum* and *Odoribacter*, associated with an enrichment in *Lactobacillus*.¹¹⁸ However, another study reported that a combination of *Bifidobacterium* and *Lactobacillus* was not able to improve 5-FU treatment, despite improving the flora composition.¹²³

The mixture of three different strains of *Lactobacillus* and one of *Bifidobacterium* decreases the incidence and number of tumors in a preclinical model of CRC associated with colitis.¹³⁰ VSL#3 is a commercial probiotic containing a mixture of different strains of *Lactobacillus*, *Bifidobacterium* and *Streptococcus*. In a murine model with a Western diet, treatment with VSL#3 and metformin decreases colonic tumor formation, through reduced cell proliferation and macrophage infiltration.¹²⁸ However, VSL#3 alone shows no protection against colonic tumor development in other studies.^{116, 128} In fact, in a model of colitis associated with CRC and inflammation, VSL#3 increases macroscopic lesions, as well as their invasiveness and histological grade.¹¹⁶ Therefore, it is important not to prescribe probiotics without scientific basis.

In addition, the *Lactobacillus* genus alone is probably the most widely tested probiotic in CRC (**Fig. 2 d&e**). A study conducted with *Lactobacillus casei* in a model of CRC associated with colitis, combined with FOLFOX, showed improvement in various inflammatory and microbiota parameters; however, it did not show reduction in tumor development.¹²⁴ In this study, FOLFOX increases apoptosis in intestinal epithelial cells, while the probiotic inhibits cell death in healthy epithelial cells.¹²⁴ Nonetheless, it is difficult to assess the effect of a given treatment in an animal model, but it is even more complicated in models that use injections of tumor cells in different (heterotopic) locations, as is the case in this study.¹²⁴ Another study with the *L. casei* BL23 strain showed a total inhibition of the incidence of colorectal tumors in mice. This mechanism could be associated with increased expression of Bik, a proapoptotic gene, and a decrease in Ki67.¹²⁶ *Lactobacillus reuteri* strain can also reduce colon tumors through the production of histamine, which is associated with a decrease in inflammatory markers at both local and systemic levels.¹²⁷

Another well-studied probiotic is *Akkermansia muciniphila* (**Fig. 2 d**). Several studies support its potential use in CRC patients with promising results. Preclinical studies have shown that it is more efficient against CRC than IL-2 therapy. The combination of *A. muciniphila* and IL-2 was significantly more effective than their separate actions in reducing tumor volume and increasing survival.¹¹⁵ The effect of *A. muciniphila* appears to be related to a membrane protein called Amuc, which is recognized by Toll-like receptor 2 (TLR2). Cellular mechanisms and immune effects mediated by TLR2 lead to increased efficacy of IL-2 immune therapy.¹¹⁵

Therefore, it seems that probiotics, especially lactic acid bacteria, SCFA producers and *A. muciniphila* could be beneficial for patients diagnosed with CRC.

Prebiotics

Many of the prebiotics used in the CRC literature are oligo- and polysaccharides (Fig. 2 e & f). Among them, pectin has shown beneficial effects such as SCFA production,¹⁷⁰ reduction of ammonia,¹⁷¹ and improve glucose tolerance.¹⁷² Pectin is the fundamental metabolite of plant walls, and its fermentation serves as a source of energy for numerous microorganisms, producing metabolites with beneficial functions for our organism.^{131, 173} Furthermore, pectin is currently used in numerous food additives, so its study is of great interest.^{131, 174} In fact, it has been observed that pectin can be beneficial against several types of tumors such as CRC, breast and prostate cancer by halting tumor development.¹⁷⁴ Its effect seems to be mediated through the increase of different SCFA-producing bacteria and in particular acetate, and the enhancement of the immune response through IFN- γ -producing TCD8+. In addition, a transplantation of fecal microbiota from PD-1 refractory patients showed enhanced anti-tumor response in PD-1 and pectin-treated mice.¹³¹

Galacto-oligosaccharides derived from lactulose, and mainly lactose, have been tested in preclinical models with promising results in terms of reducing tumor development and modulating the microbiota.¹⁴⁰ Other well-studied oligosaccharides include fructo-oligosaccharides/inulin and mucin.^{66, 132, 164} Inulin has been shown to significantly reduce the volume of subcutaneously transplanted colon tumors, while mucin did not cause any significant change.¹³² In another study, inulin was reported to strongly inhibit tumor development through modification of the microbiota and butyrate production.⁶⁶ This paper dissects the pathways leading to this effect, finding that butyrate is absorbed, but not oxidized due to the Warburg effect of cancer cells.¹⁷⁵ However, healthy colonocytes do use butyrate as a major source of energy.¹⁷⁶ In tumor cells, butyrate accumulates in the cell interior where it inhibits histone deacetylase activity, thus decreasing proliferation and increasing apoptosis.⁶⁶

Studies with compounds derived from citrus peel, such as polymethoxyflavones, have shown positive effects in preventing the development of CRC. However, mechanisms such as autophagy, which the authors highlight as possible effector pathways for these compounds, are poorly analyzed and higher quality studies are needed to draw valid conclusions.¹⁴² Other prebiotics, such as green tea extract

(EGCG), have been shown to reduce colonic tumors, but these studies are of low scientific quality (grading = 1 and 2) for the reasons mentioned above. In a rat model of colitis-associated cancer using EGCG and resistant starch, EGCG was shown to have no effect. Only resistant starch was able to decrease colonic tumor formation through a change in the microbial flora, increasing SCFA production, and decreasing inflammation.¹⁵⁴

There is great interest among scientists in Asia, but particularly in China, in demonstrating the effects of different therapeutic approaches to traditional medicine through the microbiota.^{135, 148} For example, the Canmei formula contains herbs called *Mume sieb* and *Marci Hieronymi*, whose more than 41 active ingredients include DL-arginine, L-tyrosine, L(-)-carnitine, adenine, guanine or ambrosic acid. This formulation was shown to decrease the incidence of colonic tumors by reducing inflammation and improving dysbiosis.¹³⁷ Another study using a *Boswellia serrata* resin extract has also demonstrated, in preclinical models, its ability to reduce colonic tumor formation.¹⁴⁴ Many of these studies present positive results for the inhibition of CRC development. However, in general, they are of low quality (grading = 1 and 2) due to a lack of mechanistic and statistically robust data and therefore do not allow for reliable conclusions to be drawn.

Symbiotics

One of the most interesting proposals in the literature is the possibility of using symbiotics as therapeutic options to increase the efficacy of available treatments. It has been observed that the use of *C. butyricum*, surrounded by dextran and bound to capecitabine or diclofenac, can increase the efficacy of these chemotherapeutics by >500%. This symbiotic decreases tumor invasiveness and also promotes a healthier and more diverse microbiota coupled with increased production of SCFA.⁹⁶ This work is interesting for clinical studies as both dextran and *C. butyricum* are approved for use in food applications and showed no notable side effects.⁹⁶ Another symbiotic studied is the combination of *Lactobacillus gasseri* 505 and *Cudrania tricuspidata* leaf extract in fermented milk. This symbiotic was shown to reduce tumor incidence and malignancy in a model of CRC-associated colitis.¹⁴⁶

Therefore, the use of symbiotics for improving gut microbiota in patients with CRC seems promising, although more studies are needed to clearly define which symbiotic should be used, the dose, the timing, and in which specific CRC patients.

Healthy nutrition as a source of pre- and probiotics

Most, if not all pro-, pre- and symbiotics can be obtained naturally through a healthy diet. Prebiotics have been isolated from mushrooms,¹³⁸ fruits,^{142, 151} dairy,¹⁴⁰ rice,¹⁶¹ and green tea¹³⁸ and probiotics from fruit skins,¹¹⁷ dairy,¹²² and fermented products.¹⁴⁶

From infancy, a balanced, non-processed plant-based nutrition is the best way to have a proper gut microbiota. In fact, it has been shown that a diet high in soluble fiber is not able to reverse the erosion of intestinal mucus caused by a diet without fiber. However, the same diet, but high in intact plant fiber decreased mucus degradation, strengthening the intestinal barrier.⁶⁷

Anthocyanins, compounds with a positive impact on the health of patients with CRC, are present in many foods such as tea, coffee, cocoa and red fruits.¹⁷⁷ In this regard, it has been shown that anthocyanins in blackberries can reduce the incidence of CRC, through diminished local inflammation and improved microbiota.¹⁵¹

Another food with interesting results is the marine extract of certain shellfish that may have anticarcinogenic effects. Using metabolomic analysis techniques, one can discern which compounds are toxic to tumor cells, opening the door to precision nutrition.^{147, 178} Algae-derived products such as fucoxanthin, a carotenoid present in high levels in wakame (*Undaria pinnatifida*), have also shown to decrease the number of adenocarcinomas and total tumors. This effect was also associated with an increase in the number of *Lachnospiraceae*.¹³⁴ Spices have similarly attracted a great deal of interest for their health-promoting properties. For example, nutmeg can decrease the development of APC genetically engineered tumors by modulating the metabolism of the microbial flora and improving lipid metabolism.¹⁵⁰ Numerous food extracts are also used in traditional Chinese medicine, including American ginseng. The use of this compound in a murine model prevented tumors and ameliorated dysbiosis induced by different murine models of CRC and high-fat diet.^{153, 155}

Moving from specific foods to dietary patterns, the Mediterranean diet has been shown to have beneficial effects on multiple levels, specifically in the prevention of certain types of cancer.¹⁷⁹ In the case of CRC, epidemiological studies show that adherence to a Mediterranean diet reduces the incidence of CRC by up to 11%.¹⁸⁰ Likewise, the effect of mixing a Mediterranean diet with other types of diet (low-fat or high-fat Western-type diet) has been studied in models of cancer associated with

colitis. This study showed that consumption of the Mediterranean diet, regardless of the diet with which it was mixed, decreased tumor incidence, and improved the microbiota, especially when accompanying a high-fat diet.¹⁴⁹

Randomized clinical trials

At the highest level of scientific evidence are randomized clinical trials. Within this group, those conducted with lactic acid strains stand out. Among them, a study conducted with 78 patients analyzed the role of probiotics in reducing side effects after CRC surgery. Inclusion criteria were a diagnosis of stage III adenomatous CRC and that the patients were to undergo surgery. The probiotic group received post-operative treatment with a mixture of different strains of *Lactobacillus*, *Bifidobacterium*, and *Streptococcus thermophilus* for almost one year. The most important conclusion of this study was that probiotic treatment decreased the most frequent side effects of surgery, specifically paralytic ileus. However, limitations of this study include the low number of subjects included, the apparent lack of double-blinding, active placebo, and a weak statistical analysis.⁷⁴ In another clinical trial, a probiotic consisting of *Bifidobacterium longum*, *Lactobacillus acidophilus* and *Enterococcus faecalis* was tested in CRC patients undergoing surgery. It was shown that, in addition to improving the diversity of the microbial flora, the probiotic mixture was able to decrease the levels of *Fusobacterium*, a bacterium associated with the development of CRC, by >500%.¹⁵⁶ In addition, its use decreased post-operative infections.¹⁵⁷ Another similar probiotic mixture consisting of *Lactobacillus plantarum*, *Lactobacillus acidophilus* and *Bifidobacterium longum* reduced the risk of postoperative infection and improved the intestinal epithelial barrier, resulting in a lower incidence of diarrhea, abdominal pain, and fever.¹⁵⁹ A study using a mixture of *Bifidobacterium longum* (BB536) and *Lactobacillus johnsonii* (La1) showed that only the latter (La1) colonized the gut, reducing the number of pathogenic microorganisms and reducing colonic inflammation in CRC patients undergoing surgery. This result shows that adding more probiotics will not have a greater effect and it is important to study each strain individually and its combination before it can be recommended to patients.¹⁶⁰

In another clinical trial, a probiotic mixture consisting of *Enterococcus*, *Clostridium* and *Bacillus* was used in patients undergoing CRC surgery to discern whether their consumption could ameliorate the adverse effects of surgery. It was found that peri-

surgical use of this mixture significantly reduced incisional infections and decreased the acute inflammatory response to surgery.¹⁵⁸ However, in this study, no randomization was performed, and patients were included in one group (control) or the other (probiotic) according to the date of surgery, with the control group (2009-2011) prior to the probiotic group (2011-2013). This does not allow for a clear conclusion on the effect of the probiotic, because multiple factors may have changed during those two years, including greater experience of the surgical team or better technical means. In this regard, the authors describe that a higher number of laparotomies were performed in the probiotic group, and that performing or not performing laparotomy is a key factor related to the outcome of the surgery.¹⁵⁸

As far as prebiotics are concerned, rice bran is of great interest due to its large commercial use. Numerous studies show the benefit of whole grain consumption in the prevention of CRC.¹⁸¹ In the case of rice bran, various phytochemicals in its composition have been associated with a decrease in the risk of cancer development, in particular CRC.¹⁸² Its consumption increases levels of *Firmicutes* and *Lactobacillus*¹⁶¹ and produces numerous metabolic modifications that could be related to a decrease in the risk of developing CRC.¹⁶⁶

In another clinical trial with prebiotics in CRC patients, a mixture of different non-digestible oligosaccharides was used for 7 days before patients underwent radical surgery to remove the tumor. Both groups received the same nutrition (provided by the hospital) in the 7 days before and after surgery (parenteral nutrition). Surgery decreased the diversity of the microbiota in both groups, underlining the impact of these interventions on gut health. Interestingly, a principal component analysis of microbiome diversity was able to unequivocally identify samples from postoperative patients who had received prebiotics. Furthermore, the use of these prebiotics improved systemic immune function, confirming the positive effect that microbiota can have on immune response. An important limitation of this study is that the randomization did not specify the stage or type of tumor.¹⁶²

Symbiotics have also been tested in randomized clinical trials. A commercially named symbiotic, simbioflora®, containing fructooligosaccharides, 3 strains of *Lactobacillus*, and *Bifidobacterium lactis* showed promising results in CRC patients undergoing surgery. The use of this symbiotic significantly reduced inflammation and postoperative infections.¹⁶³ However, this is a small study of patients with cancer at different stages, so the data should be analyzed with caution. In another study, a

mixture of *Bifidobacterium*, *Lactobacillus* and inulin strains was used in stage I-III patients undergoing surgery. This treatment resulted in an improvement in the composition of the microbial community, especially the SCFA-producing bacterium. However, this study does not define blind randomization of patients and there is no placebo, so the data should be also analyzed with caution.¹⁶⁴ Finally, a randomized clinical study was conducted in 20 healthy subjects with backcrossed arms with the aim of preventing CRC. This study of 4 weeks duration per intervention (switching every 4 weeks to another arm until each group completed all 4) was done with 4 different arms: placebo, *Bifidobacterium lactis*, resistant starch and their combination (*Bifidobacterium lactis* and resistant starch). It was shown that in healthy subjects the combination of *Bifidobacterium lactis* and resistant starch increased the levels of *Lachnospiraceae spp.* However, the other parameters analyzed, such as stool mass, pH or ammonium, SCFA, circulating cytokines, and markers of colonic epithelial health, did not change.¹⁶⁵

The use of pomegranate extract in patients undergoing surgical tumor resection has also been investigated. Pomegranate extract was shown to produce significant changes in the expression of numerous genes and microRNAs related to tumor development. These results underline the crucial role of diet in modifying gene expression and, therefore, in the etiology of disease.^{167, 168}

5. Conclusion

We can summarize the conclusions of this work regarding the modification of the microbiota in the CRC patient as follows:

1) **Antibiotic use** is one of the **major deleterious factors** for the prognosis of **treatment efficacy**, due to its **harmful effect on the microbiota**.

2) **Prebiotics** such as inulin and resistant starch can support the development of a healthy microbiota.

3) **Probiotics**, mainly based on **SCFA-producing lactic acid strains**, show a **favorable profile** in preclinical and clinical studies. However, not all strains are equally favorable and are therefore not recommended for use in patients without individualized study of the patient and the probiotic.

4) A **healthy** and varied **nutrition**, consisting mainly of unprocessed **plant-based foods**, seems to be one of the **best sources** to maintain a **homeostatic microbiota**.

Finally, after a detailed analysis of the literature, in a **healthy person**, there is no need to recommend the use of pre-, pro- or symbiotics in addition to a **healthy diet**. However, CRC patients often have severe dysbiosis due to their own disease, and broad-spectrum antibiotic use. In these patients, apart from a **healthy diet**, or if a healthy diet cannot be established, carefully chosen pre-, pro- and/or symbiotics may be beneficial to restore a beneficial microbiota and support the efficacy of hospital treatments more effectively.

Acknowledgements

We thank Dr. Alberto Canfran-Duque and Noemi Rotllan for their for critical reading of the manuscript. We apologize to those whose work could not be cited owing to space limitations.

Authors Contributions

JMM designed and performed the systematic search, wrote and edited the manuscript. SBE contributed to the editing and interpretation of the manuscript. All authors read and edited the final draft of the manuscript.

Competing Financial Interest

The authors declare no conflicts of interest

Correspondence and requests for materials should be addressed to: madrigal-matute@ibions.com.

Data availability

There are not restrictions on data availability in this manuscript.

616

617 6. Bibliography

- 618 1. Henrikson, N.B. *et al.* Family history and the natural history of colorectal
619 cancer: systematic review. *Genet Med* **17**, 702-712 (2015).
- 620 2. Czene, K., Lichtenstein, P. & Hemminki, K. Environmental and heritable
621 causes of cancer among 9.6 million individuals in the Swedish Family-Cancer
622 Database. *Int J Cancer* **99**, 260-266 (2002).
- 623 3. Lichtenstein, P. *et al.* Environmental and heritable factors in the causation of
624 cancer--analyses of cohorts of twins from Sweden, Denmark, and Finland. *N Engl J*
625 *Med* **343**, 78-85 (2000).
- 626 4. Jiao, S. *et al.* Estimating the heritability of colorectal cancer. *Hum Mol Genet*
627 **23**, 3898-3905 (2014).
- 628 5. Syngal, S. *et al.* ACG clinical guideline: Genetic testing and management of
629 hereditary gastrointestinal cancer syndromes. *Am J Gastroenterol* **110**, 223-262; quiz
630 263 (2015).
- 631 6. Dekker, E., Tanis, P.J., Vleugels, J.L.A., Kasi, P.M. & Wallace, M.B. Colorectal
632 cancer. *Lancet* **394**, 1467-1480 (2019).
- 633 7. Nishida, A. *et al.* Gut microbiota in the pathogenesis of inflammatory bowel
634 disease. *Clin J Gastroenterol* **11**, 1-10 (2018).
- 635 8. Murphy, N. *et al.* Lifestyle and dietary environmental factors in colorectal
636 cancer susceptibility. *Mol Aspects Med* **69**, 2-9 (2019).
- 637 9. Crosnier, C., Stamataki, D. & Lewis, J. Organizing cell renewal in the intestine:
638 stem cells, signals and combinatorial control. *Nat Rev Genet* **7**, 349-359 (2006).
- 639 10. (IARC Working Group on the Evaluation of Carcinogenic Risks to Humans.
640 Lyon (FR): International Agency for Research on Cancer; 2010. (IARC Monographs
641 on the Evaluation of Carcinogenic Risks to Humans, No. 92.); 2010).
- 642 11. Mottram, D.S., Wedzicha, B.L. & Dodson, A.T. Acrylamide is formed in the
643 Maillard reaction. *Nature* **419**, 448-449 (2002).
- 644 12. (IARC Working Group on the Evaluation of Carcinogenic Risks to Humans.
645 Chemical Agents and Related Occupations. Lyon (FR): International Agency for
646 Research on Cancer; 2012. (IARC Monographs on the Evaluation of Carcinogenic
647 Risks to Humans, No. 100F.) 2012).

13. Diggs, D.L. *et al.* Polycyclic aromatic hydrocarbons and digestive tract cancers: a perspective. *J Environ Sci Health C Environ Carcinog Ecotoxicol Rev* **29**, 324-357 (2011).
14. Tabatabaei, S.M., Heyworth, J.S., Knuiman, M.W. & Fritschi, L. Dietary benzo[a]pyrene intake from meat and the risk of colorectal cancer. *Cancer Epidemiol Biomarkers Prev* **19**, 3182-3184 (2010).
15. Coussens, L.M. & Werb, Z. Inflammation and cancer. *Nature* **420**, 860-867 (2002).
16. Medema, J.P. Cancer stem cells: the challenges ahead. *Nat Cell Biol* **15**, 338-344 (2013).
17. Cancer Genome Atlas, N. Comprehensive molecular characterization of human colon and rectal cancer. *Nature* **487**, 330-337 (2012).
18. Stappenbeck, T.S. & Virgin, H.W. Accounting for reciprocal host-microbiome interactions in experimental science. *Nature* **534**, 191-199 (2016).
19. Zitvogel, L., Ma, Y., Raoult, D., Kroemer, G. & Gajewski, T.F. The microbiome in cancer immunotherapy: Diagnostic tools and therapeutic strategies. *Science* **359**, 1366-1370 (2018).
20. Thursby, E. & Juge, N. Introduction to the human gut microbiota. *Biochem J* **474**, 1823-1836 (2017).
21. Claesson, M.J. *et al.* Gut microbiota composition correlates with diet and health in the elderly. *Nature* **488**, 178-184 (2012).
22. Wang, T. *et al.* Structural segregation of gut microbiota between colorectal cancer patients and healthy volunteers. *ISME J* **6**, 320-329 (2012).
23. Lin, Y. *et al.* NMR-based fecal metabolomics fingerprinting as predictors of earlier diagnosis in patients with colorectal cancer. *Oncotarget* **7**, 29454-29464 (2016).
24. Chen, J. & Vitetta, L. Inflammation-Modulating Effect of Butyrate in the Prevention of Colon Cancer by Dietary Fiber. *Clin Colorectal Cancer* **17**, e541-e544 (2018).
25. Zeng, H., Safratowich, B.D., Wang, T.T.Y., Hamlin, S.K. & Johnson, L.K. Butyrate Inhibits Deoxycholic-Acid-Resistant Colonic Cell Proliferation via Cell Cycle Arrest and Apoptosis: A Potential Pathway Linking Dietary Fiber to Cancer Prevention. *Mol Nutr Food Res* **64**, e1901014 (2020).

- 681 26. Tan, J. *et al.* The role of short-chain fatty acids in health and disease. *Adv*
682 *Immunol* **121**, 91-119 (2014).
- 683 27. Ganapathy, V., Thangaraju, M., Prasad, P.D., Martin, P.M. & Singh, N.
684 Transporters and receptors for short-chain fatty acids as the molecular link between
685 colonic bacteria and the host. *Curr Opin Pharmacol* **13**, 869-874 (2013).
- 686 28. Zhao, Y. *et al.* GPR43 mediates microbiota metabolite SCFA regulation of
687 antimicrobial peptide expression in intestinal epithelial cells via activation of mTOR
688 and STAT3. *Mucosal Immunol* **11**, 752-762 (2018).
- 689 29. Madara, J.L. & Pappenheimer, J.R. Structural basis for physiological
690 regulation of paracellular pathways in intestinal epithelia. *J Membr Biol* **100**, 149-164
691 (1987).
- 692 30. Eeckhaut, V. *et al.* Butyricococcus pullicaecorum in inflammatory bowel
693 disease. *Gut* **62**, 1745-1752 (2013).
- 694 31. Kang, S. *et al.* Dysbiosis of fecal microbiota in Crohn's disease patients as
695 revealed by a custom phylogenetic microarray. *Inflamm Bowel Dis* **16**, 2034-2042
696 (2010).
- 697 32. Takaishi, H. *et al.* Imbalance in intestinal microflora constitution could be
698 involved in the pathogenesis of inflammatory bowel disease. *Int J Med Microbiol* **298**,
699 463-472 (2008).
- 700 33. Karlsson, C.L. *et al.* The microbiota of the gut in preschool children with
701 normal and excessive body weight. *Obesity (Silver Spring)* **20**, 2257-2261 (2012).
- 702 34. Crovesy, L., Masterson, D. & Rosado, E.L. Profile of the gut microbiota of
703 adults with obesity: a systematic review. *Eur J Clin Nutr* **74**, 1251-1262 (2020).
- 704 35. Vigsnaes, L.K., Brynskov, J., Steenholdt, C., Wilcks, A. & Licht, T.R. Gram-
705 negative bacteria account for main differences between faecal microbiota from
706 patients with ulcerative colitis and healthy controls. *Benef Microbes* **3**, 287-297
707 (2012).
- 708 36. Wang, L. *et al.* A purified membrane protein from Akkermansia muciniphila or
709 the pasteurised bacterium blunts colitis associated tumourigenesis by modulation of
710 CD8(+) T cells in mice. *Gut* **69**, 1988-1997 (2020).
- 711 37. Ottman, N. *et al.* Genome-Scale Model and Omics Analysis of Metabolic
712 Capacities of Akkermansia muciniphila Reveal a Preferential Mucin-Degrading
713 Lifestyle. *Appl Environ Microbiol* **83** (2017).

38. Wang, G. *et al.* Role of SCFAs in gut microbiome and glycolysis for colorectal cancer therapy. *J Cell Physiol* **234**, 17023-17049 (2019).
39. Petersen, C. & Round, J.L. Defining dysbiosis and its influence on host immunity and disease. *Cell Microbiol* **16**, 1024-1033 (2014).
40. Wong, S.H. & Yu, J. Gut microbiota in colorectal cancer: mechanisms of action and clinical applications. *Nat Rev Gastroenterol Hepatol* **16**, 690-704 (2019).
41. Peters, B.A. *et al.* The gut microbiota in conventional and serrated precursors of colorectal cancer. *Microbiome* **4**, 69 (2016).
42. Dethlefsen, L., Huse, S., Sogin, M.L. & Relman, D.A. The pervasive effects of an antibiotic on the human gut microbiota, as revealed by deep 16S rRNA sequencing. *PLoS Biol* **6**, e280 (2008).
43. Henao-Mejia, J. *et al.* Inflammasome-mediated dysbiosis regulates progression of NAFLD and obesity. *Nature* **482**, 179-185 (2012).
44. Newton, D.F., Cummings, J.H., Macfarlane, S. & Macfarlane, G.T. Growth of a human intestinal *Desulfovibrio desulfuricans* in continuous cultures containing defined populations of saccharolytic and amino acid fermenting bacteria. *J Appl Microbiol* **85**, 372-380 (1998).
45. Carbonero, F., Benefiel, A.C. & Gaskins, H.R. Contributions of the microbial hydrogen economy to colonic homeostasis. *Nat Rev Gastroenterol Hepatol* **9**, 504-518 (2012).
46. Roediger, W.E., Duncan, A., Kapaniris, O. & Millard, S. Reducing sulfur compounds of the colon impair colonocyte nutrition: implications for ulcerative colitis. *Gastroenterology* **104**, 802-809 (1993).
47. Suarez, F., Furne, J., Springfield, J. & Levitt, M. Production and elimination of sulfur-containing gases in the rat colon. *Am J Physiol* **274**, G727-733 (1998).
48. Cai, W.J., Wang, M.J., Ju, L.H., Wang, C. & Zhu, Y.C. Hydrogen sulfide induces human colon cancer cell proliferation: role of Akt, ERK and p21. *Cell Biol Int* **34**, 565-572 (2010).
49. Deplancke, B. & Gaskins, H.R. Hydrogen sulfide induces serum-independent cell cycle entry in nontransformed rat intestinal epithelial cells. *FASEB J* **17**, 1310-1312 (2003).
50. Attene-Ramos, M.S., Wagner, E.D., Plewa, M.J. & Gaskins, H.R. Evidence that hydrogen sulfide is a genotoxic agent. *Mol Cancer Res* **4**, 9-14 (2006).

51. Kanazawa, K. *et al.* Factors influencing the development of sigmoid colon cancer. Bacteriologic and biochemical studies. *Cancer* **77**, 1701-1706 (1996).
52. Kostic, A.D. *et al.* Genomic analysis identifies association of *Fusobacterium* with colorectal carcinoma. *Genome Res* **22**, 292-298 (2012).
53. Castellarin, M. *et al.* *Fusobacterium nucleatum* infection is prevalent in human colorectal carcinoma. *Genome Res* **22**, 299-306 (2012).
54. Yang, Y. *et al.* *Fusobacterium nucleatum* Increases Proliferation of Colorectal Cancer Cells and Tumor Development in Mice by Activating Toll-Like Receptor 4 Signaling to Nuclear Factor-kappaB, and Up-regulating Expression of MicroRNA-21. *Gastroenterology* **152**, 851-866 e824 (2017).
55. Yu, T. *et al.* *Fusobacterium nucleatum* Promotes Chemoresistance to Colorectal Cancer by Modulating Autophagy. *Cell* **170**, 548-563 e516 (2017).
56. Campisciano, G. *et al.* The Obesity-Related Gut Bacterial and Viral Dysbiosis Can Impact the Risk of Colon Cancer Development. *Microorganisms* **8** (2020).
57. Feng, Q. *et al.* Gut microbiome development along the colorectal adenoma-carcinoma sequence. *Nat Commun* **6**, 6528 (2015).
58. Arthur, J.C. *et al.* Intestinal inflammation targets cancer-inducing activity of the microbiota. *Science* **338**, 120-123 (2012).
59. Garrett, W.S. The gut microbiota and colon cancer. *Science* **364**, 1133-1135 (2019).
60. Dejea, C.M. *et al.* Patients with familial adenomatous polyposis harbor colonic biofilms containing tumorigenic bacteria. *Science* **359**, 592-597 (2018).
61. Zhang, G., Svenungsson, B., Karnell, A. & Weintraub, A. Prevalence of enterotoxigenic *Bacteroides fragilis* in adult patients with diarrhea and healthy controls. *Clin Infect Dis* **29**, 590-594 (1999).
62. Turnbaugh, P.J. *et al.* A core gut microbiome in obese and lean twins. *Nature* **457**, 480-484 (2009).
63. Turnbaugh, P.J. *et al.* The effect of diet on the human gut microbiome: a metagenomic analysis in humanized gnotobiotic mice. *Sci Transl Med* **1**, 6ra14 (2009).
64. Carmody, R.N. *et al.* Diet dominates host genotype in shaping the murine gut microbiota. *Cell Host Microbe* **17**, 72-84 (2015).
65. De Filippis, F. *et al.* High-level adherence to a Mediterranean diet beneficially impacts the gut microbiota and associated metabolome. *Gut* **65**, 1812-1821 (2016).

66. Donohoe, D.R. *et al.* A gnotobiotic mouse model demonstrates that dietary fiber protects against colorectal tumorigenesis in a microbiota- and butyrate-dependent manner. *Cancer Discov* **4**, 1387-1397 (2014).
67. Desai, M.S. *et al.* A Dietary Fiber-Deprived Gut Microbiota Degrades the Colonic Mucus Barrier and Enhances Pathogen Susceptibility. *Cell* **167**, 1339-1353 e1321 (2016).
68. Halmos, E.P., Power, V.A., Shepherd, S.J., Gibson, P.R. & Muir, J.G. A diet low in FODMAPs reduces symptoms of irritable bowel syndrome. *Gastroenterology* **146**, 67-75 e65 (2014).
69. Sanders, M.E., Merenstein, D.J., Reid, G., Gibson, G.R. & Rastall, R.A. Probiotics and prebiotics in intestinal health and disease: from biology to the clinic. *Nat Rev Gastroenterol Hepatol* **16**, 605-616 (2019).
70. Nami, Y., Haghshenas, B., Haghshenas, M., Abdullah, N. & Yari Khosroushahi, A. The Prophylactic Effect of Probiotic *Enterococcus lactis* IW5 against Different Human Cancer Cells. *Front Microbiol* **6**, 1317 (2015).
71. Kaeid Sharaf, L. & Shukla, G. Probiotics (*Lactobacillus acidophilus* and *Lactobacillus rhamnosus* GG) in Conjunction with Celecoxib (selective COX-2 inhibitor) Modulated DMH-Induced Early Experimental Colon Carcinogenesis. *Nutr Cancer* **70**, 946-955 (2018).
72. Sharaf, L.K., Sharma, M., Chandel, D. & Shukla, G. Prophylactic intervention of probiotics (*L.acidophilus*, *L.rhamnosus* GG) and celecoxib modulate Bax-mediated apoptosis in 1,2-dimethylhydrazine-induced experimental colon carcinogenesis. *BMC Cancer* **18**, 1111 (2018).
73. Li, J. *et al.* Probiotics modulated gut microbiota suppresses hepatocellular carcinoma growth in mice. *Proc Natl Acad Sci U S A* **113**, E1306-1315 (2016).
74. Bajramagic, S. *et al.* Usage of Probiotics and its Clinical Significance at Surgically Treated Patients Suffering from Colorectal Carcinoma. *Med Arch* **73**, 316-320 (2019).
75. Azcarate-Peril, M.A., Sikes, M. & Bruno-Barcena, J.M. The intestinal microbiota, gastrointestinal environment and colorectal cancer: a putative role for probiotics in prevention of colorectal cancer? *Am J Physiol Gastrointest Liver Physiol* **301**, G401-424 (2011).

76. Zhuo, Q. *et al.* Lysates of *Lactobacillus acidophilus* combined with CTLA-4-blocking antibodies enhance antitumor immunity in a mouse colon cancer model. *Sci Rep* **9**, 20128 (2019).
77. Eeckhaut, V. *et al.* The Probiotic *Butyricococcus pullicaecorum* Reduces Feed Conversion and Protects from Potentially Harmful Intestinal Microorganisms and Necrotic Enteritis in Broilers. *Front Microbiol* **7**, 1416 (2016).
78. Everard, A. *et al.* Cross-talk between *Akkermansia muciniphila* and intestinal epithelium controls diet-induced obesity. *Proc Natl Acad Sci U S A* **110**, 9066-9071 (2013).
79. Gopalakrishnan, V. *et al.* Gut microbiome modulates response to anti-PD-1 immunotherapy in melanoma patients. *Science* **359**, 97-103 (2018).
80. Routy, B. *et al.* Gut microbiome influences efficacy of PD-1-based immunotherapy against epithelial tumors. *Science* **359**, 91-97 (2018).
81. Zhai, R. *et al.* Strain-Specific Anti-inflammatory Properties of Two *Akkermansia muciniphila* Strains on Chronic Colitis in Mice. *Front Cell Infect Microbiol* **9**, 239 (2019).
82. Paolillo, R., Romano Carratelli, C., Sorrentino, S., Mazzola, N. & Rizzo, A. Immunomodulatory effects of *Lactobacillus plantarum* on human colon cancer cells. *Int Immunopharmacol* **9**, 1265-1271 (2009).
83. Kahouli, I., Malhotra, M., Westfall, S., Alaoui-Jamali, M.A. & Prakash, S. Design and validation of an orally administrated active *L. fermentum*-*L. acidophilus* probiotic formulation using colorectal cancer Apc (Min/+) mouse model. *Appl Microbiol Biotechnol* **101**, 1999-2019 (2017).
84. Zhu, J. *et al.* *Lactobacillus salivarius* Ren prevent the early colorectal carcinogenesis in 1, 2-dimethylhydrazine-induced rat model. *J Appl Microbiol* **117**, 208-216 (2014).
85. Chen, L. *et al.* NLRP12 attenuates colon inflammation by maintaining colonic microbial diversity and promoting protective commensal bacterial growth. *Nat Immunol* **18**, 541-551 (2017).
86. Redman, M.G., Ward, E.J. & Phillips, R.S. The efficacy and safety of probiotics in people with cancer: a systematic review. *Ann Oncol* **25**, 1919-1929 (2014).
87. Gibson, G.R. *et al.* Expert consensus document: The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the

definition and scope of prebiotics. *Nat Rev Gastroenterol Hepatol* **14**, 491-502 (2017).

88. van Loo, J., Coussement, P., de Leenheer, L., Hoebregs, H. & Smits, G. On the presence of inulin and oligofructose as natural ingredients in the western diet. *Crit Rev Food Sci Nutr* **35**, 525-552 (1995).

89. Roberfroid, M.B., Van Loo, J.A. & Gibson, G.R. The bifidogenic nature of chicory inulin and its hydrolysis products. *J Nutr* **128**, 11-19 (1998).

90. Campbell, J.M., Fahey, G.C., Jr. & Wolf, B.W. Selected indigestible oligosaccharides affect large bowel mass, cecal and fecal short-chain fatty acids, pH and microflora in rats. *J Nutr* **127**, 130-136 (1997).

91. Fehlbauer, S. *et al.* In Vitro Fermentation of Selected Prebiotics and Their Effects on the Composition and Activity of the Adult Gut Microbiota. *Int J Mol Sci* **19** (2018).

92. Li, K. *et al.* Dietary inulin alleviates diverse stages of type 2 diabetes mellitus via anti-inflammation and modulating gut microbiota in db/db mice. *Food Funct* **10**, 1915-1927 (2019).

93. Fuentes-Zaragoza, E. *et al.* Resistant starch as prebiotic: A review. *Starch - Stärke* **63**, 406-415 (2011).

94. Sharma, A., Yadav, B.S. & Ritika Resistant Starch: Physiological Roles and Food Applications. *Food Reviews International* **24**, 193-234 (2008).

95. Faraj, A., Vasanathan, T. & Hoover, R. The effect of extrusion cooking on resistant starch formation in waxy and regular barley flours. *Food Research International* **37**, 517-525 (2004).

96. Zheng, D.W. *et al.* Prebiotics-Encapsulated Probiotic Spores Regulate Gut Microbiota and Suppress Colon Cancer. *Adv Mater* **32**, e2004529 (2020).

97. Swanson, K.S. *et al.* The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of synbiotics. *Nat Rev Gastroenterol Hepatol* **17**, 687-701 (2020).

98. Le Leu, R.K., Hu, Y., Brown, I.L., Woodman, R.J. & Young, G.P. Synbiotic intervention of *Bifidobacterium lactis* and resistant starch protects against colorectal cancer development in rats. *Carcinogenesis* **31**, 246-251 (2010).

99. Ganesh, K. *et al.* Immunotherapy in colorectal cancer: rationale, challenges and potential. *Nat Rev Gastroenterol Hepatol* **16**, 361-375 (2019).

100. Paulos, C.M. *et al.* Microbial translocation augments the function of adoptively transferred self/tumor-specific CD8+ T cells via TLR4 signaling. *J Clin Invest* **117**, 2197-2204 (2007).
101. Iida, N. *et al.* Commensal bacteria control cancer response to therapy by modulating the tumor microenvironment. *Science* **342**, 967-970 (2013).
102. Viaud, S. *et al.* The intestinal microbiota modulates the anticancer immune effects of cyclophosphamide. *Science* **342**, 971-976 (2013).
103. Vetizou, M. *et al.* Anticancer immunotherapy by CTLA-4 blockade relies on the gut microbiota. *Science* **350**, 1079-1084 (2015).
104. Sivan, A. *et al.* Commensal Bifidobacterium promotes antitumor immunity and facilitates anti-PD-L1 efficacy. *Science* **350**, 1084-1089 (2015).
105. Gaines, S. *et al.* Western Diet Promotes Intestinal Colonization by Collagenolytic Microbes and Promotes Tumor Formation After Colorectal Surgery. *Gastroenterology* **158**, 958-970 e952 (2020).
106. Chen, D.S. & Mellman, I. Elements of cancer immunity and the cancer-immune set point. *Nature* **541**, 321-330 (2017).
107. Matson, V. *et al.* The commensal microbiome is associated with anti-PD-1 efficacy in metastatic melanoma patients. *Science* **359**, 104-108 (2018).
108. Alberda, W.J. *et al.* The Importance of a Minimal Tumor-Free Resection Margin in Locally Recurrent Rectal Cancer. *Dis Colon Rectum* **58**, 677-685 (2015).
109. Manfredi, S. *et al.* Incidence and patterns of recurrence after resection for cure of colonic cancer in a well defined population. *Br J Surg* **93**, 1115-1122 (2006).
110. Naxerova, K. *et al.* Origins of lymphatic and distant metastases in human colorectal cancer. *Science* **357**, 55-60 (2017).
111. Fermor, B., Umpleby, H.C., Lever, J.V., Symes, M.O. & Williamson, R.C. Proliferative and metastatic potential of exfoliated colorectal cancer cells. *J Natl Cancer Inst* **76**, 347-349 (1986).
112. Shogan, B.D. *et al.* Collagen degradation and MMP9 activation by *Enterococcus faecalis* contribute to intestinal anastomotic leak. *Sci Transl Med* **7**, 286ra268 (2015).
113. Mirnezami, A. *et al.* Increased local recurrence and reduced survival from colorectal cancer following anastomotic leak: systematic review and meta-analysis. *Ann Surg* **253**, 890-899 (2011).

114. Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G. & Group, P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* **6**, e1000097 (2009).
115. Shi, L. *et al.* Combining IL-2-based immunotherapy with commensal probiotics produces enhanced antitumor immune response and tumor clearance. *J Immunother Cancer* **8** (2020).
116. Arthur, J.C. *et al.* VSL#3 probiotic modifies mucosal microbial composition but does not reduce colitis-associated colorectal cancer. *Sci Rep* **3**, 2868 (2013).
117. Chandel, D., Sharma, M., Chawla, V., Sachdeva, N. & Shukla, G. Isolation, characterization and identification of antigenotoxic and anticancerous indigenous probiotics and their prophylactic potential in experimental colon carcinogenesis. *Sci Rep* **9**, 14769 (2019).
118. Song, H. *et al.* Pretreatment with probiotic Bifico ameliorates colitis-associated cancer in mice: Transcriptome and gut flora profiling. *Cancer Sci* **109**, 666-677 (2018).
119. Xu, H. *et al.* Inhibitory Effects of Breast Milk-Derived *Lactobacillus rhamnosus* Probio-M9 on Colitis-Associated Carcinogenesis by Restoration of the Gut Microbiota in a Mouse Model. *Nutrients* **13** (2021).
120. Chang, S.C. *et al.* A gut butyrate-producing bacterium *Butyrivibrio* *pullicaecorum* regulates short-chain fatty acid transporter and receptor to reduce the progression of 1,2-dimethylhydrazine-associated colorectal cancer. *Oncol Lett* **20**, 327 (2020).
121. Hradicka, P., Beal, J., Kassayova, M., Foey, A. & Demeckova, V. A Novel Lactic Acid Bacteria Mixture: Macrophage-Targeted Prophylactic Intervention in Colorectal Cancer Management. *Microorganisms* **8** (2020).
122. Silveira, D.S.C. *et al.* *Lactobacillus bulgaricus* inhibits colitis-associated cancer via a negative regulation of intestinal inflammation in azoxymethane/dextran sodium sulfate model. *World J Gastroenterol* **26**, 6782-6794 (2020).
123. Yuan, L. *et al.* The influence of gut microbiota dysbiosis to the efficacy of 5-Fluorouracil treatment on colorectal cancer. *Biomed Pharmacother* **108**, 184-193 (2018).
124. Chang, C.W. *et al.* *Lactobacillus casei* Variety *rhamnosus* Probiotic Preventively Attenuates 5-Fluorouracil/Oxaliplatin-Induced Intestinal Injury in a Syngeneic Colorectal Cancer Model. *Front Microbiol* **9**, 983 (2018).

125. Bindels, L.B. *et al.* Increased gut permeability in cancer cachexia: mechanisms and clinical relevance. *Oncotarget* **9**, 18224-18238 (2018).
126. Jacouton, E., Chain, F., Sokol, H., Langella, P. & Bermudez-Humaran, L.G. Probiotic Strain *Lactobacillus casei* BL23 Prevents Colitis-Associated Colorectal Cancer. *Front Immunol* **8**, 1553 (2017).
127. Gao, C. *et al.* Gut Microbe-Mediated Suppression of Inflammation-Associated Colon Carcinogenesis by Luminal Histamine Production. *Am J Pathol* **187**, 2323-2336 (2017).
128. Chung, E.J. *et al.* Combination of metformin and VSL#3 additively suppresses western-style diet induced colon cancer in mice. *Eur J Pharmacol* **794**, 1-7 (2017).
129. da Silva Duarte, V. *et al.* Chemoprevention of DMH-Induced Early Colon Carcinogenesis in Male BALB/c Mice by Administration of *Lactobacillus Paracasei* DTA81. *Microorganisms* **8** (2020).
130. Mendes, M.C.S. *et al.* Microbiota modification by probiotic supplementation reduces colitis associated colon cancer in mice. *World J Gastroenterol* **24**, 1995-2008 (2018).
131. Zhang, S.L. *et al.* Pectin supplement significantly enhanced the anti-PD-1 efficacy in tumor-bearing mice humanized with gut microbiota from patients with colorectal cancer. *Theranostics* **11**, 4155-4170 (2021).
132. Li, Y. *et al.* Prebiotic-Induced Anti-tumor Immunity Attenuates Tumor Growth. *Cell Rep* **30**, 1753-1766 e1756 (2020).
133. Zhu, L.Q. *et al.* Evodiamine inhibits high-fat diet-induced colitis-associated cancer in mice through regulating the gut microbiota. *J Integr Med* **19**, 56-65 (2021).
134. Terasaki, M. *et al.* Alteration of fecal microbiota by fucoxanthin results in prevention of colorectal cancer in AOM/DSS-treated mice. *Carcinogenesis* (2020).
135. Luo, L. *et al.* The effect of menthol supplement diet on colitis-induced colon tumorigenesis and intestinal microbiota. *Am J Transl Res* **13**, 38-56 (2021).
136. Liu, X., Wang, L., Jing, N., Jiang, G. & Liu, Z. Biostimulating Gut Microbiome with Bilberry Anthocyanin Combo to Enhance Anti-PD-L1 Efficiency against Murine Colon Cancer. *Microorganisms* **8** (2020).
137. Zhang, H., Hui, D., Li, Y., Xiong, G. & Fu, X. Canmei Formula Reduces Colitis-Associated Colorectal Carcinogenesis in Mice by Modulating the Composition of Gut Microbiota. *Front Oncol* **9**, 1149 (2019).

138. Khan, I. *et al.* Mushroom polysaccharides and jiaogulan saponins exert cancer preventive effects by shaping the gut microbiota and microenvironment in Apc(Min/+) mice. *Pharmacol Res* **148**, 104448 (2019).
139. Li, S., Fu, C., Zhao, Y. & He, J. Intervention with alpha-Ketoglutarate Ameliorates Colitis-Related Colorectal Carcinoma via Modulation of the Gut Microbiome. *Biomed Res Int* **2019**, 8020785 (2019).
140. Fernandez, J. *et al.* A Galacto-Oligosaccharides Preparation Derived From Lactulose Protects Against Colorectal Cancer Development in an Animal Model. *Front Microbiol* **9**, 2004 (2018).
141. Mudd, A.M. *et al.* Chemoprevention of Colorectal Cancer by Anthocyanidins and Mitigation of Metabolic Shifts Induced by Dysbiosis of the Gut Microbiome. *Cancer Prev Res (Phila)* **13**, 41-52 (2020).
142. Wu, J.C. *et al.* Polymethoxyflavones prevent benzo[a]pyrene/dextran sodium sulfate-induced colorectal carcinogenesis through modulating xenobiotic metabolism and ameliorate autophagic defect in ICR mice. *Int J Cancer* **142**, 1689-1701 (2018).
143. Wang, X. *et al.* Structural shift of gut microbiota during chemo-preventive effects of epigallocatechin gallate on colorectal carcinogenesis in mice. *World J Gastroenterol* **23**, 8128-8139 (2017).
144. Chou, Y.C. *et al.* Boswellia serrata resin extract alleviates azoxymethane (AOM)/dextran sodium sulfate (DSS)-induced colon tumorigenesis. *Mol Nutr Food Res* **61** (2017).
145. Fukuda, M. *et al.* Prebiotic treatment reduced preneoplastic lesions through the downregulation of toll like receptor 4 in a chemo-induced carcinogenic model. *J Clin Biochem Nutr* **49**, 57-61 (2011).
146. Oh, N.S., Lee, J.Y., Kim, Y.T., Kim, S.H. & Lee, J.H. Cancer-protective effect of a synbiotic combination between Lactobacillus gasseri 505 and a Cudrania tricuspidata leaf extract on colitis-associated colorectal cancer. *Gut Microbes* **12**, 1785803 (2020).
147. Rudd, D.A. *et al.* Mapping insoluble indole metabolites in the gastrointestinal environment of a murine colorectal cancer model using desorption/ionisation on porous silicon imaging. *Sci Rep* **9**, 12342 (2019).
148. Zhang, J. *et al.* A Functional Food Inhibits Azoxymethane/Dextran Sulfate Sodium-Induced Inflammatory Colorectal Cancer in Mice. *Onco Targets Ther* **14**, 1465-1477 (2021).

149. Piazzzi, G. *et al.* A Mediterranean Diet Mix Has Chemopreventive Effects in a Murine Model of Colorectal Cancer Modulating Apoptosis and the Gut Microbiota. *Front Oncol* **9**, 140 (2019).
150. Li, F. *et al.* Modulation of colon cancer by nutmeg. *J Proteome Res* **14**, 1937-1946 (2015).
151. Chen, L. *et al.* Chemoprevention of colorectal cancer by black raspberry anthocyanins involved the modulation of gut microbiota and SFRP2 demethylation. *Carcinogenesis* **39**, 471-481 (2018).
152. Fernandez, J., Garcia, L., Monte, J., Villar, C.J. & Lombo, F. Functional Anthocyanin-Rich Sausages Diminish Colorectal Cancer in an Animal Model and Reduce Pro-Inflammatory Bacteria in the Intestinal Microbiota. *Genes (Basel)* **9** (2018).
153. Wang, C.Z. *et al.* Role of intestinal microbiome in American ginseng-mediated colon cancer prevention in high fat diet-fed AOM/DSS mice [corrected]. *Clin Transl Oncol* **20**, 302-312 (2018).
154. Hu, Y. *et al.* Manipulation of the gut microbiota using resistant starch is associated with protection against colitis-associated colorectal cancer in rats. *Carcinogenesis* **37**, 366-375 (2016).
155. Yu, C. *et al.* American ginseng significantly reduced the progression of high-fat-diet-enhanced colon carcinogenesis in Apc (Min/+) mice. *J Ginseng Res* **39**, 230-237 (2015).
156. Gao, Z. *et al.* Probiotics modify human intestinal mucosa-associated microbiota in patients with colorectal cancer. *Mol Med Rep* **12**, 6119-6127 (2015).
157. Zhang, J.W. *et al.* Preoperative probiotics decrease postoperative infectious complications of colorectal cancer. *Am J Med Sci* **343**, 199-205 (2012).
158. Aisu, N. *et al.* Impact of perioperative probiotic treatment for surgical site infections in patients with colorectal cancer. *Exp Ther Med* **10**, 966-972 (2015).
159. Liu, Z. *et al.* Randomised clinical trial: the effects of perioperative probiotic treatment on barrier function and post-operative infectious complications in colorectal cancer surgery - a double-blind study. *Aliment Pharmacol Ther* **33**, 50-63 (2011).
160. Gianotti, L. *et al.* A randomized double-blind trial on perioperative administration of probiotics in colorectal cancer patients. *World J Gastroenterol* **16**, 167-175 (2010).

161. So, W.K.W. *et al.* Effects of a Rice Bran Dietary Intervention on the Composition of the Intestinal Microbiota of Adults with a High Risk of Colorectal Cancer: A Pilot Randomised-Controlled Trial. *Nutrients* **13** (2021).
162. Xie, X. *et al.* Effects of prebiotics on immunologic indicators and intestinal microbiota structure in perioperative colorectal cancer patients. *Nutrition* **61**, 132-142 (2019).
163. Polakowski, C.B., Kato, M., Preti, V.B., Schieferdecker, M.E.M. & Ligocki Campos, A.C. Impact of the preoperative use of synbiotics in colorectal cancer patients: A prospective, randomized, double-blind, placebo-controlled study. *Nutrition* **58**, 40-46 (2019).
164. Hibberd, A.A. *et al.* Intestinal microbiota is altered in patients with colon cancer and modified by probiotic intervention. *BMJ Open Gastroenterol* **4**, e000145 (2017).
165. Worthley, D.L. *et al.* A human, double-blind, placebo-controlled, crossover trial of prebiotic, probiotic, and synbiotic supplementation: effects on luminal, inflammatory, epigenetic, and epithelial biomarkers of colorectal cancer. *Am J Clin Nutr* **90**, 578-586 (2009).
166. Brown, D.G., Borresen, E.C., Brown, R.J. & Ryan, E.P. Heat-stabilised rice bran consumption by colorectal cancer survivors modulates stool metabolite profiles and metabolic networks: a randomised controlled trial. *Br J Nutr* **117**, 1244-1256 (2017).
167. Nunez-Sanchez, M.A. *et al.* Gene expression changes in colon tissues from colorectal cancer patients following the intake of an ellagitannin-containing pomegranate extract: a randomized clinical trial. *J Nutr Biochem* **42**, 126-133 (2017).
168. Nunez-Sanchez, M.A. *et al.* MicroRNAs expression in normal and malignant colon tissues as biomarkers of colorectal cancer and in response to pomegranate extracts consumption: Critical issues to discern between modulatory effects and potential artefacts. *Mol Nutr Food Res* **59**, 1973-1986 (2015).
169. Chen, C.C. *et al.* Oral inoculation of probiotics *Lactobacillus acidophilus* NCFM suppresses tumour growth both in segmental orthotopic colon cancer and extra-intestinal tissue. *Br J Nutr* **107**, 1623-1634 (2012).
170. Bellville, J.W., Swanson, G.D., Miyake, T. & Aqleh, K.A. Respiratory stimulation observed following ethanol ingestion. *West J Med* **124**, 423-425 (1976).

171. Shinohara, K., Ohashi, Y., Kawasumi, K., Terada, A. & Fujisawa, T. Effect of apple intake on fecal microbiota and metabolites in humans. *Anaerobe* **16**, 510-515 (2010).
172. Schwartz, S.E. *et al.* Sustained pectin ingestion: effect on gastric emptying and glucose tolerance in non-insulin-dependent diabetic patients. *Am J Clin Nutr* **48**, 1413-1417 (1988).
173. Annunziata, G. *et al.* Effect of Grape Pomace Polyphenols With or Without Pectin on TMAO Serum Levels Assessed by LC/MS-Based Assay: A Preliminary Clinical Study on Overweight/Obese Subjects. *Front Pharmacol* **10**, 575 (2019).
174. Glinsky, V.V. & Raz, A. Modified citrus pectin anti-metastatic properties: one bullet, multiple targets. *Carbohydr Res* **344**, 1788-1791 (2009).
175. Warburg, O. On the origin of cancer cells. *Science* **123**, 309-314 (1956).
176. Roediger, W.E. Role of anaerobic bacteria in the metabolic welfare of the colonic mucosa in man. *Gut* **21**, 793-798 (1980).
177. Khoo, H.E., Azlan, A., Tang, S.T. & Lim, S.M. Anthocyanidins and anthocyanins: colored pigments as food, pharmaceutical ingredients, and the potential health benefits. *Food Nutr Res* **61**, 1361779 (2017).
178. Reglero, C. & Reglero, G. Precision Nutrition and Cancer Relapse Prevention: A Systematic Literature Review. *Nutrients* **11** (2019).
179. Toledo, E. *et al.* Mediterranean Diet and Invasive Breast Cancer Risk Among Women at High Cardiovascular Risk in the PREDIMED Trial: A Randomized Clinical Trial. *JAMA Intern Med* **175**, 1752-1760 (2015).
180. Bamia, C. *et al.* Mediterranean diet and colorectal cancer risk: results from a European cohort. *Eur J Epidemiol* **28**, 317-328 (2013).
181. (2018).
182. Henderson, A.J. *et al.* Chemopreventive properties of dietary rice bran: current status and future prospects. *Adv Nutr* **3**, 643-653 (2012).

1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154

Figure legends

Figure 1. Flow chart of the search and selection process.

Figure 2. Distribution of the studies analyzed. a. Preclinical and clinical studies. **b** and **c.** Pre-, pro-, symbiotics and food in **b.** preclinical and **c.** clinical studies. **d.** Strains used as probiotics in preclinical studies. **e.** Prebiotics used in preclinical studies. **f.** Strains used as probiotics in clinical studies. **g.** Prebiotics used in clinical studies. * More than 1 pro- and prebiotic is used in some studies.

1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166

Table 1. Details of the search strategy used.

Field	Key words	Boolean Operator
Title	(Cancer* OR carcino* OR tumor* OR onco*)	AND
Title/Abstract	(colon* OR colorect*)	AND
Title/Abstract	(microbi* OR dysbi*)	AND
Title/Abstract	(chemo* OR immuno* OR car-t* OR radium* OR surgery* OR therapy*)	AND
Title/Abstract	(nutrition* OR diet* OR food* OR probio* OR prebio* OR symbio*)	NOT
Type of publication	(Biography OR Comment OR Duplicate publication OR Editorial OR Guideline OR Interview OR News OR Personal Narrative OR Published Erratum OR Retracted Publication OR Retraction of Publication OR Meta-Analysis OR Review OR Systematic Review)	

Using words/prefixes with an asterisk opens the search to more results based on plurals and varying suffixes.

1167

Table 2. Summary of the results of the articles analyzed.

Intervention	Type of cancer	Type of study	Impact on microbiota	Mechanism	Effect on the tumor	Ref	Scientific quality
Preclinical studies							
Probiotics							
Probiotic: <i>Akkermansia muciniphila</i>	Subcutaneous, syngeneic model of colon cancer	Female BALB/c mice injected subcutaneously with CT26 cells in combination or not with IL-2	↑ diversity and <i>Akkermansia</i> , <i>Allistipes</i> and <i>Lactobacillus</i>	↑ necrosis, ↓ proliferation, and ↑ tumor apoptosis. ↑ CTL and ↓ Treg recruitment. ↑ IFN-γ and IL-2 in the tumor and ↑ TNF-α in serum, together with ↓ TGF-β. The Amuc protein effects through TLR2.	↓ tumor volume and weight and ↑ survival time	115	4
Probiotic: VSL#3, is composed of: <i>Lactobacillus plantarum</i> , <i>Lactobacillus delbrueckii</i> subsp. <i>Bulgaricus</i> , <i>Lactobacillus paracasei</i> , <i>Lactobacillus acidophilus</i> , <i>Bifidobacterium breve</i> , <i>Bifidobacterium longum</i> , <i>Bifidobacterium infantis</i> and <i>Streptococcus salivarius</i> subsp. <i>Thermophilus</i> .	Gnobioc model of CRC associated with colitis and inflammation	129/SvEv deficient in IL10 treated with AOM	↓ <i>Verrucomicrobia</i> and ↑ <i>Proteobacteria</i> in the mucosa. ↓ <i>Bacteroidetes</i> in feces	↓ <i>Clostridium</i> in the mucous membrane	↑ penetrance, invasiveness, histological grade, tumor volume and number of tumors	116	4
Inactivated	CRC	Syngeneic	↓ <i>Proteobacteria</i>		↓ of tumors	76	3

probiotic: <i>Lactobacillus acidophilus</i> lysates (<i>Phylum Firmicutes</i>)	associated with colitis	male BALB/C treated with AOM and DSS		↑ Th1 lymphocytes (CD3 + CD4 + IFN-γ+) and M1 macrophages (CD11b + F4/80 + CD86+) in mesenteric lymphoid nodules. In combination with anti CTLA-4 ↑ TCD8+ effector memory (CD44 + CD8 + CD62L+), ↓ Treg (CD4 + CD25 + Foxp3+) and M2 macrophages (F4/80 + CD206+) in the tumor microenvironment. In addition to ↑ IL-2 in nodules and plasma and plasma IFN-γ	and trend towards ↑ survival in combination with CTLA-4 inhibitors		
Probiotic: <i>Lactobacillus rhamnosus</i> , <i>Lactobacillus plantarum</i> and <i>Pediococcus pentosaceus</i>	Early colon carcinogenesis	Male Wistar rats treated with DHM	↑ lactic acid bacteria	↓ weight loss, fecal pH, liver transaminases and fecal enzyme normalization	↓ aberrant intestinal crypts, especially with <i>L. rhamnosus</i> .	117	3
Probiotic: <i>Enterococcus faecalis</i> , <i>Bifidobacterium longum</i> and <i>Lactobacillus acidophilus</i> (Bifico)	CRC associated with colitis	C57BL/6 male WT treated with AOM and DSS	↑ <i>Lactobacillus</i> . ↓ <i>Desulfovibrio</i> , <i>Mucispirillum</i> , and <i>Odoribacter</i>	↑ colon size. ↓ IL-1β, IL-6, COX-2 and TNF-α, in addition to PGE2 production. ↓ Cxcl1, Cxcl2, Cxcl3, Cxcl5 and Ccl7.	↓ number and volume of tumors	118	3
Probiotic: <i>Akkermansia muciniphila</i> or its specific membrane protein Amuc_1100	CRC associated with colitis	C57BL/6 male WT treated with AOM and DSS	NR	↑ cytotoxic TCD8 in mesenteric nodules. ↑ TCD8-TNF-α+ and ↓ TCD8-PD-1+. ↓ TCD8 and Ki67 in colon.	↓ malignancy score, tumor formation and area	36	2
Probiotic: <i>Lactobacillus rhamnosus</i> M9	CRC associated with colitis	Male 6-C57BL/6NcrSlc mice treated with AOM and DSS	↑ diversity of the flora and ↑ of <i>Blautia</i> , <i>Akkermansia</i> , <i>Lactobacillus</i> and <i>Bifidobacterium</i>	↓ PCNA, pSTAT3, pAKT. ↓ CD68+ and CD163+ in subserosa.	↓ number and size of tumors. ↓ inflammatory and fibrotic score.	119	2
Probiotic: <i>Butyrivibrio pullicaecorum</i>	CRC associated with colitis	male BALB/cByJNarl treated with DHM	NR	↑ SLC5A8 and GPR43	↓ infiltration, anal bleeding, and CEA levels	120	2

Probiotic: <i>Lactobacillus plantarum</i> VD23 <i>Lactobacillus plantarum</i> C28 <i>Lactobacillus plantarum</i> MS18 <i>Lactobacillus salivarius</i> MS3 <i>Lactobacillus salivarius</i> MS6 <i>Lactobacillus salivarius</i> MS16	Early colon carcinogenesis	Male Wistar rats treated with DHM	NR	↑IL-18	↓ proliferation, volume and total number of tumors. ↑ colon length (non-significant results)	121	2
Probiotic: <i>Lactobacillus bulgaricus</i>	CRC associated with colitis	C57BL/6 male WT treated with AOM and DSS	NR	↓ IL-6, TNF-α, IL-17, IL-23 and IL-1β and ↑ IFN-γ in healthy and tumor intestinal tissue	↓ mean tumor volume and size. ↓ pathological severity score and ↑ colon length.	122	2
Probiotic: <i>Bifidobacterium</i> and <i>Lactobacillus</i>	Subcutaneous, syngeneic model of colon cancer	Female BALB/c mice injected subcutaneously with CT26 cells	↑ <i>Alloprebotella</i> , <i>Citrobacter</i> , <i>Prevotellaceae_UC</i> G-001, <i>Roseburia</i> , <i>Thalassospira</i> , <i>Erysipelatoclostridium</i> and <i>Lachnospiraceae_UC</i> CG-006. ↓ <i>Desulfovibrio</i> , <i>Anaerotruncus</i> , <i>Mucispirillum</i> , <i>Odoribacter</i> . and <i>Escherichia-Shigella</i> . ↑ <i>Bacteroides_chinchillae</i> and <i>Helicobacter_Ganmani</i> . ↓ <i>Escherichia coli</i> , <i>Lachnospiraceae_bacterium_COE1</i> , <i>Bacteroides vulgatus</i> and <i>Lachnospiraceae_bacterium_10-1</i>	NR	No change in tumor volume	123	2
Probiotic: <i>Lactobacillus casei</i> variety <i>rhannosus</i> (Lcr35)	Subcutaneous, syngeneic model of colon cancer	Male BALB/c mice injected subcutaneously with CT26 cells and treated with FOLFOX (5-fluorouracil, leucovorin, and Lactobacillus casei oxaliplatin).	↑ <i>Firmicutes</i> and <i>Bacteroidetes</i>	↓ severe diarrhea and mucositis. ↓ apoptotic cells, NF-κB and BAX. ↓ TNF-α and IL6.	Unchanged	124	2
Probiotic: <i>Faecalibacterium prausnitzii</i>	Subcutaneous model, colon cancer with cachexia	Male CD2F1 mice injected subcutaneously with C26 (I understand this refers to CT26, although it is	NR	Improvement of various parameters related to the intestinal barrier, but not significant	↓ tumor mass, not significant	125	2

		not specified).					
Probiotic: <i>Lactobacillus casei</i> BL23	CRC associated with colitis	C57BL/6 WT females treated with AOM and DSS	↑ <i>Lactobacillus zeae</i>	↓ Ki67. ↑ Bik	↓ tumor volume, number of tumors and histological malignancy	126	2
Probiotic: <i>Lactobacillus reuteri</i>	CRC associated with colitis	Hdc-/- BALB/c male BALB/c treated with AOM and DSS	NR	↓ tumor zone KC, IL22, IL6, TNF-α, IL-1-α. ↓ serum KC, IL22, IL6. ↓ CD11b+Gr-1+	↓ number of tumors	127	2
Probiotic: VSL#3, is composed of: <i>Lactobacillus plantarum</i> , <i>Lactobacillus delbrueckii subsp. Bulgaricus</i> , <i>Lactobacillus paracasei</i> , <i>Lactobacillus acidophilus</i> , <i>Bifidobacterium breve</i> , <i>Bifidobacterium longum</i> , <i>Bifidobacterium infantis</i> and <i>Streptococcus salivarius subsp. Thermophilus</i> .	CRC associated with colitis	Male BALB/c WT treated with AOM and DSS with Western diet and metformin.	NR	Metformin + VSL#3 = ↓ Ki67, F4/80	Metformin and metformin+VSL#3 = ↓ number of tumors	128	2
Probiotic: <i>Lactobacillus paracasei</i> DTA81	Early colon carcinogenesis	male BALB/c WT treated with DHM	↑ <i>Ruminiclostridium</i> and ↑ fecal acetic acid* and short-chain short acids* (*poorly performed comparator groups)	↓ IL-6, IL-17, and PCNA in colonic homogenates	Show no change	129	1
Probiotic: <i>Lactobacillus rhamnosus</i>	Early colon carcinogenesis	male BALB/c WT treated with DHM	↑ <i>Romboutsia</i> and <i>Turicibacter</i>	↑ IL4, IL-10 IL-6, IL-17, TNF-α, IFN-γ, c-MYC and P-53 in colonic homogenates	Show no change	129	1
Probiotics: <i>Lactobacillus acidophilus</i> , <i>Lactobacillus rhamnosus</i> and <i>Bifidobacterium bifidum</i>	CRC associated with colitis	C57BL/6 male WT treated with AOM and DSS	↑ <i>Lactobacillus</i> , <i>Bifidobacterium</i> , <i>Allobaculum</i> , <i>Clostridium XI</i> , and <i>Clostridium X III</i>	↓ colitis, ↓ circulating levels of RANTES and eotaxin; ↓ p-IKK and TNF-α with ↑ IL-10	↓ incidence and number of tumors	130	1
Prebiotics							
Prebiotic: Fructooligosaccharide/inulin	Gnobioc model of CRC associated with colitis	BALB/C mice (gender undefined) treated with AOM and DSS	NR	↑ histone 3 and ↓ HDAC activity. ↑ levels of Fas and active caspase 3. ↓ of p21, p27 and Ki-67.	↓ histological grade, volume and number of tumors	66	5

Prebiotic: Pectin	Subcutaneous, syngeneic model of colon cancer	Female C57BL/6 WT mice injected with MC38 cells. With humanized microbiota from healthy humans or patients, anti-PD-1 treatment.	↑ diversity, ↑ <i>Lactobacillaceae</i> , <i>Bifidobacteriaceae</i> , <i>Erysipelotrichaceae</i> , <i>Ruminococcaceae</i> , <i>Faecalibacterium</i> , and <i>Holdemania</i> . ↑ percentage <i>Firmicutes/Bacteroidetes</i> . ↑ acetate and butyrate.	↑ TCD4+, TCD8+ and TCD8+IFN γ + tumor infiltration	↓ tumor volume and size	131	4
Prebiotic: Mucin and inulin	Subcutaneously transplanted colon cancer (MC-38)	Syngeneic C57BL/6 mice	↑ <i>Clostridium</i> cluster XIVa	↑ of MHC class I and MHC class II expression in CD	Inulin causes ↓ tumor volume. No change in tumor volume with Mucin	132	4
Prebiotic: food extract: Evodiamine	CRC associated with colitis	C57BL/6 WT treated with HFD, AOM and DSS	↑ <i>Campylobacter</i> , <i>Bifidobacterium</i> and <i>Lactobacillus</i> . ↓ <i>E. faecalis</i> and <i>E. coli</i>	↓ IL-1, IL-6 and circulating TNF- α and in colonic tissue ↓ p-STAT3 and p-P65	↑ survival, ↓ tumor diameter and percentage weight/length of intestine. ↓ pathological score and proliferation, ↑ apoptosis.	133	3
Prebiotic: Food extract: Fucoxanthin	CRC associated with colitis	ICR WT male treated with AOM and DSS	↑ <i>Lachnospiraceae</i> and ↓ <i>Bacteroidales</i> and <i>Rikenellaceae</i>	↑ caspase 3	↓ size and number of colorectal adenomas and total tumors	134	3
Prebiotic: Food extract: Menthol	CRC associated with colitis	C57BL/6 male WT treated with AOM and DSS	↑ in butyrate-producing bacteria (<i>Allobaculum</i> , <i>Roseburia</i> and <i>Intestinimonas</i>) and fecal ↑ butyrate	↓ expression of β -catenin, Ki67, IL-6, TNF- α and MPO. ↓ infiltration of immune cells in the spleen, ↑ IL10 in the distal colon	↓ the number of 1-3 mm adenomas in a dose-dependent manner and damage scales.	135	2
Prebiotic: Chitosan and low molecular weight citrus pectins encapsulating bilberry anthocyanins	Subcutaneously transplanted colon cancer (MC-38)	Female C57BL/6 mice	↑ <i>Lachnospiraceae</i> (Clostridiales) and diversity	They suggest a ↑ of SCFA and TCD4+ and TCD8+ infiltration of the tumor, but are not significant.	↓ tumor volume in combination with PD-L1 inhibitors.	136	2
Prebiotic: Canmei formula	CRC associated with colitis	C57BL/6 WT females treated with AOM and DSS and HFD	↑ <i>Turicibacter</i> , <i>Bacteroides</i> , <i>Bacteroidaceae</i> , <i>Faecalibaculum</i> , <i>Erysipelatoclostridium</i> and <i>Staphylococcus</i>	↓ serum levels of IL-17 and NF- κ B	↓ tumor incidence	137	2
Prebiotic: Polysaccharides from <i>Ganoderma lucidum</i> and saponin from	Prevention of colon cancer of genetic origin	ApcMin+/- male mice	↑ bacteria producing SCFA and ↓ of sulfur-reducing bacteria	Shift from M1 to M2 macrophages. Reversal of expression, E-cadherin/N-	↓ number and size of polyp	138	2

<i>Gynostemma pentaphyllum</i>				cadherin ratio and ↓ oncogenic signaling. ↑ signaling of Gprotein coupled-receptors due to modulation of histone deacetylases, PYY and PPARγ expression.			
Prebiotic α-ketoglutarate	CRC associated with colitis	C57BL/6 WT females treated with AOM and DSS	↑ diversity. ↑ <i>Akkermansia</i> , <i>Butyricoccus</i> , <i>Clostridium</i> and <i>Ruminococcus</i> . ↓ <i>Escherichia</i> and <i>Enterococcus</i> .	↓ IL-6, IL-22, TNF-α, and IL-1β	↓ tumor frequency, size, and stage.	139	2
Prebiotic: Lactulose-derived galactooligosaccharides	CRC associated with colitis	Male <i>Rattus norvegicus</i> F344 rats treated with AOM and DSS	↑ propionate. ↑ diversity of flora, anti-inflammatory: <i>Bifidobacterium</i> , <i>Bacteroidaceae</i> , <i>Prevotellaceae</i> , <i>Acidaminococcaceae</i> , <i>Bifidobacteriaceae</i> , <i>Peptococcaceae</i> , <i>Oscillospiraceae</i> and ↓ of pro-inflammatory: <i>Lachnospiraceae</i> , <i>Eubacteriaceae</i> , <i>Acholeplasmataceae</i> and <i>Firmicutes/Bacteroidetes</i> %.	↑ cecum length	↓ number and volume of polyps	140	2
Prebiotic: Anthocyanidins wrapped or not in exosomes.	CRC associated with dysbiosis	ApcMin/+ Mice Inoculated with <i>Bacteroides fragilis</i>	NR	↑ in the liver, but not in the tumor, of phase I enzymes CYP1A1 and CYP1B1 and ↓ of phase II enzymes GSTM1 and SULT1	↓ number of tumors	141	1
Prebiotic: polymethoxyflavones	CRC associated with colitis	ICR WT males treated with benzo[a]pyrene and DSS	↑ <i>Sphingobacteriaceae</i> , <i>Gammaproteobacteria</i> , <i>Ruminococcaceae</i> and <i>Blautia</i> . ↓ <i>Bacilli</i> , <i>Parabacteroides</i> , <i>Lactobacillus Ruminis</i> (without statistical analysis)	↑ colon size. ↓ hepatic and renal toxicity. ↓ BaP-7,8-dihydrodiol-9,10-epoxide-DNA binding. ↓ genes for inflammation, metastasis and ↑ antioxidants.	↓ hyperplasias, adenoma and adenocarcinomas	142	1
Prebiotic: Epigallocatechin in gallate (EGCG)	CRC associated with colitis	FVB/N WT female treated with benzo[a]pyrene and DSS	↑ <i>Lactobacillus</i> , <i>Fusobacterium</i> , <i>Ruminococcus</i> , ↓ <i>Bacteroides</i> , <i>Anaerotruncus</i>	NR	↓ volume and number of tumors and pre-cancerous lesions	143	1
Prebiotic: <i>Boswellia</i>	CRC associated	ICR WT male treated with	↑ <i>Clostridiales spp</i> and ↓ <i>Bacteroidales</i>	↓ NF-κB, PCNA. ↑	↓ number of tumors	144	1

<i>serrata</i> resin extract	with colitis	AOM and DSS		colon length			
Prebiotic: Germinated barley extract	CRC associated with colitis	AOM-treated male Fisher 344 rats	NR	↓ TLR4 and COX2	↓ aberrant crypts	145	1
Symbiotics							
Symbiotic: <i>Clostridium butyricum</i> and prebiotic dextran	Subcutaneous and orthotopic model of colon cancer	CT26luc and ApcMin+/-	↑ <i>Muribaculaceae</i> , <i>Bacteroides</i> , <i>Mucispirillum</i> , <i>Alloprevotella</i> , <i>Lachnospiraceae</i> , and <i>Ruminococcaceae</i>	↑ iso-butyrate, butyrate, iso-valerate, propionate, valerate, iso-hexanoate, and hexanoate, ↓ dysbiosis	↑ effectiveness of capecitabine and diclofenac, ↓ tumor size of 74% /in combination with diclofenac, while diclofenac only 11%, abdominal invasion, ↑ apoptosis	96	5
Symbiotic: <i>Lactobacillus gasseri</i> 505 (LG) and leaf extract of <i>Cudrania tricuspidata</i> (CT) in fermented milk called FCT	CRC associated with colitis	C57BL/6 male WT treated with AOM and DSS	↑ <i>Lactobacillus</i> , <i>Bifidobacterium</i> , and <i>Akkermansia</i> associated with ↑ SCFA and ↓ <i>Staphylococcus</i>	↓ TNF-α, IFN-γ, IL-1β, IL-6, iNOS and COX-2. ↑ IL-4 and IL-10. ↑ MUC2, TFF3 and ZO-1. ↑ p53, p21, and Bax. ↓ Bcl-2 and Bcl-xL. ↓ β-Catenin, NF-κB and ↑ IkB-α	↓ incidence of tumors and dysplasia. ↑ colon length	146	3
Food							
Western diet, antibiotics and <i>Enterococcus faecalis</i> .	Model of CRC recurrence after surgery	Male BALB/c mice inoculated with CT26 after resection and anastomosis	Western diet = ↑ <i>E. faecalis</i> . Western diet associated with tumors ↑ <i>Proteus</i> , <i>Akkermansia</i> and <i>Trabulsiella</i> . ↓ in <i>Bacteroides</i> , <i>Roseburia</i> and <i>Ruminococcus</i> . Inoculation with <i>E. faecalis</i> = ↑ <i>Proteus</i> , <i>Enterococcus</i> , <i>Trabulsiella</i> <i>Akkermansia</i> and <i>Clostridium</i> and ↓ <i>Roseburia</i> .	NR	↑ tumor formation with Western diet and <i>E. faecalis</i> . ↑ distant metastasis with Western diet (not significant).	105	4
Food extract: selfish <i>Muricidae</i> : 1. marine extract (NE) 2. metabolite of the extract 6-bromoisatin (6-Br)	CRC genotoxic	AOM-treated male C57BL/6 WT	NR	6-Br = ↓ tumor proliferation. NE = ↓ proliferation and ↑ apoptosis in crypts.	6-Br = ↓ number of aberrant crypts and tumors. NE = ↓ number of tumors	147	4
Functional food: extract of 55 different foodstuffs mixed	CRC associated with colitis	AOM- and DSS-treated male BALB/c mice	↓ <i>Desulfovibrio</i> and <i>Ruminococcaceae</i> and ↑ from <i>Bacteroides</i> , <i>Ruminiclostridium_6</i> , <i>Holdemanella</i> , <i>Clostridium_sensu_</i>	Improved intestinal barrier and body weight maintenance. ↓ IL-6 and TNF-α. ↑ in	↓ number and size of tumors. ↑ survival	148	3

			<i>stricto_1</i> , <i>Allobaculum</i> and <i>Parasutterella</i> . ↓ of deuteroporphylin IX, citrulline, and diferuloylputrescine, and ↑-acid acetic, ascorbic, palmitic, quinic, ursocholic, branched amino acids, rosmaridiphenol and scillirosidine	the proportion of total T cells, CD3+CD4+, CD3+CD8+, and IFN-γ-producing CD4+ T cells.			
Diet: - low fat (LFD) - high fat (HFD) - a mixture of one of them with Mediterranean (MD). This being a mixture of ω-3 polyunsaturated fatty acids (PUFA), eicosapentanoic acid and polyphenols.	CRC genotoxic	A/J WT male treated with AOM	MD improves dysbiosis with an HFD, but worsens it with an LFD.	MD ↑ apoptosis in HFD, and especially in LFD. MD ↑ eicosapentanoic acid (EPA), especially in LFD.	MD ↓ incidence in HFD, but especially in LFD	149	3
Food extract: Nutmeg	Prevention of colon cancer of genetic origin	C57BL/6J WT and ApcMin+/- Male Mice	NR	↓ in urine of cresol sulphate and glucuronide, indoxyl sulphate and phenyl sulphate. ↓ colon and liver IL6. ↓ ALT and amylase. ↑ LPE 18:0, LPC 18:0, azelaic acid, sebacic acid, palmitolcarnitine and oleocarnitine. ↑ lipid metabolism gene expression in the colon and liver	↓ tumor volume and number of tumors	150	3
Food extract: Anthocyanin extract from blackberries.	CRC associated with colitis	C57BL/6 male WT treated with AOM and DSS	↑ <i>Eubacterium rectale</i> , <i>Faecalibacterium prausnitzii</i> and <i>Lactobacillus</i> . ↓ <i>Desulfovibrio</i> and <i>Enterococcus</i> spp.	↑ SFRP2 triggers the ↓ of DNMT1 and p-STAT3. In addition to ↓ IL-1β, IL-6, IL-10, COX-2 and TNF-α.	↓ number of tumors	151	2
Food extract: Anthocyanins extracted from strawberries and blackberries	CRC associated with colitis	Male <i>Rattus norvegicus</i> F344 rats treated with AOM and DSS	↓ <i>Bilophila wadsworthia</i>	Improved plasma reducing capacity	↓ number of tumors	152	2
Food extract: American	CRC associated	A/J male treated with	Improving dysbiosis	NR	↓ number of tumors and	153	2

Ginseng	with colitis	AOM and DSS and HFD			histological malignancy		
Food extract and prebiotic: Green tea extract (EGCG) and resistant starch	CRC associated with colitis	Male <i>Sprague-Dawley</i> rats treated with AOM and DSS	↑ <i>Parabacteroides</i> , <i>Barnesiella</i> , <i>Ruminococcus</i> , <i>Marvinbryantia</i> and <i>Bifidobacterium</i> . ↑ acetate, butyrate and propionate.	↑ GPR43. ↓ COX-2, NF-κB, TNF-α and IL-1β.	Resistant starch = ↓ number of tumors and adenocarcinoma formation	154	2
Food extract: American Ginseng	Prevention of colon cancer of genetic origin with a Western diet	Male <i>ApcMin</i> +/- mice on high-fat diet	NR	↓ IL-1α, IL-1β, IL6, TNF-α, G-CSF and GM-CSF	↓ tumor volume and number of tumors	155	2
Clinical studies							
Probiotics							
Probiotic: <i>Bifidobacterium longum</i> , <i>Lactobacillus acidophilus</i> and <i>Enterococcus faecalis</i>	Patients with CRC who have previously undergone radical surgery and who are about to undergo further surgery.	Randomized clinical trial Peri-surgery placebo (n=11), probiotic (n=11). Healthy control patients (n=11).	↑ <i>Enterococcus</i> and ↓ <i>Peptostreptococcus</i> , <i>Comamonas</i> , <i>Fusobacterium</i>	NR	NR	156	2
Probiotic: <i>Bifidobacterium longum</i> , <i>Lactobacillus acidophilus</i> and <i>Enterococcus faecalis</i>	CRC patients whose tumor could be removed by surgery and who are scheduled for surgery	Randomized clinical study Placebo (n=30), preoperative probiotic (n=30)	↑ preoperative percentage <i>Bifidobacterium/E. coli</i>	↑ IgG and sIgA in serum. ↓ endotoxins, D-lactic acid, IL-6 and CRP	↓ incidence of post-surgery infections	157	2
Probiotic: <i>Enterococcus faecalis</i> T110, <i>Clostridium butyricum</i> TO-A and <i>Bacillus mesentericus</i>	CRC patients whose tumor could be removed by surgery and who are scheduled for surgery	Clinical study Control (n=81), Probiotic (n=75)	NR	↓ incisional infections and reduced acute inflammation after surgery	NR	158	2
Probiotic: <i>Lactobacillus plantarum</i> , <i>Lactobacillus acidophilus</i> and <i>Bifidobacterium longum</i>	CRC patients whose tumor could be removed by surgery and who are scheduled for surgery	Randomized double-blind clinical study. Placebo (n=50), perioperative probiotic (n=50).	↑ diversity. ↑ <i>Bifidobacterium</i> and <i>Lactobacilli</i> . ↓ <i>Enterobacteriaceae</i> , <i>Pseudomonas</i> and <i>Candida</i>	↑ claudin1, JAM1 and occludin	NR	159	2
Probiotic: <i>Bifidobacterium longum</i> (BB536) and <i>Lactobacillus johnsonii</i> (La1)	CRC patients whose tumor could be removed	Randomized double-blind clinical study. Placebo (n=10), probiotic	<i>La1</i> = ↑ <i>La1</i> in mucosa and feces. ↓ <i>Enterobacteriaceae</i>	↑ CD3, CD4, CD8 and naive and memory lymphocytes. ↓ CD83-123 markers,	NR	160	2

	by surgery and who are scheduled for surgery	mixture at a dose of ¹⁰⁷ (n=11) and probiotic mixture at a dose of 109 (n=11). Peri-surgery treatment		CD83-HLADR, and CD83-11c			
Probiotic: <i>Lactobacillus acidophilus</i> , <i>Lactobacillus casei</i> , <i>Lactobacillus plantarum</i> , <i>Lactobacillus rhamnosus</i> , <i>Bifidobacterium lactis</i> , <i>Bifidobacterium bifidum</i> , <i>Bifidobacterium breve</i> , <i>Streptococcus thermophilus</i>	Stage III colorectal adenocarcinoma	Randomized clinical study of 78 patients with colorectal carcinoma with surgery	NR	NR	↓ number of complications associated with surgery	74	1
Prebiotics							
Prebiotic: food extract: evodiamine	CRC patients	CRC patients undergoing colonoscopies vs. healthy patients without any colorectal disease	↑ <i>E. faecalis</i> and <i>E. coli</i> ↓ <i>Campylobacter</i> , <i>Bifidobacterium</i> and <i>Lactobacillus</i> in CRC. ↑ Endotoxin and D-lactic acid in CRC.	↑ pSTAT3 in cancerous vs. peritumor biopsies	↓ proliferation, p-STAT3 and p-P65 after incubation with evodiamine in colon cancer cell lines HCT-116 and SW-480	133	3
Prebiotic: Rice bran	Patients at high risk of developing CRC	Randomized, double-blind, placebo-controlled trial (placebo = 19, treatment = 20). Placebo = rice powder, rice bran treatment. 30 g every 24h, for 24 weeks.	↑ <i>Firmicutes</i> and <i>Lactobacillus</i>	NR	NR	161	3
Prebiotic: fructo-oligosaccharides, xylooligosaccharides, polydextrose, and resistant dextrin	CRC patients requiring radical surgery (peri-surgery: pre- and post-surgery are analyzed).	Randomized double-blind clinical study with 140 CRC patients (control = 70; prebiotics = 70)	Pre-surgery = ↑ <i>Bifidobacterium</i> and <i>Enterococcus</i> , ↓ <i>Bacteroides</i> . Post-surgery = ↑ <i>Bifidobacterium</i> and <i>Escherichia-Shigella</i> , ↓ <i>Enterococcus</i>	Pre-surgery = ↑ IgG, IgM and transferrin. Post-surgery = ↑ IgG, TCD3+CD8+ (cytotoxic/suppressive T lymphocytes) and CD19+ (B lymphocytes).	NR	162	2
Symbiotics							
Symbiotic: Maltodextrin as placebo	CRC patients to be treated	Prospective, double-blind, randomized,	NR	↓ plasma IL6 and CRP levels. ↓ post-	3 deaths in the control group and	163	2

Microbiota in CRC

Madrigal-Matute & Banon-Escandell

and treatment with Simbioflora® (6g fructooligosaccharide, plus <i>Lactobacillus acidophilus</i> NCFM, <i>Lactobacillus rhamnosus</i> HN001, <i>Lactobacillus casei</i> LPC-37 and <i>Bifidobacterium lactis</i> HN019).	with surgery. Stages I-III	placebo-controlled clinical study (placebo; n = 36. Treatment; n = 37).		surgery infections, antibiotic use and hospital stay.	none in the treatment group.		
Symbiotic: <i>Bifidobacterium lactis</i> BI-04, <i>Lactobacillus acidophilus</i> NCFM and inulin	Stage I-III CRCs undergoing surgery	Clinical trial 21 healthy controls (colonoscopy), 15 CRC patients (8 non-intervention, 7 probiotic)	↑ <i>Firmicutes</i> , <i>Clostridiales</i> spp and <i>Faecalibacterium</i>	NR	NR	164	1
Pre-, pro- and symbiotic							
Prebiotic (resistant starch) probiotic (<i>Bifidobacterium lactis</i>) or combination of both (symbiotic)	Study in healthy subjects for CRC prevention	Randomized double-blind clinical study with crossover treatments. Healthy subjects, n = 17, for 16 weeks, 4 weeks per arm (placebo, prebiotic, probiotic and symbiotic).	Symbiotic = ↑ <i>Lachnospiraceae</i> spp.	No changes in fecal ammonium, SCFA or circulating cytokines. No epithelial changes.	NR	165	3
Food							
Food extract: Rice bran	CRC patients who have undergone surgery.	Randomized clinical study (control; n = 10. 30g/day rice bran; n = 9)	NR	↑ carbohydrate, lipid, amino acid, cofactor and vitamin metabolism	NR	166	2
Food extract: Pomegranate extract with high ellagitannin content	CRC patients whose tumor could be removed by surgery	Randomized clinical study. Control (n=10), PE (n=35)	NR	↓ CD44 and CTNNB1 in adjacent healthy tissue and tumor. ↓ CDKN1A, EGFR and TYMS in tumor.	NR	167	2
Food extract: Pomegranate extract with high ellagitannin content	CRC patients whose tumor could be removed by surgery and who are scheduled for surgery	Randomized double-blind clinical study Control (n=11), pomegranate extract #1 (n=19) and #2 (n=16)	NR	Modulation of miRNAs	NR	168	2

Microbiota in CRC

Madrigal-Matute & Banon-Escandell

Increase/activation has been indicated as (↑) and reduction/inhibition as (↓). NR - Not reported. Ref = References

1168