Elevated plasma levels of CXCL16 in severe COVID-19 patients

Sandra P. Smieszek¹, Vasilios M. Polymeropoulos¹, Christos M. Polymeropoulos¹, Bartłomiej P. Przychodzen¹, Gunther Birznieks¹, Mihael H. Polymeropoulos¹

¹ Vanda Pharmaceuticals Inc., Washington, DC, USA

Corresponding author

sandra.smieszek@vandapharma.com

Declarations:

Funding: no relevant funding

Conflicts of interest/Competing interests (include appropriate disclosures): authors are employees of Vanda Pharmaceuticals Inc.

Availability of data and material (data transparency): Data available upon request

Code availability (software application or custom code): not applicable

Authors' contributions (optional: please review the submission guidelines from the journal whether statements are mandatory): not applicable

Additional declarations for articles in life science journals that report the results of studies involving humans and/or animals: not applicable

Ethics approval (include appropriate approvals or waivers): Advarra IRB; Pro00043096

Consent to participate (include appropriate statements): All participants were consented prior to participation

Consent for publication (include appropriate statements): not applicable

Abstract

Genome-wide association studies have recently identified 3p21.31, with lead variant pointing to the *CXCR6* gene, as the strongest thus far reported susceptibility risk locus for severe manifestation of COVID-19. In order to determine its role, we measured plasma levels of Chemokine (C-X-C motif) ligand 16 (CXCL16) in the plasma of COVID-19 hospitalized patients. CXCL16 interacts with CXCR6 promoting chemotaxis or cell adhesion. The CXCR6/CXCL16 axis mediates homing of T cells to the lungs in disease and hyper-expression is associated with localised cellular injury. To characterize the CXCR6/CXCL16 axis in the pathogenesis of severe COVID-19, plasma concentrations of CXCL16 collected at baseline from 115 hospitalized COVID-19 patients participating in ODYSSEY COVID-19 clinical trial were assessed together with a set of controls. We report elevated levels of CXCL16 in a cohort of COVID-19 hospitalized patients. Specifically, we report significant elevation of CXCL16 plasma levels in association with severity of COVID-19 (as defined by WHO scale) (P -value<0.02). Our current study is the largest thus far study reporting CXCL16 levels in COVID-19 hospitalized patients (with whole-genome sequencing data available). The results further support the significant role of the CXCR6/CXCL16 axis in the immunopathogenesis of severe COVID-19 and warrants further studies to understand which patients would benefit most from targeted treatments.

Introduction

Genome-wide association studies have recently identified 3p21.31, encompassing the *CXCR6* gene, as the strongest thus far reported susceptibility region associated with a severe manifestation of COVID-19 (rs73064425, OR=2.14, P -value=4.77E-30)¹. Another lead severity variant rs10490770:T>C pointing to *CXCR6* was also reported with a P -value of 1.443E-73 (hospitalization) and 2.20E-61 (critical illness) by the COVID-19 Host Genetics Initiative². Nakanishi et al., reported that risk allele (rs10490770) carriers experienced an increased risk of COVID-19 related mortality (HR 1.5, 95% CI 1.3-1.8)³. Risk allele carriers had increased odds of several COVID-19 complications: severe respiratory failure (OR 2.0, 95% CI 1.6-2.6), venous thromboembolism (OR 1.7, 95% CI 1.2-2.4)³.

Chemokine (C-X-C motif) ligand 16 (CXCL16) is synthesized as a transmembrane molecule that is expressed as a cell surface-bound molecule, and as a soluble chemokine⁴. CXCL16 interacts with CXCR6 in leukocytes and other cells promoting chemotaxis or cell adhesion. It has been shown that inflammatory cytokines such as IFN γ and TNF α promote CXCL16 expression⁵. CXCL16 has been previously implicated in the pathogenesis of lung injury upon which it is released and functions as a chemoattractant for CXCR6+ T, natural killer (NK), B, and dendritic cells⁶. CXCL16 levels were elevated in the serum of acute lung injury patients in comparison to controls⁶. CXCR6 is expressed on T cells not only in T helper type 1 (Th1) inflammation but also in Th2 inflammation, where it is increased after allergen challenge. It was previously shown that reduction in CD8+ T-lymphocytes occurred between C57BL/6 and CXCR6KO mice⁷. CXCR6 is a major coreceptor of HIV type 2 variants⁸. The CXCR6/CXCL16 axis mediates homing of T cells to the lungs in disease and when hyper expressed is associated with localized cellular injury⁷. This CXCR6/CXCL16 axis is involved in lung pathology also associated with other infections, including influenza. Antagonism was reported to result in reduced inflammation⁶. In a recent report authors have shown that knockdown of CXCR6 reversed the effect of CXCL16 and so did the treatment with the p38 inhibitor SB203580 abolished the effects of CXCL16⁶. Another recent study has shown that the absence of CXCR6 significantly decreases airway CD8 resident memory T cells due to altered trafficking of CXCR6^{-/-} cells within the lung⁹. Moreover mice lacking CXCL16 also had decreased CD8 resident memory T cells in the airways and ultimately blocking CXCL16 resulted in the inhibition of the steady-state maintenance of airway resident memory T cells⁹.

Methods

Participants

ODYSSEY is a double-blinded Phase 3 study with a planned randomization of a total of 300 hospitalized severely ill COVID-19 patients (Clinicaltrials.gov: NCT04326426). Inclusion criteria for the study comprised of: 1. Adults aged 18-90; 2. confirmed laboratory COVID-19 infection; 3. confirmed pneumonia by chest radiograph or computed tomography; 4. fever defined as temperature ≥ 36.6 °C armpit, ≥ 37.2 °C oral, or ≥ 37.8 °C rectal since admission or the use of antipyretics; 5. $\text{PaO}_2 / \text{FiO}_2 \leq 300$; 6. In patient hospitalization. Patients were to be followed for up to 28 days to record clinical outcomes. Patients' clinical progress was recorded on a 7 point clinical status ordinal scale defined as follows: 1- Death; 2- Hospitalized on mechanical ventilation or ECMO; 3- Hospitalized on non invasive ventilation or high-flow oxygen supplementation; 4- Hospitalized requiring supplemental oxygen; 5- Hospitalized not requiring supplemental oxygen, requiring continued medical care; 6- Hospitalized not requiring supplemental oxygen, not requiring continued medical care; 7- Not hospitalized. Main Exclusion Criteria included: 1. Inability to provide informed consent or to have an authorized relative or designated person provide informed consent, or to comply with the protocol requirements; 2. Known allergy to tradipitant or other neurokinin-1 antagonists; 3. Pregnancy; 4. Uncontrolled HIV, HBV, or HCV infection; 5. Other uncontrolled medically significant diseases; 6. Enrollment in another clinical trial of an investigational therapy; 7. Alanine aminotransferase $> 5 \times$ Upper Limit of Normal or Creatinine clearance $< 50 \text{ mL} / 1.73 \text{ m}^2 / \text{min}$; 8. Requiring mechanical ventilation for > 72 hours.

CXCL16 measurement

To characterize the CXCR6/CXCL16 axis in the pathogenesis of severe COVID-19, plasma concentrations of CXCL16 from 115 hospitalized COVID-19 patients participating in ODYSSEY COVID-19 clinical trial ((NCT04326426)) and additionally samples from 37 controls were assessed. CXCL16 levels in plasma were determined with an enzyme-linked immunosorbent assay (ELISA) assay (Thermofisher ID: # EHCXCL16) performed on plasma collected at baseline in cases and in a set of healthy controls (assay range 2.74-2000 pg/mL).

Genetic Analysis

Incoming nucleic acid samples are quantified using fluorescent-based assays (PicoGreen) to accurately determine whether sufficient material is available for library preparation and sequencing. DNA sample size distributions are profiled by a Fragment Analyzer (Advanced Analytics) or BioAnalyzer (Agilent Technologies), to assess sample quality and integrity. HumanCoreExome 24v1.3 array was performed on all human DNA samples sequenced. Whole genome sequencing (WGS) libraries were prepared using the Truseq DNA PCR-free Library Preparation Kit. Whole Genome data were processed on NYGC automated pipeline. Paired-end 150 bp reads were aligned to the GRCh37 human reference (BWA-MEM v0.7.8) and processed with GATK best-practices workflow (GATK v3.4.0). The mean coverage was 35.8, it reflects the samples average. All high quality variants obtained from GATK were annotated for functional effects (intronic, intergenic, splicing, nonsynonymous, stopgain and frameshifts) based on RefSeq transcripts using Annovar³¹. Additionally, Annovar was used to match general population frequencies from public databases (Exac, gnomAD, ESP6500, 1000g) and to

prioritize rare, loss-of-function variants. Linear models adjusted for PC, age and sex were conducted in PLINK.

Results

We report elevated levels of CXCL16 in a set of COVID-19 hospitalized patients. Baseline demographics of the studied cohort are presented in **Table 1**. Clinical characteristics are provided in **Table 2**. Specifically, we report significantly elevated levels of CXCL16 in a subset of COVID-19 severe (as defined by WHO scale) hospitalized patients (P -value <0.02) displayed on **Figure 1**. The most significant difference reported in plasma CXCL16 between most severe (WHO- 2) COVID-19 hospitalized cases and less severe patients (P -value <0.02) as well as most severe and controls. We furthermore report a larger variance within the cases, signifying further necessity of stratification based on clinical characteristics (ANCOVA p -value – 0.001, covariates: age, sex, BMI). The effect is not driven by age nor BMI, or sex (Figure 2 – Supplemental Material). We further investigated the correlation between IFN- γ , TNF- α . We report lack thereof at baseline (Figure 3 – Supplemental Material). We were able to replicate the lead variant association (rs10490770) in our cohort of hospitalized patients (cases n 140 control n

1900) with confirmed COVID-19 infection (OR 1.9, CI 1.2-3.5, P -value <0.002). The carriers of this variant were at higher risk, of being in the hospitalized severe group (as defined by the WHO scale), finding which is consistent across multiple COVID-19 GWAS results reported thus far. Moreover, we inspected the association between the lead variant status and directly CXCL16 plasma levels. This association between lead coding variant carrier status and CXCL16 levels did not attain statistical significance (result that is trending), a finding that warrants further replication in a larger sample size. FYCO1 variant within the extended haplotype did attain statistical significance (P -value <0.0001) and the result is displayed in Supplementary Figure 4.

Discussion

Previously it was demonstrated that circulating CD8+CXCR6+ T cells were significantly elevated with advanced age, yet virtually absent in patients with severe COVID-19. In a smaller sample set, plasma levels of CXCL16 were significantly upregulated in severe COVID-19 patients compared to mild COVID-19 patients¹⁰. Differential expression of *CXCR6* and *CXCL16* mRNA was observed in severe COVID-19 compared to mild disease and significant functional polymorphisms in *CXCR6* are linked to viral control¹.

Furthermore, based on lung tissue autopsy data, the top differentially expressed genes from GWAS hits included *FYCO1*, specifically in AT2 cells, ciliated cells, and club cells¹¹. *FYCO1* coding variant (rs33910087) is an eQTL for *CXCR6*, whereas *CXCR6* was differentially expressed in lung CD8 T cells in COVID-19 infected lung tissue¹¹.

Prior studies targeting the axis (both knockout and pharmacological blocking) support the hypothesis that antagonism may constitute potential therapies^{6,9}. It may be the case that the treatment of the hyper-expression of the CXCL16/CXCR6 axis hence reduction of the severely heightened homing signal may elicit such therapeutic effects, for example in a cytokine storm. Besides the potential implications in lung injury, both the CXCR6 and its ligand plays an important role in mediating a proinflammatory microenvironment for tumor growth in hepatocellular carcinoma¹². Here targeting the CXCR6/CXCL16 axis is needed to elucidate the mechanism whereby neutrophils are affected in the tumor environment¹². Authors propose one such orally bioavailable compound - antagonist of the CXCR6 receptor shown to decrease growth in a mouse xenograft model of hepatocellular carcinoma¹². Further studies on antagonism of the axis are needed to explain the timing of such antagonism treatment that would result in optimal outcomes. Our current study, is the largest thus far reporting on the plasma levels of CXCL16 in association with severity in COVID-19 patients. The results support the significant role of the CXCR6/CXCL16 axis in the immunopathogenesis of severe COVID-19. Further studies warrant analysis of outcomes based on the carrier status for this locus and on the levels of CXCL16.

203

204

205

206 References

207

- 208 1. Pairo-Castineira E, Clohisey S, Klaric L, et al. Genetic mechanisms of critical illness in Covid-19.
209 *medRxiv*. 2020;17(8):25. <https://doi.org/10.1101/2020.09.24.20200048>.
- 210 2. Initiative C-19 HG. Mapping the human genetic architecture of COVID-19. *Nature*. 2021.
211 doi:10.1038/s41586-021-03767-x
- 212 3. Nakanishi T, Pigazzini S, Degenhardt F, et al. Age-dependent impact of the major common genetic
213 risk factor for COVID-19 on severity and mortality. *medRxiv*. 2021.
214 doi:10.1101/2021.03.07.21252875
- 215 4. Collado A, Marques P, Escudero P, et al. Functional role of endothelial CXCL16/CXCR6-platelet-
216 leucocyte axis in angiotensin II-associated metabolic disorders. *Cardiovascular Research*.
217 2018;114(13):1764-1775. doi:10.1093/cvr/cvy135
- 218 5. Abel S, Hundhausen C, Mentlein R, et al. The Transmembrane CXC-Chemokine Ligand 16 Is
219 Induced by IFN- γ and TNF- α and Shed by the Activity of the Disintegrin-Like Metalloproteinase
220 ADAM10. *The Journal of Immunology*. 2004;172(10):6362 LP-6372.
221 doi:10.4049/jimmunol.172.10.6362
- 222 6. Tu G-W, Ju M-J, Zheng Y-J, et al. CXCL16/CXCR6 is involved in LPS-induced acute lung injury via
223 P38 signalling. *Journal of cellular and molecular medicine*. 2019;23(8):5380-5389.
224 doi:10.1111/jcmm.14419
- 225 7. Ashhurst AS, Flórido M, Lin LCW, et al. CXCR6-Deficiency Improves the Control of Pulmonary
226 Mycobacterium tuberculosis and Influenza Infection Independent of T-Lymphocyte Recruitment
227 to the Lungs. *Frontiers in Immunology*. 2019;10:339. doi:10.3389/fimmu.2019.00339
- 228 8. Blaak H, Boers PHM, Gruters RA, Schuitemaker H, van der Ende ME, Osterhaus ADME. CCR5,
229 GPR15, and CXCR6 are major coreceptors of human immunodeficiency virus type 2 variants
230 isolated from individuals with and without plasma viremia. *Journal of virology*. 2005;79(3):1686-
231 1700. doi:10.1128/JVI.79.3.1686-1700.2005
- 232 9. Wein AN, McMaster SR, Takamura S, et al. CXCR6 regulates localization of tissue-resident
233 memory CD8 T cells to the airways. *Journal of Experimental Medicine*. 2019;216(12):2748-2762.
234 doi:10.1084/jem.20181308
- 235 10. Payne DJ, Dalal S, Leach R, et al. The CXCR6/CXCL16 axis links inflamm-aging to disease severity in
236 COVID-19 patients. *bioRxiv*. January 2021:2021.01.25.428125. doi:10.1101/2021.01.25.428125
- 237 11. Delorey TM, Ziegler CGK, Heimberg G, et al. COVID-19 tissue atlases reveal SARS-CoV-2 pathology
238 and cellular targets. *Nature*. 2021;595(7865):107-113. doi:10.1038/s41586-021-03570-8

12. Peddibhotla S, Hershberger PM, Jason Kirby R, et al. Discovery of small molecule antagonists of chemokine receptor CXCR6 that arrest tumor growth in SK-HEP-1 mouse xenografts as a model of hepatocellular carcinoma. *Bioorganic & medicinal chemistry letters*. 2020;30(4):126899. doi:10.1016/j.bmcl.2019.126899

Figure 1 Baseline plasma CXCL16 levels in association with severity of COVID-19 manifestation and case control analysis. Severe (as defined by WHO scale) cases had significantly higher levels of CXCL16 – effect that is statistically significant.

257

258

259

260

261

262

263

264

Supplemental Figure 2 Scatter plot between CXCL16 and IFN γ and TNF α (baseline samples collected from COVID-19 hospitalized patients with severity determined with WHO scale)

Supplemental Figure 3 Correlation between age and CXCL16 plasma levels (collected at baseline from COVID-19 hospitalized patients)

287
288
289
290
291
292
293
294
295
296

Supplemental Figure 4 Association between genotype status and CXCL16 plasma levels (collected at baseline from COVID-19 hospitalized patients)

FYCO1 3UTR rs60431156 vs CXCL16

Table 1

Characteristic	Total (N=141)
Age (years)	64 (54 - 72)
Sex, n (%)	
Male	91 (64.5)
Female	50 (35.5)
Race, n (%)	
Asian	2 (1.4)
Black or African American	35 (24.8)
Native Hawaiian or Other Pacific Islander	1 (0.7)
White	73 (51.8)
Other	30 (21.3)
Any Comorbidities, n (%)	135 (95.7)
Hypertension	80 (56.7)
Diabetes	56 (39.7)
Coronary Heart Disease	10 (7.1)
Asthma	16 (11.3)
7 Point Ordinal Scale at Baseline, n (%)	
2 - Hospitalized on mechanical ventilation or ECMO	8 (5.7)
3 - Hospitalized on non-invasive ventilation or high-flow oxygen supplement	69 (48.9)
4 - Hospitalized requiring supplemental oxygen	61 (43.3)
5 - Hospitalized not requiring supplemental oxygen, requiring continued medical care	3 (2.1)

Table 2

Characteristic	Total (N=141)
Time from Hospitalization to Starting Study Treatment, Days	4 (2 - 7)
Early (<=10 Days from Hospitalization)	127 (90.1)
Late (>10 Days from Hospitalization)	14 (9.9)
Highest Oxygen Therapy Support	
Room Air	3 (2.1)
Nasal Cannula (NC)	46 (32.6)
Non Rebreather (NRB)	15 (10.6)
High Flow Nasal Cannula (HFNC)	52 (36.9)
CPAP Mask	7 (5.0)
BiPAP Mask	10 (7.1)
Mechanical Ventilation	8 (5.7)
Concomitant Medications, n (%) *	39 (27.7)
Azithromycin	19 (13.5)
Tocilizumab	4 (2.8)
Hydroxychloroquine	12 (8.5)
Steroids#	21 (14.9)