1 **Live wildlife trade in markets – a scoping review to inform risk assessment**

2 **of emerging infectious diseases**

3

4 V.J. Brookes^{1,2}, O. Wismandanu³, E. Sudarnika⁴, J.A. Roby^{2,5}, L. Hayes^{1,2}, M.P. Ward⁶, C.
5 Basri⁴, H. Wibawa⁷, J. Davis⁸, D. Indrawan⁹, J. Manyweathers^{1,2}, W.S. Nugroho¹⁰, S.
6 Windria¹¹, M. Hernandez-Jover^{1,2}

7

8 ¹ School of Agriculture, Environmental and Veterinary Sciences, Faculty of Science and
9 Health, Charles Sturt University, Wagga Wagga, NSW 2678, Australia

10 ² Graham Centre for Agricultural Innovation, (NSW Dept. of Primary Industries and Charles
11 Sturt University), Charles Sturt University, Wagga Wagga, NSW 2678, Australia

12 ³ Veterinary Medicine Study Program, Faculty of Medicine, Padjadjaran University,
13 Indonesia

14 ⁴ Faculty of Veterinary Medicine, IPB (Institut Pertanian Bogor) University, Indonesia

15 ⁵ School of Dentistry and Medical Sciences, Faculty of Science and Health, Charles Sturt
16 University, Wagga Wagga, NSW 2678, Australia

17 ⁶ Sydney School of Veterinary Science, The University of Sydney, Camden, 2570, Australia

18 ⁷ Disease Investigation Centre Wates, Directorate General of Livestock and Animal Health
19 Services, Ministry of Agriculture of Indonesia

20 ⁸ Australian Department of Agriculture, Water and the Environment, Canberra, Australia

21 ⁹ School of Business, IPB University, Indonesia

22 ¹⁰ Faculty of Veterinary Medicine, Universitas Gadjah Mada, Indonesia

23 ¹¹ Department of Biomedical Sciences, Division of Microbiology, Veterinary Medicine Study

24 Program, Faculty of Medicine, Padjadjaran University, Indonesia

25

26 **Corresponding author**

27 Victoria J. Brookes, victoria.brookes@sydney.edu.au. Sydney School of Veterinary Science,

28 Faculty of Science, Camperdown, NSW, 2006, Australia.

29

30

31

32

33 **Abstract**

34 Wet markets are important for food security in many regions worldwide but have come under
 35 scrutiny due to their potential role in the emergence of infectious diseases. The sale of live
 36 wildlife has been highlighted as a particular risk, and the World Health Organisation has
 37 called for the banning of live, wild-caught mammalian species in markets unless risk
 38 assessment and effective regulations are in place. Following PRISMA guidelines, we
 39 conducted a global scoping review of peer-reviewed information about the sale of live,
 40 terrestrial wildlife in markets that are likely to sell fresh food, and collated data about the
 41 characteristics of such markets, activities involving live wildlife, the species sold, their
 42 purpose, and animal, human, and environmental health risks that were identified. Of the 59
 43 peer-reviewed records within scope, only 25% (n = 14) focussed on disease risks; the rest
 44 focused on the impact of wildlife sale on conservation. Although there were some global
 45 patterns (for example, the types of markets and purpose of sale of wildlife), there was wide
 46 diversity and huge epistemic uncertainty in all aspects associated with live, terrestrial wildlife
 47 sale in markets such that the feasibility of accurate assessment of the risk of emerging
 48 infectious disease associated with live wildlife trade in markets is limited. Given the value of
 49 both wet markets and wildlife trade and the need to support food affordability and
 50 accessibility, conservation, public health, and the social and economic aspects of livelihoods
 51 of often vulnerable people, there are major information gaps that need to be addressed to
 52 develop evidence-based policy in this environment. This review identifies these gaps and
 53 provides a foundation from which information for risk assessments can be collected.

54

55 **Key words**

56 Live wildlife trade, wet markets, emerging infectious disease, EID

57 1. Introduction

58 The COVID-19 pandemic – together with the emergence of SARS, Ebola virus disease and
 59 avian influenza H5N1 – has focused global attention on the potential role played by animal
 60 markets in the emergence of novel pathogens in human populations. Following the
 61 epidemiologic association between early cases of COVID-19 and the Huanan seafood market
 62 in Wuhan, Hubei, China (Chen et al., 2020; Huang et al., 2020), the potential for spill-over of
 63 potentially pathogenic microbes from live wildlife traded at fresh food markets, or ‘wet
 64 markets,’ was again highlighted as a risk for emergence of infectious diseases. Reactions
 65 have ranged from calls to improve risk assessment at wet markets to banning the sale of live
 66 wildlife in markets (World Health Organisation, 2021; Xiao et al., 2021). Although food-
 67 borne disease risks due to inadequate sanitation and hygiene practices at wet markets are
 68 well-documented (World Health Organization, 2006), the risk of pathogen spillover
 69 associated with live wildlife sale in wet markets and the circumstances in which this could
 70 lead to emergence of a new infectious disease are not well understood (Naguib et al., 2021;
 71 Pruvot et al., 2019).

72 Wet markets are common in many parts of the world, selling fresh food such as meat,
 73 seafood, vegetables and fruits, and sometimes live animals – including wildlife – which
 74 might be slaughtered and butchered at the market (Nadimpalli and Pickering, 2020; Sigal,
 75 2020; Zhong et al., 2020). Such markets are found in China, Southeast Asia, Africa and South
 76 America (The Law Library of Congress, 2020). Food markets, including wet markets, serve
 77 local communities as social and commercial centres, and are especially important in urban
 78 settings in low income regions where they are important for food security by providing easy
 79 access to affordable food (World Health Organization, 2006).

80 Most emerging infectious diseases of humans have an animal origin (Jones et al., 2008;
81 Taylor et al., 2001). Non-specific drivers of disease emergence – such as climate change and
82 ecosystem degradation – are well-described, but exact mechanisms at interfaces between
83 wildlife and other species such as livestock and people are generally poorly defined or
84 unknown (Magouras et al., 2020; Plowright et al., 2017; Woolhouse et al., 2005). The
85 probability of emergence of a pathogen depends on interacting factors associated with
86 microorganism type, the characteristics and circumstances of the reservoir and potential new
87 hosts, as well as opportunities for spill-over between these hosts by direct or indirect routes.
88 These factors could facilitate or impede evolutionary processes and events that need to occur
89 to enable cross-species transmission, and pathogenicity and transmissibility between
90 individuals in the new host population (Plowright et al., 2017).

91 Wet markets provide an interface between live wildlife, domestic animals and people, and
92 have previously been associated with disease emergence events (Woo et al., 2006). An
93 outbreak of highly-pathogenic avian influenza (H5N1) in people in Hong Kong in 1997 was
94 in part disseminated via live-bird markets (Tam, 2002). Crowding and mixing of bird species
95 and unhygienic slaughter in markets, as well as proximity of markets to residential areas,
96 were considered to be contributing factors. Similarly, live-poultry markets were considered a
97 risk factor during an outbreak of H7N9 in people in Zhejiang, China (He et al., 2015). In the
98 outbreak of severe acute respiratory syndrome (SARS) which emerged in late 2002 in
99 Guangzhou Province, China, markets were again suggested as a risk factor because early
100 community cases were more likely to have lived within walking distance of wet markets than
101 community cases later in the outbreak (Xu et al., 2004). Supporting this, in markets in
102 Guangzhou, there was evidence of greater exposure to SARS-CoV among live animal traders
103 (seroprevalence 13.0%, N = 508), especially palm civet traders (seroprevalence 72%, N =
104 22), compared to control groups (seroprevalence range 1.2—2.9%, N = 284) (Yu et al., 2003).

SARS-CoV-like virus was also isolated from live Himalayan palm civets (*Paguma larvata*) at a market in Shenzhen municipality, China, and evidence of infection with SARS-CoV-like virus was also found in a raccoon dog (*Nyctereutes procyonoides*), a Chinese ferret-badger (*Melogale moschata*), and people working at the market (Guan et al., 2003). Bats are considered to be natural reservoirs of many *Coronaviridae* and SARS-CoV-2 is closely related to a bat coronavirus (Andersen et al., 2020; Zhou et al., 2020). Although a route between bats and people has not been identified, species that are sometimes sold live in wet markets – for example, Pholidota (pangolin), one of the most trafficked animals in the world (Aisher, 2016) – have been suggested as intermediate hosts of SARS-CoV-2, though evidence for the role of any specific intermediate species is incomplete and consequently various theories remain controversial (Lam et al., 2020; Zhang et al., 2020). Additionally, evidence of the location or time during which transmission to humans could have occurred is lacking.

Whether live wildlife traded at markets provide opportunity for infectious disease emergence and the circumstances in which this might occur is of critical importance to developing and implementing policy to prevent pandemics. Risk assessments should form the basis of such policy by informing evidenced-based regulatory and surveillance frameworks that mitigate identified risks associated with the sale of live wildlife in food markets; such tools are already being developed (Wikramanayake et al., 2021). Detailed understanding of the hazards associated with live wildlife sold in markets is needed, not only to conduct meaningful risk assessments, but also to understand the policy environment of live wildlife trade in markets so that risk mitigation strategies can be implemented whilst supporting the essential socio-economic roles of markets.

In this review, our objective was to collate and describe peer-reviewed evidence of live, terrestrial wildlife sold in wet markets, including associated factors such as market type and

130 documented risks to animal, human and environmental health. We aimed to provide a
131 baseline of current knowledge – and identify gaps in knowledge – to support risk assessment
132 and policy development regarding disease emergence associated with the sale of live,
133 terrestrial wildlife at markets.

2. Materials and Methods

2.1 Protocol

This scoping review was conducted according to the Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) guidelines (Tricco et al., 2018). An overview of this methodology is described at <http://www.prisma-statement.org/Extensions/ScopingReviews>, accessed 18 October 2020.

The review protocol included three levels: Level 1, screening on title and abstract; Level 2, screening on full record; Level 3, data extraction. The web-based review platform Sysrev[®] (Insilica LLC, 2020; www.sysrev.com, accessed 18 October 2020) was used for Levels 1 and 2, and a spreadsheet in Excel[®] (Microsoft 2020, www.microsoft.com, accessed 18 October 2020) was used for Level 3.

We use the term ‘record’ to describe any bibliographic citation captured in the searches. We use the term ‘study’ as an individual investigation of wildlife sale in a market. Whilst most records comprised only one study, if a record contained multiple relevant studies (for example, theses, books, conference proceedings and reports), each relevant study was evaluated separately within that record.

2.2 Eligibility criteria

Types of records that were eligible included theses, dissertations, and peer-reviewed literature for which the full text was available. Records were eligible if they were published in English, from 1980 onwards, and from any country, and contained primary research of interest about the sale of live terrestrial wildlife at markets likely to sell food.

Markets of interest were collections of stalls which were identified in the record as food, wet or fish markets, street markets (when the meaning was not in the context of covert

sale), medicinal markets, wildlife markets or open fairs. Pet shops, restaurants, bars or cafés were not included.

Terrestrial wildlife of interest were animals from classes Amphibia, Reptilia, Aves and Mammalia (see Supplementary Material for lists of orders in each class) which were presented live for sale at the market, and were not feral, companion or livestock animals. Vermin that lived in and around the market were not within scope unless presented live for sale at the market. Meat or other products from animals that were slaughtered prior to presentation for sale at the market were also not within scope. Live wildlife could be sold at the market for any purpose; for example, pets, medicine and consumption, including wildlife that were slaughtered at the market for these purposes.

Records which described the chain of events around wildlife trade (for example, hunting and sale of bushmeat) but did not include information about the sale of live wildlife at specific markets were excluded.

2.3 Information sources and search strategy

The literature search was conducted in July 2020, using the following combination of search terms:

- (i) market* OR sale,
- (ii) AND: wildlife OR "wild life" OR "wild-life" OR "wild animal" OR "wild bird",
- (iii) AND: food OR medicine,
- (iv) timeframe: 1980—2020, language: “English.”

Three electronic databases were searched: Web of Science, PubMed, and SCOPUS. Searches for theses and dissertations were conducted in ProQuest Thesis and Dissertations Global Database, and EBSCO Open Dissertations. A literature search was conducted via the Google

Scholar search engine, in which the first 100 results were screened. In addition, a search was conducted in Google Scholar with the addition of the term ‘TRAFFIC bulletin’ to identify relevant records from this journal, which was not available through the electronic databases. All records were exported into the citation manager software Endnote (<https://endnote.com/>, accessed 18 October 2020), and duplicates were removed. Records were then uploaded to Sysrev© for Level 1 agreement tests and screening.

2.4 Selection of relevant records – Levels 1 and 2

Agreement tests were conducted prior to screening at Levels 1 and 2, in which all 10 reviewers in Levels 1 and 2 screened the same records (50 and 30 records at Levels 1 and 2, respectively). Conflicting opinions about inclusion or exclusion of records were discussed to achieve consensus between reviewers and to refine and improve the clarity of the questions at each level (see Supplementary Material 1).

During Level 1 (screening on title and abstract), two reviewers assessed each record. To maximise the sensitivity of identification of relevant records, records progressed to Level 2 if either reviewer assessed that the record might be eligible. During Level 2 (screening on the full record), two reviewers initially assessed each record. Records were only included for charting in Level 3 if there was consensus that the record met the eligibility criteria between at least 2 reviewers. Consensus was achieved via discussion and if required, consultation with a third reviewer.

2.5 Data items and charting process – Level 3

A spreadsheet was used to standardise data charting in Level 3 (Supplementary Material 1). Data items that were extracted included the year of publication, the location and year of the study, the driving concern for the study (for example, conservation or disease investigation), and then further data within the overarching topics of interest: markets, animals, and risks.

Data that were extracted about markets included the type of market (for example, wildlife or food), and the physical structure, frequency, and types of people and activities at the markets. Hygiene practices and temporal changes (for example, increased sale of live animal species) were also recorded.

Data that were extracted about animals at the markets included the species of live animals sold (if listed as individual species), their source, the volume sold and their physical condition and housing. Classes and orders of all wildlife (including dead animals and animal products) sold at the market were also extracted, as well as the total volume of all wildlife and the purposes of wildlife sale. Any non-wildlife species sold at the market were recorded, as well as contacts that were documented between wildlife species and any other live animal species.

Data that were extracted about risks included specific pathogens and diseases, measures of their frequency and how they were identified, and any other risks associated with human and animal health.

Due to the longer form at Level 3, four additional reviewers were included, and agreement amongst all reviewers was initially assessed with three records (selected based on diversity of study type). Modifications were made to the form following reviewer feedback. The form was then tested by all reviewers with two more records to assess agreement. All conflicts were discussed. Once consensus was achieved for all five records and further modifications to the Level 3 form were made to improve clarity and increase agreement, the full Level 3 review was conducted. Conflicting opinions about extracted data were discussed between each record's pair of reviewers and a third reviewer to determine a consensus prior to synthesis of the extracted data.

2.6 Verification of the search strategy

228 Verification of the search strategy was undertaken by screening the bibliography list of three
229 records that were retained for data analysis in Level 3 (one for each decade of the search
230 window). Records which were potentially relevant to the study were included in the review
231 process using the same methods as records identified in the initial search.

232 2.7 Synthesis of extracted data

233 Topics of interest were tabulated according to region, and information about each topic was
234 then summarized and presented as a narrative.

235

236

3. Results

3.1. Screening

Initial and verification searches identified 3721 records. Following removal of 512 duplicates, 3209 records remained for screening (Figure 1). During Level 1 screening, 2795 records were excluded, leaving 414 records for screening of full text (Level 2). The most common reason for exclusion of records in Level 1 was lack of information about wildlife in markets (for example, the study investigated wildlife in the context of hunting or fishing; 98.6 %, n = 2757). During Level 2 screening, the most common reason for exclusion was lack of mention of live wildlife (92 %, n = 329). Of 133 records screened in the verification strategy, 5 were eligible for inclusion. Overall, 56 records were included for data charting and synthesis in Level 3 (Supplementary Material 2).

Figure 1: Diagram of the flow of records through the levels of a scoping review of reported research (1980–2020) associated with live, terrestrial, vertebrate wildlife sold for any purpose at markets likely to sell food.

3.2. Data charting

Of 56 records included in Level 3, all were single studies. A spreadsheet of the Level 3 studies and the data charted for each study is included in the Supporting Information. One record was sourced from a thesis (Edwards, 2012), and all other records were published in 35 peer-reviewed journals (Supplementary Material 3: Figure S1). The most frequently represented journal was *TRAFFIC Bulletin* (20%; $n = 11$), followed by *Biodiversity and Conservation* (7%; $n = 4$). Studies collected data from 28 countries (Figure 2). The most frequently represented study region was Southeast Asia ($n = 22$; mode country Lao PDR, $n = 6$), followed by China (here, classified as a region due to its large geographic size and number of studies; $n = 17$; mode province Guangdong, $n = 7$), Africa ($n = 16$; mode country Nigeria, $n = 4$), and South America ($n = 8$; mode country Brazil, $n = 6$). Two studies were conducted in more than one region: Europe and Africa (Spanish enclaves of Ceuta and Melilla, and Morocco; Nijman and Bergin (2017)), and China and Southeast Asia (Vietnam; Yiming and Dianmo (1998)).

Studies used data collected over 1–21-year periods from 1980–2019 and were published from 1998–2020. Although publications increased after the avian influenza outbreak in Hong Kong in 1997, and then subsequent to the SARS outbreak in 2003, data in the publications in the early 2000s was frequently collected prior to 1997; therefore, research intensity did not appear to be influenced by particular EID events (Figure 3).

The most common focus for studies was investigation of the wildlife trade in relation to conservation concerns such as species depletion and loss of biodiversity (75%; $n = 42$).

274 Disease risk was the primary focus of study in only 14 studies (25%), and there was no
275 association between the focus of studies and global region ($X^2 = 9.5$, d.f. = 8, $P = 0.3$).
276

277
278 Figure 2: Regions and countries (China by Province) in which data were collected in peer-reviewed records
279 identified in a scoping review of reported research (1980–2020) associated with live, terrestrial, vertebrate
280 wildlife sold for any purpose at markets likely to sell food.

Figure 3: Years in which data were collected (A) and published (B) in peer-reviewed records identified in a scoping review of research associated with live, terrestrial, vertebrate wildlife sold for any purpose at markets likely to sell food. Triangles denote major outbreaks of emerging infectious disease: brown = Avian influenza A H5N1, Hong Kong 1997; red = SARS, 2003; green = H1N1, 2009; yellow = Avian influenza A H7N9, China 2013; purple = Ebola virus disease, West Africa 2007; blue = COVID-19, 2019.

3.2.1 Markets

The most frequent market types (according to author definitions and descriptions) were general markets in South America, traditional medicine and bushmeat markets in Africa, food in Southeast Asia and China, live-bird markets in Africa, and wildlife in Southeast Asia (Figure 4, and Supplementary Material 2). The frequency of topics associated with markets are shown in Table 1.

Market physical structure was described in 20 records (36%); a range of permanent and temporary structures were reported, either indoor or outdoor (for example, on the sides of roads or in car parks), and often in locations accessible for the arrival and distribution of goods. For example, in Belém, Brazil, market trade takes advantage of riverside ports (da Nobrega Alves and Pereira Filho, 2007), and in the Niger Delta, hunters deliver goods via rivercraft to one of the largest bushmeat markets in the region which is also accessible via recently constructed roads (Akani et al., 2015). In central Lao PDR, authors reported that the most active wildlife markets were located close to major roads which facilitated trade across Southeast Asia (Schweikhard et al., 2019).

Figure 4: Market types identified in a scoping review of reported research (1980–2020) associated with the sale of live, terrestrial, vertebrate wildlife at markets likely to sell food globally.

Market frequency was highly variable although usually regular, with markets occurring every day on weekdays, on one day of the week only, on mornings only, at weekends only, at times dependent on the lunar calendar, or on special occasions. Unpredictable times and locations were a feature of some markets; for example, in Yunnan Province, China, wildlife trade could occur at irregular times and sites, and sometimes, markets appeared to be the sites of advertising and negotiation as vendors displayed their goods (including live wildlife in cages) whilst transfer of goods occurred elsewhere (Haibin and Kunming, 1999).

The types of people involved in markets were vendors and consumers – also called traders and buyers – as well as hunters and trappers, middlemen (also termed intermediaries and brokers), wildlife farmers, restaurant owners, government officials, slaughtermen, traditional healers and drivers. Middlemen could be involved in the chain of events before or after the market between wildlife acquisition and the end-user. For example, some middlemen traded between hunters and market vendors (these people could also be termed ‘collectors’ because they house wildlife as well as facilitated transactions), and others traded between market vendors and end-users such as restaurants, traditional medicine shops and tourists. Market trade was also associated with families as younger generations inherited the family business (for example, turtle collectors in Kerala, India; Krishnakumar et al. (2009)).

Illegal trade was reported in most records (68 %, $n = 38$), particularly if the focus was wildlife trade rather than disease risk (OR 17.0, 95 % CI 3.4—120.0, $P < 0.001$). Reports of illegal trade were not associated with a particular region ($X^2 = 5.1$, d.f. = 7, $P = 0.64$). Types of illegal trade included the sale of wildlife species that were prohibited for sale by national or international laws, sourced from protected areas such as National Parks (for example, in Liberia and the Democratic Republic of the Congo and Liberia, reported by Greengrass (2016) and Van Vliet et al. (2012), respectively), collected by illegal methods such as hunting (for example, pangolin in Myanmar; McEvoy et al. (2019)), and illegally imported (for

example, wildlife from Vietnam sold in Yunnan Province, China; Zhang et al. (2008)). Sometimes, laws were open to misinterpretation because they varied according to animal purpose (for example, hunting could be allowed depending on the destination and purpose of the wildlife in Indonesia; Chng et al. (2018); Janssen and Gomez (2019)) or were inconsistent between regional and national levels (for example, in Brazil; Fernandes-Ferreira et al. (2012)). Illegal trade was often conducted openly because it was either unregulated or ineffectively regulated. Examples include open trade despite signage that trade is illegal in Lao (Schweikhard et al., 2019), and turtle collectors in Kerala, India who were aware that trade was illegal, but were not deterred by small fines from local police and forest officials (Krishnakumar et al., 2009). Government officials were reportedly sometimes corrupt and therefore, not necessarily promoting regulatory activities (Haibin and Kunming, 1999). Sometimes illegal trade was clandestine. In Lao PDR, Davenport and Heatwole (2013) suspected that as live wildlife on display were sold, they were replenished from bags that were kept out of sight to avoid attention from local authorities, and in Mozambique, live wildlife and their products were concealed unless trade was likely (Williams et al., 2016). Other than trade, activities directly associated with live wildlife included slaughter and butchering. This was most frequently reported in the context of reptile and amphibian trade (for example, in Africa, South America and Southeast Asia (Catenazzi et al., 2010; da Nobrega Alves and Pereira Filho, 2007; Edwards, 2012; Gilbert et al., 2012; Kusrini and Alford, 2006; Pruvot et al., 2019)). Levels of hygiene and sanitation were reported in only 6 records (11%) and were not more associated with studies which investigated disease risk than studies of trade ($X^2 = 2.7$, d.f. = 1, $P = 0.10$). Cronin et al. (2015) observed that a bushmeat market in Bioko Island, Equatorial Guinea, Africa was occasionally closed for cleaning, and Van Vliet et al. (2012) noted that lack of refrigeration might be a reason for the high proportion of smoked bushmeat for sale at a market in Kisingani, DRC, Africa (and not an

indication of a large catchment area of wildlife as some have previously suggested). Others also noted lack of refrigeration of fresh meat in studies in Asia (Pruvot et al., 2019; Shepherd and Nijman, 2007). In an assessment of zoonotic disease risk in markets throughout Lao PDR, (Greatest et al., 2016) noted that hygiene practices such as hand-washing were rare, sanitation was poor (butchers knives and tables were rarely cleaned; blood and entrails were on the floor) and wildlife contact with fresh food was high in most markets. In a study from Nigeria and Egypt in which biosecurity compliance was evaluated (Fasanmi et al., 2016), wild animal trade was predictive for detection of highly-pathogenic avian influenza virus H5N1, and although not specific to activities involving wild animals, hygiene and sanitation measures were poor at many of the markets.

Temporal trends were mentioned in 18 records (32%), with changes in activity and species availability observed in some markets. In Yunnan Province, China, Haibin and Kunming (1999) noted increased smaller markets as well as underground wildlife trade networks. This could have been driven by increased regulation, reduced availability of some species, and increased consumption associated with development of the local economy and transport networks. Seasonality was observed in some markets; trade increased in the wet season in Nigeria (Akani et al., 2015) and in winter in Guangdong, China (Dong et al., 2007), and season was associated with species availability in Lao PDR and Brazil (Baía et al., 2010; Davenport and Heatwole, 2013; Schweikhard et al., 2019). Trade increased overall in Bioko Island (Cronin et al., 2015; Fa et al., 2000), and decreased in Togo, Mozambique and Papua New Guinea (D'Cruze et al., 2020; Eisemberg et al., 2011; Williams et al., 2016), whilst in markets in Myanmar, increases and decreases of particular species were observed over time (Min, 2012). Actors at markets also changed over time; for example, trade with non-locals has increased at Mong La market in Myanmar (Shepherd and Nijman, 2007). Trade also

sometimes increased around religious festivals; for example, in Belém, Brazil (Nobrega Alves and Rosa, 2010).

3.2.2 Wildlife Species

A total of 534 individual species of live, vertebrate, terrestrial wildlife were reported in the records from all taxonomic classes of vertebrates (Supplementary Material 3: Table S1). Of these, the most frequent orders were Testudines (37%, $n = 197$), followed by Passeriformes (20%, $n = 109$) and Squamata (12%, $n = 65$; Supplementary Material 3: Figure S2). Of mammalian orders that have been considered high risk for EID events (Delahay et al., 2021; Wikramanayake et al., 2021), no live species of Chiroptera (bats) were reported. Live Viverridae (large Indian, lesser Indian and masked palm civets) were reported in 4 records from China (Chow et al., 2014; Dong et al., 2007; Haibin and Kunming, 1999; Yiming and Dianmo, 1998). Pholidota (pangolins) were reported in 10 records from Africa ($n = 5$), Southeast Asia ($n = 4$) and China ($n = 2$) (Nijman 2016, (Akani et al., 2015; Edwards, 2012; Haibin and Kunming, 1999; Ingram et al., 2019; McEvoy et al., 2019; Shepherd and Nijman, 2007; Sodeinde and Soewu, 1999; Soewu and Ayodele, 2009; Yiming and Dianmo, 1998). Primate species were observed in markets in Africa, China and Southeast Asia, including a live chimpanzee in a bushmeat market in Liberia (Greengrass, 2016). Mustelidae species were observed in markets in China (Dong et al., 2007), and squirrels (Sciuridae) were observed in Southeast Asia and China (Privot et al., 2019; Yiming and Dianmo, 1998). Most species were reported in records from China (51%, $n = 271$) and Southeast Asia (24%, $n = 129$), and the most frequent classes sold in China and Southeast Asia were Reptilia and Aves, respectively (Figure 5). Relatively few live species were reported in records from South America ($n = 73$), or from the Middle East, Africa, Europe, the Indian subcontinent and Oceania (35, 29, 6, 2, and 1 species, respectively). In South America and the Middle East,

406 most species were also of class Aves, and in the remaining regions combined, Reptilia were
407 most frequently reported. Some records reported live wildlife by a common name only or by
408 their genus (Figure 5 inset; Supplementary Material 3: Table S2). Of these, wildlife of class
409 Aves were most frequently reported.

410 The greatest number of individual species of live, vertebrate, terrestrial wildlife reported in a
411 single study was 157 turtle species reported from a 35-month survey of turtle trade in Hong
412 Kong, Shenzhen and Guangzhou in southern China (Cheung and Dudgeon, 2006). Large
413 numbers of live wildlife species were also reported from agricultural and general markets in
414 Yunnan, China (n = 85; Haibin and Kunming (1999)), bird markets in Brazil (n = 51;
415 Fernandes-Ferreira et al. (2012)), and wildlife markets in Sumatra (n= 48; Chng et al. (2018),
416 at the Guangxi border between China and Vietnam (n = 47; Yiming and Dianmo (1998)), in
417 South China (n = 26; Chow et al. (2014)), and in Tabuk, Saudi Arabia (n = 31; Aloufi and
418 Eid (2014)). However, nearly half of the records only reported one live wildlife species (48%,
419 n = 26; Figure S3).

Figure 5: Classes of live, vertebrate, terrestrial wildlife reported in records in a scoping review of reported research (1980–2020) associated with the sale of live wildlife at markets likely to sell food globally. Main plot: classes of wildlife that were identified by species. Inset plot: classes of wildlife that were identified by common name, genus or order.

Wildlife (live or products) were mostly sold for food and medicine globally, as well as pets in Southeast Asia and China, and spiritual purposes in South America and Africa (Figure 6). The reported volume of live wildlife and total wildlife products sold was dependent on the study design (number of markets and frequency of visits), and inventories were not possible in some markets due to covert trading – information from records about volume of wildlife is included in the Supplementary Material 2. Most studies reported that wildlife were sourced locally, often from national parks and similarly protected areas; however, in markets in Africa, the Middle East, China and Southeast Asia, wildlife (not necessarily live) were sometimes imported (Figure S4). The condition of live wildlife was rarely reported although

435 Chow et al. (2014) noted tooth-like wounds of the feet of mammals that indicated they had
 436 been caught in traps, and Nekaris et al. (2010) found that the front teeth of lorises were
 437 clipped so that they made safer pets. In Lao PDR, Gilbert et al. (2012) observed animals tied
 438 with twine or bamboo through their hind limbs or foot webs, and Davenport and Heatwole
 439 (2013) observed frogs tied together, and lizards with their feet tied or their mouths wired or
 440 tied closed. In Brazil, Maloney et al. (2020) found that birds were in good condition. Live
 441 wildlife were often reported in unhygienic conditions in sacks, buckets in the case of reptiles
 442 and amphibians, or in overcrowded cages (Catenazzi et al., 2010; Nijman and Bergin, 2017;
 443 Stuart et al., 2000). The presence of other live animals was rarely reported (livestock in
 444 Mozambique; Williams et al. (2016), and poultry in Vietnam and China; Chow et al. (2014);
 445 Edmunds et al. (2011)). Likewise, proximity or contact between live wildlife and other
 446 species was rarely reported (contact with people; Pruvot et al. (2019), wild birds; Fasanmi et
 447 al. (2016), and poultry in the markets; Edmunds et al. (2011)).

Figure 6: Purpose of sale of live wildlife and wildlife products reported in records in a scoping review of reported research (1980–2020) associated with the sale of live wildlife at markets likely to sell food globally. Main plot: classes of wildlife that were identified by species. Inset plot: classes of wildlife that were identified by common name, genus or order.

3.2.2 Health risks associated with live wildlife

Twelve records identified specific diseases or microbes. Chytridiomycosis (caused by the fungus, *Batrachochytrium dendrobatidis*) was diagnosed by PCR and histological examination in 4 of 37 (10.8%) live bullfrogs in a food market in Yunnan Province, China (Bai et al., 2010). *B. dendrobatidis* was also detected in all live frogs sampled in a study in

459 Peru (N = 5; Catenazzi et al. (2010)), and in one live bullfrog in Cambodia (N = 71; Gilbert et
460 al. (2012)).

461 *Spirometra*, a zoonotic cestode that causes sparganosis, was detected with > 25% prevalence
462 in snakes sampled in Guangdong (Wang et al., 2014; Wang et al., 2011) and Indonesia
463 (Yudhana et al., 2019). In a study in markets in Brazil, the protozoan, *Blastocystis*, was
464 detected with approximately 20% prevalence in Galliformes, Struthioniformes and
465 Anseriformes, although not found in wild birds for sale (Maloney et al., 2020). Whilst some
466 subtypes of *Blastocystis* have zoonotic potential, pathogenicity in people is variable; the
467 authors noted that isolation of a subtype of *Blastocystis*, ST5, in ostriches in the study was
468 significant because ST5 has been found in pigs and pig caretakers in China and Australia.

469 Eighteen isolates of avian influenza virus (H5N1, H9N2, H7N3, H7N9, H10N6, H3N6) were
470 found in migratory Anseriformes in live-bird markets in Egypt (Kayed et al., 2019). Highly
471 pathogenic H5N1 was also detected in tracheal and cloacal swabs, post-mortem, and
472 moribund or freshly killed birds in 65% of live-bird markets surveyed in Nigeria and Egypt
473 (N= 91; Fasanmi et al. (2016)).

474 In a study to examine the ecology and evolutionary pathways of *Coronaviridae* (CoV),
475 Vijaykrishna et al. (2007) investigated viral sequences obtained from species including bats,
476 humans and livestock, as well as wild animals (raccoon dog and Chinese ferret badger)
477 sampled at live-animal markets in Guangdong Province, China. The circumstances of the
478 animals at the markets or the frequency of positive CoV identification was not described. In
479 another study in live-animal markets in Guangdong and Guangxi Province, China, 24 live
480 wildlife species were sampled (Dong et al., 2007). CoV was detected in several Asian leopard
481 cats (*Prionailurus bengalensis*; n = 35/1,453, 2.4%) and Chinese ferret badgers (*Melogale*
482 *moschata*; n = 11/934, 1.1%), and once in 6 other species: yellow-bellied weasel (*Mustela*

483 *kathiah*), Siberian weasel (*M. sibirica*), masked palm civet (*Paguma larvata*), Chinese
484 bamboo rat (*Rhizomys sinensis*), lesser Indian civet (*Viverricula indica*), and flying squirrel
485 (*Petaurista sp.*). Ferret badgers and Asian leopard cats were available in the markets all year
486 round, although most CoV positive swabs were detected in winter.
487 Zoonotic disease risks that were mentioned other than those associated with disease and
488 pathogens detected in the studies included acknowledgement of the presence of large
489 numbers of species susceptible to HPAI H5N1 in ornamental bird markets in Vietnam
490 (Edmunds et al., 2011), and concern about zoonoses such as Ebola virus disease, rabies, and
491 tuberculosis associated with handling primates in Benin, Africa (Sogbohossou et al., 2018).
492 In markets in Guangxi at the China-Vietnam border, Yiming and Dianmo (1998) noted that
493 wildlife mortality could be as high as 50%, especially during transport due to unhygienic
494 conditions and overcrowding and that subsequent disposal of animals around trading areas
495 and in forests could result in disease spread. Chow et al. (2014) commented that
496 overcrowding could increase the probability of disease spread, and Kusriini and Alford (2006)
497 noted that escaped exotic species (bullfrogs found in rice fields in West Java, Indonesia)
498 could be a disease risk to local frog species.
499
500

Table 1: Number of records (percentage) from each region, which contained information from specific topics of interest in a scoping review of reported research (1980–2020) associated with live, terrestrial, vertebrate wildlife sold for any purpose at markets likely to sell food. Bold indicates that $\geq 50\%$ of studies from that region contained information on this topic (excluding regions in which there were ≤ 2 studies).

Region		South America	Africa	Africa/Europe	Middle East	Indian subcontinent	Southeast Asia	China	China/ Southeast Asia	Oceania	Total
Median year of publication (minimum – maximum)		2010 (1999— 2020)	2015 (1999— 2020)	2017 (2017—2017)	2016 (2014— 2018)	2009 (2009— 2009)	2015 (2000— 2019)	2007 (1999— 2014)	1998 (1998— 1998)	2011 (2011— 2011)	
Broad topic	Topic										
Market	Physical structure	3 (38%)	7 (50%)	0	1 (50%)	0	8 (47%)	0	1 (100%)	0	20 (36%)
	Frequency	2 (25%)	6 (43%)	0	1 (50%)	0	6 (35%)	1 (9%)	0	1 (100%)	17 (30%)
	Activities	3 (38%)	6 (43%)	0	0	1 (100%)	7 (41%)	1 (9%)	0	0	18 (32%)
	Illegal wildlife trade	6 (75%)	11 (79%)	1 (100%)	1 (50%)	1 (100%)	11 (65%)	6 (54%)	1 (100%)	0	38 (68%)
	General hygiene	0	3 (21%)	0	0	0	2 (12%)	0	0	0	5 (9%)
	Temporal trend	2 (25%)	6 (43%)	0	0	1 (100%)	6 (35%)	2 (18%)	0	1 (100%)	18 (32%)
Live wildlife	Listed by species	8 (100%)	10 (71%)	1 (100%)	2 (100%)	1 (100%)	14 (82%)	9 (81%)	0	1 (100%)	46 (82%)
	Source	6 (75%)	11 (79%)	0	1 (50%)	1 (100%)	14 (82%)	9 (81%)	1 (100%)	1 (100%)	44 (78%)
	Volume	5 (63%)	7 (50%)	1 (100%)	0	1 (100%)	11 (65%)	3 (27%)	0	1 (100%)	29 (52%)
	Purpose of sale	7 (88%)	13 (93%)	1 (100%)	1 (50%)	1 (100%)	15 (88%)	10 (91%)	1 (100%)	1 (100%)	50 (89%)
	Condition of wildlife	3 (38%)	1 (7%)	0	0	0	5 (29%)	1 (9%)	0	0	10 (18%)
	Housing	4 (50%)	4 (28%)	1 (100%)	0	1 (100%)	8 (47%)	2 (18%)	0	0	21 (38%)
	Contact between species	0	0	0	1 (50%)	0	2 (12%)	0	0	0	3 (5%)

Study focus Risks	Other species	2 (25%)	3 (21%)	0	0	0	1 (6%)	2 (18%)	0	0	8 (14%)
	Disease	2 (25%)	1 (7%)	0	1 (50%)	0	4 (22%)	6 (54%)	0	0	14 (25%)
	Disease identified	0	0	0	0	0	1(6%)	3 (27%)	0	0	4 (7%)
	Microbe identified	2 (25%)	1 (7%)	0	1 (50%)	0	2 (12%)	5 (46%)	0	0	10 (18%)
	Other risk identified	1 (13%)	1 (7%)	0	0	0	3 (19%)	2 (18%)	1 (100%)	0	8 (14%)
Total records		8	14	1	2	1	16	11	1	1	56

4. Discussion

This review highlighted enormous global variability and epistemic uncertainty in all aspects regarding circumstances in which live, terrestrial wildlife are sold in markets that are likely to sell fresh food. Specific insights included the limited research focus on disease associated with live wildlife in markets, the wide variety of live wildlife species traded at markets especially in China and Southeast Asia, and the high accessibility of markets (both socially and geographically). These insights, as well as the variability and uncertainty related to the markets, the wildlife, and associated risks, greatly influence the feasibility of accurate assessment of the risk of emerging infectious disease associated with live wildlife trade in markets, and the development of effective, sustainable policy in this environment.

Most studies in this review focused on the impact of live wildlife trade in markets on conservation, particularly biodiversity loss. Whilst this can indirectly inform the scale of biodiversity loss as a broad driver of emerging infectious disease, focus on potential pathogens and disease is needed for risk analysis frameworks, which require information about specific hazards and exposure pathways. Many zoonotic and foodborne microorganisms have been identified in items other than live wildlife (including fresh meat, vegetables, and the environment) in wet markets worldwide, such as *Leptospira* spp., *Vibrio* spp., *Toxoplasma* spp., *Campylobacter* spp., and *Salmonella* spp. (Hamid et al., 2020; Kottawattage et al., 2017; Ngan et al., 2020; Sekoai et al., 2020), yet only three potentially zoonotic microorganisms (*Spirometra* species, *Coronaviridae*, and avian influenza virus) were reported from live wildlife in this review. There was also limited information about water, sanitation, and hygiene (WASH; <https://www.who.int/health-topics/water-sanitation-and-hygiene-wash>, accessed 20 August 2021) conditions associated with live wildlife trade, such as butchering and slaughter of wildlife, contamination of other food products or the environment, or contact between wildlife and other species, including people. Although

market regulations are often aimed at reducing the risk of zoonotic and foodborne disease, and some countries – for example China, Indonesia, and Thailand – have specific WASH regulations applied to wet markets (The Law Library of Congress, 2020), there appears to be a gap in knowledge about the range of microorganisms and WASH risks associated with live wildlife. A greater understanding of current WASH conditions around activities associated with live wildlife trade (including market infrastructure such as hand washing facilities, adequate drainage, and separation of animals) is required to inform risk assessment exposure pathways. In addition, previous recommendations to increase surveillance of wildlife to identify potential pathogens and to detect and respond to emerging infectious disease (Grogan et al., 2014; Kuiken et al., 2005; Vrbova et al., 2010) appear to have been largely overlooked in the context of live wildlife in markets and should be adopted more comprehensively.

The greatest number of live wildlife species was reported from studies in markets in China and Southeast Asia. Wide species diversity might provide more opportunity for EID events due to a greater number of potential reservoir-spillover host pairs. However, following previous patterns of disease emergence, species from classes Mammalia and Aves have been considered a higher disease-risk due to their association with potentially pathogenic microorganisms (Olival et al., 2017; White and Razgour, 2020; Woolhouse et al., 2012), and a greater focus on these classes of wildlife might be warranted. Consistent with this, the World Health Organization have recommended suspension of trade of live-caught mammalian wildlife in traditional food markets (World Health Organisation, 2021). In this review, live mammalian wildlife species that could be considered high risk were identified in studies from markets in Africa, Southeast Asia, and China, and wild birds were also reported in markets from South America and the Middle East in addition to these regions. However, the likelihood of spillover will also depend on additional characteristics of the wildlife-potential spillover host interface, including its magnitude (amount of live wildlife traded and

contact with the potential spillover species) and factors that affect transmission, such as stressors that might influence the shedding of microorganisms (including travel, overcrowded housing, comorbidities and the physical condition of the wildlife), the type of contact occurring and the susceptibility of the potential spillover host (Plowright et al., 2017; Wikramanayake et al., 2021). The scale of interfaces could not be determined in this review, and although most studies and species were identified from China and Southeast Asia, this might not be representative of the global distribution of markets or volume of live wildlife sold. Ultimately, resources for mitigation strategies will be limited, and development of effective policy will require assessment of relative risk dependent on the diverse global contexts that were apparent in this review, such as market type (for example, markets in this review were predominantly bushmeat in Africa, wildlife in Southeast Asia, and food in China) and species sold, as well as local, market-specific factors.

The scope of policy to mitigate EID risks associated with trade of live wildlife in markets is also important to define. If the goal is to prevent dissemination of disease from markets and an epidemic – and not only to prevent spillover of microorganisms from live wildlife reservoir hosts – social and geographic pathways from markets also need to be considered in risk assessments. Short value-chains that support local producers and ensure a supply of fresh food at low cost – due to the limited requirement for transport and refrigeration – have been considered an advantage of wet markets (Goldman et al., 1999; Zhang and Pan, 2013).

However, several studies in this review described how markets were located to facilitate trade and distribution, and highlighted that wildlife trade is a high-value, global activity with long value chains and movement of wildlife across many regions. In 2017, the value of the global wildlife trade was an estimated USD23 billion (van Uhm and Wong, 2019). They also described a multitude of people associated with live wildlife trade, consistent with the typologies defined by Phelps et al. (2016). Long value-chains which involve many people

might be specific features of markets selling live wildlife that could be related to the purpose and value of wildlife. Although quantitative methods can be used to characterise risks and their pathways within markets (such as species abundance and contact durations), social science and ethnographic methods, such as those described by Edwards (2012) in the context of markets in Mali, Africa, are likely to be of greater value to understand complex wildlife trading networks. Understanding the drivers and risks behind network structure is valuable to identify potential control points (Phelps et al., 2016), and will also serve to understand the socio-economic impacts of regulations to mitigate risks. Networks involving people and transport are often small-world networks and it is possible that wildlife trade networks are of this type; such networks can be robust in the face of disruption (Hu and Verma, 2011). Therefore, as well as not addressing the drivers of wildlife consumption, banning trade of live mammalian wildlife in markets might not be an effective strategy to mitigate EID risks. It has already been suggested that those whose livelihoods depend on wildlife trade are likely to find alternative trade routes that avoid regulations; trade diverted to hidden networks could inadvertently pose a greater risk to public health (Roe et al., 2020).

Overall, this review demonstrates that accurate assessment of the risk of EIDs associated with live wildlife trade in markets is currently infeasible due to massive epistemic uncertainty. The magnitude of live wildlife-potential spillover host interfaces is unknown, high-risk species and microorganism pairs are largely unknown, there is limited information about activities that could influence exposure to microorganisms, and there is limited evidence to associate EID events with wet markets. Although this review did not identify records in languages other than English, the small proportion of records which focused on disease indicates that assessing EID risks in markets has had less research focus than the conservation impacts of live wildlife trade. This does not mean that markets are not a risk for EID emergence and that a precautionary approach is unwarranted. Ultimately however, policy will need to be

evidence-based and commensurate with risk so that it can be implemented and sustained. The high-profile association of early COVID-19 cases with a wet market in Wuhan and the current recommendation to ban trade of live mammalian species in markets has catalysed the need to understand these risks. This review provides a baseline of currently available, peer-reviewed information about live wildlife trade in markets globally. Addressing the gaps in knowledge to accurately assess risks across the wide range of market types, activities and wildlife species purpose globally will provide the framework in which policy to promote biodiversity, protection of livelihoods and prevention of EIDs can be supported.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

We thank the Australian Department of Education, Skills and Employment, Enabling Growth and Innovation Program for their funding and support.

References

- Aisher A. Scarcity, alterity and value: decline of the pangolin, the world's most trafficked mammal. *Conservation and Society* 2016; 14: 317-329.
- Akani GC, Amadi N, Eniang EA, Luiselli L, Petrozzi F. Are mammal communities occurring at a regional scale reliably represented in "hub" bushmeat markets? A case study with Bayelsa State (Niger Delta, Nigeria). *Folia Zoologica* 2015.
- Aloufi A, Eid E. Conservation perspectives of illegal animal trade at markets in Tabuk, Saudi Arabia. *Traffic Bulletin* 2014.

- Andersen KG, Rambaut A, Lipkin WI, Holmes EC, Garry RF. The proximal origin of SARS-CoV-2. *Nature Medicine* 2020; 26: 450-452.
- Bai C, Garner TWJ, Li Y. First evidence of batrachochytrium dendrobatidis in China: Discovery of chytridiomycosis in introduced American bullfrogs and native amphibians in the Yunnan Province, China. *EcoHealth* 2010.
- Baía PC, Jr., Guimarães DA, Le Pendu Y. Non-legalized commerce in game meat in the Brazilian Amazon: a case study. *Rev Biol Trop* 2010.
- Catenazzi A, Vredenburg VT, Lehr E. Batrachochytrium dendrobatidis in the live frog trade of Telmatobius (Anura: Ceratophryidae) in the tropical Andes. *Diseases of Aquatic Organisms* 2010.
- Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *The Lancet* 2020; 395: 507-513.
- Cheung SM, Dudgeon D. Quantifying the Asian turtle crisis: market surveys in southern China, 2000–2003. *Aquatic Conservation: Marine and Freshwater Ecosystems* 2006.
- Chng SC, Shepherd CR, Eaton JA. In the market for extinction: birds for sale at selected outlets in Sumatra. *TRAFFIC Bulletin* 2018.
- Chow AT, Cheung S, Yip PK. Wildlife markets in south China. *Human-Wildlife Interactions* 2014.
- Cronin DT, Woloszynek S, Morra WA, Honarvar S, Linder JM, Gonder MK, et al. Long-Term Urban Market Dynamics Reveal Increased Bushmeat Carcass Volume despite Economic Growth and Proactive Environmental Legislation on Bioko Island, Equatorial Guinea. *PLoS One* 2015.
- D'Cruze N, Assou D, Coulthard E, Norrey J, Megson D, Macdonald DW, et al. Snake oil and pangolin scales: insights into wild animal use at “Marché des Fétiches” traditional medicine market, Togo. *Nature Conservation* 2020; 39: 45.
- da Nobrega Alves RR, Pereira Filho GA. Commercialization and use of snakes in North and Northeastern Brazil: Implications for conservation and management. *Biodiversity and Conservation* 2007.
- Davenport D, Heatwole H. Wild terrestrial vertebrates sold as food in open markets in Laos. *Pacific Conservation Biology* 2013.
- Delahay RJ, de la Fuente J, Smith GC, Sharun K, Snary EL, Flores Girón L, et al. Assessing the risks of SARS-CoV-2 in wildlife. *One Health Outlook* 2021; 3: 7.
- Dong BQ, Liu W, Fan XH, Vijaykrishna D, Tang XC, Gao F, et al. Detection of a novel and highly divergent coronavirus from asian leopard cats and Chinese ferret badgers in Southern China. *J Virol* 2007.

- Edmunds K, Robertson SI, Few R, Mahood S, Bui PL, Hunter PR, et al. Investigating Vietnam's Ornamental Bird Trade: Implications for Transmission of Zoonoses. Ecohealth 2011.
- Edwards IB. The social life of wild-things: Negotiated wildlife in Mali, West Africa. 2012.
- Eisemberg CC, Rose M, Yaru B, Georges A. Demonstrating decline of an iconic species under sustained indigenous harvest - The pig-nosed turtle (*Carettochelys insculpta*) in Papua New Guinea. Biological Conservation 2011.
- Fa JE, Yuste JEG, Castelo R. Bushmeat markets on Bioko Island as a measure of hunting pressure. Conservation Biology 2000.
- Fasanmi OG, Ahmed SSU, Oladele-Bukola MO, El-Tahawy AS, Elbestawy AR, Fasina FO. An evaluation of biosecurity compliance levels and assessment of associated risk factors for highly pathogenic avian influenza H5N1 infection of live-bird-markets, Nigeria and Egypt. Acta Trop 2016.
- Fernandes-Ferreira H, Mendonca SV, Albano C, Ferreira FS, Nobrega Alves RR. Hunting, use and conservation of birds in Northeast Brazil. Biodiversity and Conservation 2012.
- Gilbert M, Bickford D, Clark L, Johnson A, Joyner PH, Ogg Keatts L, et al. Amphibian pathogens in Southeast Asian frog trade. Ecohealth 2012.
- Goldman A, Krider R, Ramaswami S. The Persistent Competitive Advantage of Traditional Food Retailers in Asia: Wet Markets' Continued Dominance in Hong Kong. Journal of macromarketing 1999; 19: 126-139.
- Greatorrex ZF, Olson SH, Singhalath S, Silithammavong S, Khammavong K, Fine AE, et al. Wildlife Trade and Human Health in Lao PDR: An Assessment of the Zoonotic Disease Risk in Markets. Plos One 2016.
- Greengrass E. Commercial hunting to supply urban markets threatens mammalian biodiversity in Sapo National Park, Liberia. Oryx 2016.
- Grogan LF, Berger L, Rose K, Grillo V, Cashins SD, Skerratt LF. Surveillance for emerging biodiversity diseases of wildlife. PLoS Pathogens 2014; 10: e1004015.
- Guan Y, Zheng B, He Y, Liu X, Zhuang Z, Cheung C, et al. Isolation and characterization of viruses related to the SARS coronavirus from animals in southern China. Science 2003; 302: 276-278.
- Haibin W, Kunming G. Wildlife trade in Yunnan Province, China, at the border with Vietnam. Traffic Bulletin 1999.
- Hamid NA, Sadiq MB, Ramanoon SZ, Mansor R, Watanabe M, Nur Mahiza Md I, et al. Seroprevalence and Risk Factors of *Toxoplasma gondii* in Ruminant Meats from Wet Markets in Klang Valley and Abattoirs in Selangor, Malaysia. Animals 2020; 10: 1139.

- He F, En-Fu C, Fu-Dong L, Xin-Yi W, Xiao-Xiao W, Jun-Fen L. Human infection and environmental contamination with Avian Influenza A (H7N9) Virus in Zhejiang Province, China: risk trend across the three waves of infection. BMC Public Health 2015; 15.
- Hu Z, Verma PK. Topological resilience of complex networks against failure and attack. 2011 Fifth IEEE International Conference on Advanced Telecommunication Systems and Networks (ANTS), 2011, pp. 1-6.
- Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. The lancet 2020; 395: 497-506.
- Ingram DJ, Cronin DT, Challender DWS, Venditti DM, Gonder MK. Characterising trafficking and trade of pangolins in the Gulf of Guinea. Global Ecology and Conservation 2019.
- Janssen J, Gomez L. Common sun skink eutropis multifasciata (kuhl 1820) sold for traditional medicine in indonesia and potential conservation implications. Ethnobiology and Conservation 2019.
- Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, et al. Global trends in emerging infectious diseases. Nature 2008; 451: 990-U4.
- Kayed AS, Kandeil A, Gomaa MR, El-Shesheny R, Mahmoud S, Hegazi N, et al. Surveillance for avian influenza viruses in wild birds at live bird markets, Egypt, 2014-2016. Influenza Other Respir Viruses 2019.
- Kottawattage SAK, VanBergen MAP, Abeynayake P, Wagenaar JA, Veldman KT, Kalupahana RS. Campylobacter in Broiler Chicken and Broiler Meat in Sri Lanka: Influence of Semi-Automated vs. Wet Market Processing on Campylobacter Contamination of Broiler Neck Skin Samples. Foods 2017; 6: 105.
- Krishnakumar K, Raghavan R, Pereira B. Protected on paper, hunted in wetlands: exploitation and trade of freshwater turtles (Melanochelys trijuga coronata and Lissemys punctata punctata) in Punnamada, Kerala, India. Tropical Conservation Science 2009.
- Kuiken T, Leighton FA, Fouchier RAM, LeDuc JW, Peiris JSM, Schudel A, et al. Public Health. Pathogen Surveillance in Animals, 2005.
- Kusrini MD, Alford RA. Indonesia's exports of frogs' legs. Traffic Bulletin 2006.
- Lam TT-Y, Jia N, Zhang Y-W, Shum MH-H, Jiang J-F, Zhu H-C, et al. Identifying SARS-CoV-2-related coronaviruses in Malayan pangolins. Nature 2020; 583: 282-285.
- Magouras I, Brookes VJ, Jori F, Martin A, Pfeiffer DU, Dürr S. Emerging Zoonotic Diseases: Should We Rethink the Animal–Human Interface? Frontiers in Veterinary Science 2020; 7.
- Maloney JG, Molokin A, da Cunha MJR, Cury MC, Santin M. Blastocystis subtype distribution in domestic and captive wild bird species from Brazil using next generation amplicon sequencing. Parasite Epidemiol Control 2020.

- McEvoy JF, Connette G, Huang Q, Soe P, Pyone KHH, Valitutto M, et al. Two sides of the same coin—Wildmeat consumption and illegal wildlife trade at the crossroads of Asia. *Biological Conservation* 2019; 238: 108197.
- Min S. Impacts of Wildlife Trade on Conservation in Kachin State, Myanmar. *TRAFFIC Bulletin* 2012.
- Nadimpalli ML, Pickering AJ. A call for global monitoring of WASH in wet markets. *The Lancet Planetary Health* 2020; 4: e439-e440.
- Naguib MM, Li R, Ling J, Grace D, Nguyen-Viet H, Lindahl JF. Live and Wet Markets: Food Access versus the Risk of Disease Emergence. *Trends in Microbiology* 2021.
- Nekaris KAI, Shepherd CR, Starr CR, Nijman V. Exploring cultural drivers for wildlife trade via an ethnoprimateological approach: A case study of slender and slow lorises (*Loris* and *Nycticebus*) in South and Southeast Asia. *American Journal of Primatology* 2010.
- Ngan WY, Rao S, Chan LC, Sekoai PT, Pu Y, Yao Y, et al. Impacts of Wet Market Modernization Levels and Hygiene Practices on the Microbiome and Microbial Safety of Wooden Cutting Boards in Hong Kong. *Microorganisms* 2020; 8: 1941.
- Nijman V, Bergin D. Reptiles traded in markets for medicinal purposes in contemporary Morocco. *Contributions to Zoology* 2017.
- Nobrega Alves RR, Rosa IL. Trade of Animals Used in Brazilian Traditional Medicine: Trends and Implications for Conservation. *Human Ecology* 2010.
- Olival KJ, Hosseini PR, Zambrana-Torrel C, Ross N, Bogich TL, Daszak P. Host and viral traits predict zoonotic spillover from mammals. *Nature* 2017; 546: 646-650.
- Phelps J, Biggs D, Webb EL. Tools and terms for understanding illegal wildlife trade. *Frontiers in Ecology and the Environment* 2016; 14: 479-489.
- Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, et al. Pathways to zoonotic spillover. *Nature Reviews Microbiology* 2017; 15: 502-510.
- Pruvot M, Khammavong K, Milavong P, Philavong C, Reinharz D, Mayxay M, et al. Toward a quantification of risks at the nexus of conservation and health: The case of bushmeat markets in Lao PDR. *Science of the Total Environment* 2019.
- Roe D, Dickman A, Kock R, Milner-Gulland EJ, Rihoy E, 't Sas-Rolfes M. Beyond banning wildlife trade: COVID-19, conservation and development. *World Development* 2020; 136: 105121.
- Schweikhard J, Kasper K, Ebert C, Lehmann M, Erbe P, Ziegler T. Investigations into the illegal wildlife trade in central Lao PDR. *Traffic Bulletin* 2019; 31: 19-25.
- Sekoai PT, Feng S, Zhou W, Ngan WY, Yang P, Yao Y, et al. Insights into the Microbiological Safety of Wooden Cutting Boards Used for Meat Processing in Hong Kong's Wet Markets: A Focus on Food-Contact Surfaces, Cross-Contamination and the Efficacy of Traditional Hygiene Practices. *Microorganisms* 2020; 8: 579.

- Shepherd C, Nijman V. An assessment of wildlife trade at Mong La market on the Myanmar-China border. *Traffic Bulletin* 2007; 21: 85-88.
- Sigal S. The Coronavirus Likely Came from China's Wet Markets. They're Reopening Anyway. 2021. Vox, 2020.
- Sodeinde O, Soewu D. Pilot study of the traditional medicine trade in Nigeria. *TRAFFIC BULLETIN-CAMBRIDGE-TRAFFIC INTERNATIONAL*- 1999.
- Soewu DA, Ayodele IA. Utilisation of Pangolin (Manis sps) in traditional Yorubic medicine in Ijebu province, Ogun State, Nigeria. *J Ethnobiol Ethnomed* 2009.
- Sogbohossou EA, Kassa B, Akpona HA, Amahowe IO, Djagoun J, Sinsin B. *TRADE IN PRIMATE SPECIES FOR MEDICINAL PURPOSES IN SOUTHE IMPLICATIONS FOR CONSERVATION*. *TRAFFIC Bulletin* 2018.
- Stuart BL, Smith J, Davey K, Din P, Platt SG. Homalopsine watersnakes: the harvest and trade from Tonle Sap, Cambodia. *TRAFFIC BULLETIN-CAMBRIDGE-TRAFFIC INTERNATIONAL*- 2000; 18: 115-124.
- Tam JS. Influenza A (H5N1) in Hong Kong: an overview. *Vaccine* 2002; 20: S77-S81.
- Taylor LH, Latham SM, Woolhouse ME. Risk factors for human disease emergence. *Philos Trans R Soc Lond B Biol Sci* 2001; 356: 983-9.
- The Law Library of Congress. Regulation of Wild Animal Wet Markets in Selected Jurisdictions: Law Library of Congress, Global Legal Research Directorate, 2020.
- Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Annals of Internal Medicine* 2018; 169: 467-473.
- van Uhm DP, Wong RW. Establishing trust in the illegal wildlife trade in China. *Asian Journal of Criminology* 2019; 14: 23-40.
- Van Vliet N, Nebesse C, Gambalemoke S, Akaibe D, Nasi R. The bushmeat market in Kisangani, Democratic Republic of Congo: Implications for conservation and food security. *ORYX* 2012.
- Vijaykrishna D, Smith GJ, Zhang JX, Peiris JS, Chen H, Guan Y. Evolutionary insights into the ecology of coronaviruses. *J Virol* 2007.
- Vrbova L, Stephen C, Kasman N, Boehnke R, Doyle Waters M, Chablitt Clark A, et al. Systematic review of surveillance systems for emerging zoonoses. *Transboundary and emerging diseases* 2010; 57: 154-161.
- Wang F, Li W, Hua L, Gong S, Xiao J, Hou F, et al. Spirometra (Pseudophyllidea, Diphyllbothriidae) severely infecting wild-caught snakes from food markets in Guangzhou and Shenzhen, Guangdong, China: implications for public health. *ScientificWorldJournal* 2014.

- Wang F, Zhou L, Gong S, Deng Y, Zou J, Wu J, et al. Severe infection of wild-caught snakes with *Spirometra erinaceieuropaei* from food markets in Guangzhou, China involves a risk for zoonotic sparganosis. *J Parasitol* 2011.
- White RJ, Razgour O. Emerging zoonotic diseases originating in mammals: a systematic review of effects of anthropogenic land-use change. *Mammal Review* 2020; 50: 336-352.
- Wikramanayake E, Pfeiffer DU, Magouras I, Conan A, Ziegler S, Bonebrake TC, et al. A tool for rapid assessment of wildlife markets in the Asia-Pacific Region for risk of future zoonotic disease outbreaks. *One Health* 2021; 13: 100279.
- Williams VL, Moshoeu TJ, Alexander GJ. Reptiles sold as traditional medicine in Xipamanine and Xiquelene Markets (Maputo, Mozambique). *South African Journal of Science* 2016.
- Woo PC, Lau SK, Yuen K-y. Infectious diseases emerging from Chinese wet-markets: zoonotic origins of severe respiratory viral infections. *Current opinion in infectious diseases* 2006; 19: 401-407.
- Woolhouse M, Scott F, Hudson Z, Howey R, Chase-Topping M. Human viruses: discovery and emergence. *Philosophical Transactions of the Royal Society B: Biological Sciences* 2012; 367: 2864-2871.
- Woolhouse MEJ, Haydon DT, Antia R. Emerging pathogens: the epidemiology and evolution of species jumps. *Trends in Ecology & Evolution* 2005; 20: 238-44.
- World Health Organisation. Reducing public health risks associated with the sale of live wild animals of mammalian species in traditional food markets. In: Safety NaF, editor. World Health Organization, WHO Headquarters (HQ), 2021, pp. 8.
- World Health Organization. A guide to healthy food markets. World Health Organization, Geneva, Switzerland, 2006.
- Xiao X, Newman C, Buesching CD, Macdonald DW, Zhou Z-M. Animal sales from Wuhan wet markets immediately prior to the COVID-19 pandemic. *Scientific reports* 2021; 11: 1-7.
- Xu RH, He JF, Evans MR, Peng GW, Field HE, Yu DW, et al. Epidemiologic clues to SARS origin in China. *Emerg Infect Dis* 2004; 10: 1030-7.
- Yiming L, Dianmo L. The dynamics of trade in live wildlife across the Guangxi border between China and Vietnam during 1993–1996 and its control strategies. *Biodiversity & Conservation* 1998.
- Yu D, Li H, Xu R, MPH, He J, Lin J, et al. Prevalence of IgG antibody to SARS-associated coronavirus in animal traders--Guangdong Province, China, 2003. *MMWR. Morbidity and mortality weekly report* 2003; 52: 986-987.
- Yudhana A, Praja RN, Supriyanto A. The medical relevance of *Spirometra* tapeworm infection in Indonesian Bronzeback snakes (*Dendrelaphis pictus*): A neglected zoonotic disease. *Vet World* 2019.

- Zhang L, Hua N, Sun S. Wildlife trade, consumption and conservation awareness in southwest China. *Biodivers Conserv* 2008.
- Zhang QF, Pan Z. The transformation of urban vegetable retail in China: Wet markets, supermarkets and informal markets in Shanghai. *Journal of Contemporary Asia* 2013; 43: 497-518.
- Zhang T, Wu Q, Zhang Z. Probable Pangolin Origin of SARS-CoV-2 Associated with the COVID-19 Outbreak. *Current Biology* 2020; 30: 1346-1351.e2.
- Zhong S, Crang M, Zeng G. Constructing freshness: the vitality of wet markets in urban China. *Agriculture and Human Values* 2020; 37: 175-185.
- Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 2020; 579: 270-273.