

**Social inequalities, length of hospital stay for chronic conditions and the mediating role of comorbidity and discharge destination: A multilevel analysis of hospital administrative data linked to the population census in Switzerland**

**Short Title: Social inequalities, comorbidity and length of hospital stays**

Lucy Bayer-Oglesby<sup>1</sup>, Andrea Zumbrunn<sup>1</sup>, Nicole Bachmann<sup>1</sup> and SIHOS Team


<sup>1</sup>University of Applied Sciences and Arts Northwestern Switzerland FHNW, School of Social Work, Institute for Social Work and Health, Riggenbachstrasse 16, CH-4600 Olten, Switzerland.

Correspondence: [lucy.bayer@fhnw.ch](mailto:lucy.bayer@fhnw.ch)


# Supporting Information

**Table S1** Cluster sizes and Intra-Class-Correlation (ICC) of cluster Variables, linear CCMM (null-model with outcome length of stay)


Custer Variable	Records (N)	Clusters (N)	Cluster Size					ICC
			Min.	Max.	Median	Mean	SD	
Hospitals	141'307	188	1	5513	321.50	751.63	1085.96	0.281
Patients	141'307	92'623	1	23	1.00	1.53	1.02	0.154



**Figure S1:** Length of stay by age and social factors (top left: sex; top right: education; middle left: insurance class, middle right: type of household; bottom right: language skills; bottom left: migration background)



**Figure S2:** Comorbidity by age and social factors (top left: sex; top right: education; middle left: insurance class, middle right: type of household; bottom right: language skills; bottom left: migration background)



**Figure S3:** Discharge destination by age and social factors (top left: sex; top right: education; middle left: insurance class, middle right: type of household; bottom right: language skills; bottom left: migration background)

**Table S2: Effect estimates for significant interactions between main diagnosis and social factors**

Simple Contrast for significant interactions between main diagnosis and social factors		Contrast Estimate	Adj. Sig.	95% Confidence Interval	
Main Diagnosis	Educational Attainment			Lower	Upper
Other main diagnoses <sup>1)</sup>	Mandatory vs Tertiary	0.05	0.29	-0.04	0.14
	Upper Secondary vs Tertiary	0.04	0.38	-0.04	0.12
Colon Cancer	Mandatory vs Tertiary	0.99	0.02	0.13	1.84
COPD	Upper Secondary vs Tertiary	0.56	0.04	0.02	1.10
Asthma	Mandatory vs Tertiary	0.93	0.04	0.06	1.80
Ischaemic HD	Upper Secondary vs Tertiary	-0.17	0.04	-0.33	-0.01
<b>Main Diagnosis</b>	<b>Hospital insurance class</b>				
Other main diagnoses <sup>1)</sup>	private vs. general	0.36	<0.001	0.18	0.53
	semi-private vs general	0.15	0.03	0.02	0.29
Colon Cancer	semi-private vs general	-1.10	0.00	-1.85	-0.35
Breast Cancer	private vs. general	-0.56	0.01	-0.99	-0.13
	semi-private vs general	-0.36	0.03	-0.67	-0.04
AMI	private vs. general	0.65	0.01	0.20	1.10
	semi-private vs general	0.54	0.00	0.20	0.87
COPD	private vs. general	1.50	0.00	0.49	2.50
	semi-private vs general	0.92	0.01	0.27	1.58
Back problems	private vs. general	0.95	0.00	0.46	1.44
	semi-private vs general	0.61	0.00	0.28	0.93
<b>Main diagnosis</b>	<b>Type of household</b>				
Other main diagnoses <sup>1)</sup>	living alone vs living with others	0.28	<0.001	0.20	0.37
Lung cancer	living alone vs living with others	0.92	0.000	0.46	1.38
Colon Cancer	living alone vs living with others	1.21	0.001	0.49	1.94
Back problems	living alone vs living with others	0.56	0.000	0.30	0.82
<b>Main diagnosis</b>	<b>Langug skills</b>				
Other main diagnoses <sup>2)</sup>	allophone vs. regional language	0.34	<0.001	0.13	0.55
	no regional vs. regional language	0.00	0.98	-0.12	0.13
Colon cancer	allophone vs. regional language	-1.559	0.06	-3.16	0.04
COPD	allophone vs. regional language	-0.749	0.06	-1.54	0.04
Ischaemic HD	no regional vs. regional language	0.345	0.01	0.10	0.59
<b>Main diagnosis</b>	<b>Migration background</b>				
Other main diagnoses <sup>2)</sup>	1st Generation vs Swiss w/o migr.background	0.16	<0.001	0.05	0.27
	2nd Generation vs Swiss w/o mig. background	0.02	0.68	-0.07	0.10
Colon Cancer	2nd Generation vs Swiss w/o mig. background	-1.284	0.005	-2.167	-0.402
Osteoarthritis	1st Generation vs Swiss w/o migr.background	0.019	0.838	-0.163	0.201
Back problems	2nd Generation vs Swiss w/o mig. background	0.387	0.011	0.089	0.685
The least significant difference adjusted significance level is .05.					
<sup>1)</sup> of tabel 4; <sup>2)</sup> cf tabel 5					

**Table S3: Associations of comorbidity with social factors (Linear CCMM)**

Outcome: N° of side diagnosis+	Modell Comorbidity (N=140'903)			
	β (N)	p-value	95% CI	
			Lower	Upper
<b>Fixed Effects*</b>				
Intercept	2.70	0.109	-0.60	6.00
Educational attainment				
Compulsory	0.37	0.000	0.33	0.41
Upper secondary	0.23	0.000	0.19	0.26
Tertiary (Ref.)	0 <sup>b</sup>			
Insurance Class				
Private	0.01	0.863	-0.06	0.07
Semi-private	-0.05	0.115	-0.11	0.01
Mandatory (Ref.)	0 <sup>b</sup>			
Household type				
Living alone	0.22	0.000	0.17	0.26
Living with others (Ref.)	0 <sup>b</sup>			
Sex				
Men	0.23	0.000	0.18	0.27
Women (Ref.)	0 <sup>b</sup>			
Nationality				
Other nationality	0.32	0.000	0.21	0.42
EU/EFTA	0.02	0.507	-0.04	0.08
Swiss	0 <sup>b</sup>			

+ centred by chronic conditions  
 \*Controlling for clustering on hospital- and patient-level and adjusted for age, chronic condition, language region of hospital and year of discharge

β: Difference in average length of hospital stay to the respective reference category estimated from the mixed linear regression model containing all variables listed in the table and random effects for hospitals and patients.

**Table S4: Associations of discharge destination with social factors and factors related to hospital stay (Logistic CCMM)**

Binary Outcome: Transfer to inpatient setting (vs. discharge to home)	Model Discharge Destination (N=140'903)						
	β (N)	95% CI		OR	p-value	95% CI	
		Lower	Upper			Lower	Upper
<b>Fixed Effects*</b>							
Intercept	-1.86	-6.66	2.95	0.16	0.449	0.00	19.11
Educational attainment							
Compulsory	0.01	-0.05	0.07	1.01	0.708	0.95	1.07
Upper secondary	0.00	-0.05	0.05	1.00	0.973	0.95	1.05
Tertiary (Ref.)	0 <sup>b</sup>						
Insurance Class							
Private	-0.08	-0.21	0.04	0.92	0.196	0.81	1.04
Semi-private	-0.10	-0.20	-0.01	0.90	0.030	0.82	0.99
Mandatory (Ref.)	0 <sup>b</sup>						
Household type							
Living alone	0.45	0.40	0.50	1.56	0.000	1.49	1.64
Living with others (Ref.)	0 <sup>b</sup>						
Sex							
Men	-0.29	-0.33	-0.25	0.75	0.000	0.72	0.78
Women (Ref.)	0 <sup>b</sup>						
Nationality							
Other nationality	0.09	-0.02	0.20	1.09	0.116	0.98	1.22
EU/EFTA	0.05	0.00	0.10	1.05	0.046	1.00	1.10
Swiss	0 <sup>b</sup>						
Comorbidity							
NSD (centered by CC)	0.11	0.10	0.13	1.12	0.000	1.10	1.14
Psychic comorbidity: yes	0.34	0.28	0.40	1.40	0.000	1.33	1.49
Psychic comorbidity: no (Ref.)	0 <sup>b</sup>						
Hospital Ward							
Surgical	0.29	0.09	0.49	1.34	0.004	1.10	1.64
Internal medicine or other (Ref.)	0 <sup>b</sup>						
Need of intensive care							
Yes	0.68	0.47	0.90	1.98	0.000	1.60	2.45
No (Ref.)	0 <sup>b</sup>						

\*Controlling for clustering on hospital- and patient-level and adjusted for age, chronic condition, language region of hospital and year of discharge

β: Difference in average length of hospital stay to the respective reference category estimated from the mixed linear regression model containing all variables listed in the table and random effects for hospitals and patients.

**Table S5** Association of length of stay with demographic factors (**age implemented as categorical variable**), social factors, health status and factors related to hospital stay (linear CCMM D)

Outcome: LOS (days)	Model D - age categorical (N=140'903)			
	$\beta$ (days)	p-value	95% CI	
			Lower	Upper
<b>Fixed Effects*</b>				
Intercept	2.25	<0.001	1.34	3.15
Educational attainment				
Compulsory	0.05	0.25	-0.04	0.14
Upper secondary	0.04	0.35	-0.04	0.12
Tertiary	Ref.			
Insurance Class				
Private	0.36	<0.001	0.18	0.54
Semi-private	0.16	0.02	0.02	0.29
Mandatory	Ref.			
Household type				
Living alone	0.28	<0.001	0.20	0.37
Living with others	Ref.			
Sex				
Men	-0.30	<0.001	-0.38	-0.22
Women	Ref.			
Age				
75-84	0.67	<0.001	0.31	1.03
65-74	0.55	<0.001	0.23	0.86
55-64	0.47	<0.001	0.23	0.71
45-54	0.27	<0.001	0.09	0.46
25-44	Ref.			
Nationality				
Other nationality	0.21	0.04	0.01	0.41
EU/EFTA	0.14	0.02	0.02	0.26
Swiss	Ref.			
Chronic Condition (CC)				
Lung cancer	4.81	<0.001	4.38	5.24
Colon cancer	6.97	<0.001	6.27	7.67
Breast cancer	1.53	<0.001	1.17	1.89
Prostata cancer	2.54	<0.001	1.97	3.11
Diabetes w/o complication	4.03	<0.001	3.04	5.02
Diabetes with complication	7.49	<0.001	6.79	8.19
Acute myocardial infarction	0.70	<0.001	0.39	1.02
Acute cerebrovascular diseases	3.92	<0.001	3.41	4.43
Congestive heart failure	6.41	<0.001	5.93	6.90
COPD	5.09	<0.001	4.61	5.57
Asthma	2.00	<0.001	1.63	2.37
Osteoarthritis	2.93	<0.001	2.38	3.49
Back problems	3.50	<0.001	3.00	3.99
Disc disorder	2.51	<0.001	1.93	3.09
Ischämïc heart disease	Ref.			
Comorbidity				
NSD (centered by CC)	0.80	<0.001	0.70	0.89
Psychic comorbidity: yes	0.36	<0.001	0.14	0.57
Psychic comorbidity: no	Ref.			
Hospital Ward				
Surgical	1.72	<0.001	1.24	2.21
Internal medicine or other	Ref.			
Need of intensive care				
Yes	3.15	<0.001	2.78	3.53
No	Ref.			
Discharge destination				
Died in hospital	0.34	0.31	-0.32	1.01
Transfer to inpatient setting	1.99	0.00	1.76	2.22
Discharge to home	Ref.			
Akaike criterion, corrected	868'829			

\*The model controls for clustering on hospital- and on patient-level and is adjusted for language region of hospital and year of discharge

The regression coefficients  $\beta$  are the estimated differences in average length of stay between the respective category and the reference category obtained from the respective model.