New insights from GWAS on longitudinal and cross-sectional BMI and related phenotypes in admixed children with Native American and European ancestries

Lucas Vicuña^{1,2}, Esteban Barrientos^{1,2}, Tomás Norambuena^{1,2}, Danilo Alvares³, Juan Cristobal Gana⁴, Valeria Leiva³, Veronica Mericq⁵, Cristian Meza⁶, Ana Pereira⁷, José L. Santos⁸, Susana Eyheramendy^{1,2*}

1 Faculty of Engineering and Sciences, Universidad Adolfo Ibáñez, Santiago, Chile.

2 Instituto Milenio de Investigación Sobre los Fundamentos de los Datos (IMFD), Santiago, Chile.

3 Department of Statistics, Pontificia Universidad Católica de Chile, Santiago, Chile.

4 Department of Pediatric Gastroenterology and Nutrition. Division of Pediatrics. School of Medicine. Pontificia Universidad Católica de Chile, Santiago, Chile

5 Institute of Maternal and Child Research, Faculty of Medicine, University of Chile, Santiago, Chile.

6 INGEMAT-CIMFAV, Faculty of Engineering, Universidad de Valparaíso, Valparaíso, Chile.

7 Institute of Nutrition and Food Technology, University of Chile, Santiago, Chile.

8 Department of Nutrition, Diabetes and Metabolism, School of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile.

* Corresponding author: susana.eyheramendy@uai.cl

Abstract

Body-mass index (BMI) is a well-known marker of adiposity across all ages. The genetic architecture of BMI has been thoroughly studied among adults. In contrast, there are a few genome-wide association studies (GWAS) on children. Further, GWAS on children have been performed almost exclusively in Europeans at single ages. We aimed to better understand the genetic architecture of BMI trajectory across ages and how BMI is affected by Native American genetic ancestry. We performed cross-sectional and longitudinal GWAS for BMI-related traits on 904 admixed Chilean children with mostly European and Mapuche Native American genetic ancestry. We focused on BMI and two traits that occur at the minimum of the childhood BMI growth trajectory, namely, age at adiposity rebound (Age-AR) and BMI at adiposity rebound (BMI-AR). We found several variants in the immune gene *HLA-DQB3* that are strongly associated with BMI at ages 1.5-2.5 years old, but not at other ages. We also identified a variant in the sex-determining gene *DMRT1* significantly associated with Age-AR ($P = 9.8 \times 10^{-9}$). Further, BMI was significantly higher in Mapuche than in European children at all ages between 5.5 and 16.5 years old, but not before. Finally, Age-AR was significantly lower ($P = 0.013$) by 1.64 years in the Mapuche children compared with Europeans.

Introduction

Childhood obesity is a major public health problem across the world. One well-studied indicator of obesity is the body mass index (BMI), whose alterations in children have been associated with risk of adult obesity, as well as related diseases, including type 2

diabetes [1] and cardiometabolic diseases [2]. Even though several environmental causes explain an important part of childhood obesity, there is still a lack of knowledge about the genetic factors underlying susceptibility to this disease.

BMI follows a nonlinear trajectory over time during childhood. BMI trajectory is characterized by three main phases: i) a rapid increase with an adiposity peak (AP) at ~ 9 month of age; ii) a decline reaching its lowest value at ~ 5.5 years, also called the adiposity rebound (AR); and iii) a subsequent peak at early adulthood [3].

The heritability of adult obesity has been estimated to be 40 – 70% [34]. Approximately 22.4% of adult BMI heritability is explained by 941 SNPs, according to a genome-wide association study (GWAS) meta-analysis performed in $\sim 700,000$ individuals of European ancestry [5]. Such studies have been mostly performed in adults, but for most of these loci it is unknown whether or not and to which extent they also affect BMI in children. Some cross-sectional GWA studies on European populations have shown that several loci influence BMI in adulthood and childhood, whereas other loci seem to act only during adulthood or childhood. For instance, a GWAS found variants in the *FTO* and *MC4R* genes significantly associated with BMI in children (< 6 years old) as well as in adults, but a locus in *MAF* only associated in adults [6]. A GWAS meta-analysis found significant associations with *FTO*, *MC4R*, *TMEM18*, *SDCCAG8* and *TNKS/MSRA* loci, but associations for the later gene were limited to children and adolescents [7]. Another GWAS meta-analysis identified 15 loci significantly associated with BMI (they considered a single BMI value at the oldest age between 2 and 10 years). Among them, 12 were previously associated with adult BMI or childhood obesity, suggesting that the 3 remaining loci, located near *ELP3*, *RAB27B* and *ADAM23* genes, act only during childhood [8]. A further GWAS meta-analysis identified 18 SNPs significantly associated with BMI in paediatric cohorts from diverse ancestries pooled together, although most of these loci only reached the genome-wide significance threshold in Europeans when same-ancestry cohorts were analyzed separately. Most of these loci had been previously associated with adult BMI, childhood BMI or childhood obesity [9].

A few GWAS on European children have identified loci affecting BMI in distinct phases of child growth. A recent GWAS meta-analysis identified 4 loci significantly associated with childhood BMI-related traits: one *LEPR/LEPROT* gene variant associated with BMI-AP; one *FTO* variant and one *TFAP2B* variant associated with Age-AR; and a *GNPDA2* variant associated with BMI-AR. Among them, the *TFAP2B*, *GNPDA2* and *FTO* variants are also associated with adult BMI, whereas the *LEPR/LEPROT* variant only associates with a BMI-related trait in the infant phase. The authors suggested that adult BMI-related variants start influencing BMI by the time of AR [3]. A candidate gene study on a European multi-cohort found that a *FTO* locus is positively associated with BMI from 5.5 years onward, but is inversely associated below age 2.5 years old. They also confirmed this findings using a longitudinal linear mixed model [13]. Warrington *et al* 2015 performed a longitudinal GWAS to identify associations between genotypes and BMI trajectories across childhood [14]. They found 4 loci previously associated with BMI in adults or children as well as a novel locus in the *FAM120AOS* gene significantly associated with BMI at 8 years of age. To our knowledge, Warrington *et al* 2015 is the only longitudinal GWAS on childhood BMI.

As mentioned before, the vast majority of these GWA studies have been performed in Europeans, but we do not know to what extent these loci affect BMI in populations with other continental ancestries, in particular those with Native American ancestry. This is relevant, as obesity and obesity-related disorders are markedly affected by genetic ancestry [34]. For instance, Latinos, which usually have mixed European, Native American and African ancestries, are more susceptible to lipid-related disorders than any other US group, in part due to their Native-American genetic heritage [15].

The aims of this study are twofold: i) to estimate how much does Native American ancestry affect childhood BMI trajectory, Age-AR as well as BMI-AR; and ii) to identify genetic variants associated with these traits.

Materials and methods

Sample collection and genotyping.

We analyzed individuals of the “Growth and Obesity Chilean Cohort Study” (GOCS; [22]), who were genotyped using the Infinium® Multi-Ethnic Global BeadChip (Illumina), as described in [21]. After quality controls were applied, we obtained a filtered data set of 904 individuals and 774,433 autosomal SNPs. Further details can be found in [21].

Weight and height measurements.

Weight and height measurements were taken since 2006 at Instituto de Nutrición y Tecnología de los Alimentos (INTA), Santiago, Chile, on children/adolescents in barefoot and light clothes by a trained dietitian following standardized protocols; ICC for all measurements was ≥ 0.75 [41]. Weight and height were measured twice at each individual visit, and the average was considered as the final value. Before and after puberty, weight and height were measured once per year; during puberty they were measured every 6 months. Weight was measured with a portable electronic scale (Seca 770), precision of 0.1 kg, and height was measured with a portable stadiometer (Harpender 603) to the nearest 0.1 cm. BMI (kg/m^2) and Z-scores (for < 18 years old) were calculated based on the World Health Organization 2007 growth references. Weight and height data prior to 2006 were retrieved from medical records.

Local Ancestry Estimation.

We used RFMIX [48] to infer the local ancestry of each SNP allele from our Chilean sample. We used reference populations from the 1000 Genomes Project [49], namely, Yoruba (YRI, $n=108$) for the African ancestry, Utah Residents with Northern and Western European Ancestry (CEU, $n=36$) for the European ancestry, and Peruvian (PEL) individuals with $> 95\%$ Native American ancestry ($n=29$). We excluded individuals with $> 5\%$ SNP missing rate. We inferred the gametic phase of individuals with the Beagle 5 software [24], using the HapMap37 human genome build 37 recombination map. To obtain the local ancestry, we used the `--forward-backward` parameter, as recommended in the manual. Local ancestry estimation identified three continental ancestries: European, African and Native American.

Global Ancestry Estimation.

Global ancestry proportions of Chilean children were estimated with Admixture 1.3 [23], using $K=4$ ancestral populations. This K value was chosen because it was able to distinguish the two main Native American subcomponents of Chileans, namely, Mapuche and Aymara [21]. In order to distinguish Peruvian (PEL) individuals with $> 95\%$ Native American ancestry, we used $K=3$ ancestral populations and the reference populations mentioned in the previous section. Global ancestry estimation identified four proportion of ancestries corresponding to European, African, Aymara and Mapuche. We used the Mapuche proportion ancestry in the GWAS to identify its effect on the phenotypes.

Variant and gene annotations.

Variant annotations [GRCh37 (hg19) assembly] were retrieved with the web tool Variant Effect Predictor (VEP) from Ensembl [25]. Upstream and downstream variants were defined as those located 10 Kb upstream or downstream of the gene, respectively. Intergenic variants were defined as those located > 100 Kb upstream or downstream of the closest gene. We used HaploReg v4.1 [26] to identify the genes located closest to associated intergenic variants. Reported GWAS associations were retrieved from the NHGRI GWAS Catalog [27]. When more than one variant in a gene has been associated with the same phenotype, we reported the strongest association.

Derivation of Age-AR and BMI-AR

Before modeling BMI trajectory over time, we applied some filters on the individuals. We excluded 51 individuals who had no sufficient measurements between 2 and 10 years old. We also excluded 156 individuals with all their measurements taken after 4.5 years old. We obtained a final dataset of 696 individuals with measurements between 2 and 10 years old. To estimate Age-AR, we implemented a longitudinal statistical model of BMI (in logarithm scale) on this filtered data. The model equation is:

$$\begin{aligned} \log(\text{BMI})_{ij} = & (\beta_1 S_i + \beta_0) + (\beta_2 S_i + \beta_3) \text{Age}_{ij} + (\beta_4 S_i + \beta_5) \text{Age}_{ij}^2 + (\beta_6 S_i + \beta_7) \text{Age}_{ij}^3 + \\ & + \beta_8 i + \beta_9 i \text{Age}_{ij} + \beta_{10i} \text{Age}_{ij}^2 + \beta_{11i} \text{Age}_{ij}^3 + \epsilon_{ij} \\ & j = 1, 2, \dots, n_i \text{ and } i = 1, \dots, 696. \end{aligned} \quad (1)$$

where n_i corresponds to the number of Age/BMI measurements for individual i , S_i represents the gender of the individual (0 female, 1 male). The parameters β_k for $k = 1, \dots, 7$ are fixed effects whereas the parameters β_{ki} for $k = 8, \dots, 11$ are random effects for each individual i (i.e. $\beta_{ki} \sim N(0, \sigma_k^2)$). The errors $\{\epsilon_{ij}\}$ are assumed independent across the different individuals i but dependent between observations for the same individual (index j), and are normally distributed with variance σ^2 .

The predicted trajectory for individual i can then be written as

$$\begin{aligned} \log(\hat{\text{BMI}})_{ij} = & (\hat{\beta}_1 S_i + \hat{\beta}_0) + (\hat{\beta}_2 S_i + \hat{\beta}_3) \text{Age}_{ij} + (\hat{\beta}_4 S_i + \hat{\beta}_5) \text{Age}_{ij}^2 + (\hat{\beta}_6 S_i + \hat{\beta}_7) \text{Age}_{ij}^3 + \\ & + \hat{\beta}_8 i + \hat{\beta}_9 i \text{Age}_{ij} + \hat{\beta}_{10i} \text{Age}_{ij}^2 + \hat{\beta}_{11i} \text{Age}_{ij}^3 \\ & j = 1, 2, \dots, n_i \text{ and } i = 1, \dots, 696. \end{aligned} \quad (2)$$

where $\hat{}$ (hat) represents the estimators of the parameters. From this trajectory, the Age of Adiposity rebound (Age-AR) and the BMI at Age-AR are obtained by finding the Age at which the minimum BMI is found.

A local minimum of BMI is found by solving the following equation:

$$3(\hat{\beta}_6 S_i + \hat{\beta}_7 + \hat{\beta}_{11i}) x_i^2 + 2(\hat{\beta}_4 S_i + \hat{\beta}_5 + \hat{\beta}_{10i}) x_i + (\hat{\beta}_2 S_i + \hat{\beta}_3 + \hat{\beta}_9 i) = 0 \quad (3)$$

which has as its solution:

$$\begin{aligned} \text{Age-AR}_i = & \frac{1}{3(\hat{\beta}_6 S_i + \hat{\beta}_7 + \hat{\beta}_{11i})} (-(\hat{\beta}_4 S_i + \hat{\beta}_5 + \hat{\beta}_{10i}) + \\ & \pm 2\sqrt{(\hat{\beta}_4 S_i + \hat{\beta}_5 + \hat{\beta}_{10i})^2 - 3(\hat{\beta}_6 S_i + \hat{\beta}_7 + \hat{\beta}_{11i})(\hat{\beta}_2 S_i + \hat{\beta}_3 + \hat{\beta}_9 i)}) \end{aligned} \quad (4)$$

where the minimum obtained above needs to satisfy:

$$6(\hat{\beta}_6 S_i + \hat{\beta}_7 + \hat{\beta}_{11i}) \text{Age-AR}_i + 2(\hat{\beta}_4 S_i + \hat{\beta}_5 + \hat{\beta}_{10i}) > 0 \quad (5)$$

To identify BMI-AR, we replace the Age term from equation (1) by the Age-AR value obtained from equation (3).

Cross-sectional GWAS on BMI.

For the cross-sectional GWAS, we analyzed individuals between 1.5 and 16.5 years old, across 16 strata of 1 year interval each (S1 Table). BMI was expressed as kg/m² and age in years. Individuals with no weight or height measurements in a particular stratum were removed. When > 1 measurement was taken in the same individual and time interval, we took the mean of the BMI measurements. To get a normal distribution of BMI values, we applied Box-Cox transformations (λ values for each stratum are shown in S1 Table). These transformed values were standardized to get mean = 0 and standard deviation = 1, obtaining zBMI scores [13] [14]. These zBMI scores were used to test for genome-wide associations. We applied the following linear regression model:

$$zBMI_i = \beta_0 + \beta_1 GT_{ij} + \beta_2 S_i + \beta_3 GA_i + \beta_4 LA_{ij} + \epsilon_i \quad (6)$$

where β_1 represents the effect of the additive genotype (GT) of SNP j ; β_2 represents the effect of gender (S); β_3 represents the effect of the global Mapuche Native American ancestry proportion (GA); and β_4 represents the effect of local Native American ancestry, where LA_j takes the values 0, 1 or 2 depending on the number of Native American alleles. Of note, while our global ancestry analysis is able to separately estimate the Mapuche and Aymara ancestral Native American subcomponents of Chileans, the local ancestry analysis only considers a general Native American population.

GWAS on longitudinal BMI data.

To estimate longitudinal genotype-phenotype associations, we implemented a linear mixed model for BMI. We adjusted for gender, age, global ancestry, and local ancestry. We assumed an additive inheritance model. The model equation is as follows:

$$\begin{aligned} \log(\text{BMI})_{ij} = & (\beta_1 S_i + \beta_0) + (\beta_2 S_i + \beta_3) \text{Age}_{ij} + \beta_4 \text{Age}_{ij}^2 + \beta_5 \text{Age}_{ij}^3 + \beta_6 GA_i + \\ & + \beta_7 LA_i + \beta_8 GT_i + \beta_9 i + \beta_{10i} \text{Age}_{ij} + \beta_{11i} \text{Age}_{ij}^2 + \epsilon_{ij} \quad (7) \\ & j = 1, 2, \dots, m_i \text{ and } i = 1, \dots, 901. \end{aligned}$$

where m_i corresponds to the number of Age/BMI measurements for individual i , S_i represents the gender of the individual (0 female, 1 male).

GWAS on Age-AR and BMI-AR.

Similarly as with BMI, we obtained standardized Age-AR and BMI-AR values by applying Box-Cox transformations followed by standardization. We obtained estimates of $\lambda = 0.841$ for Age-AR and $\lambda = -0.759$ for BMI-AR. We performed genome-wide associations for standardized Age-AR, using the following linear regression model:

$$\begin{aligned} \text{zAge-AR}_i = & (\beta_1 S_i + \beta_0) + (\beta_2 S_i + \beta_3) \text{zBMI-AR}_i + (\beta_4 S_i + \beta_5) \text{zBMI-AR}_i^2 + \\ & (\beta_6 S_i + \beta_7) \text{zBMI-AR}_i^3 + \beta_8 G A_i + (\beta_9 S_i + \beta_{10}) G T_i + \beta_{11} L A_i + \epsilon_i \end{aligned} \quad (8)$$

where β_9 represents the interaction effect between gender and genotype; $(\beta_2 S_i + \beta_3), (\beta_4 S_i + \beta_5), (\beta_6 S_i + \beta_7)$ = interaction effect between sex and a third degree polynomial of zBMI-AR. For the GWAS on BMI-AR, we used a similar model:

$$\begin{aligned} \text{zBMI-AR}_i = & (\beta_1 S_i + \beta_0) + (\beta_2 S_i + \beta_3) \text{zAge-AR}_i + (\beta_4 S_i + \beta_5) \text{zAge-AR}_i^2 + \\ & (\beta_6 S_i + \beta_7) \text{zAge-AR}_i^3 + \beta_8 G A_i + (\beta_9 S_i + \beta_{10}) G T_i + \beta_{11} L A_i + \epsilon_i \end{aligned} \quad (9)$$

Results

Estimation of Age-AR and BMI-AR

We first characterized BMI trajectories in 904 children from the “Growth and Obesity Chilean Cohort Study” (GOCS) [22], from age 2 years to age 10 years (Fig 1). We implemented longitudinal mixed models (see details in Methods) in boys and girls pooled together as well as separately. Fig 1 shows the expected BMI trajectories for the three analyses. We observed a strong fit between observed and expected curves, as revealed by a Pearson’s correlation coefficient of $\rho = 0.97$. Using the same longitudinal model, we estimated BMI-AR and Age-AR (see Methods). The mean Age-AR was 4.5 (SD = 1.5) years in pooled children, 4.35 (SD = 1.35) years in boys and 4.55 (SD = 1.71) years in girls. The mean BMI-AR was 16.3 (SD = 1.4) kg/m² in pooled children, 16.4 (SD = 1.29) kg/m² in boys and 16.2 (SD = 1.42) kg/m² in girls.

Effect of global ancestry on BMI, Age-AR and BMI-AR.

The children of this cohort have on average 52.1% European, 43.8% Mapuche Native American, 2.6% Aymara Native American, and 1.5% African global ancestry proportions [21] (S1 Fig). To quantify how much Mapuche and European ancestries affect BMI at each age stratum, we applied linear regressions with BMI as the dependent variable and global Mapuche ancestry as the independent variable. We estimated BMI values for hypothetical individuals with 100% Mapuche ancestry and with 100% European ancestry, hereafter referred to as “Mapuche” and “Europeans”. When compared with Europeans, Mapuche individuals have significantly higher BMI at all age strata between 5.5 and 16.5 years old, but not before 5.5 years old (Fig 2A). Indeed, intercept P -values were almost all < 0.01 except at 7.5-8.5 years old ($P < 0.05$; Fig 2A and S1 Table). Further, BMI differences between Mapuche and European individuals tend to increase with age (Fig 2A). Similarly, global ancestry had a mild but significant effect in the whole longitudinal BMI trajectory ($P = 0.045$; effect size = 0.094).

We explored whether individual Mapuche and European global ancestries have an effect on Age-AR and BMI-AR. By regressing Age-AR onto Mapuche global ancestry, our model predicted that Age-AR is significantly lower by 1.64 years in a Mapuche child (mean = 3.53 years) than in a European child (mean = 5.17 years; $P = 0.013$; Fig 2B). On the other hand, BMI-AR was 16.9 and 15.9 kg/m² in a Mapuche and a European child, respectively (Fig 2C). However, this difference was not significant.

Identification of variants associated with BMI.

We performed GWAS on BMI in each age stratum by performing linear regression models (see Methods for details; S1 Table shows the number of individuals in each

Fig 1. Estimation of Age-AR and BMI-AR. Fitted BMI (kg/m^2) curves over time (years) in the whole cohort as well as in boys and girls considered separately. Vertical and horizontal dotted lines pinpoint the observed mean Age-AR and mean BMI-AR in boys and girls.

stratum). Besides gender and age, we adjusted for the global ancestry proportion of each individual to account for population substructure as well as for local ancestry to adjust for the ancestry of each haplotype (i.e. SNP). S3 Fig S4 Fig S5 Fig S6 Fig S7 Fig S8 Fig S9 Fig S10 Fig S11 Fig S12 Fig S13 Fig S14 Fig S15 Fig S16 Fig S17 Fig show the corresponding Manhattan plots. We found a significant association for the *rs75964957* variant at 4.5 - 5.5 years old stratum ($P < 3.5 \times 10^{-8}$; S1 Table). The second strongest association ($P < 1.6 \times 10^{-7}$) was achieved by a genetic signal at chromosome 6 harboring several variants in high linkage disequilibrium, as revealed by a clear peak in the Manhattan plot of Fig 3A. This genetic signal was apparent in the 1.5 - 2.5 years old stratum, but not at other strata (Fig 3B).

The strongest variants of this peak, namely, *rs9275582*, *rs9275593* and *rs9275595*, map a promoter flanking region of the *HLA-DQB3* pseudogene, which is part of the human leukocyte antigen (HLA) region (see Discussion). S2 Table shows all variants from the HLA peak achieving nominal genome-wide significance ($P < 1 \times 10^{-6}$), with their corresponding annotations. We identified additional association peaks at chromosomes 4 and 20 (S18 Fig). The strongest peak variant at chromosome 4, *rs12501266*, maps the gene *SORCS2*, whereas the strongest peak variant at chromosome 20, *rs474169*, maps the gene *SNAP25-AS1*. Of note, *rs474169* is also associated with BMI at age stratum 10.5 - 11.5 (S1 Table). In contrast to the HLA peak, peak variants at chromosomes 4 and 20 show associations across several strata during childhood and/or adolescence (S18 Fig).

In order to identify variants associated with the whole BMI trajectory during body growth, we performed a longitudinal GWAS on BMI (see Methods for details). The strongest association was achieved by *rs35266519* ($P = 2.9 \times 10^{-7}$), which is a missense variant of the *GSDMB* gene. This gene has been GWAS-associated with

Fig 2. Effect of global Mapuche ancestry on BMI trajectory, Age-AR and BMI-AR. A: BMI (kg/m²) estimated at each age stratum for predicted individuals with 100% Mapuche (red line) and 100% European global ancestry (blue line). B: Linear model of Age-AR with Mapuche ancestry as covariate. C: Linear model of BMI-AR with Mapuche ancestry as covariate.

obesity-related traits (see Discussion). [S19 Fig](#) shows the corresponding Manhattan plot, whereas [S3 Table](#) shows the associations achieving nominal genome-wide significance ($P < 1 \times 10^{-6}$), as well as their annotations.

Identification of variants associated with Age-AR and BMI-AR.

To identify variants associated with Age-AR, we implemented a regression model described in the Methods section. We detected one variant significantly associated with Age-AR ($P < 9.8 \times 10^{-9}$; [S2 Table](#) and [S20 Fig](#)). This variant, namely *rs445398*, maps an intron of the *DMRT1* gene. *DMRT1* is a key gene involved in sex differentiation (see Discussion). In addition, we identified 33 variants achieving nominal significance ($P < 1 \times 10^{-6}$; [S2 Table](#)).

We also performed a GWAS for BMI-AR (see Methods for details). We detected 8 variants achieving nominal significance ([S3 Table](#)). [S21 Fig](#) shows the Manhattan plot.

SNP ID-Allele	Location	Consequence	Gene	β -GT	P-GT	Age stratum	Frequency
rs75964957-G	1:239032385	intergenic	-	1.56	4.7E-08	4.5-5.5	0.0077
rs269511-A	5:114688671	intergenic	-	-0.27	2.2E-07	1.5-2.5	0.4646
rs9275582-T	6:32770853	promoter flanking region	HLA-DQB3	0.30	2.5E-07	1.5-2.5	0.2865
rs9275593-A	6:32771630	promoter flanking region	HLA-DQB3	0.30	2.5E-07	1.5-2.5	0.2865
rs9275595-C	6:32772142	promoter flanking region	HLA-DQB3	0.30	2.5E-07	1.5-2.5	0.2865
rs7134291-A	12:14128322	intron	GRIN2B	-0.33	3.0E-07	8.5-9.5	0.1726
rs7896870-T	10:127827684	intron	ADAM12	0.25	4.0E-07	0.5-1.5	0.4906
rs474169-T	20:10081800	intron/NCT	SNAP25-AS1	-0.29	4.8E-07	10.5-11.5	0.3031
rs1495271-C	15:101962723	intron	PCSK6	0.26	6.0E-07	4.5-5.5	0.4060
rs11244839-G	10:127834046	intron	ADAM12	0.25	6.7E-07	0.5-1.5	0.4751
rs13257360-G	8:6117252	upstream	RP11-124B13.1	0.28	7.2E-07	14.5-15.5	0.3335
rs10150218-A	14:73174338	intron	DPF3	0.35	8.3E-07	5.5-6.5	0.4629
rs77468591-T	10:12937704	downstream	CCDC3	-1.84	9.2E-07	4.5-5.5	0.0039

Table 1. Top cross-sectional associations for BMI. Shown is the SNP rs ID with the associated allele, physical location, SO consequence/feature type, gene, effect size of the genotype (β_{GT}), association P -value of the genotype (P_{GT}), age stratum and frequency of associated allele. NCT: noncoding transcript variant.

SNP ID-Allele	Location	Consequence	Gene	β -GTxSex	P-GTxSex	Frequency
rs445398-C	9:954336	intron	DMRT1	5.81	9.8E-09	0.165
rs73952010-G	18:37721370	downstream	RPL17P45	-5.45	7.3E-08	0.007
rs78106261-G	6:110655312	intron	METTL24	-5.13	3.8E-07	0.008
rs73139959-T	3:13815160	intergenic	-	-4.97	8.4E-07	0.003
rs969092-G	12:42234778	intron/NCT	RP11-630C16.2	4.96	8.9E-07	0.238
rs11066997-T	12:114661524	intergenic	-	4.94	9.9E-07	0.358

Table 2. Strongest associations for Age-AR. Shown is the SNP rs ID with the associated allele, physical position in the chromosome, consequence type, gene, effect size of the interaction between genotype and sex (β_{GTxSex}) with the corresponding association P -value (P_{GTxSex}), and the frequency of the associated allele. NCT: noncoding transcript variant.

Discussion

In this study, we investigated the genetic architecture of BMI-related traits on admixed children with mainly European and Mapuche Native American ancestry. There are a few studies that have analyzed the genetic architecture of longitudinal growth traits during childhood and adolescence [6, 7, 8, 9, 14], and most of them have been performed in European populations. To our knowledge, Vicuña *et al.*, 2021 [21] is the only study that has quantified the relationship between Native American genetic ancestry and growth traits. The authors showed that the pubertal age where maximum height growth occurs (peak height velocity), is significantly older by 0.73 years in Europeans than in Mapuche adolescents on average. We are the first to quantify how Native American ancestry affects childhood growth traits. One of our most important findings is the observation that increases in Mapuche ancestry associate with significant increases in BMI between 5.5 and 16.5 years old as well as with BMI trajectory between 2 – 10 years old. Moreover, the difference in BMI between Mapuche and Europeans

Fig 3. Genome-wide associations with BMI. A: GWAS at age stratum 1.5 – 2.5. B: Association peak of chromosome 6 across age strata. The region shown has physical coordinates 6 : 32676320 – 32683820 and was centered at the variant with the strongest association.

increases steadily after 5.5 years old. This makes sense, since it is known that the contribution of heritable genetic variation to BMI increases from childhood (4 years) to young adulthood (19 years) [11]. We did not find significant differences at age stratum 4.5 – 5.5 years old, likely due to a lack of statistical power derived from the sample size of our cohort. Another important finding was the significantly lower Age-AR in the Mapuche compared with Europeans on average. A higher adiposity in the Mapuche during childhood could explain in part why this population is particularly susceptible to developing metabolic disorders such as insulin resistance, obesity, cholesterol gallstones and metabolic syndrome during adulthood, since all of these disorders are strongly associated with elevated lipid levels [20]. Moreover, Mapuche ancestry among Chileans is distinctly associated with heart diseases, hypertension and diabetes mellitus [19]. Further, the prevalence of type II diabetes and obesity among the Mapuche increased significantly following the change from a rural to an urban life style [33], suggesting that

genotype-environment interactions may lead to a higher genetic susceptibility to developing cardiometabolic diseases.

It is unknown whether the effect of Mapuche genetic ancestry on BMI-related traits was originated by random genetic drift in ancient Native American or European populations, by adaptation to selective pressures (see [16]) or by a combination of adaptation, drift and/or other evolutionary forces. It is also possible that environmental factors could partially contribute to such differences. However, the children of our cohort belong to the same urban district in Santiago and to the same middle-lower socioeconomic group. Therefore, we do not expect high variation in environmental factors such as nutrition, exposure to pollutants or incidence of pathogenic diseases. However, it is worth noting that in this same cohort, mothers of boys with high Mapuche ancestry (≥ 3 Mapuche last names: 4.8% of total boys) have a lower educational level than mothers of boys with low Mapuche ancestry (0 Mapuche last names: 80% of boys) [35], which in turn could potentially have an effect on their son's BMI.

Our GWAS on BMI detected a significant association for the intergenic variant *rs75964957* at age stratum 4.5 – 5.5 years. This was despite the sample size at this stratum ($n = 708$). The second stronger association was for the *rs9275582*, *rs9275593* and *rs9275595* variants, which map a promoter flanking region of the *HLA-DQB3* unprocessed pseudogene. Noteworthy, *rs9275595* has been previously associated with BMI in adults ($P = 6 \times 10^{-6}$; 0.016 kg/m² increase; [28]). These three variants constitute the top of an association peak that was observed at age stratum 1.5 – 2.5 years, but not at other age strata. The *HLA-DQB3* gene belongs to the human leucocyte antigen (HLA) gene cluster, which harbors hundreds of genes that are fundamental for immune function. HLA genes encode proteins of the major histocompatibility complex, which present antigenic peptides to immune cells, in order to distinguish between “self” and “non-self” agents [29]. It is unknown why these *HLA-DQB3* variants have such a distinct effect on BMI at 1.5 – 2.5 years old, but the observation that they map a promoter flanking region suggests that they affect early BMI by regulating the expression of *HLA-DQB3* or additional HLA genes in high linkage disequilibrium with *HLA-DQB3*. Future studies will be needed to validate this finding. It is worth noting that the HLA region shows high variability among ethnic groups [37] and that the HLA-DR/DQ region is the major determinant of susceptibility to childhood type 1 diabetes [38]. The strongest association of the longitudinal GWAS on BMI was for *rs35266519*, a missense variant of the *GSDMB* gene. Among other phenotypes, *GSDMB* variants have been GWAS-associated with type 1 diabetes ($P < 6 \times 10^{-13}$) [39] and subcutaneous adipose tissue ($P < 3 \times 10^{-8}$) [40], two traits that are strongly related with obesity.

We identified a variant of the *DMRT1* gene significantly associated with Age-AR. Remarkably, this association was captured using a sample size of $n = 696$. *DMRT1* is a hallmark gene involved in sex differentiation by maintaining the fates of testes or ovaries in adult mammals [30]. Indeed, deletion or inactivation of *DMRT1* in humans causes XY male-to-female sex reversal [31]. Thus, it is possible that genetic variation in this gene could affect endocrine mechanisms of infant growth. Moreover, this effect on infant growth may lead to an altered pubertal timing, since it has been observed that Mapuche boys have a higher incidence of precocious puberty [35].

We could not replicate our findings in an independent cohort, due to the lack of longitudinal paediatric growth cohorts with enough longitudinal BMI measurements and relatively high Native American ancestry. To our knowledge, the Salvador-SCAALA cohort from Brazil [32] and the COIPIS cohort from Mexico [10] are the only paediatric cohorts with genome-wide genotype information and Native American admixture. However, these cohorts present severe limitations for replication. In the

Salvador-SCAALA cohort, the mean Native American global ancestry is too low (6%), in contrast to high mean African and European ancestries (51% and 46%, respectively; [32]). Also, in this cohort only a single BMI measurement per individual was taken in a broad age window (4 – 11 years old). Regarding the COIPIS cohort, while the level of Native American ancestry is appropriate (36%; [10]), it has a similar problem, namely, that only a single BMI measurement per individual was taken in children aged 3 – 18 years old.

Conclusion

Our study contributes to better understand the biological mechanisms underlying BMI and Age-AR differences between Native Americans and Europeans. Also, it highlights the importance of performing serial cross-sectional GWAS for complex longitudinal traits like BMI trajectory, since longitudinal studies alone might not be well-suited to capture effects that occur at short time intervals. Finally, our results have medical relevance, since they could be useful to identify genetic risk factors for infant obesity based on the individual's proportion of Native American genetic ancestry.

Consent to Participate

Informed consents from participants were obtained from parents or guardians. Children agreed to participate when they turned 7 years old. This study was approved by the Scientific Ethics Committees of Instituto de Nutrición y Tecnología en Alimentos (INTA) and Pontificia Universidad Católica de Chile.

Acknowledgments

This work was supported by the Fondo Nacional de Ciencia y Tecnología (FONDECYT) [1200146 to S.E., L.V. and E.B.; 1190346 to V.M.; 1150416 and 1150486 to J.L.S., 1190801 to C.M.]. S.E., T.N. and L.V. were additionally supported by the Instituto Milenio de Investigación Sobre los Fundamentos de los Datos (IMFD). S.E. conceived the project. S.E. and L.V. designed experiments. E.B., L.V., T.N., C.M., D.A. and V.L. analyzed the data. A.P. and V.M. collected phenotype data. S.E., J.L. S. and J.C. G. raised funds for the genotyping data. L.V., S.E., A.P. E.B. wrote the manuscript. All authors critically reviewed and accepted the final version.

References

1. Can, B. & Can, B. Change in overweight from childhood to early adulthood and risk of type 2 diabetes. *N Engl J Med* **378**, 2537 (2018).
2. Sovio, U. *et al.* How do changes in body mass index in infancy and childhood associate with cardiometabolic profile in adulthood? findings from the northern finland birth cohort 1966 study. *Int J Obes (Lond)* **38**, 53–9 (2014).
3. Couto Alves, A. *et al.* Gwas on longitudinal growth traits reveals different genetic factors influencing infant, child, and adult bmi. *Sci Adv* **5**, eaaw3095 (2019).
4. Llewellyn, C. H., Trzaskowski, M., van Jaarsveld, C. H. M., Plomin, R. & Wardle, J. Satiety mechanisms in genetic risk of obesity. *JAMA Pediatr* **168**, 338–344 (2014).

5. Yengo, L. *et al.* Meta-analysis of genome-wide association studies for height and body mass index in ~700000 individuals of european ancestry. *Hum Mol Genet* **27**, 3641–3649 (2018). 354
355
356
6. Meyre, D. *et al.* Genome-wide association study for early-onset and morbid adult obesity identifies three new risk loci in european populations. *Nat Genet* **41**, 157–9 (2009). 357
358
359
7. Scherag, A. *et al.* Two new loci for body-weight regulation identified in a joint analysis of genome-wide association studies for early-onset extreme obesity in french and german study groups. *PLoS Genet* **6**, e1000916 (2010). 360
361
362
8. Felix, J. F. *et al.* Genome-wide association analysis identifies three new susceptibility loci for childhood body mass index. *Hum Mol Genet* **25**, 389–403 (2016). 363
364
365
9. Bradfield, J. P. *et al.* A trans-ancestral meta-analysis of genome-wide association studies reveals loci associated with childhood obesity. *Hum Mol Genet* **28**, 3327–3338 (2019). 366
367
368
10. Costa-Urrutia, P. *et al.* Genome-wide association study of body mass index and body fat in mexican-mestizo children. *Genes (Basel)* **10** (2019). 369
370
11. Dubois, L. *et al.* Genetic and environmental contributions to weight, height, and bmi from birth to 19 years of age: an international study of over 12,000 twin pairs. *PLoS One* **7**, e30153 (2012). 371
372
373
12. Jacob, R. *et al.* The role of eating behavior traits in mediating genetic susceptibility to obesity. *Am J Clin Nutr* **108**, 445–452 (2018). 374
375
13. Sovio, U. *et al.* Association between common variation at the fto locus and changes in body mass index from infancy to late childhood: the complex nature of genetic association through growth and development. *PLoS Genet* **7**, e1001307 (2011). 376
377
378
379
14. Warrington, N. M. *et al.* A genome-wide association study of body mass index across early life and childhood. *Int J Epidemiol* **44**, 700–12 (2015). 380
381
15. Ko, A. *et al.* Amerindian-specific regions under positive selection harbour new lipid variants in latinos. *Nat Commun* **5**, 3983 (2014). 382
383
16. NEEL, J. V. Diabetes mellitus: a "thrifty" genotype rendered detrimental by "progress"? *Am J Hum Genet* **14**, 353–62 (1962). 384
385
17. Amorim, C. E. *et al.* Genetic signature of natural selection in first americans. *Proc Natl Acad Sci U S A* **114**, 2195–2199 (2017). 386
387
18. Eyheramendy, S., Martinez, F. I., Manevy, F., Vial, C. & Repetto, G. M. Genetic structure characterization of chileans reflects historical immigration patterns. *Nat Commun* **6**, 6472 (2015). 388
389
390
19. Lorenzo Bermejo, J. *et al.* Subtypes of native american ancestry and leading causes of death: Mapuche ancestry-specific associations with gallbladder cancer risk in chile. *PLoS Genet* **13**, e1006756 (2017). 391
392
393
20. Miquel, J. Salud y enfermedad en nuestras poblaciones originarias. *Rev Chil Cardiol* **31**, 129–133 (2012). 394
395

21. Vicuña, L. *et al.* Novel loci and mapuche genetic ancestry are associated with pubertal growth traits in chilean boys. *Hum Genet* (2021). 396
397
22. Pereira, A. *et al.* Breast bud detection: a validation study in the chilean growth obesity cohort study. *BMC Womens Health* **14**, 96 (2014). 398
399
23. Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in unrelated individuals. *Genome Res* **19**, 1655–64 (2009). 400
401
24. Browning, S. R. & Browning, B. L. Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. *Am J Hum Genet* **81**, 1084–97 (2007). 402
403
404
25. McLaren, W. *et al.* The ensembl variant effect predictor. *Genome Biol* **17**, 122 (2016). 405
406
26. Ward, L. D. & Kellis, M. Haploreg: a resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants. *Nucleic Acids Res* **40**, D930–4 (2012). 407
408
409
27. Welter, D. *et al.* The nhgri gwas catalog, a curated resource of snp-trait associations. *Nucleic Acids Res* **42**, D1001–6 (2014). 410
411
28. Locke, A. E. *et al.* Genetic studies of body mass index yield new insights for obesity biology. *Nature* **518**, 197–206 (2015). 412
413
29. Lokki, M.-L. & Paakkanen, R. The complexity and diversity of major histocompatibility complex challenge disease association studies. *HLA* **93**, 3–15 (2019). 414
415
416
30. Huang, S., Ye, L. & Chen, H. Sex determination and maintenance: the role of dmrt1 and foxl2. *Asian J Androl* **19**, 619–624 (2017). 417
418
31. Raymond, C. S. *et al.* A region of human chromosome 9p required for testis development contains two genes related to known sexual regulators. *Hum Mol Genet* **8**, 989–96 (1999). 419
420
421
32. Scliar, M. O. *et al.* Admixture/fine-mapping in brazilians reveals a west african associated potential regulatory variant (rs114066381) with a strong female-specific effect on body mass and fat mass indexes. *Int J Obes (Lond)* **45**, 1017–1029 (2021). 422
423
424
425
33. Pérez-Bravo, F. *et al.* Prevalence of type 2 diabetes and obesity in rural mapuche population from chile. *Nutrition* **17**, 236–8 (2001). 426
427
34. Stryjecki, C., Alyass, A. & Meyre, D. Ethnic and population differences in the genetic predisposition to human obesity. *Obes Rev* **19**, 62–80 (2018). 428
429
35. Fernández, M., Pereira, A., Corvalán, C. & Mericq, V. Precocious pubertal events in chilean children: ethnic disparities. *J Endocrinol Invest* **42**, 385–395 (2019). 430
431
36. Vicuña, L. *et al.* Novel loci and mapuche genetic ancestry are associated with pubertal growth traits in chilean boys. *Hum Genet* (2021). 432
433
37. Degenhardt, F. *et al.* Construction and benchmarking of a multi-ethnic reference panel for the imputation of hla class i and ii alleles. *Hum Mol Genet* **28**, 2078–2092 (2019). 434
435
436

38. Ilonen, J. *et al.* Genetic susceptibility to type 1 diabetes in childhood - estimation of hla class ii associated disease risk and class ii effect in various phases of islet autoimmunity. *Pediatr Diabetes* **17 Suppl 22**, 8–16 (2016). 437–439
39. Barrett, J. C. *et al.* Genome-wide association study and meta-analysis find that over 40 loci affect risk of type 1 diabetes. *Nat Genet* **41**, 703–7 (2009). 440–441
40. Chu, A. Y. *et al.* Multiethnic genome-wide meta-analysis of ectopic fat depots identifies loci associated with adipocyte development and differentiation. *Nat Genet* **49**, 125–130 (2017). 442–444
41. Kain, J. *et al.* [evolution of the nutritional status of chilean children from preschool to school age: anthropometric results according to the source of the data]. *Arch Latinoam Nutr* **60**, 155–9 (2010). 445–447
42. Helgeland, Ø. *et al.* Genome-wide association study reveals dynamic role of genetic variation in infant and early childhood growth. *Nat Commun* **10**, 4448 (2019). 448–450
43. Berkey, C. S. & Reed, R. B. A model for describing normal and abnormal growth in early childhood. *Hum Biol* **59**, 973–87 (1987). 451–452
44. Stergiakouli, E. *et al.* Genome-wide association study of height-adjusted bmi in childhood identifies functional variant in adcy3. *Obesity (Silver Spring)* **22**, 2252–9 (2014). 453–455
45. Cole, T. J. Weight/height² compared to weight/height³ for assessing adiposity in childhood: influence of age and bone age on p during puberty. *Ann Hum Biol* **13**, 433–51 (1986). 456–458
46. Cole, T. J. & Green, P. J. Smoothing reference centile curves: the lms method and penalized likelihood. *Stat Med* **11**, 1305–19 (1992). 459–460
47. Tobar, H. E. *et al.* Identification and functional analysis of missense mutations in the lecithin cholesterol acyltransferase gene in a chilean patient with hypoalphalipoproteinemia. *Lipids Health Dis* **18**, 132 (2019). 461–463
48. Maples, B. K., Gravel, S., Kenny, E. E. & Bustamante, C. D. Rfmix: a discriminative modeling approach for rapid and robust local-ancestry inference. *Am J Hum Genet* **93**, 278–88 (2013). 464–466
49. 1000 Genomes Project Consortium *et al.* A global reference for human genetic variation. *Nature* **526**, 68–74 (2015). 467–468
50. Chang, C. C. *et al.* Second-generation plink: rising to the challenge of larger and richer datasets. *Gigascience* **4**, 7 (2015). 469–470
51. Delaneau, O., Zagury, J.-F. & Marchini, J. Improved whole-chromosome phasing for disease and population genetic studies. *Nat Methods* **10**, 5–6 (2013). 471–472