Stress signaling and STAT1 activation characterize the keratinocytic gene expression pattern in Hidradenitis suppurativa

VG Frings¹, L Jopp¹, M Srivastava², D Presser¹, M Goebeler¹, M Schmidt¹

¹Department of Dermatology, Venereology and Allergology, University Hospital Würzburg, Germany

²Core Unit Systemmedizin (SysMed), Medical Faculty, University of Würzburg, Würzburg, Germany

Text word count: 3.312

Figures: 6

Tables: 0

Supplemental Material: 2 suppl. Fig. and 6 suppl. Tab.

Corresponding author:

Prof. Dr. Marc. Schmidt

Department of Dermatology, Venereology and Allergology

University Hospital Würzburg

Josef-Schneider-Straße 2

97076 Würzburg

Germany

Mail: schmidt_m11@ukw.de

Phone: +49-931-20126710

Funding sources: This work was supported by the *Interdisziplinäres Zentrum für Klinische Forschung (IZKF)*, University Hospital Würzburg, Germany (ZZ/25).

Conflict of interest: D.P. is member of the European Hidradenitis Suppurativa Foundation (EHSF) e. V. The Department of Dermatology, Venereology and Allergology, University Hospital Würzburg (with active members D. P. and M. G.) is a health care provider center of the European Network for Rare and Low Prevalence Complex Skin diseases (*ERN Skin*).

ABSTRACT

Background: The underlying pathogenetic factors generating the innate immune signal necessary for T cell activation, initiation and chronification of *Hidradenitis suppurativa* (HS, also known as *Acne inversa*) are still poorly understood. Emerging evidence suggests that defective keratinocyte function critically contributes to HS disease development and progression.

Objectives: To elucidate the role of keratinocytes in HS lesion formation, we compared the transcriptomes of isolated lesional and perilesional HS epidermis by RNA sequencing.

Methods: Lesional and perilesional HS skin samples of at least 3 different donors were obtained. Isolated epidermal keratinocytes were further processed for cell culture, protein extraction, immunostaining procedures or RNA isolation and RNA sequencing. For large scale promotor site analysis, DEGs were analyzed for overrepresented transcription factor binding sites. Functional annotation clustering for analyzing enriched functional-related gene groups was performed employing the DAVID Bioinformatics Resources.

Results: We show that HS is characterized by a strong epidermal stress state as evident by a significant overrepresentation of an AP-1-driven stress signature in the overall gene expression pattern of lesional keratinocytes and a substantial activation of the stress-activated cJun N-terminal kinase (JNK) pathway in lesional HS epidermis. Additionally, our data reveal a strong induction of STAT1 activation in lesional HS epidermis that likely results from IFN γ production and governs the expression of key inflammatory genes that coordinate activation of innate immunity and the adaptive T cell response in HS.

Conclusions: Taken together, these data implicate a new role of combined stress signaling and JAK/STAT1 pathway activation in disease progression of HS suggesting interference with JAK/STAT1 signaling as a potentially promising therapeutic approach for HS.

INTRODUCTION

Hidradenitis suppurativa (HS, also known as *Acne inversa*)¹⁻³ is a chronic inflammatory skin disease characterized by recurrent deep-seated and painfully inflamed lesions in the intertriginous apocrine gland-bearing areas of the body⁴. Both a T cell-dependent adaptive response and innate immune activation play key roles in disease progression. Those two processes are tightly interwoven since activation of the innate immune system is a prerequisite for the generation of T cells and promotes further adaptive immune responses⁵. In HS, both Th17 cell activation⁶, regulatory T cell (Treg) deficiency⁷, and a strong non-specific inflammation^{3,8} correlate with disease progression. However, the underlying pathogenetic factors of HS generating the seminal innate immune signal that bridges T cell activation, initiation and chronification of the disease remain to be elucidated.

There is accumulating evidence that keratinocyte dysfunction plays an important role in HS. For instance, a functional defect of hair follicle keratinocytes⁹ and imbalances in epithelial cell differentiation³ were proposed to sensitize for inflammatory responses thereby contributing to HS evolution. Additionally, stimuli implicated to promote disease persistence involve bacterial overgrowth^{9,10} and release of ruptured hair follicle contents (including bacteria, damage-associated molecular patterns (DAMPs), keratin fibers and sebum) into the surrounding tissue¹. Both signals stimulate other cells, including keratinocytes to produce inflammatory cytokines¹¹. Moreover, we demonstrated a role of epidermal inflammasome activation in the pathogenesis of HS¹². The inflammasome complex controls generation of active interleukin (IL)-1 β , which is a critical regulator of inflammation and driver of antigen-specific adaptive immune responses¹³. In HS, enhanced IL-1 β production from both epidermal keratinocytes¹² and keratinocytes isolated from hair follicles⁹ has been reported. These data raise the possibility that increased inflammation during disease progression may

critically depend on dysregulation of keratinocyte-dependent gene responses fostering a vicious cycle of proinflammatory events that culminate in persistent chronic inflammation in HS.

To gain better insight into the exact contribution of keratinocytes to lesional gene expression, we performed RNA sequencing of lesional versus perilesional HS epidermis. Unlike previous whole-skin proteome, transcriptome, exome or mRNA microarray analyses ¹⁴⁻¹⁹, which did not distinguish between keratinocytic and non-keratinocytic responses, our study provides a first comprehensive overview of the isolated transcriptional pattern of lesional epidermal keratinocytes without major obscuring effects of other immunologically active cells. Our data identify a complex epidermal stress response to external and internal stimuli that is characterized by a significant overrepresentation of an AP-1-driven stress signature in the overall gene expression pattern of lesional keratinocytes. Additionally, we observe a strong induction of STAT1 activation in lesional HS epidermis that likely results from IFN γ production and governs the expression of key inflammatory genes orchestrating activation of innate immunity and the adaptive T cell response. Our data implicate a new role of combined stress signaling and JAK/STAT1 pathway activation in disease progression of HS.

MATERIALS AND METHODS

Isolation of HS skin samples and epidermal sheets

Human skin samples were obtained by in-house therapeutic skin surgery from HS patients after informed consent (ethical vote was given by the local Ethics Committee, University of Würzburg, Bavaria, Germany; No. 306/12). Skin samples of at least 3 different donors were used for experiments. Lesional and perilesional (excised at 2 cm distance from the next active HS lesion) HS skin biopsies of 8 mm in diameter were taken and epidermis and dermis separated by dispase treatment and subsequent mechanical separation as described ²⁰. After trypsinization isolated epidermal keratinocytes were further processed for cell culture, protein extraction or RNA isolation.

RNA-sequencing and bioinformatic analysis

mRNA was extracted from five pairwise-matched lesional and perilesional epidermal HS pellets using the Qiagen RNeasy Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions and RNA sequencing was performed as detailed in the supplementary informations. For large scale promotor site analysis, DEGs were analyzed for overrepresented transcription factor binding sites represented in the Jasper 2018 database using Pscan (v1.4 ²¹). Functional annotation clustering for analyzing enriched functional-related gene groups was performed employing the Data base for Annotation, Visualization, and Integrated Discovery (DAVID Bioinformatics Resources) (v6.8; david.abcc.ncifcrf.gov) as described ²².

Cell culture

Human epithelial keratinocytes (hEK) isolated from lesional epidermal HS sheets were expanded and cultured as previously described²³. Cells were stimulated with tumor necrosis factor (TNF) α (50 ng/ml), type I interferon (IFN) (10 U/ml; both from R&D Systems, Wiesbaden, Germany) or IFN γ (100 ng/ml; Peprotech, Hamburg, Germany) by diluting them directly from appropriate stock solutions into the medium. The selective JAK1 inhibitor upadacitinib (250nM; #S8162; Selleckchem, Munich, Germany) was preincubated for 30 min at 37°C prior to addition of the stimuli.

Immunoblotting analysis of secreted proteins

Cell pellets from lesional and perilesional HS epidermis were lysed and protein expression analyzed by immunoblot as described before (Frings et al., 2019) using the following primary antibodies: p-SAPK/JNK (#9251) and p-STAT1 (#9167; both from Cell Signaling Technology, Frankfurt, Germany), Anti- α -tubulin (#T5168, Sigma-Aldrich, Darmstadt, Germany). Proteins were visualized by enhanced chemiluminescence using appropriate horseradish-peroxidase (HRP)-coupled secondary antibodies.

Quantitative real-time polymerase chain reaction (qRT-PCR)

Total RNA isolation, cDNA synthesis and qRT-PCR of lesional and perilesional epidermal HS cell isolates was performed as described previously²⁰ using the SYBR Green method and appropriate gene-specific primers (Sigma-Aldrich; Table S1). Expression of TLR4 and Keratin 14 was investigated to ensure purity of isolated epidermal sheets²⁰. Test gene expression was normalized to the expression of the housekeeping gene GAPDH (Table S1) and related to an experimental control using the $\Delta\Delta$ CT method.

Immunostaining procedures

3 μ m-skin sections from formalin-fixed and paraffin-embedded biopsies from healthy donor skin (excess tissue after tumor surgery) or perilesional and lesional areas from skin samples of HS patients were selected for immunohistological analysis and stained for p-SAPK/JNK (#4668, Cell Signaling), p-c-Jun (#3270, Cell Signaling), p-STAT1 (#9167, Cell Signaling), and c-Jun (#ab218576, Abcam, Cambridge, UK) proteins using antibodies from Cell Signaling Technology. Antibodies were used at 1:100 (pJNK, pJun), 1:200 (c-Jun) or 1:500 (p-STAT1) dilution and visualized using a HRP system (#K5003; Dako, Hamburg, Germany)

12.

Statistical analysis

If not indicated otherwise, data are depicted as averages $\pm$ s.d. from at least three independent experiments. Statistical significance of differences was determined by paired t-test or one-way analysis of variance (ANOVA) with post-hoc Dunnett correction and considered significant at $p \leq 0.05$ (*). Higher significances are indicated accordingly (** ≤ 0.01 , *** ≤ 0.001). Statistical analysis was performed using GraphPad Prism 5.0 (GraphPad software Inc., La Jolla, CA).

RESULTS

HS epidermis is characterized by a complex stress response that may influence innate and adaptive immune activation

To gain insight into the repertoire of *in situ* expressed keratinocyte genes in HS we studied pairwise matched perilesional and lesional epidermis isolated from tissue after HS surgery by RNA sequencing. Principal component analysis (Figure S1) showed clustering of expression profiles across patients but clear differences between the transcriptomes of lesional and perilesional HS epidermis. This was unlikely to result from incomplete isolation of basal keratinocytes since both qRT-PCR and sequencing analysis consistently verified high expression of the basal keratinocyte marker K14 and lack of Toll-like receptor (TLR)-4 (Table S2). Expression of the latter is prominent in dermal fibroblasts and macrophages²⁴ but absent in epidermal keratinocytes²⁰. In addition, there was no major contamination with non-keratinocytic immune cells since sequencing counts for immunological markers such as CD3 for T-cells, CD19 for B-cells, CD68 for macrophages, or MPO for neutrophils were extremely low and expression ranged close to the detection limit (Table S2). By statistical evaluation we totally identified 183 up- and 49 downregulated genes in lesional HS epidermis with significantly altered mRNA abundance of ≥ 2 -fold (full dataset publicly available at EGA European Genome-Phenome Archive, <https://ega-archive.org>, ID EGAS00001005544). We next performed functional annotation cluster (FAC) analysis on the differentially expressed genes (DEGs) to identify statistically enriched groups of lesionally regulated functional genes (Figure 1). Consistent with previous reports demonstrating disturbances of the cutaneous microbiome and a role of neutrophils in HS^{12,25-29}, we observed a significant overrepresentation of various gene ontology (GO) groups related to cellular responses to bacterial products and neutrophil chemotaxis among the top-upregulated FACs. Additionally,

we found enrichment of several GO groups related to leukocyte differentiation, T cell activation, cytokine signaling and innate immune activation. To confirm our findings by an alternative method we performed qRT-PCR for various statistically upregulated gene products from selected functional groups overrepresented in the HS lesions (Figure 2). Since perifollicular deficiencies in Treg numbers or function and imbalance of the Th17/Treg ratio⁷ appear to play a central role in HS inflammation, we focused our analysis on genes of the functional cluster “T cell regulation” (*IL4R*, *PDL1*, *JAK3*; Table S3). Additionally, we chose representative genes of the cluster “activation of immune response” (*TLR2*, *TLR8*, *TNIP3*, *cGAS*; Table S3) containing several genes regulating innate immunity since an altered expression of innate immunity markers may explain the development of chronic inflammatory lesions² or contribute to initiation of the dominant T-cell immune response in HS¹. qRT-PCR confirmed significant lesional induction of all tested DEGs (*IL4R*, *PDL1*, *JAK3*, *TLR2*, *TLR8*, *TNIP3*, *cGAS*; Figure 2) except of *TLR2*, which expression in lesional skin was increased without reaching statistical significance.

Epidermal HS lesions show prominently enriched AP-1- and STAT1-binding sites in the promoters of upregulated genes

To gain better insight into the regulatory pathways underlying the observed transcriptional changes in lesional HS epidermis, we next performed large scale promoter analysis of all ≥ 2.0 -fold statistically DEGs (Table S4). These studies revealed a prominent enrichment of promoter binding sites for multiple AP-1 transcription factors and members of the JAK-STAT signaling pathway among the lesionally upregulated DEGs (Figure 3). DEGs with well conserved AP-1 sites in their promoters involved several of the top upregulated genes such as the antibacterial defensin genes *DEFB4B* and *DEFB4A*, the A20-binding protein *TNIP3* or the neutrophil-attracting chemokine *CXCL8* (Table S5). These genes also consistently appear in

the top-regulated FACs (Table S3). Surprisingly, binding sites for NF- κ B transcription factors were less prominent among the lesionally induced genes. In particular, we failed to detect statistical overrepresentation of the NF- κ B transcription factor cREL (Table S4), which commonly mediates expression of key proinflammatory genes in response to stimulation with HS-relevant cytokines such as TNF and IL-1 β ³⁰. Moreover, binding sites for NF κ B subunits NFKB1 or NFKB2 were not statistically overrepresented and Rel B sites were only enriched in the lesionally downregulated DEGs (Table S4 and Figure 3). Only binding sites for Rel A showed statistical enrichment in the promoters of the upregulated DEGs albeit with a >500 fold lower enrichment score (ES: 114.81, equivalent to $p \leq 8.7E-3$; Table S4) than e.g. STAT1 (ES: 61784.46, $p \leq 1.6E-5$) or the AP-1 factor FOSL1 (ES: 784473.70, $p < 1.3E-6$), suggesting a more prominent contribution of AP-1 factors and JAK/STAT signaling to the keratinocytic gene expression profile in HS lesions.

Activation of AP-1 factors is a well-known consequence of cytokine- and microbially induced stress signaling and is usually triggered via phosphorylation by stress-activated protein kinases such as c-Jun N-terminal kinase (JNK) ³¹. Additionally, AP-1 activation is regulated at the transcriptional level e.g. by activation of the mitogen-activated protein kinases ERK1/2 ³². Initial analysis of our RNA sequencing data revealed no overt differences in the expression of individual AP-1 factors in lesional vs. perilesional HS epidermis (data not shown). However, we observed a strongly increased epidermal phosphorylation of c-Jun and JNK in lesional HS epidermis (Figure 4), suggesting that enhanced JNK-mediated stress signaling may account for the observed overrepresentation of AP-1 sites in the regulated transcriptome of lesional HS epidermis.

While a prominent JAK/STAT signature was already obvious from our large scale promoter analysis of all lesionally upregulated genes (Figure 3), a separate promoter investigation with smaller subsets of upregulated DEGs from individual overrepresented FACs uncovered an

even more striking enrichment of STAT transcription factor sites. This was particularly true for STAT1 or STAT1:STAT2 sites that were highly enriched ($p < 0.001$) in the promoters of the top 10 enriched FACs “Activation of immune response” (STAT1:STAT2; $p \leq 5.2E-5$; STAT1; $p \leq 1.7E-4$), “Neutrophil chemotaxis” (STAT1; $p \leq 8.0E-05$), “Response to bionic stimulus” (STAT1; $p \leq 1.2E-4$) “Cellular response to molecules of bacterial origin” (STAT1; $p \leq 1.0E-4$), and “Response to cytokine” (STAT1; $p \leq 7.3E-4$) (data not shown) but were also prominent in other top-enriched clusters such as “Innate immune response” (STAT1; $p \leq 2.4E-5$; Figure 5, a). This apparent STAT1 signature in lesional HS epidermis could also be confirmed by immunohistochemistry (Figure 5, b-c) and immunoblot analysis (Figure 5, d). DEGs with well conserved IFN γ -responsive STAT-1 sites in their promoters also involved several of the lesionally most increased genes such as *TNIP3*, *TLR8* and *CXCL8* (Table S6), which consistently appeared in the top-regulated FACs (Table S3). STAT1 can be activated by several ligands including type I IFNs (IFN α and IFN β) or IFN γ ³³, which both have been implicated in the pathogenesis of HS^{6,28}. Western blot analysis of lysates from human epidermal keratinocytes (hEKs) isolated from lesional HS biopsies revealed a strong and sustained STAT1 phosphorylation upon IFN γ treatment and a transient activation with type I IFN (Figure 5, e). Thus, principally either IFN γ production or release of type I IFNs may explain the observed STAT1 activation in lesional HS epidermis.

IFN γ -mediated STAT1 activation controls expression of immune-regulatory genes in lesional HS epidermis

Our results raised the possibility that STAT1 activation may particularly influence induction of T cell-modulating genes or gene products controlling innate immune responses in HS lesions. To test whether stimulation with STAT1-activating IFNs may initiate expression of at least some of our previously confirmed lesional-specific DEGs, we isolated hEKs from

lesional HS epidermis and stimulated them with either type I IFN or IFN γ . Additionally, we performed stimulations with TNF α , which has previously been implicated in the pathogenesis of HS³⁴ and included combinatory treatments of TNF α with either of the two IFNs to explore a potential cooperativity of both signals. Strikingly, the majority of our lesionally enriched test genes were transcriptionally induced by IFN γ but barely responsive to mono-treatment with TNF α or type I IFN. Only *TNIP3* induction was clearly TNF-restricted (Figure 6, a). Notably, most test genes except for *TNIP3*, showed a trend towards increased mRNA abundance upon co-treatment with IFN γ and TNF α , suggesting that both cytokines may work in concert to induce a certain set of immune-modulating genes in HS epidermis (Figure 6, a). Treatment with the selective JAK1 inhibitor upadacitinib efficiently suppressed induction of all investigated genes but *TNIP3* by IFN γ or IFN γ /TNF α cotreatment (Figure 6, b). Consistent with the prominent STAT1 signature characterizing several lesionally induced functional clusters of our RNA seq analysis, parallel Western blots confirmed efficient suppression of IFN γ - or IFN γ /TNF α -induced STAT1 phosphorylation by upadacitinib (Figure 6, c). This indicates that STAT1 activation critically contributed to the induction of our newly discovered HS-related keratinocytic IFN γ target genes.

DISCUSSION

Here we provide evidence that keratinocytes of lesional HS skin crucially orchestrate stress, innate and adaptive immune responses by inducing a complex AP-1- and STAT1-dominated gene expression pattern. DEGs with well conserved AP-1 or STAT1 sites in their promoters involved several of the top upregulated genes such *DEFB4B*, *DEFB4A*, *TNIP3* and *CXCL8*, suggesting a prominent role of AP-1 and STAT1 signaling in epidermal HS responses. The observed AP-1 signature is likely a consequence of increased JNK activation in lesional keratinocytes, which may result from a general keratinocytic stress response due to excessive bacterial colonization. Accumulation of bacteria has previously been proposed to elicit a strong stress signal affecting T cell responses¹ and innate immunity³⁵ in HS and may trigger activation of epidermal innate immune receptors such as TLR2 and/or production of TNF or IL-1 β , which all are established triggers of JNK activation³⁶.

Unlike the rather unspecific AP-1 stress signature, the observed STAT1 phosphorylation probably indicates a specific response to high IFN levels in lesional HS epidermis. Both IFN γ released by effector T cells and type I IFN produced in response to neutrophil-induced damage have been implicated in the pathogenesis of HS^{6,28} and could principally account for the observed STAT1 phosphorylation. Yet, two findings of our study argue against a major role of type I IFN in the epidermal response in HS. First, our large scale promoter analysis showed a strong overrepresentation of the more IFN γ -specific STAT1 sites in most regulated FACs but a less pronounced overrepresentation of STAT1::STAT2 sites that are more type I IFN-restricted³⁷⁻³⁹. Secondly, we did not observe a sustained STAT1 phosphorylation and an appreciable induction of potential STAT1 genes by type I IFN stimulation of lesionally derived keratinocytes. In contrast, IFN γ stimulation induced both a sustained STAT1 phosphorylation and an upregulation of most of our test genes *in vitro*. Thus,

overrepresentation of STAT1 sites in lesionally upregulated genes likely indicates a predominant response of lesional HS keratinocytes to IFN γ .

Strikingly, our RNA seq analysis and the subsequent bioinformatic evaluation revealed a clear overrepresentation of various T cell-relevant transcripts among the lesionally upregulated epidermal DEGs, which was confirmed by qRT-PCR analysis. While the keratinocytic function of many of those genes is unclear, we could show that most of them were strongly co-regulated by IFN γ and TNF α . Hence, their regulation in epidermal HS lesions above all may reflect a conserved epidermal responsiveness to high levels of IFN γ and TNF α contributing to T cell dysregulation, which has been proposed to initiate and sustain chronic inflammation in HS (Moran et al., 2017). On the other hand, the response pattern might also mirror the control of inflammatory processes in the dermis by lesional HS keratinocytes, which may inhibit T cell proliferation and cytokine production as recently suggested for allergic contact dermatitis ⁴⁰. Particularly remarkable in this context is the observed strong induction of *PDL1* in lesional HS epidermis. PDL1 is an established target of IFN γ -JAK1/2-STAT1/2/3-IRF1 signaling ⁴¹ and suppresses T cell immunity (proliferation, differentiation, cytokine secretion, cytolytic function) by binding to programmed cell death (PD)-1 ^{42,43}.

Besides the observed regulation of T cell-relevant genes, our RNA seq study also revealed a significant lesional overrepresentation of the functional gene cluster “activation of immune response” in HS epidermis. This cluster comprises several genes involved in innate immune activation that may instigate the apparently dominant T cell immune response in HS. Statistical upregulation of such genes might sensitize cells to endogenous and exogenous danger signals thereby contributing to disease perpetuation and progression ⁴⁴.

While NF- κ B-dependent gene expression is critical for induction of an efficient immune response to bacterial infection ⁴⁵, excessive or prolonged NF- κ B signaling can foster inflammatory diseases ⁴⁶. Therefore, the NF- κ B pathway is tightly regulated by several

intracellular proteins^{47,48}. These include *TNIP3*, an interaction partner of the ubiquitin-editing enzyme A20/ TNFAIP3, which counteracts inflammation by restricting NF-κB pathway signaling downstream of TNF and TLR4^{49,50}. NF-κB transcription factors were not among the top hits in the unfiltered promoter analysis of all regulated DEGs, albeit TNFα is associated with (auto-) inflammatory mechanisms in the pathogenesis of HS⁵¹. The overexpression of *TNIP3* might explain the low overrepresentation of NF-κB transcription factors in our upregulated gene set and contribute to chronification in HS by dampening NF-κB activity allowing innate immune-activating bacteria to persist in HS lesions to trigger a new cycle of recurrent inflammation.

The sensitivity of our verified IFNγ- and TNFα-inducible DEGs to JAK1/2 inhibition suggests a striking dependency of the epidermal gene response on JAK1/2 signaling. Considering that the currently only approved targeted therapy with the TNFα-antagonist adalimumab often results in unsatisfactory responses⁵² these data raise the question whether JAK inhibitors might improve the success of non-surgical therapeutic approaches. Intriguingly, it was recently shown that anti-TNFα therapy resulted in a preferential attenuation of B cell-specific gene expression, whereas other parameters including IFNγ levels were unchanged after this treatment⁵³. Accordingly, suboptimal clinical response to anti-TNFα therapy in HS⁵² might perhaps be due to a persistent IFNγ response making HS patients refractory to TNFα antagonists. At any rate, our results suggest that treatment regimens with JAK inhibitors - either alone or in combination with other agents – could be a promising new treatment option in HS.

ACKNOWLEDGMENTS

We thank Andrea Knorz, Katharina Meder and Helga Sennefelder for excellent technical assistance. This work was supported by the *Interdisziplinäres Zentrum für Klinische Forschung (IZKF)*, University Hospital Würzburg, Germany (ZZ/25). We thank the Core Unit Systems Medicine at the Medical Faculty, University of Würzburg, that is co-funded by the *Interdisziplinäres Zentrum für Klinische Forschung (IZKF)* Würzburg for support of this work.

DATA AVAILABILITY STATEMENT

Datasets related to this article can be found under EGA ID EGAS00001005544, hosted at EGA European Genome-Phenome Archive (<https://ega-archive.org>).

ORCIDS

V. G. F.: <https://orcid.org/0000-0002-8256-7961>

L. J.: <https://orcid.org/0000-0002-0982-7499>

D. P.: <https://orcid.org/0000-0001-8535-1309>

M. Srivastava <https://orcid.org/0000-0002-7449-9911>

M. G.: <https://orcid.org/0000-0001-7095-9848>

M. Schmidt.: <https://orcid.org/0000-0002-8643-1615>

AUTHOR CONTRIBUTION STATEMENT

Conceptualization: MS*, VGF; Formal analysis: LJ, VGF; Bioinformatical analysis: MS*, MS[#]; Funding Acquisition: VGF; Collection of clinical specimens: DP; Investigation: VGF, LJ; Methodology: MG, MS*; Project Administration: MS*; Resources: MG; Supervision: MS*; Visualization: VGF, LJ, MS*; Writing - Original Draft Preparation: MS*, VGF; Writing - Review and Editing: MS*, VGF, MG, LJ, DP, MS[#].

*Marc Schmidt

[#]Mughda Srivastava

REFERENCES

- 1 Wolk K, Join-Lambert O, Sabat R. Aetiology and pathogenesis of hidradenitis suppurativa. *Br J Dermatol* 2020; **183**: 999-1010.
- 2 Dreno B, Khammari A, Brocard A *et al*. Hidradenitis suppurativa: the role of deficient cutaneous innate immunity. *Arch Dermatol* 2012; **148**: 182-6.
- 3 Zouboulis CC, Nogueira da Costa A, Makrantonaki E *et al*. Alterations in innate immunity and epithelial cell differentiation are the molecular pillars of hidradenitis suppurativa. *J Eur Acad Dermatol Venereol* 2020; **34**: 846-61.
- 4 Zouboulis CC, Desai N, Emtestam L *et al*. European S1 guideline for the treatment of hidradenitis suppurativa/acne inversa. *J Eur Acad Dermatol Venereol* 2015; **29**: 619-44.
- 5 Chaplin DD. Overview of the immune response. *J Allergy Clin Immunol* 2010; **125**: S3-23.
- 6 Wolk K, Warszawska K, Hoeflich C *et al*. Deficiency of IL-22 contributes to a chronic inflammatory disease: pathogenetic mechanisms in acne inversa. *J Immunol* 2011; **186**: 1228-39.
- 7 Moran B, Sweeney CM, Hughes R *et al*. Hidradenitis Suppurativa Is Characterized by Dysregulation of the Th17:Treg Cell Axis, Which Is Corrected by Anti-TNF Therapy. *J Invest Dermatol* 2017; **137**: 2389-95.
- 8 Witte-Handel E, Wolk K, Tsaousi A *et al*. The IL-1 Pathway Is Hyperactive in Hidradenitis Suppurativa and Contributes to Skin Infiltration and Destruction. *J Invest Dermatol* 2019; **139**: 1294-305.
- 9 Hotz C, Boniotto M, Guguin A *et al*. Intrinsic Defect in Keratinocyte Function Leads to Inflammation in Hidradenitis Suppurativa. *J Invest Dermatol* 2016; **136**: 1768-80.
- 10 Hofmann SC, Saborowski V, Lange S *et al*. Expression of innate defense antimicrobial peptides in hidradenitis suppurativa. *J Am Acad Dermatol* 2012; **66**: 966-74.
- 11 Akira S, Uematsu S, Takeuchi O. Pathogen recognition and innate immunity. *Cell* 2006; **124**: 783-801.
- 12 Lima AL, Karl I, Giner T *et al*. Keratinocytes and neutrophils are important sources of proinflammatory molecules in hidradenitis suppurativa. *Br J Dermatol* 2016; **174**: 514-21.
- 13 Dinarello CA. Immunological and inflammatory functions of the interleukin-1 family. *Annu Rev Immunol* 2009; **27**: 519-50.
- 14 Shanmugam VK, Jones D, McNish S *et al*. Transcriptome patterns in hidradenitis suppurativa: support for the role of antimicrobial peptides and interferon pathways in disease pathogenesis. *Clin Exp Dermatol* 2019; **44**: 882-92.
- 15 Gauntner TD. Hormonal, stem cell and Notch signalling as possible mechanisms of disease in hidradenitis suppurativa: a systems-level transcriptomic analysis. *Br J Dermatol* 2019; **180**: 203-4.
- 16 Brandao L, Moura R, Tricarico PM *et al*. Altered keratinization and vitamin D metabolism may be key pathogenetic pathways in syndromic hidradenitis suppurativa: a novel whole exome sequencing approach. *J Dermatol Sci* 2020; **99**: 17-22.

- 17 Blok JL, Li K, Brodmerkel C *et al.* Gene expression profiling of skin and blood in hidradenitis suppurativa. *Br J Dermatol* 2016; **174**: 1392-4.
- 18 Gudjonsson JE, Tsoi LC, Ma F *et al.* Contribution of plasma cells and B cells to hidradenitis suppurativa pathogenesis. *JCI Insight* 2020; **5**.
- 19 Rumberger BE, Boarder EL, Owens SL *et al.* Transcriptomic analysis of hidradenitis suppurativa skin suggests roles for multiple inflammatory pathways in disease pathogenesis. *Inflamm Res* 2020; **69**: 967-73.
- 20 Frings VG, Muller D, Storz G *et al.* Improved metal allergen reactivity of artificial skin models by integration of Toll-like receptor 4-positive cells. *Contact Dermatitis* 2019; **81**: 254-61.
- 21 Zambelli F, Pesole G, Pavesi G. Pscan: finding over-represented transcription factor binding site motifs in sequences from co-regulated or co-expressed genes. *Nucleic Acids Res* 2009; **37**: W247-52.
- 22 Ohnesorge N, Viemann D, Schmidt N *et al.* Erk5 activation elicits a vasoprotective endothelial phenotype via induction of Kruppel-like factor 4 (KLF4). *J Biol Chem* 2010; **285**: 26199-210.
- 23 Schmidt M, Goebeler M, Posern G *et al.* Ras-independent activation of the Raf/MEK/ERK pathway upon calcium-induced differentiation of keratinocytes. *J Biol Chem* 2000; **275**: 41011-7.
- 24 Ling GS, Bennett J, Woollard KJ *et al.* Integrin CD11b positively regulates TLR4-induced signalling pathways in dendritic cells but not in macrophages. *Nat Commun* 2014; **5**: 3039.
- 25 Ring HC, Thorsen J, Saunte DM *et al.* The Follicular Skin Microbiome in Patients With Hidradenitis Suppurativa and Healthy Controls. *JAMA Dermatol* 2017; **153**: 897-905.
- 26 Ring HC, Sigsgaard V, Thorsen J *et al.* The microbiome of tunnels in hidradenitis suppurativa patients. *J Eur Acad Dermatol Venereol* 2019; **33**: 1775-80.
- 27 Langan EA, Recke A, Bokor-Billmann T *et al.* The Role of the Cutaneous Microbiome in Hidradenitis Suppurativa-Light at the End of the Microbiological Tunnel. *Int J Mol Sci* 2020; **21**.
- 28 Byrd AS, Carmona-Rivera C, O'Neil LJ *et al.* Neutrophil extracellular traps, B cells, and type I interferons contribute to immune dysregulation in hidradenitis suppurativa. *Sci Transl Med* 2019; **11**.
- 29 Narla S, Azzam M, Townsend S *et al.* Identifying key components and therapeutic targets of the immune system in hidradenitis suppurativa with an emphasis on neutrophils. *Br J Dermatol* 2020.
- 30 Vallabhapurapu S, Karin M. Regulation and function of NF-kappaB transcription factors in the immune system. *Annu Rev Immunol* 2009; **27**: 693-733.
- 31 Minden A, Karin M. Regulation and function of the JNK subgroup of MAP kinases. *Biochim Biophys Acta* 1997; **1333**: F85-104.
- 32 Gilley R, March HN, Cook SJ. ERK1/2, but not ERK5, is necessary and sufficient for phosphorylation and activation of c-Fos. *Cell Signal* 2009; **21**: 969-77.
- 33 Baris S, Alroqi F, Kiykim A *et al.* Severe Early-Onset Combined Immunodeficiency due to Heterozygous Gain-of-Function Mutations in STAT1. *J Clin Immunol* 2016; **36**: 641-8.
- 34 van der Zee HH, de Ruiter L, van den Broecke DG *et al.* Elevated levels of tumour necrosis factor (TNF)-alpha, interleukin (IL)-1beta and IL-10 in hidradenitis suppurativa skin: a rationale for targeting TNF-alpha and IL-1beta. *Br J Dermatol* 2011; **164**: 1292-8.

- 35 van der Zee HH, de Ruiter L, Boer J *et al.* Alterations in leucocyte subsets and histomorphology in normal-appearing perilesional skin and early and chronic hidradenitis suppurativa lesions. *Br J Dermatol* 2012; **166**: 98-106.
- 36 Li DQ, Luo L, Chen Z *et al.* JNK and ERK MAP kinases mediate induction of IL-1beta, TNF-alpha and IL-8 following hyperosmolar stress in human limbal epithelial cells. *Exp Eye Res* 2006; **82**: 588-96.
- 37 Levy DE, Darnell JE, Jr. Stats: transcriptional control and biological impact. *Nat Rev Mol Cell Biol* 2002; **3**: 651-62.
- 38 Banoth B, Cassel SL. Mitochondria in innate immune signaling. *Transl Res* 2018; **202**: 52-68.
- 39 Borden EC, Sen GC, Uze G *et al.* Interferons at age 50: past, current and future impact on biomedicine. *Nat Rev Drug Discov* 2007; **6**: 975-90.
- 40 Seiringer P, Eyerich S, Eyerich K *et al.* Keratinocytes Regulate the Threshold of Inflammation by Inhibiting T Cell Effector Functions. *Cells* 2021; **10**.
- 41 Garcia-Diaz A, Shin DS, Moreno BH *et al.* Interferon Receptor Signaling Pathways Regulating PD-L1 and PD-L2 Expression. *Cell Rep* 2017; **19**: 1189-201.
- 42 Latchman YE, Liang SC, Wu Y *et al.* PD-L1-deficient mice show that PD-L1 on T cells, antigen-presenting cells, and host tissues negatively regulates T cells. *Proc Natl Acad Sci U S A* 2004; **101**: 10691-6.
- 43 Mallett G, Laurence A, Amarnath S. Programmed Cell Death-1 Receptor (PD-1)-Mediated Regulation of Innate Lymphoid Cells. *Int J Mol Sci* 2019; **20**.
- 44 Schell SL, Schneider AM, Nelson AM. Yin and Yang: A disrupted skin microbiome and an aberrant host immune response in hidradenitis suppurativa. *Exp Dermatol* 2021.
- 45 Gilmore TD. Introduction to NF-kappaB: players, pathways, perspectives. *Oncogene* 2006; **25**: 6680-4.
- 46 Kumar A, Takada Y, Boriek AM *et al.* Nuclear factor-kappaB: its role in health and disease. *J Mol Med (Berl)* 2004; **82**: 434-48.
- 47 Lang T, Mansell A. The negative regulation of Toll-like receptor and associated pathways. *Immunol Cell Biol* 2007; **85**: 425-34.
- 48 Wullaert A, Heyninck K, Janssens S *et al.* Ubiquitin: tool and target for intracellular NF-kappaB inhibitors. *Trends Immunol* 2006; **27**: 533-40.
- 49 Ma A, Malynn BA. A20: linking a complex regulator of ubiquitylation to immunity and human disease. *Nat Rev Immunol* 2012; **12**: 774-85.
- 50 Verstrepen L, Adib-Conquy M, Kreike M *et al.* Expression of the NF-kappaB inhibitor ABIN-3 in response to TNF and toll-like receptor 4 stimulation is itself regulated by NF-kappaB. *J Cell Mol Med* 2008; **12**: 316-29.
- 51 Vossen A, van der Zee HH, Prens EP. Hidradenitis Suppurativa: A Systematic Review Integrating Inflammatory Pathways Into a Cohesive Pathogenic Model. *Front Immunol* 2018; **9**: 2965.
- 52 Kimball AB, Okun MM, Williams DA *et al.* Two Phase 3 Trials of Adalimumab for Hidradenitis Suppurativa. *N Engl J Med* 2016; **375**: 422-34.
- 53 Lowe MM, Naik HB, Clancy S *et al.* Immunopathogenesis of hidradenitis suppurativa and response to anti-TNF-alpha therapy. *JCI Insight* 2020; **5**.
- 54 Huang da W, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 2009; **4**: 44-57.

TABLES

None.

FIGURE LEGENDS

Figure 1. Biological processes enriched amongst up- and downregulated DEGs in lesional vs. perilesional epidermal HS. Table with all statistically enriched FACs as obtained using DAVID FAC analysis (v6.8; <https://david.abcc.ncifcrf.gov>; ⁵⁴) with mean collective p-values of the included GO groups ≤ 0.01 (as calculated by Fisher's exact test). Designated functional clusters are ranked according to their enrichment score defined as $-\log$ of the collective mean p-value of the included GO groups. The top ten upregulated FACs are presented in bold script, the further analyzed gene clusters "*T cell regulation*" and "*Activation of immune response*" are highlighted in red and blue respectively. GO group= gene ontology group.

Figure 2. Confirmation of lesional upregulation of epidermal genes in HS. Comparative qRT-PCR analysis of mRNA expression of selected DEGs from the overrepresented FACs "*T cell regulation*" (**a**, framed in red) and "*activation of immune response*" (**b**, framed in blue) in lesional vs. perilesional epidermis of HS patients. Data represent mean ΔCT values $\pm$ s.d. determined for the respective gene in relation to GAPDH expression. Asterisks indicate statistically significant mRNA induction in n=10 experiments (* $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$ paired t-test; CT values for the indicated gene < 40 cycles). CT, cycle threshold; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; ns, not significant; PLE, perilesional epidermis; LE, lesional epidermis.

Figure 3. Dominance of AP-1- and JAK-STAT-sites within the promoters of regulated genes in lesional HS epidermis. Overrepresented TF binding sites within the promoters of all ≥ 2.0 fold differentially regulated epidermal genes in lesional vs. perilesional HS samples as

identified by large scale promoter analysis (PSCAN; ²¹). Blue colored bars represent overrepresented AP-1 TF sites, red colored ones sites corresponding to the JAK-STAT signaling pathway and purple ones TF sites of the NF- κ B signaling pathway. Grey bars denote overrepresented sites belonging to other TF families. The overrepresentation score corresponds to $1/p$ -value. Cut-off for upregulated TF binding sites was arbitrarily set at an overrepresentation score >1000.00 (corresponding to $p < 0.001$), for downregulated ones at >20 (corresponding to p -value < 0.05). TF, transcription factor.

Figure 4. Lesional HS epidermis exhibits a JNK-mediated AP-1 stress response. (a)

Immunohistochemistry shows an increased JNK and c-Jun phosphorylation in lesional as compared to intraindividual perilesional HS skin. Healthy donor skin served as negative control (not shown). Pictures are representative for 4 individual donors per group. Scale bars indicate 50 μ m. **(b)** Western blot analysis from perilesional and lesional HS epidermis of three individual patients confirming stronger lesional p-JNK activation in lesional HS. LS, lesional skin; PLS, perilesional skin; PLE, perilesional epidermal lysates; LE, lesional epidermal lysates.

Figure 5. Lesional HS epidermis shows a prominent STAT1 signature. (a)

Overrepresented TF binding sites in the FAC “Innate immune response”. Red colored bars represent overrepresented STAT binding sites. Cut-off was set at an overrepresentation score >100.00 ($p < 0.01$). **(b, c)** Anti-p-STAT1 immunohistochemistry of **(b)** perilesional or **(c)** lesional HS epidermis. Two representative images for stainings of $n = 4$ individual donors per group are shown; scale bars indicate 50 μ m. **(d, e)** pSTAT1 immunoblots of total lysates from **(d)** perilesional and lesional HS epidermis of four individual patients, or **(e)** type I IFN- (10 U/ml) and IFN γ - (100 ng/ml) stimulated cultured hEK derived from lesional HS epidermis.

hEK, human epidermal keratinocytes; PLS, perilesional skin; LS, lesional skin; PLE, perilesional epidermal lysates; LE, lesional epidermal lysates.

Figure 6. IFN responsiveness and JAK1 inhibitor sensitivity of selected lesionally induced epidermal genes. (a, b) qPCR analysis of selected DEGs upon stimulation of HS-derived hEKs with (a) diluent (dil), IFN type I (10 U/ml), IFN γ (100 ng/ml), TNF α (50 ng/ml) or a combination of the respective IFNs with TNF α for 48 h, or (b) upon 48 h mono- or co-stimulation with IFN γ (100 ng/ml) and TNF α (50 ng/ml) +/- 250 nM of the JAK1-inhibitor upadacitinib. Data represent mean fold mRNA induction $\pm$ s.d. of the DEG in relation to GAPDH expression (at least n = 3 independent experiments with cells from different donors). *p $\leq$ 0.05, **p $\leq$ 0.01, ***p $\leq$ 0.001; one-way ANOVA with post-hoc Dunnett correction. (c) Representative pSTAT1 Western blot with lysates of hEK stimulated in parallel.

Clusters of upregulated DEGs in epidermal HS	Enrichment Score
Response to external biotic stimulus	11,59
Cytokine response	6,15
Cellular response to molecules of bacterial origin	5,38
Cell migration	5,25
Neutrophil chemotaxis	4,80
Blood vessel morphogenesis	4,16
Leukocyte differentiation	4,08
T cell regulation	3,71
Activation of immune response	3,65
Blood coagulation	3,47
Regulation of TNF family cytokine production	3,05
Regulation of protein transport	3,04
Regulation of protein kinase activity	3,02
Negative regulation of leukocyte differentiation	3,01
Positive regulation of lymphocyte activation	2,86
Mast cell activation	2,84
Myeloid leukocyte differentiation	2,82
Regulation of wound healing	2,80
Leukocyte degranulation	2,80
Innate immune response	2,75
Negative regulation of leukocyte activation	2,73
Cell death	2,70
Regulation of protein phosphorylation	2,55
Positive regulation of leukocyte chemotaxis	2,42
Leukocyte degranulation	2,40
Negative regulation of signal transduction	2,39
Entry into other organisms involved in symbiotic interaction	2,35
(Metal) ion homeostasis	2,35
Positive regulation of cell migration	2,34
Mast cell degranulation	2,34
Cation homeostasis	2,24
Positive regulation of T cell proliferation	2,22
Positive regulation of chemokine production	2,20
Regulation of MAPK cascade	2,20
Regulation of locomotion	2,17
Response to insulin	2,13
Cell differentiation involved in embryonic placenta development	2,12
Positive regulation of phosphate metabolic process	2,12
JAK/STAT cascade	2,11
Positive regulation of protein serine/threonine kinase activity	2,06
Positive regulation of adaptive immune response based on somatic recombination of immune receptors built from immunoglobulin	2,02
Negative regulation of insulin receptor signaling pathway	2,02
Clusters of downregulated DEGs in epidermal HS	Enrichment Score
Neurogenesis	3,03
Positive regulation of nervous system development	2,43
Cardiovascular system development	2,09

Figure 1

(a) “T cell regulation”

(b) “Activation of immune response”

Figure 2

Transcription factor binding sites

Figure 3

Figure 4

Figure 5

(a)

“T cell regulation”

“Activation of immune response”

(b)

“T cell regulation”

“Activation of immune response”

(c)

Figure 6