1 **Title Page:**

2 **Type of the article:** Research article

3 **Title:** Prevalence of SARS-CoV-2: An age-stratified, population-based, sero-epidemiological survey
4 in Islamabad, Pakistan

5 **Running title:** Seroprevalence of SARS-CoV-2 in Islamabad-Pakistan

6

7 Mirza Amir Baig, Jamil Ahmed Ansari, Aamer Ikram, Mumtaz Ali Khan, Muhammad Salman, Zakir
8 Hussain, Mirza Zeeshan Iqbal Baig, Ambreen Chaudhry, Muhammad Wasif Malik, Khurram Shahzad
9 Akram, Asim Saeed, Muazzam Abbas Ranjha, Faisal Sultan, Sohail Sabir

10

11 **Authors' Affiliation:** Field Epidemiology and Disease Surveillance Division, National Institute of
12 Health, Islamabad, Pakistan

13

14 **Scope of this study:**

15 This relatively small and focused study in context of severe local transmission gives an estimation of
16 burden of disease in Islamabad Capital Territory of Pakistan and will aid as a reference for national
17 large-scale surveys. In the context of an overburdened health system and high mortality attributable
18 to COVID-19, regional seroprevalence data will deliver valuable information to customize public
19 health interpolations against this epidemic.

20

21 **Corresponding Author:**

22 Ambreen Chaudhry (MSPH)
23 ORCID: 0000-0002-1549-2445

24 National Institute of Health
25 Park Road, Chak Shehzad,
26 Islamabad, Pakistan

27 Phone: +92 3339949969

28 Email: ambr.chaudhry@gmail.com

29 Abstract

30 **Introduction:**

31 Serological surveys are valuable tools to evaluate the extent of disease transmission,
32 measuring preventive effectiveness and proportion of asymptomatic individuals. This age-
33 stratified, serological survey was aimed to measure the COVID-19 distribution and
34 determinants in district Islamabad of Pakistan.

35 **Methodology:**

36 Three-stage cluster sampling, using population proportionate to size technique, starting with a
37 random number was used. A structured, pretested questionnaire was used after taking
38 informed written consent, to gather demographic, risk factor information.

39 **Results:**

40 Seroprevalence was found 16.5% (AR: 16.5%/100,000). The mean age was 35 (± 16 Years).
41 The majority were male (64%), self-employed (29%), and had primary level education
42 (33%). The highest seroprevalence was found in the 21-30 years age group (24.8%) while the
43 41-50 years age group showed the highest attack rate (112.9/100,000 population). The
44 proportion of the population tested that were asymptomatic was 69% (n=711) while the most
45 frequently reported sign/symptom was cough (99%) followed by fever (20%). No known co-
46 morbidity was reported in 86% (n=884) of respondents while hypertension remained the most
47 reported condition (8%). High seroprevalence was observed in urban areas (12.3%)
48 compared to rural union councils (6.4%). Visiting a house where COVID-19 case was
49 isolated (OR 2, CI 1.38-2.84, $P < 0.001$), history of contact with a known case of COVID-19
50 (OR 1.42, CI 1.11-1.82, $P = 0.005$), and attending a mass gathering (OR 1.21, CI: 1.02-1.42,
51 $p = 0.02$) were significant risk factors associated with contracting an infection. A Chi-Square
52 test of independence showed significant protection while using regular hand hygiene
53 practices (6.5; $p < 0.05$) and regular usage of face masks (8.6; $p < 0.05$).

54 **Conclusion:**

55 Seroprevalence gives a direct estimation of population groups exposed to the virus. A
56 remarkable difference in prevalence is found in urban and rural areas, extreme age groups,
57 and socioeconomic statuses, suggesting targeted public health interventions. Sero-studies are
58 affordable counterparts of molecular testing where quick estimation, prevention
59 effectiveness, and data-driven public health policies are priorities.

60 **Keywords:** COVID-19, Epidemiology, Pakistan, Serology, Survey

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81 Introduction:

82 In Pakistan, the first case of COVID-19 was identified on February 26, 2020 and by June 16, 2020,
83 148,921 cases and 2,839 deaths had been reported ¹. Highest incidence was observed in Karachi,
84 while Islamabad (capital of Pakistan) had a toll of 8,857 cases and 83 deaths during this time period ².
85 Confirmatory test for cases and their contacts (symptomatic or asymptomatic) is reverse transcriptase-
86 polymerase chain reaction (real time RT-PCR), a gold standard confirmatory test. The detected
87 number was largely influenced by the country's daily testing capacity and active surveillance of
88 contacts ³.
89 Epidemiological surveillance of confirmed COVID-19 cases, using RT-PCR, has provided a view of
90 tip of the iceberg, referring to the spectrum of the disease from severe infection to asymptomatic
91 transmission, and has been proven suitable for acute infections with signs and symptoms ⁴. Immune
92 response to SARS-CoV-2 is varied and largely dependent on the severity of infection, host antibody
93 response and is usually detectable after 7-14 days ^{5 6}. Serological testing has advantage in disease
94 surveillance in quantifying proportion of population exposed in a given period of time, where
95 antibodies are used as a marker of partial or total immunity.
96 During this pandemic, seroprevalence studies has been carried out in confined populations, such as
97 hospitals and nursing homes, health care workers or blood donors ⁷ where humoral immune response
98 is measured but results are not generalizable. Sub national seroprevalence study results has been
99 dissimilar in diverse areas, depending on magnitude of outbreak like; 1.5% of seropositivity was
100 observed in Santa Clara, California while it was 32% in Boston which was considered to be a hot
101 spot for COVID-19 outbreak ⁸, likewise, 7.1% in Atlanta USA, 9.6% in Wuhan, China and 21% in
102 Guilan, Iran ⁵.
103 In Pakistan, attempts have been made to estimate the seroprevalence of COVID-19 in confined
104 populations (police squads) ⁹ and in low and high transmission areas only ¹⁰ which are quite small and
105 contained. A larger population based, age and gender stratified survey was needed to estimate the
106 population at risk and factors associated with the exposures to SARS-CoV-2 ¹¹. Since Islamabad
107 Capital Territory (ICT) was the 4th most affected city of Pakistan and to combat the current scenario,
108 an enhanced surveillance strategy, based on population-based methods, was required ¹². In an

extensive literature review, use of serological testing of SARS-CoV-2 antibodies in general population, its role in estimation of burden of disease in different demographics during an epidemic, has been done but we could not find any evidence on the comparison of serology with molecular testing. Therefore, Field Epidemiology and Laboratory Training Program (FELTP) of Pakistan in collaboration with National Institute of Health (NIH) conducted a survey in district Islamabad, the capital of Pakistan, where age and gender stratification were used to help identify the exposed and at-risk population.

The primary objectives of the study were:

1. To estimate the extent of exposure to SARS-CoV-2 on demographic parameters
2. To estimate the proportion of asymptomatic infection in the population and prevalence of co-morbid conditions with SARS-CoV-2 exposure
3. To estimate the factors associated with SARS-CoV-2 exposure

Methods:

A community-based, age-stratified cross-sectional survey was conducted in district Islamabad in June 2020. Islamabad is divided into two administrative units; urban and rural. The World Health Organization's (WHO) "*Population-based age-stratified sero-epidemiological investigation protocol for COVID-19 virus infection*" was followed to conduct this study¹³.

Referring to 2017-18 census¹⁴, total population of Islamabad is 200,657 with 336,182 households.

The following formula was used to calculate the sample size:

$$n = [Z^{2\alpha} (P) * (1-P)] \div d^2 = 6333$$

A three-staged stratified sampling procedure was adopted, using Population Proportionate to Size (PPS) at each stage. Union councils (UCs) are the lowest administrative units in a district. At the first stage, as per Pakistan Bureau of Statistics (PBS)¹⁴ all urban and rural union councils (UCs) were enrolled (n=44) according to their population two each strata. The sample size was equally divided between rural (n=31, 49%), and urban (n=13, 51%) union councils for equal representation of both populations where urban UCs contain sectors and rural UCs contain villages in them. At the second stage, all the villages and sectors in these strata (rural and urban) were listed according to their

population and again PPS sampling techniques was used. The sample was converted into households, and according to PBS, average five individuals live in a household. Finally, the number of households were enlisted, according to the data available at respective Town Management Office and Pakistan Bureau of Statistics, based on PPS techniques. A household was defined as “*a dwelling with a common kitchen or common opening onto a shared space.*” Using a systematic random sampling approach, a sampling interval was calculated by dividing the total number of households by number of households allocated per village/sector. All persons living in a household were included in the study for representation of all ages and genders (Figure 1).

A structured questionnaire was designed to collect demographics, disease status, clinical features, exposure information and other epidemiological parameters. Pilot testing was conducted for ensuring reliability and validity of the tool, in the neighboring district of Rawalpindi which guided further editing and improvement of the questionnaire.

Survey teams were trained on key components of the survey i.e., questionnaire administration, infection prevention and control (IPC) measures, sample collection, labelling, transportation and risk communication.

An institutional review board (IRB) approval was obtained from National Institute of Health (NIH). A consent form was developed and translated into local language for the purpose of enabling the respondents to understand their autonomous right to decide to participate in the study and announcing the beneficence and justice rendered in the study¹⁵.

Strict infection control measures set forth by Public Health Laboratories Division, NIH were adopted by trained laboratory staff. Blood samples were collected from participants upon recruitment in the investigation for antibody testing with chemiluminescent immunoassay (CMI). CE/FDA approved in vitro diagnostic kits for ELISA were used (Abbot’s Diagnostic IgG/IgM RTD). Nasopharyngeal swabs were also collected from symptomatic and/or high-risk respondents (age>60 years or have contact with COVID-19 confirmed case).

Data was coded to mask identities of respondents, double entered, validated and password protected at central office. All hard copies were kept in the lock and key. Response rate was 100% and the missed entries on participants' information were completed by follow-up calls.

Statistical analyses were done using Epi Info-7 software. Frequencies, rates and ratios were calculated for demographics, and prevalence was computed. Crude attack rates, age, gender and area-specific attack rates were determined. Bivariate and multivariate analyses were carried out for risk factors at 95% confidence interval (CI) and a p-value less than 0.05. Chi-Square test was used to evaluate the significance of protective variables.

Results:

A total of 6,333 individuals were approached at their doorsteps, 100% provided consent and blood samples. No test results were inconclusive. Of 6,333 individuals enrolled, 1033 (16.3%) showed a reactive antibody test. Mean age of all seropositive individuals was 35 years (± 16 years). Majority of seropositive individuals were male (64%), had at least primary level education (88%) and were self-employed (59%) (Table 1).

Highest seroprevalence was found in 21-30 years age group (24.8%) while highest attack rate was found in 41-50 years (11.3/10,000 population) (Table 2).

Of all seropositive individuals, 711 (69%) were found asymptomatic within previous 14 days. Among symptomatic individuals, cough was the most reported symptom (99%) followed by fever (20%) and sore throat (12%). Among all respondents, 149 (14%) reported at least one known co-morbid condition and the most frequently reported co-morbidity was Hypertension (8%) followed by Diabetes Mellitus (7%) (Table 3).

Survey was conducted in 44 geographic localities of district Islamabad. High seroprevalence was found in urban union councils i.e., 12.3% in Charge 13 (Sector: H-10 & I-10), 6.9% in charge 6 (Sector: F-6, 7 and Pir Sohawa) compared to lower seroprevalence in rural UCs (Alipur-6.4%, Tarlai Khurd-6.4%).

Out of 1033 seropositive respondents, 42/1033 (4 %) reported that they have visited a house where COVID-19 case was isolated (OR 2.0, CI: 1.38-2.84, p=0.001), 88/1033 (9%) had contact with a known case (OR 1.4, CI: 1.11-1.82, p=0.004), 216/1033 (21%) responded that they attended a mass gathering (OR 1.2, CI: 1.1-1.4, p=0.02) (Table 4).

Multi-variate analysis showed that visiting a house where COVID-19 case was isolated (aOR 2, CI: 1.38-2.84, p-value 0.001) was a significant risk factor associated with contracting an infection (Table 5).

A Chi-Square test of independence indicated that the practices of regular use of a face mask and regular hand washing and/or hand sanitizing were correlated with lower seroprevalence (Table 6).

Among all respondents 458/1033 (44%) were tested for RT-PCR and positivity rate was 14% (n=64).

Discussion:

Our results showed that the antibody response against COVID-19 was 16%, which translates to about 300,000 individuals being exposed to SARS-CoV-2 in district Islamabad. This should be constructed as this is not a reflection of cases and only depicts exposed individuals. It should be realized that these results were obtained through detecting antibodies against the SARS-CoV-2 virus which is not qualified as a diagnostic test for COVID-19. Confirmed cases can only be labelled through RT-PCR testing.

The results are in concordance with surveys conducted in Spain¹⁶ where similar patterns has been documented in the general population and few other surveys conducted in Pakistan in localized and confined communities¹¹. Some small scale studies have also shown low seroprevalence with certain sampling biases such as enrolling only adult population¹⁷. Our study displayed higher positivity rate among males and entrepreneurs which explains the outdoor activities related to these population groups even amidst series of smart lock downs in the district as well as skewed participation of males. On the other hand, low prevalence in extreme age groups has been observed (lower than 10 years and more than 60 years of age) which has been indicated in other studies also, conducted in developed countries including China, which are comparable with our findings (4.5 % and 6.6% respectively)¹⁸.

We found 69% of seropositive individuals were asymptomatic, endorsing the outcome of various researches on stealth disease transmission¹⁹ and a varied spectrum of signs and symptoms in the community and hospitalized individuals. This wide spectrum of clinical presentation has led to delayed diagnosis, delay in reaching health care facility and seeking medical care on one hand while undue fear, stigma and panic in society on the other hand²⁰. This high seroprevalence in asymptomatic individuals confirms the need for a greater serosurvey.

Likewise, our study has shown that more than half of responders did not declare a known co-morbid condition other than COVID-19, contrary to many studies done in hospitalized and older aged individuals who presented with a co-morbid condition with high fatality outcomes. Since we conducted this survey on general population so the results are difficult to compare with researches done in hospitalized and nursing home settings²¹. We identified that among reported co-morbid conditions, hypertension and type-2 diabetes mellitus remained the most frequently reported one. A high prevalence of these non-communicable diseases in our communities has been documented with a maximum prevalence rate of 40% whereas some researchers suggest that more than half of them go undiagnosed due to low health services utility and health care access²². Hence forth, coexistence of COVID-19 with aforementioned NCDs is not unusual. Rural population, though appears less affected, is comprised of a comparatively larger number of elderly people who are disproportionately impacted by the disease. The most germane space has been identified in urban areas but the rural areas of the district Islamabad seem more vulnerable in relative to age distribution.

Our results afford a strong evidence for smart lock down success instigated by government of Pakistan²³ where a significant association of exposure to COVID-19 with visiting a house where a COVID-19 case has been isolated, attending a mass gathering and having an epidemiological link, has been discerned. Equally, the strong association with the much advertised and advocated preventive measures has also been established²⁴. Therefore, this population based, age stratified, sero-epidemiological study has provided a robust evidence on the proportion of Islamabad (ICT) population exposed.

As RT-PCR is gold standard for the detection of disease, sero-testing can only complement the much resource limited and patchy surveillance of the low and middle income countries like Pakistan in an outbreak settings²⁵. Here, it is pertinent to mention that, a serological test cannot replace the confirmatory test, neither the results of one district be generalized across the country. Based on population distribution in terms of age, literacy rate and access to media and health care facilities, results of such sero-surveys may contrast. It has been recognized that the recommendations issued by WHO in March 2020 on “test, test, and test”²⁶ led to a huge burden on health system. Alternatively, rapid antigen/antibody detection tests were opted for many countries irrespective of their economic status, and are now recommended for quick and affordable point-of-care testing to triage exposed individuals. Antibody test detects the host response to virus which may take several days to develop, also it cannot distinguish between acute or previous infection²⁷. Some researchers have documented that high antibody response is associated with the severe disease. Low concurrent antigen and antibody detection, determined in our survey, suggests an exclusion of severely diseased from general population as they require hospitalization²⁸.

Limitations:

This survey, conducted in one district, shows seroprevalence of the same population and surveys using similar techniques in different populations may project different seroprevalence rates depending on the varied demographics, preventive measures taken, lock down policies and extent of exposure to SARS-CoV-2. Serological surveys give only the exposure status of population to SARS-CoV-2 and hence, should not be considered as their disease status. It is important to note that a greater number of males participated in the study compared to females.

Conclusion:

Diagnostics play a crucial role in epidemic response and so is the case with COVID-19. Rapid serology test is a scalable and affordable, population-based testing to estimate the extent of SARS-CoV-2 circulation in the community. Not only it gauges the population at risk, it fortifies the surveillance system to monitor trends over time. Based on our findings, rapid serological diagnostic

testing is recommended in early and recession phase of outbreak to identify hot spots as well as measure the prevention effectiveness during high transmission phase. For astute distribution of limited public health resources, and to slacken off the burden on health care facilities, a rapid point-of-care testing will support in identifying target population and areas for smart lock down. Henceforth, implementation of testing tracking and quarantine (TTQ) strategy and achieving a data driven outbreak containment policies will be facilitated.

Financial support: Authors declare no financial support

Conflict of interest: Authors declare no conflict of interest

Ethical Approval: Institutional review Board of National Institute of Health Pakistan

References:

1. COVID-19 Health Advisory Platform by Ministry of National Health Services Regulations and Coordination . <http://covid.gov.pk/>. Accessed August 24, 2020.
2. Pakistan COVID-19 Corona Tracker. <https://www.coronatracker.com/country/pakistan/>. Accessed September 1, 2020.
3. Sethuraman N, Jeremiah SS, Ryo A. Interpreting Diagnostic Tests for SARS-CoV-2. *JAMA - J Am Med Assoc*. 2020;323(22):2249-2251. doi:10.1001/jama.2020.8259
4. Corman VM, Landt O, Kaiser M, et al. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Eurosurveillance*. 2020. doi:10.2807/1560-7917.ES.2020.25.3.2000045
5. Lai AL, Millet JK, Daniel S, Freed JH, Whittaker GR. Testing for SARS-CoV-2 (COVID-19): a systematic review and clinical guide to molecular and seological in-vitro diagnostic assay. *Lancet*. 2020;395(April):1315.
6. Zhao J, Yuan Q, Wang H, et al. Antibody responses to SARS-CoV-2 in patients of novel coronavirus disease 2019. *Clin Infect Dis*. March 2020. doi:10.1093/cid/ciaa344
7. Valenti L, Bergna A, Pelusi S, et al. SARS-CoV-2 seroprevalence trends in healthy blood donors during the COVID-19 Milan outbreak. *medRxiv*. 2020:2020.05.11.20098442. doi:10.1101/2020.05.11.20098442
8. Madsen T, Levin N, Niehus K, et al. Prevalence of IgG antibodies to SARS-CoV-2 among emergency department employees. *Am J Emerg Med*. 2020. doi:10.1016/j.ajem.2020.04.076
9. Chughtai OR, Batool H, Khan MD, Chughtai AS. Frequency of COVID-19 IgG Antibodies among Special Police Squad Lahore, Pakistan. *J Coll Physicians Surg Pak*. 2020;30(7):735-739. doi:10.29271/jcpsp.2020.07.735
10. Nisar I, Ansari Msc N, Amin Msc M, et al. Serial population based serosurvey of antibodies to SARS-CoV-2 in a low and high transmission area of Karachi, Pakistan. *medRxiv*. August

2020:2020.07.28.20163451. doi:10.1101/2020.07.28.20163451

11. Javed W, Bin Baqar J, Hussain S, Abidi B, Farooq W. Sero-prevalence Findings from Metropolises in Pakistan: Implications for Assessing COVID-19 Prevalence and Case-fatality within a Dense, Urban Working Population. *medRxiv*. August 2020:2020.08.13.20173914. doi:10.1101/2020.08.13.20173914

12. World Health Organization. *Covid 19 Strategy Update.*; 2020.

13. World Health Organization. Population-based age-stratified seroepidemiological investigation protocol for COVID-19 virus infection. *World Heal Organ*. 2020.

14. Pakistan Bureau of Statistics. <http://www.pbs.gov.pk/>. Accessed August 26, 2020.

15. Belmont Report - an overview | ScienceDirect Topics. <https://www.sciencedirect.com/topics/medicine-and-dentistry/belmont-report>. Accessed August 25, 2020.

16. Pollán M, Pérez-Gómez B, Pastor-Barriuso R, et al. Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide, population-based seroepidemiological study. *Lancet*. 2020;0(0). doi:10.1016/S0140-6736(20)31483-5

17. Sood N, Simon P, Ebner P, et al. Seroprevalence of SARS-CoV-2-Specific Antibodies among Adults in Los Angeles County, California, on April 10-11, 2020. *JAMA - J Am Med Assoc*. 2020. doi:10.1001/jama.2020.8279

18. Li W, Zhang B, Lu J, et al. The characteristics of household transmission of COVID-19. *Clin Infect Dis*. April 2020. doi:10.1093/cid/ciaa450

19. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395(10223):497-506. doi:10.1016/S0140-6736(20)30183-5

20. Rothe C, Schunk M, Sothmann P, et al. Transmission of 2019-NCOV infection from an

- 334 asymptomatic contact in Germany. *N Engl J Med*. 2020;382(10):970-971.
- 335 doi:10.1056/NEJMc2001468
- 336 21. Moccia F, Gerbino A, Lionetti V, et al. COVID-19-associated cardiovascular morbidity in
- 337 older adults: a position paper from the Italian Society of Cardiovascular Researches.
- 338 *GeroScience*. 2020;42(4):1021-1049. doi:10.1007/s11357-020-00198-w
- 339 22. Shah N, Shah Q, Shah AJ. The burden and high prevalence of hypertension in Pakistani
- 340 adolescents: A meta-analysis of the published studies. *Arch Public Heal*. 2018;76(1).
- 341 doi:10.1186/s13690-018-0265-5
- 342 23. Pakistan among pioneers of “smart lockdown” approach, says PM Imran - Pakistan -
- 343 DAWN.COM. <https://www.dawn.com/news/1561766>. Accessed August 25, 2020.
- 344 24. Coronavirus Disease 2019 (COVID-19) | CDC. [https://www.cdc.gov/coronavirus/2019-](https://www.cdc.gov/coronavirus/2019-ncov/index.html)
- 345 [ncov/index.html](https://www.cdc.gov/coronavirus/2019-ncov/index.html). Accessed August 24, 2020.
- 346 25. Nkengasong J. Let Africa into the market for COVID-19 diagnostics. *Nature*.
- 347 2020;580(7805):565-565. doi:10.1038/d41586-020-01265-0
- 348 26. WHO Director-General’s opening remarks at the media briefing on COVID-19 - 16 March
- 349 2020. [https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-](https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---16-march-2020)
- 350 [media-briefing-on-covid-19---16-march-2020](https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---16-march-2020). Accessed August 25, 2020.
- 351 27. He X, Lau EHY, Wu P, et al. Temporal dynamics in viral shedding and transmissibility of
- 352 COVID-19. *Nat Med*. 2020;26(5):672-675. doi:10.1038/s41591-020-0869-5
- 353 28. Qu J, Wu C, Li X, et al. Profile of IgG and IgM antibodies against severe acute respiratory
- 354 syndrome coronavirus 2 (SARS-CoV-2). *Clin Infect Dis*. April 2020. doi:10.1093/cid/ciaa489

355

356

357

358 Figure 1: Flow diagram showing three-staged stratified sampling technique

Figure 1: Flow diagram showing sampling Procedure (n=6333).

360

361 Table-1: Summary statistics of seropositive individuals for SARS-CoV-2 in district Islamabad (June
362 2020; n=1033)

	Cases (n)	Frequency (%)	95% CI
Gender			
Male	665	64	61-67
Female	368	36	33-38
Employment status			
Self employed	605	59	55.5-61.5
Unemployed	294	29	25.8-31.3
Government employed	134	13	11-15.2
Education			
No formal education	125	12	10-14
Primary	338	33	30-36
Secondary	15	1	0.9-2.4
Under graduate	295	29	25.9-31.4
Graduate	6	1	0.3-1.3
Post graduate	250	24	22-27

363

364

365

366

367

368

369

370 Table-2: Age distribution of sero-positivity and attack rates for SARS-CoV-2 among residents of
371 district Islamabad, June 2020 (n=1033)

Age group	Seropositive	Positivity rate		Attack rate	
(years)	(n)	%	CI	Per 10,000	CI
≤10	47	4.5	3.4-5.9	0.8	0.7-1.2
11-20	149	14.4	12.4-16.6	3.8	2.7-3.8
21-30	257	24.8	22.3-27.6	7.3	6.4-8.2
31-40	234	22.6	20.2-25.3	9.5	8.3-10.8
41-50	174	16.8	14.6-19.2	11.2	9.7-13.1
51-60	103	9.9	8.2-11.9	7.9	6.5-9.6
≥61	69	6.6	5.3-8.3	5.1	4.0-6.4
CI= 95% Confidence interval					

372

373

374

375

376

377

378

379

380

381

382

383 Table-3: Clinical characteristics of sero-positive individuals for SARS-CoV-2 in district Islamabad,

384 June 2020 (n=1033)

	Numbers	Frequency	95% CI
	(n)	(%)	
Signs and symptoms ☐			
Cough	319	99	97.7-99.2
Fever	202	20	17.2-22.1
Sore throat	119	12	9.7-13.6
Headache	109	11	8.8-12.5
Shortness of breath	67	7	5.1-8.1
Diarrhea	58	6	4.3-7.1
Runny nose	49	5	3.6-6.2
Co-morbidities			
No known co-morbidity	884	86	83.3-87.5
Hypertension	79	8	6.2-9.43
Diabetes Mellitus	74	7	5.7-8.9
IHD	30	3	2.0-4.1
CLD	14	1	0.2-2.2
CRD	13	1	0.7-2.1
Asthma	10	1	0.5-1.8
Chronic Lung Disease	5	0.5	0.2-1.1

CI=confidence interval, IHD=Ischemic health disease, CLD=Chronic liver disease,

CRD=Chronic renal disease

☐ Self-reported signs and symptoms

385

386 Table-4: Association of risk factors with seroprevalence for SARS-CoV-2 among residents of district
387 Islamabad, June 20202. (n=1033)

	Seropositive (N)	Seronegative (N)	OR	95% CI	P-value
Visited a house where COVID-19 case was isolated					
Yes	42	111	2.0	1.38-2.84	0.001
No	991	5189			
Hospitalization					
Yes	4	12	1.7	0.50-5.30	0.4
No	1029	5288			
Contact with a known case					
Yes	88	325	1.4	1.11-1.82	0.02
No	945	4975			
Visited COVID-19 treatment facility					
Yes	27	110	1.3	0.82-1.93	0.27
No	1006	5190			
Attended a mass gathering					
Yes	216	952	1.2	1.11-1.41	0.34
No	817	4348			
Travelled abroad within 14 days					
Yes	32	196	0.8	0.52-1.23	0.34
No	1006	5190			

OR=Odds ratio, CI=confidential interval,

Table-5: Multi-variate analysis of risk factors associated with seroprevalence for SARS-CoV-2 among residents of district Islamabad June 2020 (n=6333)

Risk factors	Seropositive (N)	Seronegative (N)	aOR	95% CI	p-value
Visited a house where COVID-19 case was isolated					
Yes	42	111	2.0	1.38-2.84	0.001
No	991	5189			
History of contact with a known case					
Yes	88	325	1.4	1.11-1.82	0.005
No	945	4975			
Attended a mass gathering					
Yes	216	952	1.2	1.02-1.42	0.02
No	817	4348			

aOR=Adjusted odds ratio,

395 Table-6: Association of preventive measures and SARS-CoV-2 exposure among residents of district
396 Islamabad, June 2020 (n=6333)

Preventive measures	SARS-CoV-2 Positive (N)	SARS-CoV-2 Negative (N)	Chi-Square Value	P-value
Regular use of face mask				
Regularly	716	3423	8.6	0.03
Occasionally	176	1014		
No	141	863		
Hand Washing/Hand Sanitizing				
Regularly	812	4292	6.5	0.01
Occasionally	131	661		
No	90	347		

397