1 Risk of cardiovascular disease after a diagnosis of common

2 psychiatric disorders: a matched cohort study of disease

3 susceptibility and progression trajectory in the UK Biobank

4 Xin Han^{1,2}, MD; Yu Zeng^{1,2}, MS; Yannan Shang^{1,2}, PhD; Yao Hu^{1,2}, BS; Can Hou^{1,2}, PhD;
 5 Huazhen Yang^{1,2}, MS; Wenwen Chen³, MS; Zhiye Ying^{1,2}, MS; Yajing Sun^{1,2}, MS; Yuanyuan
 6 Qu^{1,2}, MA; Junren Wang^{1,2}, MS; Wei Zhang^{1,2,4}, MD; Fang Fang⁵, MD, PhD; Unnur A.
 7 Valdimarsdóttir^{6,7,8}, PhD; Huan Song^{1,2,6*}, MD, PhD

8 1. West China Biomedical Big Data Center, West China Hospital, Sichuan University,
 9 Chengdu, China
 10 2. Med-X Center for Informatics, Sichuan University, Chengdu, China
 11 3. Division of Nephrology, Kidney Research Institute, State Key Laboratory of Biotherapy
 12 and Cancer Center, West China Hospital, Sichuan University, Chengdu, China
 13 4. Mental Health Center, West China Hospital of Sichuan University, Chengdu, China
 14 5. Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden
 15 6. Center of Public Health Sciences, Faculty of Medicine, University of Iceland, Reykjavík,
 16 Iceland
 17 7. Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm,
 18 Sweden
 19 8. Department of Epidemiology, Harvard T H Chan School of Public Health, Boston, MA,
 20 USA

21 * Corresponding to: Huan Song, MD, PhD, West China Biomedical Big Data Center, West
 22 China Hospital, Sichuan University, Guo Xue Lane 37, Chengdu, China (telephone:

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

23 86-28-85164176 e-mail: songhuan@wchscu.cn)

24 **Word count:** abstract 245; main text 3341

25 Abstract

26 **Background:** Whether associations between psychiatric disorders and cardiovascular
 27 diseases (CVDs) can be modified by disease susceptibility and the temporal pattern of these
 28 associated CVDs remain unknown.

29 **Methods:** We conducted a matched cohort study of UK Biobank including 35,227 patients
 30 with common psychiatry disorders (anxiety, depression, and stress-related disorders) between
 31 1997 and 2019, together with 176,135 sex- and birth year- individually matched unexposed
 32 individuals.

33 **Results:** The mean age at the index date was 51.76 years, and 66.0% of participants were
 34 females. During a mean follow-up of 11.94 years, we observed an elevated risk of CVD
 35 among patients with studied psychiatry disorders, compared with matched unexposed
 36 individuals (hazard ratios [HRs]=1.16, 95% confidence interval [CI]: 1.14-1.19), especially
 37 during the first six months of follow-up (HR=1.59 [1.42-1.79]). To assess the modification
 38 role of disease susceptibility, we stratified analyses by family history of CVD and by CVD
 39 PRS, which obtained similar estimates between subgroups with different susceptibilities to
 40 CVD. We conducted trajectory analysis to visualize the temporal pattern of CVDs after
 41 common psychiatry disorders, identifying primary hypertension, acute myocardial infarction,
 42 and stroke as three main intermediate steps leading to further increased risk of other CVDs.

43 **Conclusions:** The association between common psychiatry disorders and subsequent CVD is
 44 not modified by predisposition to CVD. Hypertension, acute myocardial infarction, and stroke
 45 are three initial CVDs linking psychiatric disorders to other CVD sequelae, highlighting a need
 46 of timely intervention on these targets to prevent further CVD sequelae among all individuals

47 with common psychiatric disorders.

48 **Key words:** Psychiatry disorders, Cardiovascular disease, Disease susceptibility, Disease

49 trajectory

50 **Funding:** This work is supported by the National Natural Science Foundation of China (No.

51 81971262 to HS), 1.3.5 project for disciplines of excellence, West China Hospital, Sichuan

52 University (No. ZYYC21005 to HS), EU Horizon2020 Research and Innovation Action Grant

53 (847776 to UV and FF).

Introduction

Both psychiatry disorders and cardiovascular disease (CVD) are leading causes of disability, morbidity, and mortality worldwide according to the World Health Organization (WHO)(DALYs et al., 2015; Whiteford et al., 2013). During the last decades, accumulating evidence indicate a link between psychiatry disorders and CVD(Han et al., 2021; Kollia et al., 2017; Momen et al., 2020; Song et al., 2019). Specifically, elevated risks of multiple CVDs, including ischemic heart disease(Janszky, Ahnve, Lundberg, & Hemmingsson, 2010; Wium-Andersen et al., 2019), hypertension(Stein et al., 2014), heart failure(Song et al., 2019), atherosclerosis(Yu, Ho, Lam, Woo, & Ho, 2010), and stroke(Momen et al., 2020; Swain et al., 2015), have been reported among patients with psychiatry disorders. Potential mechanisms linking psychiatric disorders and cardiovascular abnormalities are multifactorial, including chronic inflammation(Miller, Maletic, & Raison, 2009), alterations in hypothalamic–pituitary–adrenal (HPA) axis(Jokinen & Nordstrom, 2009), and unhealthy lifestyle (e.g., smoking)(Weinberger et al., 2017).

Recent genome wide association studies (GWAS) have suggested a genetic predisposition to different CVDs such as coronary artery disease(Consortium et al., 2013; Nikpay et al., 2015) and stroke(Network & International Stroke Genetics, 2016), including multiple single nucleotide polymorphisms (SNPs) significantly associated with these traits. Likewise, familial factors, as a proxy for the joint impact of genetic background and shared environmental and lifestyle factors, have been shown to play an important role in the development of CVD(Forsdahl, 1979; Iannotti, Zuckerman, & Rifai, 2000). However, data are scarce regarding whether predisposition to CVD can modify the association between psychiatry

disorders and subsequent CVD. Findings from a Danish twin study did not support a role of shared familial factors on the associations of depression with CVD(Wium-Andersen et al., 2020). In our previous study, however, we noted attenuated risk increase of CVD among patients with stress-related disorders when comparing them with their full siblings, relative to comparison with age- and sex-matched unexposed individuals from the general populations(Song et al., 2019). These inconsistent results call for further investigations of this question, ideally with direct assessment on disease susceptibility to CVD such as individual-level genotype data.

Moreover, as there are shared etiologies and comorbidity between different CVDs(Collaboration, 2015; Laurent & Boutouyrie, 2015), a further question is whether there are specific CVDs that more directly related to psychiatric disorders and may lead to other CVDs downstream. We therefore applied disease trajectory analysis to map out the temporal pattern of CVDs after psychiatric disorders (Han et al., 2021; Siggaard et al., 2020). As the illustration of the progression network of CVDs can aid the identification of key CVDs first affected by psychiatric disorders, results of such analysis can be of importance in terms of helping identify targets for preventive measures with the goal of reducing the risk of subsequent CVDs.

In the present study, taking advantage of enriched phenotypic information, complete follow-up and individual-level genotyping data in the UK Biobank, we aimed to clarify whether disease susceptibility to CVD modifies the association of common psychiatry disorders with subsequent CVDs. Further, we used trajectory network analysis to identify the key CVDs directly associated with a prior diagnosis of common psychiatric disorders.

Methods

Study design

UK biobank is a community-based cohort study which enrolled a half million participants aged 40 to 69 years at recruitment between 2006 and 2010 across England, Scotland, and Wales. Baseline questionnaires and biological samples were collected at recruitment with detailed description elsewhere (Sudlow et al., 2015). Health-related outcomes were obtained by linkages to a range of health records (Sudlow et al., 2015). UK Biobank inpatient hospital data were mapped from the Hospital Episode Statistics database, the Scottish Morbidity Record, and the Patient Episode Database, covering 89% of UK Biobank participants since 1997. The primary care data were derived from various general practitioner computer system suppliers, providing information on primary care visit for ~45% participants since 1985 (Sudlow et al., 2015).

Based on these data, we conducted a matched cohort study. After the exclusion of 19 individuals who withdrew their informed consent forms, and 3 individuals who lost to follow-up before a diagnosis of common psychiatry disorders, we identified an exposed cohort of 39,960 participants who received their first diagnosis of anxiety, depression, or stress-related disorders between 1 January 1997 and 31 December 2019 (**Figure 1**). Individuals with those common psychiatry disorders were included in the exposed cohort from the date of their diagnosis (i.e., the index date). We then excluded individuals with a history of any CVD before the index date ($n=4,733$), leaving 35,227 exposed individuals in the final analysis. For each exposed individual (i.e., the index patient), we randomly selected five unexposed individuals per index patient, individually matched by sex and birth year (± 2

years), from individuals who were free of those common psychiatry disorders and CVD at the diagnosis date of the index patient.

Follow-up

For all participants, follow-up was from the index date until the diagnosis of any or specific CVDs, death, loss to follow-up, or the end of follow-up (31 December 2019), whichever occurred first. The follow-up for matched unexposed individuals was additionally censored at the time of their first diagnosis of studied psychiatry disorders, if any, during the follow-up.

Ascertainment of common psychiatric disorders

Common psychiatry disorders were defined as a diagnosis of anxiety, depression, or stress-related disorders, based on the primary diagnoses from UK Biobank inpatient hospital and primary care data. **eTable 1 in the supplement** lists details of diagnosis codes, including the 9th to 10th versions of the International Classification of Diseases (ICD) codes for inpatient hospital data, and version 2 and version 3 read codes (i.e., Read v2 and Read v3) for primary care data. We considered a diagnosis of other psychiatry disorders before the index date as history of other psychiatry disorders. Inpatient mental health diagnosis from the Hospital Episode Statistics database was validated with a median accuracy of 73% against a gold-standard research diagnosis(Davis et al., 2018), while the positive predictive value (PPV) of Read codes for anxiety and depression from primary care data was 61%-76% against the five-item Mental Health Inventory(John et al., 2016).

Ascertainment of CVD

With a specific focus on severe CVD consequences that require hospital care, we defined an incident CVD (any or six major subtypes including hypertensive diseases, ischemic heart

diseases, embolism and thrombosis, arrhythmia/conduction disorders, heart failure, and cerebrovascular disease) as a corresponding diagnosis documented in the UK Biobank inpatient hospital data, according to ICD-9 and ICD-10 codes (**eTable 1 in the supplement**). For disease trajectory analysis, we used the 3-digit ICD-10 codes for CVD identification based on the UK Biobank inpatient hospital data and combined CVDs with clinical or biological similarities(Han et al., 2021; Yang et al., 2019) (**eTable 2 in the supplement**). The inpatient diagnosis of CVDs in UK Biobank has been validated, showing a satisfactory accuracy (i.e., PPV= 72% for coronary heart disease and >90% for stroke(Kivimaki et al., 2017; Woodfield et al., 2015)).

Disease susceptibility of CVD

In the present study, we assessed the disease susceptibility to CVD in two ways. First, we defined family history of CVD by a self-reported history of heart diseases and stroke among first-degree relatives (i.e., father, mother and siblings), obtained from baseline questionnaires (Field ID: 20107, 20110, 20111). Second, a polygenic risk score (PRS) for CVD was calculated for 376,019 individuals with eligible genotyping data (6,099,107 SNPs available after a standard quality control(Marees et al., 2018)) using LDpred2(Prive, Arbel, & Vilhjalmsen, 2020), a method for PRS computation based on combination of summary statistics and a matrix of correlations between genetic variants. The independent SNPs associated with CVD were derived from meta-analyzed GWAS summary data from CARDIoGRAMplusC4D Consortium(Nikpay et al., 2015). In a validation step, the generated CVD PRS was shown to be significantly associated with an increased risk of the CVD

phenotype in our dataset (odds ratio [95% confidence intervals (CIs)]: 1.43 [1.40-1.45], per one unite increase of PRS).

Covariates

We retrieved information on educational levels, ethnicity, smoking status, Townsend deprivation index (TDI), and body mass index (BMI) through the baseline questionnaires. Based on the UK Biobank inpatient hospital data, we calculated the Charlson comorbidity index (CCI) on the index date for each individual, as a proxy of their baseline comorbidity level(Quan et al., 2005).

Statistical analysis

We used Cox model to estimate hazard ratios (HRs) with 95% CIs of any CVD, in relation to a previous diagnosis of common psychiatry disorders, using time since the index date as the underlying time scale. The models were stratified by matched factors (sex and birth year), and partially or fully adjusted for educational levels, ethnicity, smoking status, TDI, BMI, history of other psychiatry disorders, family history of CVD, and CCI. As a high-risk time window for CVD has been indicated after diagnosis of psychiatric disorders(Song et al., 2019), we first visualized the changes of HRs over time by stratifying the Cox models to different follow-up periods (≤ 3 , 3-6, 6-12, 12-18, 18-24, 24-60, 60-120, 120-240, and >240 months follow-up). As the HR during the first six months of follow-up period was greater than those of the later times (**eFigure 1 in the supplement**), we assessed the risk of CVD within and beyond the first six months (i.e., ≤ 6 or >6 months of follow-up) separately in later analyses. In addition, we did separate analyses for six main types of CVDs as well as for anxiety, depression and stress-related disorders.

To determine the impact of disease susceptibility to CVD on the association between common psychiatric disorders and CVD, we conducted stratified analyses by family history of CVD (yes or no) and by level of CVD PRS (high, moderate, or low by tertile distribution).

The temporal progression of CVD subsequent to a diagnosis of psychiatry disorders was explored using trajectory network analysis as described in our previous study (Han et al., 2021). In brief, a phenome-wide association analysis (PheWAS) was performed to identify specific CVDs associated with a prior diagnosis of psychiatry disorders. In this step, the p-value for statistical significance was set to 0.05/number of analyses performed (Bonferroni corrections), to account for multiple testing. We then used binomial tests to identify pairs of CVD events with a temporal order, and assessed the magnitude of the associations between CVD pairs through a nested case-control study design and conditional logistic regression (eFigure 2 in the supplement).

To test the robustness of our results to the definition of CVD, we reassessed these associations by identifying CVD solely based on the primary diagnosis from UK Biobank inpatient hospital data. All statistical analyses were conducted by R (version 4.0.2), PLINK (version 1.9), Python (version 3.8) and Cytoscape (version 3.8.0). Codes script used in the primary analyses are available (Source code 1).

Results

In total, we identified 35,227 patients with studied common psychiatry disorders, together with 176,135 sex- and birth year- matched unexposed individuals (Figure 1). Table 1 shows the basic characteristics of the study population. Mean age at index date was 51.76 years, and

66.0% of participants were females. Compared with matched unexposed individuals, patients with common psychiatry disorders were more likely to be smokers (either previous or current, 47.4% vs 41.6%), have a history of other psychiatric disorders (16.0% vs 4.1%), have lower educational level (41.0% vs 44.9% for university level), but higher BMI (26.6% vs 22.2% for more than 30 Kg/M²).

During a mean follow-up time of 11.94 years, 10,619 (crude incidence rate: 25.41 per 1000 person-years) and 43,192 (20.52 per 1000 person-years) incident cases of CVD were identified among the exposed and matched unexposed individuals, respectively. According to the fully adjusted Cox model, the risk of CVD was higher for patients with common psychiatry disorders compared with matched unexposed individuals (HR [95% CIs]: 1.16 [1.14-1.19]), especially during the first six months of follow-up (1.59 [1.42-1.79], **Table 2**). Except for heart failure, all other five main categories of CVD were associated with psychiatry disorders, with the strongest association observed for cerebrovascular disease (1.28 [1.19-1.38], **Table 2**). The result patterns were similar for the periods beyond six months of follow-up, while the highest HR was for heart failure within six months of follow-up (2.66 [1.42-4.97], **Table 2**). Comparable results were found for anxiety, depression, and stress-related disorders (**eTable 3 in the supplement**).

The role of disease susceptibility

We observed similar associations among individuals with and without a family history of CVD (1.54 [1.31-1.80] vs 1.61 [1.25-2.07], $P_{\text{interaction}} = 0.7703$ within six months of follow-up, and 1.15 [1.11-1.18] vs 1.21 [1.16-1.27], $P_{\text{interaction}} = 0.0683$ beyond six months follow-up, **Table 3**). Likewise, comparable risk elevations of CVD were observed among individuals

with different levels of CVD PRS, within six months of follow-up (2.29 [1.58-3.32] and 1.41 [0.98-2.02] for low and high PRS, $P_{\text{interaction}} = 0.0667$) and beyond (1.25 [1.16-1.34] and 1.18 [1.10-1.25] for low and high PRS, $P_{\text{interaction}} = 0.2412$) (**Table 3**). Analyses on subtypes of CVD by family history of CVD and CVD PRS showed similar result patterns (**eFigure 3 in the supplement**).

Trajectory network analysis

15 CVDs with more than 150 cases among patients with common psychiatry disorders were involved in the PheWAS analysis (**eFigure 2 in the supplement**). Among these, eight CVDs showed significant associations with a prior diagnosis of common psychiatry disorders (**eTable 5 in the supplement**), forming 56 CVD pairs among which 21 passed the threshold of prevalence (i.e., experienced by at least 75 individuals). Finally, 12 pairs contributed to the visualized trajectory networks (**Figure 2 and eTable 5 in the supplement**), presenting as a CVD disease tree originated from primary hypertension, acute myocardial infarction and stroke, which further connected to a wide range of other CVDs.

In the sensitivity analysis, defining the CVD cases solely by primary diagnoses, instead of all diagnoses, in UK Biobank inpatient data led to largely similar estimates (**Tables 6 and 7 in the supplement**). However, the number of CVDs involved in the trajectory network was substantially reduced, due to insufficient data power, where we had only 2 CVD pairs in the disease tree with an initial node of acute myocardial infarction (**eFigures 4 and 5 in the supplement**).

Discussion

In this matched cohort study of UK Biobank, we found that individuals with common psychiatry disorders (i.e., anxiety, depression, and stress-related disorders) were at elevated risk of multiple CVDs, especially within first six months of follow-up. With mutually verified results obtained from stratified analyses by family history of CVD and by CVD PRS, the observed associations seemed to be constant across individuals with different predisposition to CVD. Furthermore, trajectory network analysis indicated that primary hypertension, acute myocardial infarction, and stroke were the CVD types firstly affected by a diagnosis of common psychiatric disorders which then further led to other CVDs downstream. This result highlights the need of timely intervention on these targets for preventing or interrupting further CVD sequelae among individuals with common psychiatric disorders.

In line with our results, multiple large-scale cohort studies have reported an association between a diagnosis of psychiatry disorders and subsequently elevated risk of CVD, despite often insufficient control of important confounders, such as lifestyle (Han et al., 2021; Song et al., 2019; Stein et al., 2014; Wium-Andersen et al., 2019) and familial factors (Han et al., 2021; Kollia et al., 2017; Momen et al., 2020; Stein et al., 2014; Swain et al., 2015; Yu et al., 2010). Our finding that the association was not modified by family history of CVD gains support from two recent Danish twin studies (Wium-Andersen et al., 2020; Wium-Andersen et al., 2019), both of which concluded that genetic factors and early-life environment did not explain the association of depression with ischemic heart disease and stroke using a study sample of 6714 twins (Wium-Andersen et al., 2019) and 63,038 twins (Wium-Andersen et al., 2020), respectively. The Nurses' Health Study II and the Normative Aging Study also revealed no attenuation in the magnitude of association between psychiatric disorders and CVD after

additional adjustment for family history of CVD(Kubzansky, Koenen, Spiro, Vokonas, & Sparrow, 2007; Sumner et al., 2015). Our analysis on CVD PRS expanded this knowledgebase further and, for the first time, showed similar associations across groups with different levels of disease susceptibility to CVD. Taken together, these results underscore the importance of surveillance for CVD events among individuals with psychiatry disorders, regardless of disease liability.

Based on the trajectory network analysis, we identified three specific CVDs first affected by a prior diagnosis of common psychiatric disorders (i.e., primary hypertension, followed by acute myocardial infarction and stroke). Although there are no existing studies on CVD trajectory networks after psychiatry disorders, the direct effect of psychiatric disorders on specifically these CVDs has been implied in previous studies. Mendelian randomization analyses showed associations of genetic predisposition to depression with ischemic stroke(Cai et al., 2019) and myocardial infarction(Lu, Wang, Georgakis, Lin, & Zheng, 2021), suggesting a causal link between depression and these CVDs. Also, findings of experimental studies indicated direct physiological responses to mental stress, including endothelial dysfunction (monkeys(Strawn et al., 1991) and human(Ghiadoni et al., 2000)), dysregulation of blood pressure(Nas, Riese, van Roon, & Rijdsdijk, 2020), and activated HPA axis(Jokinen & Nordstrom, 2009). Moreover, similar downstream CVDs (e.g., chronic ischemic heart diseases, other cerebrovascular diseases, and other cardiac arrhythmias) after primary hypertension, and acute myocardial infarction can be identified using a disease trajectory browser that was developed based on national data from inpatient wards, outpatient clinics, and emergency room visits of Denmark(Siggaard et al., 2020). Collectively, these data

295 suggest that these key CVDs may be potential targets for preventing or interrupting further
296 CVD sequels among individuals with psychiatric disorders.

297 The major strength of our study includes the multidimensional data from UK Biobank, i.e.,
298 rich phenotypic information on family history of CVD, health-related outcomes, and
299 individual-level genotype information, which enabled a comprehensive assessment on the
300 modification role of CVD susceptibility on the studied associations and the temporal patterns
301 of CVD development following a diagnosis of common psychiatry disorders. This rich data
302 source also enabled vigorous control of a wide range of important confounders, including
303 socio-demographic and behavioral factors, as well as baseline health status. Other strengths
304 include the long and complete follow-up (mean follow-up time of 11.94 years), providing
305 sufficient surveillance period for CVD outcomes, and the independent collections of
306 diagnoses for psychiatry disorders and CVD, which minimizes the risk of information bias.

307 Our study has several limitations. First, as both primary and secondary diagnoses from UK
308 Biobank inpatient hospital data were used for CVD identification, the diagnosis date for some
309 mild forms of CVDs (e.g., primary hypertension) may not be accurate, leading to a risk of
310 reverse causality. Consequently, the appearance of primary hypertension as the initial node in
311 the CVD trajectory tree of the main analysis may rather imply the importance of primary
312 hypertension as a comorbidity of psychiatric disorders on other subsequent CVDs, instead of
313 a key progression pathway. Nevertheless, our analyses on acute and severe CVDs (e.g., acute
314 myocardial infarction) should be less affected by such concern. Second, the SNPs used in the
315 calculation for CVD PRS was obtained from GWAS studies which mainly focused on
316 coronary arterial diseases, not all CVDs. However, a common genetic basis has been

demonstrated between different CVDs(Dichgans et al., 2014; Jafaripour, Sasanejad, Dadgarmoghaddam, & Sadr-Nabavi, 2019), and the generated PRS has been confirmed to be associated with the CVD phenotype of our dataset in our validation step. Furthermore, we obtained largely similar result patterns in the analyses of six major categories of CVDs. Third, as the trajectory network analysis is a data-driven approach which requires a large sample size, our analysis might have missed some rare CVDs in the trajectory networks after psychiatric disorders. Further, because some important confounders, such as BMI and smoking status, were measured at baseline, they might not accurately reflect the real status of these factors at the time of psychiatric disorder diagnosis. Last, generalization of our findings to the entire UK population, or other populations, should be done with caution because the UK Biobank participants were overall healthier than the general UK population(Sudlow et al., 2015).

In conclusion, individuals with common psychiatry disorders have an elevated risk of CVD regardless of their predisposition to CVD. Hypertension, acute myocardial infarction, and stroke represent the three initial CVDs that link psychiatric disorders to further CVDs, highlighting a need of a timely intervention to prevent further CVD sequels among individuals with common psychiatric disorders.

Acknowledgments

This research has been conducted using the UK Biobank Resource under Application 54803. We thank the team members and colleagues involved in West China Biomedical Big Data Center- UK Biobank project for their support.

339 **Contributions**

340 HS, FF, and UAV were responsible for the study concept and design. XH, YH, YajingS, ZY,
341 YQ, and JW did the data and project management. XH, YZ, HY, WC, and CH did the data
342 cleaning and analysis. XH, YZ, HS, and FF interpreted the data. XH, YananS, UAV, FF, WZ,
343 and HS drafted the manuscript. All authors approved the final manuscript as submitted and
344 agree to be accountable for all aspects of the work. The corresponding author attests that all
345 listed authors meet authorship criteria and that no others meeting the criteria have been
346 omitted.

347

348 **Funding**

349 This work is supported by the National Natural Science Foundation of China (No. 81971262
350 to HS), 1.3.5 project for disciplines of excellence, West China Hospital, Sichuan University
351 (No. ZYYC21005 to HS), EU Horizon2020 Research and Innovation Action Grant (847776
352 to UV and FF).

353

354 **Disclosure**

355 We declare no competing interests.

356

357 **Data availability**

358 Source data for Figure 2 has been provided in the supplementary (eTable 5). Original Data is
359 held by UK Biobank Limited. Due to the ethical approval of UK Biobank from the NHS
360 National Research Ethics Service and the ethical approval of the current study, we cannot

361 make the data publicly available. However, all researchers can access the data upon making
 362 an application to the UK Biobank. Detailed information on data application can be found at
 363 <http://www.ukbiobank.ac.uk/>.

364

365 **Ethics approval**

366 The UK Biobank has full ethical approval from the NHS National Research Ethics Service
 367 (reference number: 16/NW/0274), and this study was also approved by the biomedical
 368 research ethics committee of West China Hospital (reference number: 2019-1171).

369

370 **Additional information**

371 Supplementary Tables and Figures are available at online website.

References

- Cai, H., Cai, B., Zhang, H., Sun, W., Wang, Y., Zhou, S., . . . Liang, J. (2019). Major depression and small vessel stroke: a Mendelian randomization analysis. *J Neurol*, 266(11), 2859-2866. doi:10.1007/s00415-019-09511-w
- Collaboration, T. E. R. F. (2015). Association of Cardiometabolic Multimorbidity With Mortality. *JAMA*, 314(1), 52-60. doi:10.1001/jama.2015.7008
- Consortium, C. A. D., Deloukas, P., Kanoni, S., Willenborg, C., Farrall, M., Assimes, T. L., . . . Samani, N. J. (2013). Large-scale association analysis identifies new risk loci for coronary artery disease. *Nat Genet*, 45(1), 25-33. doi:10.1038/ng.2480
- DALYs, G. B. D., Collaborators, H., Murray, C. J., Barber, R. M., Foreman, K. J., Abbasoglu Ozgoren, A., . . . Vos, T. (2015). Global, regional, and national disability-adjusted life years (DALYs) for 306 diseases and injuries and healthy life expectancy (HALE) for 188 countries, 1990-2013: quantifying the epidemiological transition. *Lancet (London, England)*, 386(10009), 2145-2191. doi:10.1016/S0140-6736(15)61340-X
- Davis, K. A. S., Bashford, O., Jewell, A., Shetty, H., Stewart, R. J., Sudlow, C. L. M., & Hotopf, M. H. (2018). Using data linkage to electronic patient records to assess the validity of selected mental health diagnoses in English Hospital Episode Statistics (HES). *PLoS One*, 13(3), e0195002. doi:10.1371/journal.pone.0195002
- Dichgans, M., Malik, R., Konig, I. R., Rosand, J., Clarke, R., Gretarsdottir, S., . . . International Stroke Genetics, C. (2014). Shared genetic susceptibility to ischemic stroke and coronary artery disease: a genome-wide analysis of common variants. *Stroke*, 45(1), 24-36. doi:10.1161/STROKEAHA.113.002707

394 Forsdahl, A. (1979). Are poor living conditions in childhood and adolescence and important
395 risk factor for arteriosclerotic heart disease? *Int J Rehabil Res*, 2(2), 238-239.
396 doi:10.1097/00004356-197905000-00008

397 Ghiadoni, L., Donald, A. E., Cropley, M., Mullen, M. J., Oakley, G., Taylor, M., . . . Deanfield, J.
398 E. (2000). Mental stress induces transient endothelial dysfunction in humans.
399 *Circulation*, 102(20), 2473-2478. doi:10.1161/01.cir.102.20.2473

400 Han, X., Hou, C., Yang, H., Chen, W., Ying, Z., Hu, Y., . . . Song, H. (2021). Disease
401 trajectories and mortality among individuals diagnosed with depression: a
402 community-based cohort study in UK Biobank. *Mol Psychiatry*.
403 doi:10.1038/s41380-021-01170-6

404 Iannotti, R. J., Zuckerman, A. E., & Rifai, N. (2000). Correlations of cardiovascular disease risk
405 factors between African American siblings. *J Pediatr*, 136(4), 511-519.
406 doi:10.1016/s0022-3476(00)90015-5

407 Jafaripour, S., Sasanejad, P., Dadgarmoghaddam, M., & Sadr-Nabavi, A. (2019). ADAMTS7
408 and ZC3HC1 Share Genetic Predisposition to Coronary Artery Disease and Large
409 Artery Ischemic Stroke. *Crit Rev Eukaryot Gene Expr*, 29(4), 351-361.
410 doi:10.1615/CritRevEukaryotGeneExpr.2019028209

411 Janszky, I., Ahnve, S., Lundberg, I., & Hemmingsson, T. (2010). Early-onset depression,
412 anxiety, and risk of subsequent coronary heart disease: 37-year follow-up of 49,321
413 young Swedish men. *J Am Coll Cardiol*, 56(1), 31-37. doi:10.1016/j.jacc.2010.03.033

414 John, A., McGregor, J., Fone, D., Dunstan, F., Cornish, R., Lyons, R. A., & Lloyd, K. R. (2016).
415 Case-finding for common mental disorders of anxiety and depression in primary care:

416 an external validation of routinely collected data. *BMC Med Inform Decis Mak*, 16, 35.

417 doi:10.1186/s12911-016-0274-7

418 Jokinen, J., & Nordstrom, P. (2009). HPA axis hyperactivity and cardiovascular mortality in

419 mood disorder inpatients. *J Affect Disord*, 116(1-2), 88-92.

420 doi:10.1016/j.jad.2008.10.025

421 Kivimaki, M., Batty, G. D., Singh-Manoux, A., Britton, A., Brunner, E. J., & Shipley, M. J. (2017).

422 Validity of Cardiovascular Disease Event Ascertainment Using Linkage to UK Hospital

423 Records. *Epidemiology*, 28(5), 735-739. doi:10.1097/EDE.0000000000000688

424 Kollia, N., Panagiotakos, D., Georgousopoulou, E., Chrysohoou, C., Yannakoulia, M.,

425 Stefanadis, C., . . . investigators, A. S. (2017). Exploring the path between depression,

426 anxiety and 10-year cardiovascular disease incidence, among apparently healthy

427 Greek middle-aged adults: The ATTICA study. *Maturitas*, 106, 73-79.

428 doi:10.1016/j.maturitas.2017.09.005

429 Kubzansky, L. D., Koenen, K. C., Spiro, A., 3rd, Vokonas, P. S., & Sparrow, D. (2007).

430 Prospective study of posttraumatic stress disorder symptoms and coronary heart

431 disease in the Normative Aging Study. *Arch Gen Psychiatry*, 64(1), 109-116.

432 doi:10.1001/archpsyc.64.1.109

433 Laurent, S., & Boutouyrie, P. (2015). The structural factor of hypertension: large and small

434 artery alterations. *Circ Res*, 116(6), 1007-1021.

435 doi:10.1161/CIRCRESAHA.116.303596

436 Lu, Y., Wang, Z., Georgakis, M. K., Lin, H., & Zheng, L. (2021). Genetic Liability to Depression

437 and Risk of Coronary Artery Disease, Myocardial Infarction, and Other Cardiovascular

438 Outcomes. *J Am Heart Assoc*, 10(1), e017986. doi:10.1161/JAHA.120.017986

439 Marees, A. T., de Kluiver, H., Stringer, S., Vorspan, F., Curis, E., Marie-Claire, C., & Derks, E.
440 M. (2018). A tutorial on conducting genome-wide association studies: Quality control
441 and statistical analysis. *Int J Methods Psychiatr Res*, 27(2), e1608.
442 doi:10.1002/mpr.1608

443 Miller, A. H., Maletic, V., & Raison, C. L. (2009). Inflammation and its discontents: the role of
444 cytokines in the pathophysiology of major depression. *Biol Psychiatry*, 65(9), 732-741.
445 doi:10.1016/j.biopsych.2008.11.029

446 Momen, N. C., Plana-Ripoll, O., Agerbo, E., Benros, M. E., Borglum, A. D., Christensen, M.
447 K., . . . McGrath, J. J. (2020). Association between Mental Disorders and Subsequent
448 Medical Conditions. *N Engl J Med*, 382(18), 1721-1731. doi:10.1056/NEJMoa1915784

449 Nas, Z., Riese, H., van Roon, A. M., & Rijdsdijk, F. V. (2020). Higher Anxiety Is Associated with
450 Lower Cardiovascular Autonomic Function in Female Twins. *Twin Res Hum Genet*,
451 23(3), 156-164. doi:10.1017/thg.2020.47

452 Network, N. S. G., & International Stroke Genetics, C. (2016). Loci associated with ischaemic
453 stroke and its subtypes (SiGN): a genome-wide association study. *Lancet Neurol*,
454 15(2), 174-184. doi:10.1016/S1474-4422(15)00338-5

455 Nikpay, M., Goel, A., Won, H. H., Hall, L. M., Willenborg, C., Kanoni, S., . . . Farrall, M. (2015).
456 A comprehensive 1,000 Genomes-based genome-wide association meta-analysis of
457 coronary artery disease. *Nat Genet*, 47(10), 1121-1130. doi:10.1038/ng.3396

458 Prive, F., Arbel, J., & Vilhjalmsen, B. J. (2020). LDpred2: better, faster, stronger.
459 *Bioinformatics*. doi:10.1093/bioinformatics/btaa1029

460 Quan, H., Sundararajan, V., Halfon, P., Fong, A., Burnand, B., Luthi, J. C., . . . Ghali, W. A.
461 (2005). Coding algorithms for defining comorbidities in ICD-9-CM and ICD-10
462 administrative data. *Med Care*, 43(11), 1130-1139.
463 doi:10.1097/01.mlr.0000182534.19832.83

464 Siggaard, T., Reguant, R., Jorgensen, I. F., Haue, A. D., Lademann, M., Aguayo-Orozco,
465 A., . . . Brunak, S. (2020). Disease trajectory browser for exploring temporal,
466 population-wide disease progression patterns in 7.2 million Danish patients. *Nat*
467 *Commun*, 11(1), 4952. doi:10.1038/s41467-020-18682-4

468 Song, H., Fang, F., Arnberg, F. K., Mataix-Cols, D., Fernandez de la Cruz, L., Almqvist, C., . . .
469 Valdimarsdottir, U. A. (2019). Stress related disorders and risk of cardiovascular
470 disease: population based, sibling controlled cohort study. *BMJ*, 365, l1255.
471 doi:10.1136/bmj.l1255

472 Stein, D. J., Aguilar-Gaxiola, S., Alonso, J., Bruffaerts, R., de Jonge, P., Liu, Z., . . . Scott, K. M.
473 (2014). Associations between mental disorders and subsequent onset of hypertension.
474 *Gen Hosp Psychiatry*, 36(2), 142-149. doi:10.1016/j.genhosppsy.2013.11.002

475 Strawn, W. B., Bondjers, G., Kaplan, J. R., Manuck, S. B., Schwenke, D. C., Hansson, G.
476 K., . . . Clarkson, T. B. (1991). Endothelial dysfunction in response to psychosocial
477 stress in monkeys. *Circ Res*, 68(5), 1270-1279. doi:10.1161/01.res.68.5.1270

478 Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., . . . Collins, R. (2015). UK
479 biobank: an open access resource for identifying the causes of a wide range of
480 complex diseases of middle and old age. *PLoS Med*, 12(3), e1001779.
481 doi:10.1371/journal.pmed.1001779

482 Sumner, J. A., Kubzansky, L. D., Elkind, M. S., Roberts, A. L., Agnew-Blais, J., Chen, Q., . . .

483 Koenen, K. C. (2015). Trauma Exposure and Posttraumatic Stress Disorder

484 Symptoms Predict Onset of Cardiovascular Events in Women. *Circulation*, *132*(4),

485 251-259. doi:10.1161/CIRCULATIONAHA.114.014492

486 Swain, N. R., Lim, C. C., Levinson, D., Fiestas, F., de Girolamo, G., Moskalewicz, J., . . . Scott,

487 K. M. (2015). Associations between DSM-IV mental disorders and subsequent

488 non-fatal, self-reported stroke. *J Psychosom Res*, *79*(2), 130-136.

489 doi:10.1016/j.jpsychores.2015.05.008

490 Weinberger, A. H., Kashan, R. S., Shpigel, D. M., Esan, H., Taha, F., Lee, C. J., . . . Goodwin,

491 R. D. (2017). Depression and cigarette smoking behavior: A critical review of

492 population-based studies. *Am J Drug Alcohol Abuse*, *43*(4), 416-431.

493 doi:10.3109/00952990.2016.1171327

494 Whiteford, H. A., Degenhardt, L., Rehm, J., Baxter, A. J., Ferrari, A. J., Erskine, H. E., . . . Vos,

495 T. (2013). Global burden of disease attributable to mental and substance use

496 disorders: findings from the Global Burden of Disease Study 2010. *Lancet (London,*

497 *England)*, *382*(9904), 1575-1586. doi:10.1016/S0140-6736(13)61611-6

498 Wium-Andersen, M. K., Villumsen, M. D., Wium-Andersen, I. K., Jorgensen, M. B., Hjelmberg,

499 J. V. B., Christensen, K., & Osler, M. (2020). The familial and genetic contribution to

500 the association between depression and cardiovascular disease: a twin cohort study.

501 *Mol Psychiatry*. doi:10.1038/s41380-020-00954-6

502 Wium-Andersen, M. K., Wium-Andersen, I. K., Jorgensen, M. B., McGue, M., Jorgensen, T. S.

503 H., Christensen, K., & Osler, M. (2019). The association between depressive mood

504 and ischemic heart disease: a twin study. *Acta Psychiatr Scand*, 140(3), 265-274.

505 doi:10.1111/acps.13072

506 Woodfield, R., Grant, I., Group, U. K. B. S. O., Follow-Up, U. K. B., Outcomes Working, G., &

507 Sudlow, C. L. (2015). Accuracy of Electronic Health Record Data for Identifying Stroke

508 Cases in Large-Scale Epidemiological Studies: A Systematic Review from the UK

509 Biobank Stroke Outcomes Group. *PLoS One*, 10(10), e0140533.

510 doi:10.1371/journal.pone.0140533

511 Yang, H., Pawitan, Y., He, W., Eriksson, L., Holowko, N., Hall, P., & Czene, K. (2019). Disease

512 trajectories and mortality among women diagnosed with breast cancer. *Breast Cancer*

513 *Res*, 21(1), 95. doi:10.1186/s13058-019-1181-5

514 Yu, R. H., Ho, S. C., Lam, C. W., Woo, J. L., & Ho, S. S. (2010). Psychological factors and

515 subclinical atherosclerosis in postmenopausal Chinese women in Hong Kong.

516 *Maturitas*, 67(2), 186-191. doi:10.1016/j.maturitas.2010.06.014

517

518 **Figure legends:**

519 **Figure 1** Flow chart the participants selection

520 This figure shows the study design, the inclusion and exclusion process of study population selection.

521

522 **Figure 2** Trajectory progression of cardiovascular disease following a diagnosis of common
523 psychiatry disorders

524 This figure illustrates following trajectory progression of cardiovascular disease identified in our
525 analysis. The combined cardiovascular diseases are shown within the circle. The color of the circle
526 represents the hazard ratios of this cardiovascular disease when comparing patients with common
527 psychiatry disorders to matched unexposed individuals. The number above the arrow connecting two
528 circles corresponds to the number of pairs of two cardiovascular disease events among patients with
529 common psychiatry disorders. The color of the arrows indicates the odds ratio of the sequential
530 association between the two cardiovascular disease events among patients with common psychiatry
531 disorders.

532 **Table 1** Baselines characteristics of the study cohort

	Total individuals	Individuals with psychiatry disorders (n=35,227)	Matched individuals without psychiatry disorders^b (n=176,135)
Follow-up time (years), mean (SD)	11.94 (5.76)	11.86 (5.81)	11.95 (5.75)
Age at index date, mean (SD)	51.76 (8.96)	51.72 (9.00)	51.77 (8.96)
Sex (%)			
Female	139446 (66.0)	23241 (66.0)	116205 (66.0)
Male	71916 (34.0)	11986 (34.0)	59930 (34.0)
Educational levels (%)			
University	93568 (44.3)	14452 (41.0)	79116 (44.9)
High school	24712 (11.7)	3926 (11.1)	20786 (11.8)
Middle school	59307 (28.1)	10219 (29.0)	49088 (27.9)
Unknown	33775 (16.0)	6630 (18.8)	27145 (15.4)
Ethnicity (%)			
White	189912 (89.9)	32142 (91.2)	157770 (89.6)
Others	20304 (9.6)	2899 (8.2)	17405 (9.9)
Unknown	1146 (0.5)	186 (0.5)	960 (0.5)
BMI (%)			
< 18.5	1320 (0.6)	216 (0.6)	1104 (0.6)
18.5-25	74888 (35.4)	11180 (31.7)	63708 (36.2)
25-30	85499 (40.5)	14257 (40.5)	71242 (40.4)
> 30	48488 (22.9)	9353 (26.6)	39135 (22.2)
Unknown	1167 (0.6)	221 (0.6)	946 (0.5)
Smoking status (%)			
Never	120348 (56.9)	18319 (52.0)	102029 (57.9)
Previous smokers	66836 (31.6)	11593 (32.9)	55243 (31.4)
Current smokers	22991 (10.9)	5101 (14.5)	17890 (10.2)
Unknown	1187 (0.6)	214 (0.6)	973 (0.6)
Townsend deprivation index^a, mean (SD)	-1.25 (3.10)	-1.02 (3.16)	-1.29 (3.09)
History of other psychiatry disorders (%)			
No	198511 (93.9)	29594 (84.0)	168917 (95.9)
Yes	12851 (6.1)	5633 (16.0)	7218 (4.1)
Family history of			

CVD (%)			
No	93808 (44.4)	14973 (42.5)	78835 (44.8)
Yes	117554 (55.6)	20254 (57.5)	97300 (55.2)
CVD polygenic risk score			
Low	54110 (25.60%)	9243 (26.24%)	44867 (25.47%)
Intermediate	54110 (25.60%)	9285 (26.36%)	44825 (25.45%)
High	54125 (25.61%)	9400 (26.68%)	44725 (25.39%)
Unknown	49017 (23.19%)	7299 (20.72%)	41718 (23.69%)
CCI score, mean (SD)	0.01 (0.19)	0.01 (0.22)	0.01 (0.19)

533 This table shows the basic characteristics of the patients with common psychiatry disorders and their
534 matched unexposed individuals.
535 Abbreviation: CCI: Charlson comorbidity index; SD: standard deviation; BMI: Body Mass Index;
536 CVD: cardiovascular disease.
537 ^a Townsend deprivation index was assigned to each individual based on their postcode locations, with a
538 greater index score implying a greater degree of deprivation; missing data of TDI (n=300) was replaced
539 by mean TDI of all individuals.
540 ^b At most 5 participants who were alive, retained in the cohort and free of psychiatry disorders at the
541 corresponding index date were individually matched to each individual with psychiatry disorders based
542 on sex, and year of birth.

543 **Table 2** Hazard ratios (HRs) with 95% confidence intervals (CIs) of cardiovascular disease among patients with common psychiatry disorders compared to
544 matched unexposed individuals

	All individuals		<= 6 months follow-up		> 6 months follow-up	
	No. of cases (incidence ^a) in patients/match ed individuals	HR (95% CI) ^b	No. of cases (incidence*) in patients/matched individuals	HR (95% CI) [#]	No. of cases (incidence*) in patients/matched individuals	HR (95% CI) [#]
Model information						
Controlled for sex, birth year, educational level, ethnicity, smoking status, and TDI		1.23 (1.21-1.26)		1.70 (1.52-1.91)		1.22 (1.19-1.25)
As above + BMI	10619 (25.41)/43192	1.20 (1.17-1.23)	412 (23.88)/1183 (13.68)	1.66 (1.48-1.86)	10207 (25.47) /41403 (20.71)	1.18 (1.16-1.21)
As above + history of other psychiatry disorders	(20.52)	1.17 (1.14-1.20)		1.61 (1.43-1.81)		1.16 (1.13-1.18)
As above + family history of CVDs		1.16 (1.14-1.19)		1.59 (1.41-1.79)		1.15 (1.12-1.18)
As above + CCI		1.16 (1.14-1.19)		1.59 (1.42-1.79)		1.15 (1.12-1.18)
Subtypes of cardiovascular diseases^c						

Hypertensive diseases	8249 (19)/33112 (15)	1.16 (1.13-1.19)	281 (16)/819 (9.5)	1.55 (1.34-1.79)	7968 (19)/31959 (15)	1.15 (1.12-1.18)
Ischemic heart disease	2684 (5.7)/9760 (4.2)	1.22 (1.17-1.28)	88 (5.1)/250 (2.9)	1.66 (1.29-2.15)	2597 (5.7)/9461 (4.3)	1.21 (1.16-1.27)
Embolism and thrombosis	497 (1)/1938 (0.82)	1.12 (1.01-1.24)	15 (0.86)/44 (0.51)	1.71 (0.95-3.07)	482 (1)/1892 (0.84)	1.11 (1.00-1.23)
Arrhythmia/conduction disorder	2354 (4.9)/9747 (4.2)	1.10 (1.05-1.16)	45 (2.6)/145 (1.7)	1.55 (1.08-2.21)	2309 (5)/9587 (4.3)	1.10 (1.05-1.15)
Heart failure	637 (1.3)/2382 (1)	1.08 (0.98-1.19)	15 (0.86)/28 (0.32)	2.66 (1.42-4.97)	622 (1.3)/2349 (1)	1.07 (0.98-1.18)
Cerebrovascular disease	1039 (2.2)/3524 (1.5)	1.28 (1.19-1.38)	29 (1.7)/68 (0.78)	1.90 (1.18-3.08)	1010 (2.2)/3450 (1.5)	1.27 (1.18-1.37)

^a Per 1000 person years.

^b Hazard ratios (with 95% confidence intervals) of cardiovascular disease in patients with common psychiatry disorders compared to sex and year of birth matched unexposed individuals, derived from Cox regression models and adjusted for covariates listed in model information column. Time since index date was used as time scale.

^c Hazard ratios (with 95% confidence intervals) of subtypes of cardiovascular diseases in individual with common psychiatry disorders compared to sex and year of birth matched unexposed individuals, derived from Cox regression models and adjusted for sex, birth year, educational level, ethnicity, smoking status, TDI, BMI, history of other psychiatry disorders, family history of CVD, and CCI score. For subtypes of cardiovascular diseases with less than 20 cases in patients with common psychiatry disorders, hazard ratios were derived from Cox regression models and adjusted for sex, birth year. Definition of subtypes cardiovascular diseases by ICD-10 codes can be found in the eTable 1 in the supplement. Time since index date was used as time scale.

Table 3 Hazard ratios (HRs) with 95% confidence intervals (CIs) of cardiovascular disease among patients with common psychiatry disorders compared to matched unexposed individuals, by different disease susceptibility

Disease susceptibility	<= 6 months follow-up		> 6 months follow-up	
	No. of cases (incidence ^a) in patients/matched individuals	HR (95% CI) ^b	No. of cases (incidence ^a) in patients/matched individuals	HR (95% CI) ^b
Family history of cardiovascular disease				
No	139 (18.94)/172 (10.03)	1.61 (1.25-2.07)	3675 (21.2)/6229 (15.56)	1.21 (1.16-1.27)
Yes	73 (27.55)/475 (16.88)	1.54 (1.31-1.80)	6532 (28.74)/15882 (24.65)	1.15 (1.11-1.18)
P _{interaction} ^c		0.7703		0.0683
Level of polygenic risk score (PRS) of cardiovascular disease^d				
Low PRS (<the lower tertile)	105 (23.2)/62 (10.74)	2.29 (1.58-3.32)	2581 (24.14)/2665 (19.65)	1.25 (1.16-1.34)
Moderate PRS (lower-upper tertiles)	106 (23.31)/94 (16.18)	1.38 (0.96-1.98)	2723 (25.49)/2902 (21.32)	1.16 (1.09-1.24)
High PRS (>the upper tertile)	117 (25.42)/88 (14.88)	1.41 (0.98-2.02)	2936 (27.65)/3136 (22.86)	1.18 (1.10-1.25)
P _{interaction}		0.0667		0.2412

^a Per 1000 person years.

^b Cox regression models were stratified by disease susceptibility and adjusted for sex, birth year, educational level, ethnicity, smoking status, TDI, BMI, history of other psychiatry disorders, and CCI score.

^c The differences in hazard ratios for CVD polygenic risk score were assessed between low and high subgroups by Wald test.

^d Polygenic risk score of cardiovascular disease were calculated based on published GWAS summary statistics data.

Hazard ratios compared to matched unexposed individuals

Odd ratios for pairs of CVDs

Supplementary tables and figures

eTable 1 International Classification of Diseases (ICD) codes and primary care codes for diseases identifications

eTable 2 Mapping between ICD-10 codes and Combined ICD codes for PheWAS analysis

eTable 3 Hazard ratios (HRs) with 95% confidence intervals (CIs) of cardiovascular disease among patients with common psychiatry disorders compared to matched unexposed individuals, by different psychiatry disorders

eTable 4 PheWAS results of hazard ratios (HRs) with 95% confidence intervals (CIs) of different individual cardiovascular diseases among patients with common psychiatry disorders compared to matched unexposed individuals

eTable 5 Odds ratios (ORs) with 95% confidence intervals (CIs) for the significant pairs of cardiovascular diseases following a diagnosis of common psychiatry disorders

eTable 6 Hazard ratios (HRs) with 95% confidence intervals (CIs) of primary diagnosis of cardiovascular disease among patients with common psychiatry disorders compared to matched unexposed individuals

eTable 7 Hazard ratios (HRs) with 95% confidence intervals (CIs) of primary diagnosis of cardiovascular disease among patients with common psychiatry disorders compared to matched unexposed individuals, by different disease susceptibility

eTable 8 Odds ratios (ORs) with 95% confidence intervals (CIs) for the significant pairs of primary diagnosis of cardiovascular diseases following a diagnosis of common psychiatry disorders

eFigure 1 Hazard ratios (95% CIs) of cardiovascular disease among patients with common psychiatry disorders compared with their matched unexposed individuals, stratified by time of follow-up

eFigure 2 Flow chart of identifying trajectory progression of cardiovascular disease following a diagnosis of common psychiatry disorder

eFigure 3 Relative risks of six subtypes of cardiovascular diseases among patients with common psychiatry disorders compared with their matched unexposed individuals, stratified by disease susceptibility, by time of follow-up (≤ 6 or > 6 months)

eFigure 4 Flow chart of identifying trajectory progression of primary diagnosis of cardiovascular disease following a diagnosis of common psychiatry disorders

eFigure 5 Trajectory progression of primary diagnosis of cardiovascular disease following a diagnosis of common psychiatry disorders

eTable 1 International Classification of Diseases (ICD) codes and primary care codes for diseases identifications

	ICD-9	ICD-10	Primary care READ V2 code	Primary care READ V3 code
Anxiety	3000, 3002	F40, F41	E2000, E2001, E2002, E2004, E2005, E200z, E202., E2020, E2021, E2022, E2023, E2024, E2025, E2026, E2027, E2028, E2029, E202E, Eu40., Eu400, Eu402, Eu40y, Eu40z, Eu41., Eu410, Eu413, Eu41y, Eu41z, Eu931, Ua1qa, Ua1qc, Ua1qd, Ua1qe, Ua1qf, Ua1qg, Ua1qh, Ua1qi, Ua1qj, Ua1qk, Ua1ql, , Ua1qm, Ua1qn, Ua1qo, Ua1qp, Ua1qs, Ua1qt, Ua1qU, Ua1qV, Ua1qW, Ua1qX, Ua1qY, X00RP, X00Sa, X00Sb, X00Sc, X00Sd, X00Se, X00Sr, X00SV, X00SW, X00SX, X00SY, X00SZ, X50G2, X50G3, X50G5, X50G6, X50GI, X75YV, X761d, X761n, X761q, X761t, X761u, X761y, X7627, X7628, X7629, X762a, X762C, X762E, X762F, X762G, X762H, X762T, X762Z, X78wp, Xa00r, Xa00s, Xa1a8, Xa1Ev, Xa3Vj, Xa3Vk, Xa3VI, Xa3WH, Xa3WI, Xa3WJ, Xa7k9, Xa7kB, Xalo7, XaKVA, XE0rb, XE1Y7, XE1YA, XE1YB, XE1Zj, XM0Ak	E200., E2000, E2002, E2004, E200z, E2020, E2021, E2022, E2023, E2024, E2025, E2026, E2027, E2028, E2029, E202E, Eu40., Eu400, Eu402, Eu40y, Eu40z, Eu41., Eu410, Eu413, Eu41y, Eu41z, Eu931, Ua1qa, Ua1qc, Ua1qd, Ua1qe, Ua1qf, Ua1qg, Ua1qh, Ua1qi, Ua1qj, Ua1qk, Ua1ql, , Ua1qm, Ua1qn, Ua1qo, Ua1qp, Ua1qs, Ua1qt, Ua1qU, Ua1qV, Ua1qW, Ua1qX, Ua1qY, X00RP, X00Sa, X00Sb, X00Sc, X00Sd, X00Se, X00Sr, X00SV, X00SW, X00SX, X00SY, X00SZ, X50G2, X50G3, X50G5, X50G6, X50GI, X75YV, X761d, X761n, X761q, X761t, X761u, X761y, X7627, X7628, X7629, X762a, X762C, X762E, X762F, X762G, X762H, X762T, X762Z, X78wp, Xa00r, Xa00s, Xa1a8, Xa1Ev, Xa3Vj, Xa3Vk, Xa3VI, Xa3WH, Xa3WI, Xa3WJ, Xa7k9, Xa7kB, Xalo7, XaKVA, XE0rb, XE1Y7, XE1YA, XE1YB, XE1Zj, XM0Ak
Depression	2961	F32, F33	E112., E1120, E1121, E1122, E1123, E1124, E1125, E1126, E112z, E113., E1130, E1131, E1132, E1133, E1134, E1135, E1136, E1137, E113z, E135., Eu32., Eu320, Eu321, Eu322, Eu323, Eu324, Eu325, Eu326,	2257, E0043, E1120, E1121, E1122, E1123, E1124, E1125, E1126, E112z, E1130, E1131, E1132, E1133, E1134, E1135, E1136, E1137, E113z, E11y2, E130, E2B0., E2B1., Eu320, Eu321, Eu322, Eu323, Eu32y, Eu32z, Eu330, Eu331, Eu332, Eu333, Eu334, Eu33y, Eu33z, X00SO, X00SQ, X00SS, X00SU, Xa0wV, XaB9J, XaCHo, XaCHr, XaCHs, XaCIs, XaCIr, XaClu,

Stress disorders	related	308, 309	F43	Eu327, Eu328, Eu329, Eu32A, Eu32B, Eu32y, Eu32z, Eu33., Eu330, Eu331, Eu332, Eu333, Eu334, Eu33y, Eu33z	XaX53, XaX54, XaY2C, XE1Y0, XE1Y1, XE1YC, XE1Za, XE1Zb, XE1Zc, XE1Zd, XE1Ze, XE1Zf, XE1ZY, XE1ZZ, XM1GC, XSEGJ, XSGok, XSGol, XSGom, XSGon
				E280., E281., E282., E283., E2830, E2831, E283z, E284., E28z., E2900, E2925, E292y, E292z, E293., E2930, E2931, E2932, E293z, E294., E29y., E29y1, E29y2, E29y4, E29yz, E29z., Eu43., Eu430, Eu431, Eu432, Eu433, Eu434, Eu435, Eu43y, Eu43z	1B1L., 1BE., E280., E281., E282., E283., E2830, E2831, E283z, E284., E29., E290., E290z, E291., E292., E2920, E2921, E2922, E2924, E2925, E292y, E292z, E293., E2930, E2931, E2932, E293z, E294., E29y., E29y0, E29y1, E29y2, E29y3, E29y4, E29y5, E29yz, E29z., Eu430, Eu432, Eu43y, Eu43z, Eu930, Ry15., Ua18k, Ua18L, Ub1T9, X00Sf, X00TT, X40Js, Xa028, Xa18j, Xa18v, XaC2u, XaX55, XaX56, XaX58, XE1Ym, XE1Yn, XE1Yo, XE1Yp, XE2uz, XM0As, XM1Q3
Any cardiovascular disease		390-438, 444, 445	440, I00-I70, I730, I74-I75		
Hypertensive diseases			I10-I15		
Ischemic heart disease			I20-I25		
Embolism and thrombosis			I26, I74, I75		
Arrhythmia/conduction disorder			I44-I49		
Heart failure			I50		
Cerebrovascular disease			I60-I69		

Covariates: history of other psychiatry disorders	F00-F31, F34- F39, F42, F44- F99
--	--

eTable 2 Mapping between ICD-10 codes and Combined ICD codes for PheWAS analysis

ICD-10	Combined ICD codes	DESCRIPTION
I00	I00	Acute rheumatic fever
I01	I00	Acute rheumatic fever
I02	I00	Acute rheumatic fever
I05	I05	Chronic rheumatic heart disease
I06	I05	Chronic rheumatic heart disease
I07	I05	Chronic rheumatic heart disease
I08	I05	Chronic rheumatic heart disease
I09	I05	Chronic rheumatic heart disease
I10	I10	Primary hypertension
I11	I11	Other hypertensive disorders
I12	I11	Other hypertensive disorders
I13	I11	Other hypertensive disorders
I15	I11	Other hypertensive disorders
I20	I20	Angina pectoris
I21	I21	Acute myocardial infarction
I22	I22	Other ischaemic heart disease
I23	I21	Acute myocardial infarction
I24	I21	Acute myocardial infarction
I25	I25	Chronic ischaemic heart disease
I26	I26	Embolism and thrombosis
I27	I27	Other pulmonary heart diseases
I28	I27	Other pulmonary heart diseases
I30	I30	Diseases of pericardium
I31	I30	Diseases of pericardium
I32	I30	Diseases of pericardium
I33	I33	Endocarditis
I34	I34	Non-rheumatic valve disorders
I35	I34	Non-rheumatic valve disorders
I36	I34	Non-rheumatic valve disorders
I37	I34	Non-rheumatic valve disorders
I38	I33	Endocarditis
I39	I33	Endocarditis
I40	I40	Myocarditis
I41	I40	Myocarditis
I42	I42	Cardiomyopathy
I43	I42	Cardiomyopathy
I44	I49	Other cardiac arrhythmias
I45	I49	Other cardiac arrhythmias

I46	I46	Cardiac arrest
I47	I49	Other cardiac arrhythmias
I48	I48	Atrial fibrillation and flutter
I49	I49	Other cardiac arrhythmias
I50	I50	Heart failure
I51	I51	Complications and ill-defined descriptions of heart disease
I52	I52	Other heart disorders in diseases classified elsewhere
I60	I60	Stroke
I61	I60	Stroke
I62	I60	Stroke
I63	I60	Stroke
I64	I60	Stroke
I65	I65	Other cerebrovascular diseases
I66	I65	Other cerebrovascular diseases
I67	I65	Other cerebrovascular diseases
I68	I65	Other cerebrovascular diseases
I69	I69	Sequelae of cerebrovascular disease
I70	I70	Atherosclerosis
I730	I73	Peripheral vascular disease
I74	I26	Embolism and thrombosis
I75	I26	Embolism and thrombosis

eTable 3 Hazard ratios (HRs) with 95% confidence intervals (CIs) of cardiovascular disease among patients with common psychiatry disorders compared to matched unexposed individuals, by different psychiatry disorders

		<= 6 months follow-up		> 6 months follow-up	
Subtype of psychiatry disorders	Cardiovascular diseases	No. of cases (incidence*) in patients/matched individuals	HR (95% CIs)#	No. of cases (incidence*) in patients/matched individuals	HR (95% CIs) #
Anxiety	Any cardiovascular diseases	158 (24.64)/419 (13.02)	2.29 (1.58-3.32)	3838 (25.62)/15807 (21.28)	1.16 (1.11-1.20)
	Hypertensive diseases	99 (15)/288 (8.9)	1.66 (1.30-2.12)	3003 (19)/12245 (16)	1.15 (1.11-1.21)
	Ischemic heart disease	40 (6.2)/91 (2.8)	2.21 (1.48-3.29)	952 (5.6)/3637 (4.4)	1.19 (1.11-1.29)
	Embolic and thrombosis	6 (0.93)/13 (0.4)	2.37 (0.89-6.28)	163 (0.93)/725 (0.86)	0.96 (0.80-1.15)
	Arrhythmia/conduction disorder	17 (2.6)/52 (1.6)	1.66 (0.96-2.88)	886 (5.2)/3673 (4.4)	1.13 (1.05-1.22)
	Heart failure	5 (0.77)/5 (0.15)	5.69 (1.46-22.16)	214 (1.2)/911 (1.1)	0.97 (0.82-1.14)
	Cerebrovascular disease	10 (1.5)/27 (0.84)	1.81 (0.87-3.74)	362 (2.1)/1311 (1.6)	1.24 (1.09-1.40)
Depression	Any cardiovascular diseases	171 (23.76)/517 (14.34)	1.42 (1.18-1.71)	4310 (25.79)/17033 (20.24)	1.15 (1.12-1.20)
	Hypertensive diseases	124 (17)/358 (9.9)	1.44 (1.16-1.80)	3345 (19)/13154 (15)	1.14 (1.09-1.19)
	Ischemic heart disease	34 (4.7)/108 (3)	1.41 (0.93-2.15)	1158 (6.1)/3906 (4.2)	1.25 (1.16-1.34)
	Embolic and thrombosis	5 (0.69)/24 (0.66)	1.03 (0.39-2.71)	242 (1.2)/765 (0.8)	1.35 (1.16-1.58)
	Arrhythmia/conduction disorder	16 (2.2)/63 (1.7)	1.26 (0.73-2.18)	956 (4.9)/4003 (4.3)	1.06 (0.98-1.14)
	Heart failure	7 (0.97)/17 (0.47)	2.06 (0.85-5.00)	287 (1.5)/962 (1)	1.17 (1.01-1.35)
	Cerebrovascular disease	11 (1.5)/22 (0.61)	2.53 (1.22-5.26)	467 (2.4)/1444 (1.5)	1.37 (1.23-1.53)
Stress-related disorders	Any cardiovascular diseases	83 (22.8)/247 (13.55)	1.56 (0.64-2.03)	2059 (24.57)/8563 (20.64)	1.28 (1.07-1.19)

Hypertensive diseases	58 (16)/173 (9.5)	1.55 (1.13-2.15)	1620 (19)/6560 (15)	1.15 (1.08-1.21)
Ischemic heart disease	14 (3.8)/51 (2.8)	1.38 (0.76-2.51)	487 (5.2)/1918 (4.2)	1.17 (1.05-1.30)
Embolism and thrombosis	4 (1.1)/7 (0.38)	2.99 (0.85-10.48)	77 (0.79)/402 (0.86)	0.90 (0.70-1.17)
Arrhythmia/conduction disorder	12 (3.3)/30 (1.6)	2.00 (1.02-3.91)	467 (4.9)/1911 (4.1)	1.13 (1.01-1.25)
Heart failure	3 (0.82)/6 (0.33)	2.51 (0.63-10.09)	121 (1.2)/476 (1)	1.09 (0.88-1.36)
Cerebrovascular disease	8 (2.2)/19 (1)	2.12 (0.92-4.87)	181 (1.9)/695 (1.5)	1.12 (0.94-1.34)

*Per 1000 person years.

#Hazard ratios (with 95% confidence intervals) of any or subtypes of cardiovascular diseases in individual with common psychiatry disorders compared to sex and year of birth matched unexposed individuals, derived from Cox regression models and adjusted for sex, birth year, educational level, ethnicity, smoking status, TDI, BMI, history of other psychiatry disorders, family history of CVD, and CCI score. For subtypes of cardiovascular diseases with less than 20 cases in patients with common psychiatry disorders, hazard ratios were derived from Cox regression models and adjusted for sex, birth year. Definition of subtypes cardiovascular diseases by ICD-10 codes can be found in the eTable 1. Time since index date was used as time scale.

eTable 4 PheWAS results of hazard ratios (HRs) with 95% confidence intervals (CIs) of different individual cardiovascular diseases among patients with common psychiatry disorders compared to matched unexposed individuals

Cardiovascular diseases	Codes*	Case numbers [#]	HR (95% CIs) ^{&}	p
Sequelae of cerebrovascular disease	I69	155	1.25 (1.02-1.54)	0.0296
Peripheral vascular disease	I73	198	1.29 (1.09-1.52)	0.00307
Chronic rheumatic heart disease	I05	275	0.84 (0.73-0.96)	0.0106
Embolism and thrombosis	I26	502	1.12 (1.01-1.24)	0.037
Other cerebrovascular diseases	I65	551	1.38 (1.24-1.53)	1.77E-09
Non-rheumatic valve disorders	I34	566	1.04 (0.94-1.15)	0.411
Complications and ill-defined descriptions of heart disease	I51	605	1.10 (1.00-1.21)	0.041
Heart failure	I50	637	1.08 (0.98-1.19)	0.103
Stroke	I60	694	1.20 (1.09-1.32)	0.000129
Acute myocardial infarction	I21	871	1.18 (1.09-1.29)	0.000117
Other cardiac arrhythmias	I49	1370	1.16 (1.09-1.24)	7.83E-06
Atrial fibrillation and flutter	I48	1404	1.03 (0.97-1.09)	0.357
Angina pectoris	I20	1521	1.30 (1.22-1.38)	5.03E-17
Chronic ischaemic heart disease	I25	2208	1.20 (1.14-1.26)	5.96E-13
Primary hypertension	I10	8239	1.16 (1.13-1.19)	5.67E-28

*Combined ICD-10 code; mapping between the original ICD-10 code and the combined ICD-10 code can be found in the eTable 2;

[#]Numbers of the individuals of common psychiatry disorders diagnosed with the corresponding cardiovascular diseases after the diagnosis of common psychiatry disorders;

[&]Hazard ratios (with 95% confidence intervals) of different cardiovascular diseases more than 150 cases in patients with common psychiatry disorders compared to sex and year of birth matched individuals without psychiatry disorders, derived from Cox regression models and adjusted for sex, birth year, educational level, ethnicity, smoking status, TDI, BMI, history of other psychiatry disorders, family history of CVD, and CCI score.

eTable 5 Odds ratios (ORs) with 95% confidence intervals (CIs) for the significant pairs of cardiovascular diseases following a diagnosis of common psychiatry disorders

Codes of CVD pairs*	No.#	OR (95% CI) ^{&}	Percentage [§]
I10-I20	505	5.01 (4.17-6.01)	30.98%
I10-I21	224	4.02 (3.10-5.20)	31.45%
I10-I25	739	4.60 (3.97-5.33)	26.55%
I10-I49	419	2.97 (2.48-3.56)	19.88%
I10-I60	203	3.81 (2.88-5.05)	32.85%
I10-I65	226	5.10 (3.80-6.86)	21.47%
I20-I49	115	3.12 (2.33-4.18)	24.00%
I21-I20	164	24.89 (14.15-43.79)	25.00%
I21-I25	193	29.32 (16.72-51.42)	55.94%
I25-I20	430	21.65 (15.53-30.17)	36.83%
I25-I49	174	4.46 (3.41-5.84)	26.34%
I25-I65	79	2.84 (1.99-4.07)	17.65%

A total of 12 CVD pairs with N ≥ 75 and statistically significant odds ratios after Bonferroni correction for patients with common psychiatry disorders were shown in this table;

Abbreviation: CVD, cardiovascular disease;

*Combined ICD-10 code for two CVDs; mapping between the original ICD-10 code and the combined ICD-10 code can be found in the eTable 2;

#Number of patients with common psychiatry disorders who experienced the corresponding pairs of two CVDs;

[&]Odds ratios (with 95% confidence intervals) of pairs of two CVDs;

[§]Number of individuals with psychiatry disorders having the same diagnosis date of two CVDs divided by the number of patients with common psychiatry disorders diagnosed with both two CVDs.

eTable 6 Hazard ratios (HRs) with 95% confidence intervals (CIs) of primary diagnosis of cardiovascular disease among patients with common psychiatry disorders compared to matched unexposed individuals

	All individuals		<= 6 months follow-up		> 6 months follow-up	
	No. of cases (incidence*) in patients/matched individuals	HR (95% CI)#	No. of cases (incidence*) in patients/matched individuals	HR (95% CI) #	No. of cases (incidence*) in patients/matched individuals	HR (95% CI) #
Model information						
Controlled for sex, birth year, education level, ethnic, smoke, and TDI	1890 (4.522)/7982 (3.792)	1.18 (1.12-1.24)	92 (5.333)/273 (3.157)	1.66 (1.31-2.11)	1798 (4.487)/7601 (3.802)	1.16 (1.10-1.22)
As above + BMI		1.16 (1.10-1.22)		1.65 (1.30-2.10)		1.14 (1.08-1.20)
As above + history of other psychiatry disorders		1.13 (1.08-1.20)		1.67 (1.31-2.13)		1.11 (1.05-1.18)
As above + family history of CVDs		1.13 (1.07-1.19)		1.67 (1.30-2.13)		1.11 (1.05-1.17)
As above + CCI		1.13 (1.07-1.19)		1.67 (1.30-2.13)		1.11 (1.05-1.17)

Subtypes of cardiovascular diseases^{&}						
Hypertensive diseases	135 (0.28)/546 (0.23)	1.13 (0.93-1.38)	6 (0.35)/15 (0.17)	2.31 (0.86-6.23)	129 (0.27)/531 (0.23)	1.11 (0.91-1.36)
Ischemic heart disease	1796 (3.8)/6763 (2.9)	1.19 (1.13-1.26)	61 (3.5)/178 (2.1)	1.68 (1.23-2.30)	1736 (3.8)/6557 (2.9)	1.18 (1.12-1.25)
Embolism and thrombosis	352 (0.72)/1354 (0.58)	1.13 (1.00-1.28)	12 (0.69)/29 (0.33)	2.07 (1.05-4.05)	340 (0.73)/1325 (0.58)	1.12 (0.98-1.27)
Arrhythmia/conduction disorder	935 (1.9)/4141 (1.8)	1.07 (0.99-1.15)	25 (1.4)/72 (0.83)	1.84 (1.12-3.03)	910 (2)/4062 (1.8)	1.05 (0.98-1.14)
Heart failure	177 (0.36)/691 (0.29)	1.08 (0.90-1.30)	5 (0.29)/9 (0.1)	2.81 (0.94-8.42)	172 (0.37)/681 (0.3)	1.07 (0.88-1.29)
Cerebrovascular disease	644 (1.3)/2352 (1)	1.17 (1.07-1.29)	17 (0.98)/51 (0.59)	1.64 (0.95-2.84)	627 (1.3)/2297 (1)	1.17 (1.06-1.28)

*Per 1000 person years.

[#]Hazard ratios (with 95% confidence intervals) of cardiovascular disease in patients with common psychiatry disorders compared to sex and year of birth matched unexposed individuals without, derived from Cox regression models and adjusted for covariates listed in model information column. Time since index date was used as time scale.

[&]Hazard ratios (with 95% confidence intervals) of subtypes of cardiovascular diseases in individual with psychiatry disorders compared to sex and year of birth matched individuals without psychiatry disorders, derived from Cox regression models and adjusted for sex, birth year, educational level, ethnicity, smoking status, TDI, BMI, history of other psychiatry disorders, family history of CVD, and CCI score. For subtypes of cardiovascular diseases with less than 20 cases in patients with common psychiatry disorders, hazard ratios were derived from Cox regression models and adjusted for sex, birth year. Definition of subtypes cardiovascular diseases by ICD-10 codes can be found in the eTable 1. Time since index date was used as time scale.

eTable 7 Hazard ratios (HRs) with 95% confidence intervals (CIs) of primary diagnosis of cardiovascular disease among patients with common psychiatry disorders compared to matched unexposed individuals, by different disease susceptibility

Disease susceptibility	<= 6 months follow-up		> 6 months follow-up	
	No. of cases (incidence*) in patients/matched individuals	HR (95% CI) [#]	No. of cases (incidence*) in patients/matched individuals	HR (95% CI) [#]
Family history of cardiovascular disease				
No	29 (3.951)/35 (2.041)	1.88 (1.04-3.38)	636 (3.668)/1087 (2.714)	1.25 (1.12-1.40)
Yes	63 (6.357)/105 (3.732)	1.94 (1.38-2.72)	1162 (5.112)/2963 (4.598)	1.09 (1.01-1.17)
P _{interaction}		0.9283		0.0445
Level of polygenic risk score (PRS) of cardiovascular disease[§]				
Low (<the lower tertile)	17 (3.756)/17 (2.943)	1.35 (0.57-3.20)	448 (4.19)/415 (3.06)	1.48 (1.25-1.75)
Moderate (lower-upper tertile)	31 (6.817)/20 (3.442)	2.02 (1.14-3.56)	440 (4.12)/525 (3.857)	1.02 (0.87-1.19)
High (>the upper tertile)	29 (6.3)/22 (3.72)	1.93 (0.62-6.02)	579 (5.453)/656 (4.782)	1.19 (1.04-1.37)
P _{interaction} [*]		0.6241		0.0493

*Per 1000 person years.

[#]Cox regression models were stratified by disease susceptibility and adjusted for sex, birth year, educational level, ethnicity, smoking status, TDI, BMI, history of other psychiatry disorders, and CCI score.

[§]Polygenic risk score of cardiovascular disease were calculated based on published GWAS summary statistics data.

* The differences in hazard ratios for CVD polygenic risk score were assessed between low and high subgroups by Wald test.

eTable 8 Odds ratios (ORs) with 95% confidence intervals (CIs) for the significant pairs of primary diagnosis of cardiovascular diseases following a diagnosis of common psychiatry disorders

Codes of CVD pairs**	No.#	OR (95% CI) ^{&}	Percentage [§]
I21-I20	89	19.77 (9.96-39.24)	7.87%
I21-I25	184	29.99 (16.73-53.76)	15.52%

A total of 2 CVDs pairs with N ≥ 75 and statistically significant odds ratios after Bonferroni correction for individuals with psychiatry disorders were show in this table;

Abbreviation: CVD, cardiovascular disease;

*Combined ICD-10 code for two CVDs; mapping between the original ICD-10 code and the combined ICD-10 code can be found in the eTable 2;

#Number of patients with common psychiatry disorders who experienced the corresponding pairs of two CVDs;

&Odds ratios (with 95% confidence intervals) of pairs of two CVDs;

§Number of individuals with psychiatry disorders having the same diagnosis date of two CVDs divided by the number of patients with common psychiatry disorders diagnosed with both two CVDs.

eFigure 1 Hazard ratios (95% CIs) of cardiovascular disease among patients with common psychiatry disorders compared with their matched unexposed individuals, stratified by time of follow-up

Abbreviation: CI: confidence interval.

The X axis shows the different follow-up period: ≤3, 3-6, 6-12, 12-18, 18-24, 24-60, 60-120, 120-240, and >240 months follow-up. The Y axis shows the significant hazard ratios of cardiovascular disease among patients with common psychiatry disorders compared with their matched unexposed individuals, derived from Cox models adjusted for sex, birth year and ethnicity.

eFigure 2 Flow chart of identifying trajectory progression of cardiovascular disease following a diagnosis of common psychiatry disorders

eFigure 3 Relative risks of six subtypes of cardiovascular diseases among patients with common psychiatry disorders compared with their matched unexposed individuals, stratified by disease susceptibility, by time of follow-up (≤ 6 or > 6 months)

Abbreviation: N: number of cases of specific cardiovascular disease; I: incidence.

Cox models were stratified by family history of cardiovascular disease (yes or no) and level of cardiovascular disease PRS (low, middle, and high) and adjusted for sex, birth year, educational level, ethnicity, smoke, TDI, BMI, history of other psychiatry disorders, and CCI score. For subtypes of cardiovascular diseases with less than 20 cases in patients with common psychiatry disorders, hazard ratios were derived from Cox regression models and adjusted for sex, birth year.

eFigure 4 Flow chart of identifying trajectory progression of primary diagnosis of cardiovascular disease following a diagnosis of common psychiatry disorders

eFigure 5 Trajectory progression of primary diagnosis of cardiovascular disease following a diagnosis of common psychiatry disorders

This figure illustrates following trajectory progression of primary diagnosis of cardiovascular disease identified in our analysis. The combined cardiovascular diseases are shown within the circle. The color of the circle represents the hazard ratios of this cardiovascular disease when comparing patients with common psychiatry disorders to matched unexposed individuals. The number above the arrow connecting two circles corresponds to the number of pairs of two cardiovascular disease events among patients with common psychiatry disorders. The color of the arrows indicates the odds ratio of the sequential association between the two cardiovascular disease events among patients with common psychiatry disorders.